

Hypothalamic *Pomc* expression restricted to GABAergic neurons suppresses *Npy* overexpression and restores food intake in obese mice



Milagros Trotta¹, Estefanía Pilar Bello¹, Ramiro Alsina¹, María Belén Tavella², José Luis Ferrán³, Marcelo Rubinstein^{2,4}, Viviana Florencia Bumaschny^{1,5,*}

ABSTRACT

Objective: Hypothalamic arcuate proopiomelanocortin (Arc-POMC) neurons are involved in different physiological processes such as the regulation of energy balance, glucose homeostasis, and stress-induced analgesia. Since these neurons heterogeneously express different biological markers and project to many hypothalamic and extrahypothalamic areas, it is proposed that Arc-POMC neurons could be classified into different subpopulations having diverse physiological roles. The aim of the present study was to characterize the contribution of the subpopulation of Arc-POMC neurons cosecreting gamma-aminobutyric acid (GABA) neurotransmitter in the control of energy balance.

Methods: Arc-*Pomc* expression restricted to GABAergic-POMC neurons was achieved by crossing a reversible *Pomc*-deficient mouse line (*arcPomc*^{-/-}) with a tamoxifen-inducible *Gad2*-CreER transgenic line. *Pomc* expression was rescued in the compound *arcPomc*^{-/-}:*Gad2*-CreER female and male mice by tamoxifen treatment at postnatal days 25 (P25) or 60 (P60), and body weight, daily food intake, fasting glycemia, and fasting-induced hyperphagia were measured. POMC recovery was quantified by immunohistochemistry and semiquantitative RT-PCR. Neuropeptide Y (NPY) and GABAergic neurons were identified by in situ hybridization. Arc-POMC neurons projecting to the dorsomedial hypothalamic nucleus (DMH) were studied by stereotactic intracerebral injection of fluorescent retrobeads into the DMH.

Results: Tamoxifen treatment of *arcPomc*^{-/-}:*Gad2*-CreER mice at P60 resulted in *Pomc* expression in ~23–25% of Arc-POMC neurons and ~15–23% of *Pomc* mRNA levels, compared to *Gad2*-CreER control mice. *Pomc* rescue in GABAergic-POMC neurons at P60 normalized food intake, glycemia, and fasting-induced hyperphagia, while significantly reducing body weight. Energy balance was also improved in *arcPomc*^{-/-}:*Gad2*-CreER mice treated with tamoxifen at P25. Distribution analysis of rescued POMC immunoreactive fibers revealed that the DMH is a major target site of GABAergic-POMC neurons. Further, the expression of the orexigenic neuropeptide Y (NPY) in the DMH was increased in *arcPomc*^{-/-} obese mice but was completely restored after *Pomc* rescue in *arcPomc*^{-/-}:*Gad2*-CreER mice. Finally, we found that ~75% of Arc-POMC neurons projecting to the DMH are GABAergic.

Conclusions: In the present study, we show that the expression of *Pomc* in the subpopulation of Arc-GABAergic-POMC neurons is sufficient to maintain normal food intake. In addition, we found that DMH-NPY expression is negatively correlated with *Pomc* expression in GABAergic-POMC neurons, suggesting that food intake may be regulated by an Arc-GABAergic-POMC → DMH-NPY pathway.

© 2020 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords Proopiomelanocortin; GABA; Energy balance; Dorsomedial hypothalamic nucleus; Arcuate hypothalamic nucleus; Obesity

¹Grupo de Neurociencia de Sistemas, Instituto de Fisiología y Biofísica “Bernardo Houssay” (IFIBIO HOUSSAY), Facultad de Medicina, Universidad de Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas, Paraguay 2155, C1121ABG, Ciudad Autónoma de Buenos Aires, Argentina ²Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Consejo Nacional de Investigaciones Científicas y Técnicas, Vuelta de Obligado 2490, 1428, Ciudad Autónoma de Buenos Aires, Argentina ³Department of Human Anatomy, School of Medicine, University of Murcia and IMIB-Arrixaca Institute, Carretera Buenavista s/n, 30120, El Palmar, Murcia, Spain ⁴Departamento de Fisiología, Biología Molecular y Celular, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Intendente Güiraldes 2160, Ciudad Universitaria, C1428EGA, Ciudad Autónoma de Buenos Aires, Argentina ⁵Departamento de Ciencias Fisiológicas, Facultad de Medicina, Universidad de Buenos Aires, Paraguay 2155, C1121ABG, Ciudad Autónoma de Buenos Aires, Argentina

*Corresponding author. Universidad de Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto de Fisiología y Biofísica “Bernardo Houssay” (IFIBIO HOUSSAY), Grupo de Neurociencia de Sistemas, Facultad de Medicina, 2155 Calle Paraguay, Buenos Aires, C1121ABG, Argentina.

E-mails: mili.trotta@gmail.com (M. Trotta), estefania.bello@gmail.com (E.P. Bello), ramiroalsina@gmail.com (R. Alsina), belentavella@gmail.com (M.B. Tavella), jferran@um.es (J.L. Ferrán), mrubins@dna.uba.ar (M. Rubinstein), vbumaschny@fmed.uba.ar (V.F. Bumaschny).

Abbreviations: POMC, Proopiomelanocortin; GABA, Gamma-aminobutyric acid; GLU, Glutamate; Arc, Arcuate nucleus; DMH, Dorsomedial hypothalamic nucleus; PVH, Paraventricular hypothalamic nucleus; LH, Lateral hypothalamic nucleus; NPY, Neuropeptide Y; TAM, Tamoxifen

Received January 22, 2020 • Revision received March 17, 2020 • Accepted March 23, 2020 • Available online 18 April 2020

<https://doi.org/10.1016/j.molmet.2020.100985>

1. INTRODUCTION

Overweight is caused by multiple factors leading to the disruption of cerebral circuits that sense energy availability and regulate energy balance [1]. Proopiomelanocortin (POMC) anorexigenic neurons, located in the arcuate nucleus (Arc) of the hypothalamus, are the main regulators of food intake and energy expenditure [2]. Recent studies have shown that Arc-POMC neurons are heterogeneous and can be divided into subpopulations expressing diverse biomarkers and neurotransmitters [3–5]. However, the physiological roles of the different subpopulations, their anatomical distribution and targets, remain unknown.

POMC is a propeptide predominantly expressed in the pituitary gland and the hypothalamus. After posttranslational processing, POMC gives rise to bioactive products with different actions. In particular, hypothalamic melanocortins α - and β -melanocyte-stimulating hormones (α - and β -MSH) suppress food intake, stimulate energy expenditure, and enhance insulin sensitivity through melanocortin receptor 4 (MC4R) [6]. Mutations in *Pomc* or *Mc4r* genes lead to severe hyperphagia and early-onset obesity in both humans and mice [7–13]. Moreover, diet-induced obesity triggers hypothalamic inflammation and endoplasmic reticulum stress which, in turn, impairs POMC processing, leading to decreased production of α -MSH and, consequently, to hyperphagia and overweight [14–16]. Notably, the characterization of POMC neurons and their circuits in mouse models has contributed to the design and study of different drugs for the treatment of obesity and type 2 diabetes, some of which are already available for patients [17–22].

Arc-POMC neurons project to different hypothalamic areas involved in energy balance such as the paraventricular (PVH), ventromedial, dorsomedial (DMH), and lateral (LH) hypothalamic nuclei, as well as extrahypothalamic regions involved in other functions like autonomic control, reward, and analgesia [23–25]. The DMH is known to be involved in the regulation of food intake and energy expenditure [26]. It receives afferent projections from the Arc and, in turn, sends efferent projections to the PVH and the LH [26]. Lesions of the DMH result in hypophagia and weight loss, supporting a role for this region in the stimulation of food intake [27]. Moreover, it has been shown that the orexigenic neuropeptide Y (NPY) is involved in DMH induction of food intake in lactating rats and in some models of obese mice [28–31]. Finally, there is some evidence showing that melanocortin receptors mediate an inhibitory tone onto DMH-NPY expression [32]. However, a link between Arc-POMC expression and DMH-NPY expression remains to be established.

It has been shown that POMC neurons are heterogeneous regarding their electrophysiological response to peripheral signals (e.g., glucose) and the receptors they express (e.g., leptin, serotonin, insulin, and estrogen receptors) [33–35]. Moreover, subsets of hypothalamic POMC neurons express and corelease either gamma-aminobutyric acid (GABA) or glutamate (Glu) [36–38], encompassing 45–54% and 7–43% of POMC neurons, respectively [39,40]. Given the antagonistic responses elicited by GABA and Glu in neuronal excitability, it is speculated that these subpopulations carry out different functions. Recent studies applying single-cell RNA sequencing showed that Arc-POMC neurons can be classified into subpopulations with different transcriptomic profiles [3–5]. In particular, Campbell showed that Arc-POMC neurons can be classified into three clusters. While two of these clusters express mainly GABAergic markers (*Gad1* and *Gad2*), the third one expresses principally a glutamatergic marker (*Vglut2*). Interestingly, there is a preponderant response to the fasting of GABAergic-POMC clusters, evidenced by changes in their gene

expression profiles [3]. However, the physiological roles of these subpopulations in the regulation of energy homeostasis remain to be elucidated. In the present study, we aimed to dissect the contribution of the subpopulation of Arc-POMC neurons cosecreting GABA in the control of energy balance by expressing *Pomc* exclusively in GABAergic-POMC neurons of *Pomc*-deficient mice.

2. MATERIALS AND METHODS

2.1. Animal care

Mice were kept under standard laboratory conditions, with controlled photoperiod (lights on from 7 a.m. to 7 p.m.), tap water, and standard lab chow available ad libitum. Mice were weaned at P21. All procedures were approved by the Institutional Animal Care and Use Committee of the Facultad de Medicina, Universidad de Buenos Aires.

2.2. Mouse lines

Ai14:Gad2-CreER. To characterize the expression pattern of Cre recombinase in the knock-in *Gad2-CreERT²* mouse line (*Gad2-CreER*, for simplicity), we used *Ai14:Gad2-CreER* mice obtained by crossing heterozygous *Gad2-CreER* (*Gad2tm1^{Cre/ERT2}Z^h*; The Jackson Laboratory, stock: 010,702 [41]) and homozygous *Ai14* mice (B6. Cg-Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze/J}; The Jackson Laboratories, stock 7914 [42]). *Gad2-CreER* mice express a tamoxifen-inducible Cre in GABAergic neurons. Cre-expressing neurons were reported by the fluorescent protein tdTomato after tamoxifen treatment at P60 (see Section 2.3).

arcPomc^{-/-}:Gad2-CreER. *arcPomc^{+/-}* mice [11] were crossed with *Gad2-CreER* mice to obtain *arcPomc^{+/-}:Gad2-CreER* mice. Thereafter, *arcPomc^{+/-}:Gad2-CreER* mice were mated with *arcPomc^{+/-}* mice to obtain *arcPomc^{-/-}:Gad2-CreER* reversible POMC-deficient mice and obese *arcPomc^{-/-}* or normal weight *Gad2-CreER* control groups.

arcPomc^{-/-}:CreER mice were obtained as described in [11]. Briefly, the *arcPomc^{-/-}* line was crossed with CreER transgenic mouse line (B6. Cg-Tg[cre/Esr1]5Amc/J; The Jackson Laboratory, stock: 004682), in which Cre is driven by a ubiquitously active CAAG promoter and induced by tamoxifen.

2.3. CRE induction by tamoxifen

Tamoxifen (Sigma) was prepared in sesame oil (Sigma) as described previously [43]. *Gad2-CreER* mice were i. p. injected with 150 mg/kg/day tamoxifen at p60 or with 100 mg/kg/day at P25, during five consecutive days, with a solution of 15 or 10 mg tamoxifen/mL, respectively. CreER mice were i. p. injected with one dose of tamoxifen (50 mg/kg).

2.4. General study design

Two cohorts of *arcPomc^{-/-}:Gad2-CreER* mice and their control littermates were generated by breeding strategies described above and treated with tamoxifen at postnatal days 25 or 60 \pm 3 days (P25 and P60 cohorts, respectively). Mice were individually housed within the first week after weaning, and body weight and food intake were registered twice a week. Basal glycemia and glucose tolerant tests (GTTs) were measured after overnight fasting (6 pm–10 am) at the 7th (P60-PRE) and 14th weeks of age (P25 and P60-POST) by collecting blood samples from the tail with a OneTouch® glucometer (LifeScan, Johnson & Johnson). For GTTs, mice received an i. p. injection of glucose (2 g/kg; Sigma) as described previously [44]. Fasting-induced hyperphagia was measured in P60-treated mice at the 15th week of age, as described [11]. Briefly, after a 24-hour fast (10 am–10 am) ad libitum food intake of the next 24 h was measured and expressed




Figure 1: Generation of a mouse line expressing *Pomc* restricted to GABAergic-POMC neurons. (A) *Pomc*⁻ allele contains an insertion of a neomycin resistance cassette (neo), flanked by *loxP* sites (triangles), interrupting *Pomc* neuronal enhancer activity (blue circle: nPE1 enhancer; gap after neo: Deleted nPE2 enhancer). *Gad2-CreER* line drives *Pomc* reactivation restricted to GABAergic-POMC neurons after i. p. injection of tamoxifen (TAM) in *arcPomc*^{-/-}:*Gad2-CreER* mice (rescued). The POMC-cellular specificity of *Pomc* reactivation is due to the endogenous *Pomc* regulatory elements in the knock-in *arcPomc*^{-/-} allele. Gray circle: *Pomc* pituitary promoter. Arc: arcuate *Pomc* transcription. Black rectangle: *Pomc* exon 1. (B) Representative coronal brain sections of *Gad2-CreER*, *arcPomc*^{-/-}, and rescued *arcPomc*^{-/-}:*Gad2-CreER* mice showing ACTH immunopositive neurons. The arrows indicate rescued POMC expression in GABAergic neurons. 3V, third ventricle; Arc: arcuate nucleus. Magnification bars: 200 μ m. (C) Percentage of POMC neurons in *arcPomc*^{-/-}:*Gad2-CreER*-rescued mice (GABAergic-POMC neurons), relative to POMC neurons in *Gad2-CreER* control mice, at different coronal levels of the hypothalamus. The rostrocaudal analysis shows no significant differences (OWA). Error bars: \pm SEM; n = 3. (D) Representative coronal brain section of an *arcPomc*^{-/-}:*Gad2-CreER*-rescued mouse subjected to fluorescent hybridization for *Gad1* (GABAergic neurons, green) and immunohistochemistry for ACTH (POMC neurons, red). The white rectangle depicts the area of magnified images shown below. Magnification bars: 200 μ m (upper picture) and 50 μ m (bottom pictures). The white arrowheads point to GABAergic-POMC neurons. (E) Representative coronal brain section of Ai14:*Gad2-CreER* mice showing recombination in GABAergic cells (red, "*Gad2CreER-Tom*") and ACTH immunopositive neurons (green, "POMC"). The white rectangle depicts the area of the magnified image shown on the right. The white arrows point to GABAergic-POMC neurons. Magnification bars: 200 μ m (left picture) and 50 μ m (right picture).

relative to the average daily food intake of the previous week. At the 16th week of age, mice were anesthetized with 5% chloral hydrate at 3 pm and perfused with 4% paraformaldehyde (PFA) for subsequent immunohistochemistry or in situ hybridization. Alternatively, mice were killed by cervical dislocation at 11 am, and fresh hypothalami were removed and fast-frozen for RNA extraction, while inguinal (unilateral) and gonadal (bilateral) fat pads as well as livers were weighed.

2.5. Hypothalamic *Pomc* mRNA expression

Hypothalamic total RNA was prepared following phenol-chloroform extraction using TriPure (Roche) and then treated with RNase-free DNase I (Ambion). First-strand cDNA was synthesized with random primers using High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). *Pomc* mRNA was identified by relative quantitative RT-PCR using PowerUp SYBR Green Master Mix (Thermo Fisher Scientific) with specific primers (F- 5'-CTCCTGCTTCAGACCTCCAT-3' and R- 5'-CAGTCAGGGGCTGTTCATCT-3'; amplicon size: 169 bp), relative to endogenous β -actin (primers F-5'-AGAGGAAATCGTGCCTGAC-3' and R- 5'-CAATAGTGATACCTGGCCGT-3'; amplicon size: 138bp). Samples were run on the 7500 Real-Time PCR machine (Applied Biosystem),

and the results were analyzed by the $2^{-\Delta\Delta CT}$ relative quantitation method [45].

2.6. Immunohistochemistry

PFA fixed brains were collected and cut into 35 μ m coronal sections with a frozen microtome (Leica). For immunohistochemistry, floating brain sections were treated with 1% H₂O₂ for 30 min and, after PBS washing, sections were incubated with rabbit polyclonal anti-rat-ACTH antibody (1:1000, A.F. Parlow, National Hormone and Peptide Program, Harbor-UCLA Medical Center, Torrance, CA, USA), overnight at 4 $^{\circ}$ C. The following day, sections were incubated first with the secondary antibody (biotinylated anti-rabbit IgG made in goat, 1:200, Vector Labs) and then with avidin-radish peroxidase complex (Vectastain ABC Kit, Vector Labs). Finally, slices were developed with diaminobenzidine (DAB) (Vector Labs), mounted on 1% gelatin (Sigma) coated glass slides with Canada balsam, and analyzed with an Olympus BX53 microscope, coupled with a Q-Color5 digital camera and Capture Q software. The quantitative rostrocaudal analysis was performed manually and blinded by two investigators, between -1.22 mm and -2.06 mm from bregma with Fiji software




Figure 2: *Pomc* restoration in GABAergic-POMC neurons improves energy balance and glycemia. (A, H) Average of daily food intake from the 6th to 8th and from the 12th to 14th weeks of age (before (PRE) and after (POST) treatment, respectively) of mice treated i. p. with tamoxifen (TAM) at P60. N = 9–14. (B, I) Body weight curves of mice treated with TAM at P60. Graphs on the right show comparisons of body weights before (PRE, 7 weeks old) and after (POST, 16 weeks old) TAM treatment. N = 9–14. (C, J) Fast induced hyperphagia of mice previously treated with TAM at P60. Bars correspond to averages of 24-hour refeeding data expressed as a percentage of prefasting food intake. The degree of hyperphagia can be estimated above the dashed lines. N = 9–14. (D, K) Basal glycemia measured before (PRE, 7 weeks old) and after (POST, 14 weeks old) TAM treatment. N = 5–9. (E–G and L–N) Gonadal and inguinal fat pad weights and liver weights of mice treated with TAM at P60. N = 4–10. (O, R) Average of daily food intake (from the 12th to 14th weeks of age) of mice treated i. p. with TAM at P25. N = 4–11. (P–S) Body weight curves of mice treated with TAM at P25. Graphs on the right show comparisons of body weights before (PRE, 3 weeks old) and after (POST, 16 weeks old) TAM treatment. N = 4–12. (Q, T) Basal glycemia measured at the 14th weeks of age of mice treated with TAM at P25. N = 3–9. (A–G) and (O–Q) correspond to females and (H–N) and (R–T) correspond to males. In all cases, error bars: \pm SEM. (A, B, D, H, I, K, P, S) RMA: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; # $P < 0.05$, ## $P < 0.01$, and ### $P < 0.001$: pairwise comparisons within the same group, before (PRE) versus after (POST) *Pomc* restoration (Bonferroni). (C, E–G, J, L–N, O, Q, R, T) OWA: * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ (Bonferroni).




Figure 3: Partial nonspecific rescue of *Pomc* fails to normalize food intake and body weight. (A, B) Average of daily food intake from the 6th to 8th and from the 12th to 14th weeks of age (before (PRE) and after (POST) treatment, respectively) in female (A) and male (B) mice treated i. p. with a single low dose of tamoxifen (TAM) at P60. N = 4–7. (C, D) Body weight curves of female (C) and male (D) mice. The graphs on the right show comparisons of body weights before (PRE, 7 weeks old) and after (POST, 16 weeks old) TAM treatment. N = 4–7. (E, F) Comparison of body weights of female (E) and male (F) mice before and after GABA-specific or nonspecific *Pomc* rescue at P60. The data of *arcPomc*^{-/-}:*Gad2-CreER* and *Gad2-CreER* phenotypes correspond to the same mice shown in Figure 2B (females) and Figure 2I (males). N = 4–14. Error bars: \pm SEM. RMA: * $P < 0.05$, and ** $P < 0.01$, *** $P < 0.001$. Pre versus Post: # $P < 0.05$, ## $P < 0.01$, and ### $P < 0.001$ (Bonferroni).

[46]. Results were expressed as the percentage of POMC + neurons in *arcPomc*^{-/-}:*Gad2-CreER*- or *arcPomc*^{-/-}:*CreER*-rescued mice relative to POMC + neurons in *Gad2-CreER* or *CreER* control mice, respectively. For anti-ACTH immunofluorescence, sections were incubated with AlexaFluor488-coupled anti-rabbit IgG (1:1000, Invitrogen) and colocalized with endogenous red fluorescent signal (tdTomato) of Ai14 mice. In this case, images were obtained with an Olympus FV1000 confocal microscope, and slices between -1.34 mm and -1.7 mm from bregma were analyzed with Fiji software.

2.7. Chromogenic (ISH) and fluorescent in situ hybridization (FISH)

For *Npy* probe cloning, an insert of 493 bp (NM_023456.3, 32–524 bp) was amplified by RT-PCR from RNA of mouse embryos using gene-specific primers: mNpyF: 5' TCTCACAGAGGCCACCCAGAG 3' and mNpyR: 5' CAACAACAACAAGGGAAATG 3'. The PCR product was cloned (pGEM-T Easy Vector, Promega, Cat. A1360) and sequenced (SAI, University of Murcia). *Gad1* probe (NM_008077.5, 2198–2949 bp) was kindly provided by Dr. Z. Katarova. For both ISH and FISH, PFA-fixed brains were embedded in 15% gelatin (Sigma) in PBS and then collected and cut into 20 μ m coronal sections with a frozen cryostat (Thermo Fisher Scientific). ISH was performed as described [47] with minor modifications. Briefly, *Npy* probe was synthesized using a digoxigenin- (DIG-) labeling kit (Roche), incubated with ON at 72 °C, and developed with NBT/BCIP (Roche). For FISH, mounted sections were incubated with DIG-labeled *Gad1* RNA probe overnight at 55 °C. DIG probe was detected with an anti-Dig-POD antibody (1:150, Roche)

and developed with Alexa-488 Tyramide Signal Amplification Kit (1:100, Life Technologies). After FISH protocol, sections were incubated with rabbit polyclonal anti-rat-ACTH antibody (1:100, A.F. Parlow, National Hormone and Peptide Program, Harbor-UCLA Medical Center, Torrance, CA, USA) followed either by an anti-rabbit-Cy3 antisera (1:500, Jackson ImmunoResearch, Figure 1) or by a biotinylated anti-rabbit IgG antisera (1:200, Vector Labs) combined with streptavidin-Alexa Fluor-647 (1:1000, Jackson ImmunoResearch, Figure 6). Sections were mounted with VECTASHIELD (Vector Labs) and analyzed with Axio Imager M2 motorized fluorescent microscope with Apotome2 structured illumination (Zeiss). Photographs were obtained with a monochromatic camera, and fluorophores were artificially colored by Neurolucida Software (MBF Bioscience). For double-labeling analysis (*Gad1*+/*POMC*+), one image per hemisphere was taken and manually analyzed with Fiji software. Results were reported as the percentage of POMC + neurons expressing *Gad1*. For triple-labeling analysis (*retrobeads*+/*Gad1*+/*ACTH*+), z-stack images were taken every 5 μ m for a total of 15 μ m (one per hemisphere of arcuate nucleus). Stack images were collapsed in a unique plane and analyzed with Fiji software. Quantitative analysis was manually performed by two investigators in sections between -1.4 and -1.9 mm distance to bregma. 32.4 ± 2.5 POMC neurons per hemisphere of the hypothalamus were quantified. Results were expressed as the percentage of cells labeled with red (*retrobeads*) and blue (*POMC*) signals (presumably POMC neurons projecting to the DMH) that were also green (*Gad1*+).




Figure 4: POMC fibers projecting to the dorsomedial hypothalamic nucleus (DMH) of POMC rescued mice. (A) Schematic coronal brain section adapted from [57]. The red rectangle signalizes the brain area shown in the upper pictures of (B) and (C). (B) Upper panels: representative coronal brain sections of *Gad2-CreER* and rescued *arcPomc^{-/-}:Gad2-CreER* female mice showing POMC immunopositive neurons in the arcuate nucleus (Arc) and POMC immunoreactive fibers (POMC-IRFs) in the DMH. Magnification bars: 200 μm . (C) Upper panels: representative coronal brain sections of *CreER* and rescued *arcPomc^{-/-}:CreER* female mice showing POMC immunopositive neurons in the Arc and POMC-IRFs in the DMH. Magnification bars: 200 μm . (B and C) Bottom panels: magnified areas of DMH taken from the upper pictures showing POMC-IRsF. Magnification bars: 50 μm .

2.8. Stereotactic surgery

Red fluorescent latex microspheres (Lumaflores Inc.) were used as a retrograde neuronal tracer to track projections to DMH. Adult wild-type C57Bl/6 (10–12 weeks old) female mice ($n = 4$) were anesthetized using isoflurane in O_2 (2.5% induction, 0.8% maintenance) and placed in stereotactic head apparatus (Kopf Instruments). Ophthalmic ointment was applied to both eyes to prevent drying, and 2% lidocaine hydrochloride was injected subcutaneously for local analgesia. Multiple rounds of betadine and 70% ethanol wipes were applied to the scalp and then a midline 8 mm incision was made to access the skull. Then, a 1 mm diameter craniotomy was drilled. The retrobeads solution (500 nl, 1:200 in saline) was injected bilaterally at 100 nl/min at the following stereotaxic coordinates: 1.7 mm anteroposterior, ± 1.3 mm mediolateral relative to bregma, and -4.80 mm dorsoventral from the cortical surface using a 10° angle to avoid ventricles and damage of the superior sagittal sinus. After surgery, the skin was sutured and postoperative care included analgesia (i.p. administration of flunixin 5 mg/kg) for three consecutive days. Two weeks after surgery (14–19 days), mice were perfused and brains were removed and fixed in 4% PFA to prepare them for in situ hybridization and immunohistochemistry.

2.9. Statistical analysis

All data are presented as the mean \pm SEM and were analyzed by Student's unpaired two-tailed *t*-test, one-way or two-way ANOVA (OWA, TWA), or repeated measures ANOVA (RMA), using GraphPad Prism version 6.00 for Windows (GraphPad Software, La Jolla California, USA). Post hoc Bonferroni's test was used when necessary. $P < 0.05$ was considered significant.

3. RESULTS

3.1. *Pomc* expression restricted to GABAergic neurons

In order to elucidate the physiological role of POMC expressed in the subpopulation of arcuate (Arc) GABAergic-POMC neurons, we used a previously developed mouse model carrying a mutant conditional *Pomc* allele (*arcPomc^{-/-}*), which harbors a loxP-flanked neomycin cassette within *Pomc* neuronal enhancer region, which selectively prevents neuronal transcription in the Arc but not in the pituitary gland or the nucleus of the solitary tract (Figure 1A and [11]). Homozygous *arcPomc^{-/-}* mice are hyperphagic and develop early-onset extreme obesity [11]. However, eutopic *Pomc* expression can be achieved by crossing *arcPomc^{-/-}* mouse line with another one expressing Cre recombinase [11]. Interestingly, we previously found that food intake and body weight can be greatly improved after *Pomc* recovery by a ubiquitously expressed Cre recombinase [11]. In the present study, in order to assure *Pomc* expression restricted to GABAergic neurons, *arcPomc^{-/-}* mouse line was crossed with another *knock-in* line expressing a tamoxifen-inducible Cre driven by the *Gad2* promoter (*Gad2-CreER* mouse line [41], Figure 1A).

In order to selectively recover *Pomc* expression in GABAergic neurons, we treated *arcPomc^{-/-}:Gad2-CreER* mice with tamoxifen at P60, which led to POMC expression in $25.1 \pm 3.3\%$ and $23.2 \pm 2.3\%$ (mean \pm SEM) of POMC neurons in female and male mice, respectively, compared to *Gad2-CreER* control mice (Figure 1B,C). Moreover, hypothalamic *Pomc* rescue restricted to GABAergic neurons resulted in $23.3 \pm 7.1\%$ and $14.8 \pm 1.3\%$ of *Pomc* mRNA compared to *Gad2-CreER* control mice (mean \pm SEM) in females and males, respectively. POMC immunopositive neurons in rescued mice showed homogeneous distribution across the anterior–posterior axis of the arcuate nucleus (OWA: $P > 0.05$ for both sexes, Figure 1C). In order to




Figure 5: DMH-NPY expression is normalized after POMC expression restricted to GABAergic neurons. (A, C) Coronal brain sections of mice subjected to in situ hybridization, showing *Npy* mRNA (purple) in the dorsomedial hypothalamic nucleus (DMH) and arcuate nucleus (Arc) of the hypothalamus. 3V, third ventricle. ME: median eminence. Magnification bars: 200 μ m. (B, D) Average of DMH-*Npy* and Arc-*Npy* neurons per hemisection for each genotype. Error bars: \pm SEM. OWA: * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ (Bonferroni). N = 3–4.

establish the specificity of the *Gad2-CreER* driver, we performed double labeling by in situ hybridization and immunohistochemistry in brain slices of *arcPomc*^{-/-}:*Gad2-CreER* mice which revealed that at least $94.2 \pm 0.7\%$ of rescued POMC neurons are GABAergic (Figure 1D).

Since, according to previous quantifications of *Gad2*⁺-POMC neurons [39], we expected *Pomc* recovery in $\sim 45\%$ of POMC neurons, to determine the extension of *Gad2-CreER* activity, we generated Ai14:*Gad2-CreER* reporter mice. After tamoxifen administration, we found tdTomato expression in $26.7 \pm 1.9\%$ (mean \pm SEM, n = 3) of POMC neurons (Figure 1E), which suggests incomplete recombination induced by the *Gad2-CreER* driver.

3.2. GABAergic-POMC neurons regulate energy balance

With the aim of studying the role of Arc-POMC expressed in GABAergic neurons in the regulation of energy balance, we rescued *Pomc* expression in *arcPomc*^{-/-}:*Gad2-CreER* mice. After tamoxifen treatment at P60, we found that both female and male *arcPomc*^{-/-}:*Gad2-CreER*-rescued mice normalized food intake (Figure 2A,H). Remarkably, female *arcPomc*^{-/-}:*Gad2-CreER* mice greatly improved body weight after *Pomc* rescue: they were $\sim 70\%$ overweight before treatment and, after losing weight, they remain only $\sim 29\%$ overweight, compared to *Gad2-CreER* control littermates (Figure 2B). Furthermore, while *arcPomc*^{-/-} female mice showed decreased fasting-induced hyperphagia, rescued *arcPomc*^{-/-}:*Gad2-CreER* mice displayed normal response (Figure 2C). Fasting glycemia and glucose tolerance, which were increased and decreased, respectively, in *Pomc*-deficient female mice, were completely restored by *Pomc*

recovery in GABAergic-POMC neurons (Figure 2D and Supplementary Figures 1A–1D). In females, body weight improvement was due to fat loss, since inguinal and gonadal fat pads as well as liver weights were significantly lower in *arcPomc*^{-/-}:*Gad2-CreER*-rescued mice compared to obese *arcPomc*^{-/-} control mice (Figure 2E–G).

ArcPomc^{-/-}:*Gad2-CreER* male mice were rescued at P60 completely normalized food intake (Figure 2H). However, they showed mild body weight loss, with body weights that become significantly different from those of *arcPomc*^{-/-} mice seven weeks after *Pomc* recovery (RMA, Bonferroni post hoc test: $P < 0.01$; Figure 2I). Regarding fasting-induced hyperphagia, while *arcPomc*^{-/-} male mice showed decreased food intake in the refeeding period, rescued *arcPomc*^{-/-}:*Gad2-CreER* mice showed no significant differences with either *arcPomc*^{-/-} or *Gad2-CreER* control mice (Figure 2J). Fasting glycemia was not increased in *Pomc*-deficient male mice, at least until the 14th week of age (Figure 2K). However, these mice were intolerant to a glucose overload, a condition that was reverted by *Pomc* rescue (Supplementary Figures 1E–1H). Finally, weights of inguinal and gonadal fat pads of rescued *arcPomc*^{-/-}:*Gad2-CreER* male mice were not significantly different from those of *arcPomc*^{-/-} or *Gad2-CreER* mice, while liver weights were higher than those of *Gad2-CreER* mice (Figure 2L–N). The above results show that the *Pomc* expression in GABAergic-POMC neurons is sufficient to maintain normal food intake. However, despite *Pomc* rescue in obese mice at P60 leads to body weight loss, *arcPomc*^{-/-}:*Gad2-CreER*-rescued mice remain heavier than *Gad2-CreER* littermates. Since we previously found that *Pomc* recovery by a ubiquitously expressed CreER only leads to normal body weight when tamoxifen is injected to normal weight mice [11], we sought to




Figure 6: Arcuate POMC neurons projecting to DMH are mainly GABAergic. (A) Representative image of coronal brain sections showing bilateral injection of retrobeads (red) in the DMH of a wild-type female mouse. Magnification bar: 200 μ m. (B) Representative fluorescent microphotograph of a coronal brain section at -1.58 mm from bregma, subjected to POMC immunostaining (blue) and in situ hybridization against *Gad1* (green). Red: retrobeads coming from the DMH. Magnification bar: 200 μ m. (C–F) Magnified images of the square shown in (B). Arc: arcuate nucleus. 3V: third ventricle. Magnification bar: 50 μ m.

Table 1 — Brain areas with GABAergic POMC-immunopositive fibers (POMC-IRF). Numbers indicate the quantity of *arcPomc*^{-/-}:*Gad2*-CreER female and male mice rescued at P60 showing POMC-IRF projecting to a specific brain area, from a total of 6 analyzed mice.

BRAIN AREAS WITH POMC-IRF	FEMALES	MALES
Dorsomedial hypothalamic nucleus	6	6
Paraventricular hypothalamic nucleus	4	4
Posterior hypothalamic nucleus	3	4
Bed nucleus of the stria terminalis	2	3
Periventricular hypothalamic nucleus	0	2
Paraventricular thalamic nucleus	2	0
Anterior commissure	2	0
Suprachiasmatic nucleus	0	2
Preoptic nucleus	1	1
Tuber cinereum area	1	0
Lateroanterior hypothalamic nucleus	1	0
Retrochiasmatic area	1	0
Premammillary nucleus	1	0

elucidate if *Pomc* rescue restricted to GABAergic-POMC neurons in juvenile mice has similar consequences. Therefore, another cohort of mice was injected with tamoxifen at P25, before the onset of obesity. The magnitude of *Pomc* recovery in *arcPomc*^{-/-}:*Gad2*-CreER mice at P25 was similar to that of mice treated at P60: $22.31 \pm 1.83\%$ and $26.17 \pm 3.94\%$ (mean \pm SEM) of POMC + cells compared to control *Gad2*-CreER mice in females and males, respectively. We found that *Pomc* rescue in GABAergic-POMC neurons at P25 prevented hyperphagia in females and decreased hyperphagia in male *arcPomc*^{-/-}:*Gad2*-CreER mice compared to *arcPomc*^{-/-} controls (Figure 20,R). Moreover, *Pomc* rescue at P25 delayed the onset of obesity in both female and male *arcPomc*^{-/-}:*Gad2*-CreER mice: while

arcPomc^{-/-} mice became significantly heavier than *Gad2*-CreER control mice since the 7th and 4th week of age (RMA, Bonferroni post hoc test: $P < 0.001$ and $P < 0.05$, female and male, respectively), both female and male *arcPomc*^{-/-}:*Gad2*-CreER mice rescued at P25 became overweight only since the 9th week of age (RMA, Bonferroni post hoc test: $P < 0.05$ for both sexes; Figure 2P,S). Finally, *Pomc* rescue at P25 prevented hyperglycemia in both female and male *arcPomc*^{-/-}:*Gad2*-CreER mice (Figure 2Q,T). Altogether, these results suggest a key role of GABAergic-POMC neurons in the control of energy balance.

3.3. Partial nonspecific *Pomc* rescue leads to mild body weight improvement

In order to emphasize the significance of GABAergic-POMC neurons in the regulation of energy balance, we compared the improvement of obesity after *Pomc* rescue, specifically in GABAergic neurons, with that achieved by rescuing POMC in a similar number of neurons but in a nonspecific manner. With the purpose of inducing partial *Pomc* rescue driven by a ubiquitously expressed Cre recombinase, we generated *arcPomc*^{-/-}:CreER mice and treated them with a single dose of tamoxifen (50 mg/kg) at P60. Tamoxifen treatment led to $32 \pm 3.6\%$ and $35.2 \pm 7.2\%$ (mean \pm SEM; females and males, respectively) of POMC + neurons in *arcPomc*^{-/-}:CreER mice compared to CreER controls (Supplementary Figure 2). Contrary to *Pomc* rescue in GABAergic neurons, nonspecific *Pomc* rescue failed to improve food intake in both female and male *arcPomc*^{-/-}:CreER mice (Figure 3A,B). Furthermore, nonspecific POMC rescue led to a mild loss of body weight in both female and male *arcPomc*^{-/-}:CreER mice compared to *arcPomc*^{-/-} mice (Figure 3C,D). Remarkably, body weight was significantly lower after *Pomc* rescue in GABAergic neurons compared to the nonspecific rescue (Figure 3E,F). Moreover,

while *arcPomc*^{-/-}:*Gad2*-CreER mice maintained or decreased body weight compared to their pretreatment stage (female and male, respectively), *arcPomc*^{-/-}:CreER mice further increased body weight after *Pomc* nonspecific rescue (Figure 3E,F). Altogether, the comparison of obesity improvement after GABAergic-specific versus nonspecific *Pomc* expression suggests a preponderant role of GABAergic-POMC neurons in the regulation of food intake and body weight.

3.4. Distribution of GABAergic-POMC immunoreactive fibers

In order to dissect the anatomical distribution of GABAergic-POMC immunoreactive fibers (POMC-IRFs), coronal and sagittal brain sections of rescued *arcPomc*^{-/-}:*Gad2*-CreER mice were subjected to immunohistochemistry against ACTH. As expected, since POMC is rescued in only ~25% of the total POMC neurons in these mice, in general, few isolated POMC-IRFs were detected. Remarkably, we consistently found POMC-IRFs in the DMH of all analyzed female and male rescued *arcPomc*^{-/-}:*Gad2*-CreER mice (Table 1, Figure 4B, and Supplementary Figure 3). To a lesser extent, paraventricular hypothalamic nucleus and the posterior hypothalamic nucleus, which are highly innervated by POMC neurons in wild-type mice, were also the main target sites of GABAergic-POMC neurons (Table 1 and Supplementary Figure 3). Surprisingly, other areas that are highly innervated by POMC neurons in wild-type mice showed POMC-IRFs only in a few proportion of *arcPomc*^{-/-}:*Gad2*-CreER-rescued mice (e.g., the bed nucleus of the stria terminalis, retrochiasmatic area, and the periventricular hypothalamic nucleus) (Table 1). On the contrary, all male and female *Gad2*-CreER control mice showed POMC-IRFs in the brain areas listed in Table 1 (*n* = 3 and 6, males and females, respectively; data not shown).

3.5. *Pomc* rescue in GABAergic-POMC neurons prevents DMH-*Npy* overexpression

Since we found that the DMH is a major target site of GABAergic-POMC neurons and there is some evidence suggesting that DMH-NPY neurons induce food intake in different models of obese mice, including MC4-RKO [28–31], we hypothesized that POMC expression restricted to GABAergic neurons could normalize food intake by inhibiting DMH-NPY expression. In order to study if there is a correlation between DMH-*Npy* expression and Arc-POMC expression, we performed in situ hybridization (ISH) for *Npy* in the brains of female mice previously treated with tamoxifen at P60 (Figure 5). As expected for normal-weight mice, *Gad2*-CreER mice showed very low expression in the DMH (Figure 5A,B). However, DMH-*Npy* expression raised more than 30 times in obese *arcPomc*^{-/-} mice compared to *Gad2*-CreER siblings. Interestingly, DMH-*Npy* expression was reestablished by *Pomc* expression restricted to GABAergic-POMC neurons in *arcPomc*^{-/-}:*Gad2*-CreER-rescued mice (Figure 5A,B).

Partial nonspecific *Pomc* rescue also led to DMH-*Npy* reduction (Figure 5C,D) and POMC-IRFs in the DMH of *arcPomc*^{-/-}:CreER mice (Figure 4C). However, contrary to GABAergic-specific *Pomc* rescue (which leads to DMH-*Npy* normalization), the expression remains higher in *arcPomc*^{-/-}:CreER mice compared to CreER controls (Figure 5D).

It is important to note that while the number of DMH-NPY neurons is increased in *arcPomc*^{-/-} mice, the number of Arc-NPY neurons is not significantly altered (Figure 5B,D). This finding suggests a preponderant role for POMC in the regulation of DMH-NPY expression compared to Arc-NPY expression.

In order to characterize Arc-POMC → DMH projections, we performed a triple labeling experiment in coronal brain sections of wild-type mice.




Figure 7: Model of an arcuate GABAergic-POMC → DMH-NPY circuit for the regulation of food intake. Upper panel: in wild-type mice (*Pomc*⁺), GABAergic-POMC neurons (red-filled circles with G) of the arcuate nucleus (Arc) suppress food intake by inhibiting NPY expression in dorsomedial hypothalamic nucleus (DMH) neurons (green open circles), maintaining a normal weight. Middle panel: in *Pomc* knockout mice (*Pomc*⁻), lack of *Pomc* expression in Arc neurons (red-open circles) results in DMH-NPY expression (green-filled circles) and increased food intake and obesity. Lower panel: *Pomc* expression restricted to GABAergic neurons (*Pomc*^{GABA}) recovers NPY inhibition resulting in normal food intake and weight loss.

First, neurons projecting to the DMH were labeled by stereotactic guided injections of retrobeads (RBs) to the DMH of anesthetized mice (*n* = 4). Two weeks later, POMC neurons were identified by immunohistochemistry and GABAergic neurons, by in situ hybridization with a *Gad1* probe, in fixed coronal brain sections of injected mice (Figure 6A). Our results show that 16.5 ± 2.0% (mean ± SEM) of Arc-POMC neurons are RB⁺, which presumably project to the DMH. Notably, 74.9 ± 5.4% (mean ± SEM) of POMC⁺/RB⁺ neurons are *Gad1*⁺, revealing that most of Arc-POMC → DMH projections are GABAergic (Figure 6B–F). Altogether, these results support the hypothesis of an inhibitory tone of arcuate GABAergic-POMC neurons on

DMH-NPY expression which may contribute to normal control of food intake.

4. DISCUSSION

In the present study, we show that the subpopulation of arcuate GABAergic-POMC neurons is involved in body weight regulation. Moreover, we found that the DMH is a major target site of GABAergic-POMC neurons followed by the PVN. In addition, we found that DMH-NPY expression is highly increased in POMC-deficient mice and it can be restored by *Pomc* expression restricted to GABAergic-POMC neurons. Finally, we determined that Arc-POMC neurons projecting to the DMH are mainly GABAergic.

Two previous studies showed the physiological consequences of arcuate *Pomc* expression restricted to well-defined subpopulations. Burke et al. showed that congenital *Pomc* expression in the subpopulation of neurons expressing 5-hydroxytryptamine 2c receptor (5-HT2CR), which encompasses about 40% of POMC neurons, prevents hyperphagia in both sexes and obesity in male mice [48]. In another study, it was shown that constitutive *Pomc* expression restricted to neurons expressing the leptin receptor (50%–80% of POMC neurons) leads to normal food intake and body weight [49]. In these studies, *Pomc* expression was recovered in 40%–70% of POMC neurons leading to more than 40% of *Pomc* mRNA expression, which is consistent with previous data showing that there is a threshold of 30% of *Pomc* mRNA and 30% of POMC + neurons above which mice maintain normal food intake and body weight [50,51]. In this regard, here we show the first example of food intake normalization in adult mice with less than 25% of hypothalamic *Pomc* mRNA expression. It is important to note that, in the present work, we show improvement of obesity phenotype by *Pomc* rescue in postnatal life rather than prevention of hyperphagia and overweight by *Pomc* expression during embryological development. Moreover, by inducing Cre activity in adult mice, we bypassed the developmental stage in which shifts between glutamate and GABA expression can occur, as has been previously shown for POMC neurons [52,53]. Most importantly, we found that *Pomc* expression in only 23%–25% of POMC neurons restricted to the GABAergic subpopulation has a similar impact as the nonspecific rescue of *Pomc* in ~75% of POMC neurons, in mice treated at P60 (compared to [11]).

Although it has been previously shown that *Gad2*+ POMC cells encompass ~50% of Arc-POMC neurons [39,40], *Gad2*-CreER driver induced *Pomc* expression only in ~23–25% of POMC neurons. Since we found similar results in Ai14:*Gad2*-CreER (only ~26% of POMC neurons were dtTomato+), the incomplete *Pomc* recovery may be attributed to low Cre expression in the knock-in mouse line, or to incomplete tamoxifen bioavailability, or both. In either case, it is interesting to note that *Pomc* expression restricted to half of GABAergic-POMC neurons is enough to normalize food intake, while partial nonspecific POMC rescue in *arcPomc*^{-/-}:CreER mice, with ~32–35% of POMC neurons expressing *Pomc*, is not. These results suggest that GABAergic-POMC neurons have a key role in the regulation of food intake.

We previously found that *Pomc* recovery in *arcPomc*^{-/-}:CreER at P25 completely prevented obesity [11]. Conversely, *arcPomc*^{-/-}:*Gad2*-CreER mice treated at P25, despite showing lower body weight and food intake than *arcPomc*^{-/-} mice, become overweight by the 9th week of age. However, since only half of GABAergic-POMC neurons recovered *Pomc* expression in these

mice, we cannot exclude that this subpopulation could be sufficient to maintain normal body weight in wild-type mice. Moreover, the participation of glutamatergic-POMC neurons in the regulation of energy balance cannot be ruled out. In this regard, Jones et al. rescued *Pomc* expression in *arcPomc*^{-/-} mice with a *Vglut2*-Cre driver constitutively expressed since the embryological developmental period [53]. However, they found *Pomc* expression also in GABAergic neurons, probably because of the postnatal shift from glutamate to GABAergic subpopulation.

Regarding glucose homeostasis, female *arcPomc*^{-/-} and *arcPomc*^{-/-}:*Gad2*-CreER mice show fasting hyperglycemia when they are 7 weeks old. Interestingly, only rescued *arcPomc*^{-/-}:*Gad2*-CreER mice normalized glycemia after tamoxifen treatment. In addition, both female and male *Pomc*-deficient mice were intolerant to a glucose overload, a condition that was reverted by *Pomc* rescue. These results can be attributed not only to body weight loss but also to *Pomc* recovery in GABAergic-POMC neurons, since we previously showed that normal-weight *arcPomc*^{-/-}:CreER female mice are intolerant to glucose overload and display decreased insulin sensitivity, both conditions that can be reverted by *Pomc* rescue [44].

Our results show that a major target site of GABAergic-POMC neurons is the DMH, which is known to be involved in the regulation of food intake and energy expenditure. In mice, DMH-*Npy* mRNA expression is only found in some models of genetic or high-fat diet-induced obesity and in mice subjected to chronic energy restriction. For example, A^y obese mice, which have an ectopic expression of the orexigenic agouti-signaling protein, show *Npy* mRNA expression in DMH [29]. It has been suggested that DMH-*Npy* mRNA expression may be caused by agouti inhibition of MC3/4R in DMH neurons. This hypothesis is supported by results showing that targeted disruption of the MC4-R gene leads to increased expression of *Npy* mRNA in the DMH [29]. Moreover, injection of MC3/4R-selective agonist melanotan II (MTII) in the DMH suppresses fasting or suckling-induced hyperphagia and suckling-induced *Npy* mRNA expression in the DMH of lactating rats [54]. Finally, α -MSH analog decreases food intake 1 h after injection in the DMH [55].

Although it is clear that MC3/4R stimulation inhibits DMH-*Npy* mRNA expression, it remained to be elucidated if Arc-POMC neurons mediate this regulation. Here, we show that obese *arcPomc*^{-/-} mice exhibit highly increased DMH-*Npy* mRNA expression. We propose that in wild-type mice α -MSH and/or other neuropeptides derived from POMC are secreted by Arc-POMC neurons and, by interacting with postsynaptic receptors (either directly on DMH-NPY neurons or indirectly on other DMH-intermediate neurons), inhibit DMH-NPY expression assuring normal food intake and body weight (Figure 7, upper panel). Accordingly, *Pomc* deficiency in *arcPomc*^{-/-} mice would lead to DMH-*Npy* mRNA overexpression which, in turn, would contribute to hyperphagia and obesity (Figure 7, middle panel). Finally, given that, as we found, ~75% of Arc-POMC neurons projecting to the DMH are GABAergic, it is probable that the normalization of DMH-NPY expression after *Pomc* restoration in GABAergic neurons accounts for food intake normalization leading to body weight loss (Figure 7, lower panel). This model could also explain the reduction in DMH-*Npy* expression in partially rescued *arcPomc*^{-/-}:CreER mice, since we expect ~40% of neurons recovering POMC with the nonspecific driver to be GABAergic. However, we cannot exclude a contribution of non-GABAergic neurons to the inhibition of DMH-*Npy* expression either in these mice or in wild-type mice. In addition, the Arc-POMC → DMH-NPY inhibition may also be involved in the regulation of fasting-induced hyperphagia, which is also disrupted in *arcPomc*^{-/-} mice and restored by *Pomc* rescue in GABAergic-POMC neurons. Although close appositions

between α -MSH-immunoreactive fibers and DMH-NPY neurons were previously observed in female rats [54], it remains to be elucidated if Arc-POMC neurons synapse with DMH-NPY neurons in mice. Alternatively, Arc-POMC-derived peptides may exert an indirect inhibitory regulation of DMH-Npy mRNA expression. Although here we propose that food intake may be regulated by an Arc-GABAergic-POMC \rightarrow DMH-NPY pathway, an Arc-GABAergic-POMC \rightarrow PVN circuit may also be involved in the normalization of food intake in rescued mice. In this regard, we found POMC-IRFs in the PVN of rescued mice and it has been demonstrated that POMC projections in the PVN increase the activity of MC4R neurons leading to a decrease in food intake [56].

In summary, our results show that the subpopulation of Arc-GABAergic-POMC neurons is a major regulator of food intake and body weight. Since *Pomc* reactivation in these neurons normalizes DMH-NPY expression and food intake, the circuit between Arc-POMC and DMH-NPY neurons could be a potential target for the treatment of obesity.

AUTHORS' CONTRIBUTIONS

VFB designed the study. MT, RA, EPB, MBT, and VFB performed the experiments. VFB, MR, EPB, and JLF contributed new reagents and analytic tools. MT, EPB, RA, MBT, and VFB analyzed the data. VB and MT wrote the paper. All authors revised and edited the manuscript and approved the final article.

ACKNOWLEDGMENTS

We thank Dr. Juan Belforte, Dr. Lorena Rela, Dr. Gustavo Murer, Dr. Guillermo Lanuza, and Dr. Lucía Franchini for valuable mice and reagents; Jessica Unger, Verónica Riso, Analía López Díaz, and Germán N. La Iacona, for technical assistance; Dr. Mario C. Perelló and Pablo N. De Francesco, for technical advice; and Johnson&Johnson Medical S.A., Argentina, for donating glucometer strips. This work was supported in part by Agencia Nacional de Promoción Científica y Tecnológica (PICT2014-2000), Consejo Nacional de Investigaciones Científicas y Técnicas (PIP 2014-487), Universidad de Buenos Aires, Argentina, The Committee for Aid and Education in Neurochemistry (CAEN-ISN), 2015-2016, and NARSAD Young Investigator Grant Award from Brain and Behavior Research Foundation (#23983, EPB), The Pew Charitable Trusts Repatriation Grant (EPB), Mario Hirsch from Fundación Bunge y Born/Fundación Williams (EPB), and The Spanish Ministry of Science and Innovation and European Regional Fund (MICINN and FEDER-PGC2018-098229-B-100, JLF). This study was awarded by the Fundación Gador (2017) and Fundación Florencio Fiorini (2018).

CONFLICT OF INTEREST

None.

APPENDIX A. SUPPLEMENTARY DATA

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.molmet.2020.100985>.

REFERENCES

[1] Berthoud, H.R., Münzberg, H., Morrison, C.D., 2017. Blaming the brain for obesity: integration of hedonic and homeostatic mechanisms. *Gastroenterology*. <https://doi.org/10.1053/j.gastro.2016.12.050>.

[2] Waterson, M.J., Horvath, T.L., 2015. Neuronal regulation of energy homeostasis: beyond the hypothalamus and feeding. *Cell Metabolism* 22(6):962–

970. <https://doi.org/10.1016/j.cmet.2015.09.026>. S1550-4131(15)00483-0 [pii].

[3] Campbell, J.N., Macosko, E.Z., Fenselau, H., Pers, T.H., Lyubetskaya, A., Tenen, D., et al., 2017. A molecular census of arcuate hypothalamus and median eminence cell types. *Nature Neuroscience*. <https://doi.org/10.1038/nn.4495>.

[4] Chen, R., Wu, X., Jiang, L., Zhang, Y., 2017. Single-cell RNA-seq reveals hypothalamic cell diversity. *Cell Reports* 18(13):3227–3241. <https://doi.org/10.1016/j.celrep.2017.03.004>.

[5] Lam, B.Y.H., Cimino, I., Poxel-Wolf, J., Nicole Kohnke, S., Rimmington, D., Iyemere, V., et al., 2017. Heterogeneity of hypothalamic pro-opiomelanocortin-expressing neurons revealed by single-cell RNA sequencing. *Molecular Metabolism*. <https://doi.org/10.1016/j.molmet.2017.02.007>.

[6] Cawley, N.X., Li, Z., Loh, Y.P., 2016. Biosynthesis, trafficking, and secretion of pro-opiomelanocortin-derived peptides. *Journal of Molecular Endocrinology*. <https://doi.org/10.1530/JME-15-0323>.

[7] Biebermann, H., Castaneda, T.R., van Landeghem, F., von Deimling, A., Escher, F., Brabant, G., et al., 2006. A role for beta-melanocyte-stimulating hormone in human body-weight regulation. *Cell Metabolism* 3(2):141–146. <https://doi.org/10.1016/j.cmet.2006.01.007>. S1550-4131(06)00031-3 [pii].

[8] Krude, H., Biebermann, H., Luck, W., Horn, R., Brabant, G., Gruters, A., 1998. Severe early-onset obesity, adrenal insufficiency and red hair pigmentation caused by POMC mutations in humans. *Nature Genetics* 19(2):155–157. <https://doi.org/10.1038/509>.

[9] Yaswen, L., Diehl, N., Brennan, M.B., Hochgeschwender, U., 1999. Obesity in the mouse model of pro-opiomelanocortin deficiency responds to peripheral melanocortin. *Nature Med* 5(9):1066–1070. <https://doi.org/10.1038/12506>.

[10] Yeo, G.S., Farooqi, I.S., Aminian, S., Halsall, D.J., Stanhope, R.G., O'Rahilly, S., 1998. A frameshift mutation in MC4R associated with dominantly inherited human obesity. *Nature Genetics* 20(2):111–112. <https://doi.org/10.1038/2404>.

[11] Bumashny, V.F., Yamashita, M., Casas-Cordero, R., Otero-Corchón, V., De Souza, F.S.J., Rubinstein, M., et al., 2012. Obesity-programmed mice are rescued by early genetic intervention. *Journal of Clinical Investigation* 122(11):4203–4212, 62543 [pii] 10.1172/JCI62543.

[12] Huszar, D., Lynch, C.A., Fairchild-Huntress, V., Dunmore, J.H., Fang, Q., Berkemeier, L.R., et al., 1997. Targeted disruption of the melanocortin-4 receptor results in obesity in mice. *Cell*. [https://doi.org/10.1016/S0092-8674\(00\)81865-6](https://doi.org/10.1016/S0092-8674(00)81865-6).

[13] Challis, B.G., Coll, A.P., Yeo, G.S., Pinnock, S.B., Dickson, S.L., Thresher, R.R., et al., 2004. Mice lacking pro-opiomelanocortin are sensitive to high-fat feeding but respond normally to the acute anorectic effects of peptide-YY(3-36). *Proceedings of the National Academy of Sciences of the U S A* 101(13):4695–4700. <https://doi.org/10.1073/pnas.0306931101>, 0306931101 [pii].

[14] Cakir, I., Nilini, E.A., 2019. Endoplasmic reticulum stress, the hypothalamus, and energy balance. *Trends in Endocrinology and Metabolism*. <https://doi.org/10.1016/j.tem.2019.01.002>.

[15] Araujo, E., Moraes, J., Cintra, D., Velloso, L., 2016. Mechanisms IN endocrinology: hypothalamic inflammation and nutrition. *European Journal of Endocrinology* 175(3):R97–R105.

[16] Schneeberger, M., Dietrich, M.O., Sebastián, D., Imbernón, M., Castaño, C., Garcia, A., et al., 2013. Mitofusin 2 in POMC neurons connects ER stress with leptin resistance and energy imbalance. *Cell*. <https://doi.org/10.1016/j.cell.2013.09.003>.

[17] D'Agostino, G., Lyons, D., Cristiano, C., Lettieri, M., Olarte-Sanchez, C., Burke, L.K., et al., 2018. Nucleus of the solitary tract serotonin 5-HT2C receptors modulate food intake. *Cell Metabolism*. <https://doi.org/10.1016/j.cmet.2018.07.017>.

[18] Burke, L.K., Doslikova, B., D'Agostino, G., Garfield, A.S., Farooq, G., Burdakow, D., et al., 2014. 5-HT obesity medication efficacy via POMC

- activation is maintained during aging. *Endocrinology*. <https://doi.org/10.1210/en.2014-1223>.
- [19] Secher, A., Jelsing, J., Baquero, A.F., Hecksher-Sorensen, J., Cowley, M.A., Dalboge, L.S., et al., 2014. The arcuate nucleus mediates GLP-1 receptor agonist liraglutide-dependent weight loss. *Journal of Clinical Investigation* 124(10):4473–4488, 75276 [pii] 10.1172/JCI75276.
- [20] Kühnen, P., Clément, K., Wiegand, S., Blankenstein, O., Gottesdiener, K., Martini, L.L., et al., 2016. Proopiomelanocortin deficiency treated with a melanocortin-4 receptor agonist. *New England Journal of Medicine*. <https://doi.org/10.1056/NEJMoa1512693>.
- [21] Zhou, L., Sutton, G.M., Rochford, J.J., Semple, R.K., Lam, D.D., Oksanen, L.J., et al., 2007. Serotonin 2C receptor agonists improve type 2 diabetes via melanocortin-4 receptor signaling pathways. *Cell Metabolism* 6(5):398–405. <https://doi.org/10.1016/j.cmet.2007.10.008>.
- [22] Doslíkova, B., Garfield, A.S., Shaw, J., Evans, M.L., Burdakov, D., Billups, B., et al., n.d. 5-HT2C receptor agonist anorectic efficacy potentiated by 5-HT1B receptor agonist coapplication: an effect mediated via increased proportion of pro-opiomelanocortin neurons activated. *Journal of Neuroscience* 33(23): 9800–9804, Doi: 33/23/9800 [pii] 10.1523/JNEUROSCI.4326-12.2013.
- [23] O'Donohue, T.L., Miller, R.L., Jacobowitz, D.M., 1979. Identification, characterization and stereotaxic mapping of intraneuronal alpha-melanocyte stimulating hormone-like immunoreactive peptides in discrete regions of the rat brain. *Brain Research* 176(1):101–123, 0006-8993(79)90873-4 [pii].
- [24] Bagnol, D., Lu, X.Y., Kaelin, C.B., Day, H.E., Ollmann, M., Gantz, I., et al., 1999. Anatomy of an endogenous antagonist: relationship between Agouti-related protein and proopiomelanocortin in brain. *Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 19(August 2016):RC26.
- [25] King, C.M., Hentges, S.T., 2011. Relative number and distribution of Murine hypothalamic proopiomelanocortin neurons innervating distinct target sites. *PLoS One* 6(10). <https://doi.org/10.1371/journal.pone.0025864>.
- [26] Branch, A., Shen, P., 2017. *Appetite and food intake*. CRC Press.
- [27] Bellinger, L.L., Williams, F.E., 2017. Aphagia and adipsia after kainic acid lesioning of the dorsomedial hypothalamic area. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*. <https://doi.org/10.1152/ajpregu.1983.244.3.r389>.
- [28] Li, C., Chen, P., Smith, M.S., 1998. The acute suckling stimulus induces expression of neuropeptide Y (NPY) in cells in the dorsomedial hypothalamus and increases NPY expression in the arcuate nucleus. *Endocrinology*. <https://doi.org/10.1210/endo.139.4.5905>.
- [29] Kesterson, R.A., Huszar, D., Lynch, C.A., Simerly, R.B., Cone, R.D., 1997. Induction of neuropeptide Y gene expression in the dorsal medial hypothalamic nucleus in two models of the agouti obesity syndrome. *Molecular Endocrinology* 11(5):630–637. <https://doi.org/10.1210/mend.11.5.9921>.
- [30] Guan, X.-M., Yu, H., Van der Ploeg, L.H., 1998. Evidence of altered hypothalamic pro-opiomelanocortin/neuropeptide Y mRNA expression in tubby mice. *Molecular Brain Research* 59(2):273–279. [https://doi.org/10.1016/S0169-328X\(98\)00150-8](https://doi.org/10.1016/S0169-328X(98)00150-8).
- [31] Guan, X.M., Yu, H., Trumbauer, M., Frazier, E., Van Der Ploeg, L.H.T., Chen, H., 1998. Induction of neuropeptide Y expression in dorsomedial hypothalamus of diet-induced obese mice. *NeuroReport*. <https://doi.org/10.1097/00001756-199810260-00015>.
- [32] Grove, K.L., Smith, M.S., 2003. Ontogeny of the hypothalamic neuropeptide Y system. *Physiology & Behavior* 79(1):47–63.
- [33] Sohn, J.W., Xu, Y., Jones, J.E., Wickman, K., Williams, K.W., Elmquist, J.K., 2011. Serotonin 2C receptor activates a distinct population of arcuate proopiomelanocortin neurons via TRPC channels. *Neuron* 71(3):488–497. <https://doi.org/10.1016/j.neuron.2011.06.012>.
- [34] Williams, K.W., Margatho, L.O., Lee, C.E., Choi, M., Lee, S., Scott, M.M., et al., 2010. Segregation of acute leptin and insulin effects in distinct populations of arcuate proopiomelanocortin neurons. *Journal of Neuroscience* 30(7):2472–2479. <https://doi.org/10.1523/JNEUROSCI.3118-09.2010>, 30/7/2472 [pii].
- [35] Hu, J., Jiang, L., Low, M.J., Rui, L., 2014. Glucose rapidly induces different forms of excitatory synaptic plasticity in hypothalamic POMC neurons. *PLoS One*. <https://doi.org/10.1371/journal.pone.0105080>.
- [36] Hentges, S.T., Nishiyama, M., Overstreet, L.S., Stenzel-Poore, M., Williams, J.T., Low, M.J., 2004. GABA release from proopiomelanocortin neurons. *Journal of Neuroscience* 24(7):1578–1583. <https://doi.org/10.1523/JNEUROSCI.3952-03.2004>, 24/7/1578 [pii].
- [37] Hentges, S.T., Otero-Corchon, V., Pennock, R.L., King, C.M., Low, M.J., 2009. Proopiomelanocortin expression in both GABA and glutamate neurons. *Journal of Neuroscience* 29(43):13684–13690. <https://doi.org/10.1523/JNEUROSCI.3770-09>, 29/43/13684 [pii].
- [38] Dicken, M.S., Tooker, R.E., Hentges, S.T., 2012. Regulation of GABA and glutamate release from proopiomelanocortin neuron terminals in intact hypothalamic networks. *Journal of Neuroscience* 32(12):4042–4048. <https://doi.org/10.1523/JNEUROSCI.6032-11.2012>, 32/12/4042 [pii].
- [39] Jarvie, B.C., Hentges, S.T., 2012. Expression of GABAergic and glutamatergic phenotypic markers in hypothalamic proopiomelanocortin neurons. *The Journal of Comparative Neurology* 520(17):3863–3876. <https://doi.org/10.1002/cne.23127>.
- [40] Wittmann, G., Hrabovszky, E., Lechan, R.M., 2013. Distinct glutamatergic and GABAergic subsets of hypothalamic pro-opiomelanocortin neurons revealed by in situ hybridization in male rats and mice. *The Journal of Comparative Neurology* 521(14):3287–3302. <https://doi.org/10.1002/cne.23350>.
- [41] Taniguchi, H., He, M., Wu, P., Kim, S., Paik, R., Sugino, K., et al., 2011. A resource of cre driver lines for genetic targeting of GABAergic neurons in cerebral cortex. *Neuron* 71(6):995–1013. <https://doi.org/10.1016/j.neuron.2011.07.026>.
- [42] Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., et al., 2010. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nature Neuroscience* 13(1):133–140 nn.2467 [pii] 10.1038/nn.2467.
- [43] Metzger, D., Chambon, P., 2001. Site- and time-specific gene targeting in the mouse. *Methods* 24(1):71–80. <https://doi.org/10.1006/meth.2001.1159>. S1046-2023(01)91159-4 [pii].
- [44] Alsina, R., Trotta, M., Bumacshny, V.F., 2018. Hypothalamic proopiomelanocortin is necessary for normal glucose homeostasis in female mice. *Frontiers in Endocrinology* 9:554. <https://doi.org/10.3389/fendo.2018.00554>.
- [45] Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25(4):402–408. <https://doi.org/10.1006/meth.2001.1262>. S1046-2023(01)91262-9 [pii].
- [46] Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., et al., 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods*. <https://doi.org/10.1038/nmeth.2019>.
- [47] Hauptmann, G., 2015. *In situ hybridization methods*.
- [48] Burke, L.K., Doslíkova, B., D'Agostino, G., Greenwald-Yarnell, M., Georgescu, T., Chianese, R., et al., 2016. Sex difference in physical activity, energy expenditure and obesity driven by a subpopulation of hypothalamic POMC neurons. *Molecular Metabolism* 5(3):245–252. <https://doi.org/10.1016/j.molmet.2016.01.005>.
- [49] Lam, D.D., Attard, C.A., Mercer, A.J., Myers, M.G., Rubinstein, M., Low, M.J., 2015. Conditional expression of Pomc in the Lepr-positive subpopulation of POMC neurons is sufficient for normal energy homeostasis and metabolism. *Endocrinology* 156(4):1292–1302. <https://doi.org/10.1210/en.2014-1373>.
- [50] Lam, D.D., de Souza, F.S.J., Nasif, S., Yamashita, M., López-Leal, R., Otero-Corchon, V., et al., 2015. Partially redundant enhancers cooperatively maintain mammalian Pomc expression above a critical functional

- threshold. *PLoS Genetics* 11(2):1–21. <https://doi.org/10.1371/journal.pgen.1004935>.
- [51] Zhan, C., Zhou, J., Feng, Q., Zhang, J. -e., Lin, S., Bao, J., et al., 2013. Acute and long-term suppression of feeding behavior by POMC neurons in the brainstem and hypothalamus, respectively. *Journal of Neuroscience*. <https://doi.org/10.1523/jneurosci.2742-12.2013>.
- [52] Dennison, C.S., King, C.M., Dicken, M.S., Hentges, S.T., 2016. Age-dependent changes in amino acid phenotype and the role of glutamate release from hypothalamic proopiomelanocortin neurons. *Journal of Comparative Neurology* 524(6):1222–1235. <https://doi.org/10.1002/cne.23900>.
- [53] Jones, G.L., Wittmann, G., Yokosawa, E.B., Yu, H., Mercer, A.J., Lechan, R.M., et al., 2019. Selective restoration of Pomc expression in glutamatergic POMC neurons: evidence for a dynamic hypothalamic neurotransmitter network. *Eneuro*. <https://doi.org/10.1523/eneuro.0400-18.2019>.
- [54] Chen, P., Williams, S.M., Grove, K.L., Smith, M.S., 2004. Melanocortin 4 receptor-mediated hyperphagia and activation of neuropeptide Y expression in the dorsomedial hypothalamus during lactation. *Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 24(22):5091–5100. <https://doi.org/10.1523/JNEUROSCI.0588-04.2004>.
- [55] Kim, M.S., Rossi, M., Abusnana, S., Sunter, D., Morgan, D.G.A., Small, C.J., et al., 2000. Hypothalamic localization of the feeding effect of agouti-related peptide and α -melanocyte-stimulating hormone. *Diabetes*. <https://doi.org/10.2337/diabetes.49.2.177>.
- [56] Baldini, G., Phelan, K.D., 2019. The melanocortin pathway and control of appetite-progress and therapeutic implications. *Journal of Endocrinology*. <https://doi.org/10.1530/JOE-18-0596>.
- [57] Paxinos, G., Franklin, K.B.J., 2001. *The mouse brain in stereotaxic coordinates*, 2nd ed.