

PARASITIC NEMATODES OF REPTILES (LIZARDS AND SNAKES) IN THE MONTE DESERT OF ARGENTINA

GABRIEL NATALIO CASTILLO^{1,2}, JUAN CARLOS ACOSTA^{1,2}
CYNTHIA JESSICA GONZÁLEZ-RIVAS¹, GERALDINE RAMALLO³

¹Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de San Juan
Av. Ignacio de la Roza 590, 5402, San Juan, Argentina

E-mails: nataliocastillo@gmail.com; <https://orcid.org/0000-0003-1519-9457>

jacostasantuan@gmail.com; <https://orcid.org/0000-0003-3539-8568>

cynthiajesica.gr@gmail.com

²Gabinete de Investigación DIBIOVA (Diversidad y Biología de Vertebrados del Árido)
Universidad Nacional de San Juan. Av. Ignacio de la Roza 590, 5402, San Juan, Argentina

³Instituto de Invertebrados. Fundación Miguel Lillo, San Miguel de Tucumán, Argentina

E-mail: gramallo@lillo.org.ar; <https://orcid.org/0000-0002-6953-4743>

Nematodes are little known in the Argentine herpetofauna. In order to increase and contribute to the knowledge of parasitism in reptiles, we studied nematodes found in three species of lizards (*Aurivela longicauda*, *Liolaemus darwini*, and *L. riojanus*) and one species of snake (*Philodryas trilineata*) from the Monte desert of center-west Argentina. We registered generalist nematodes commonly found in reptiles, belonging to three taxa: *Physaloptera* sp. (larvae), *Physaloptera retusa* (adults) (Physalopteridae) and *Parapharyngodon riojensis* (Pharyngodontidae) (adults). *Liolaemus darwini* had the lowest prevalence of *Physaloptera* sp. (larvae) (30%) and a mean intensity of 1.3 ± 0.4 (1–2). The lizard *A. longicauda* had the highest parasitic diversity (2 taxa) with prevalence (50%) and mean intensity (4 ± 3.5) of *Physaloptera retusa* (adults), also with prevalence (12.5%) and mean intensity (20 ± 0) of *Parapharyngodon riojensis* (adults). Due to the low number of studied specimens, precise conclusions cannot be drawn for *Liolaemus riojanus* ($n = 2$) and *P. trilineata* ($n = 1$). However, because the hosts were previously fixed, the results probably may do not represent real infection patterns. The four reptile species correspond to new host records from Argentina, and the information provided contributes to the knowledge of endoparasitism in reptiles of the Argentine Monte region.

Key words: parasitic nematodes, *Parapharyngodon riojensis*, *Physaloptera retusa*, *Liolaemus*, *Aurivela*, *Philodryas*, lizard, snake.

INTRODUCTION

Until 2004, information about endoparasite metazoans in Argentine lizards has been scarce (GOLDBERG *et al.* 2004). However, in the last years an increase in the knowledge about parasitism in reptile species has occurred. Such increases are mainly due to the contributions about nematodes in reptiles from Monte and Puna regions in Argentina (CASTILLO *et al.* 2019a, b).

Although the Monte region covers an extensive area of Argentina (KARLIN *et al.* 2017), to date, parasites were only studied in a reduced number of

reptile species; namely *Salvator rufescens*, *Teius teyou* (Teiidae), *Homonota underwoodi* (Phyllodactylidae) and *Liolaemus olongasta* (Liolaemidae) (CASTILLO *et al.* 2019a, b).

Information about parasite biodiversity is essential, since parasites may play an important role in the ecology of reptiles. It is well-known in the literature that nematode communities are influenced by attributes of the host, such as phylogenetic historical factors (BRITO *et al.* 2014), diet (RIBAS *et al.* 1995, O'GRADY & DEARING 2006, PEREIRA *et al.* 2012, 2014, BRITO *et al.* 2014), foraging strategy (active and passive search) (AHO 1990, BRITO *et al.* 2014), microhabitat use (BRITO *et al.* 2014); reproduction (BRITO *et al.* 2014, GALDINO *et al.* 2014), sexes (RIBAS *et al.* 1995, VRCIBRADIC *et al.* 1999, BRITO *et al.* 2014), age (PEREIRA *et al.* 2012), body size (VAN SLUY *et al.* 1994, RIBAS *et al.* 1995, VRCIBRADIC *et al.* 1999, ANJOS *et al.* 2005, 2012, PEREIRA *et al.* 2012) and environmental characteristics such as humidity or type of habitat (CASTILLO *et al.* 2018).

In order to contribute to the knowledge of the parasitic nematodes of reptiles in the Monte region of Argentina, we analyzed the gastrointestinal content of four species of reptiles (*A. longicauda*, *L. darwini*, *L. riojanus* and *P. trilineata*) captured in the previously mentioned area.

MATERIAL AND METHODS

The Monte phytogeographic region encompasses large arid areas, with an average annual precipitation lower than 100 mm/year, including absence of annual precipitation. Xerophyte plants adapted to dry and warm weather predominate. The sector is characterized by scrublands dominated by Zigophyllaceae, Malpigiaceae and Fabaceae (CABRERA 1976, MÁRQUEZ *et al.* 2016).

Sampling was carried out in different periods and sectors of the Monte desert of San Juan province, Argentina (Figs 1 & 2). During December 2017 and January 2018, pitfall traps and manual capture were used, and the following species were collected and examined: the lizards *Liolaemus darwini* (Liolaemidae), *Liolaemus riojanus* (Liolaemidae) (Liolaemidae) and *Aurivela longicauda* (Teiidae), and the snake *Philodryas trilineata* (Dipsadidae) (Table 1). Two lizards and the snake species (i.e., *L. darwini*, *A. longicauda* and *Philodryas trilineata*) have wide distribution in the Monte region and or other provinces from Argentina, and are not under any conservation threat (GIRAUDO *et al.* 2012). While *L. riojanus* is categorized as vulnerable and has more restricted distribution (ABDALA *et al.* 2012).

All host specimens were deposited in the Herpetological Collection, Biology Department, School of Exact, Physical, and Natural Sciences, National University of San Juan (*Philodryas trilineata*: UNSJ 4010; *L. riojanus*: UNSJ 4011, 4012; *A. longicauda*: UNSJ 4013–4032; *L. darwini*: 4033–4052).

Captured specimens were euthanized with an intraperitoneal administration of sodium thiopental, fixed in 10% formalin, and preserved in 70% ethanol. In the laboratory, the dissection and analysis of the digestive tract was carried out. The lizards were dissected through a ventral incision from mouth to anus, digestive tract was extracted and revised using a binocular stereomicroscope. Nematodes found were conserved in 70% ethanol.

Table 1. Reptiles studied from the Monte desert, central-western Argentina.

Host	Specimens	Locality	Geographical coordinates
Teiidae			
<i>Aurivela longicauda</i>	♂ = 4; ♀ = 4	Encón, Dept. 25 de Mayo	32.182839°S, 67.824371°W
Liolaemidae			
<i>Liolaemus darwini</i>	♂ = 16; ♀ = 24	Encón, Dept. 25 de Mayo	32.182839°S, 67.824371°W
<i>Liolaemus riojanus</i>	♀ = 2	Encón, Dept. 25 de Mayo	32.182839°S, 67.824371°W
Dipsadidae			
<i>Philodryas trilineata</i>	juvenile	Urban Sector, Dept. Rivadavia	31°31'00"S, 68°36'00"W

For identification of parasites, specimens were cleared in lactophenol, observed using an *Arcano* light microscope, following specific literature (ANDERSON *et al.* 2009, PEREIRA *et al.* 2012, 2014). Prevalence, mean intensity and mean abundance of parasites were according to BUSH *et al.* (1997). All nematodes were deposited in the Colección Parasitológica, Department of Biology, Fac. of Exact, Physical and Natural Sciences, National University of San Juan, Argentina, (*Physaloptera retusa* (Adults): UNSJPar260; *Physaloptera* sp. (Larvae): UNSJPar 261, 262, 263; *Parapharyngodon riojensis*: UNSJPar 264).




Fig. 1. Province of San Juan, Argentina, showing the location of the present study




Fig. 2. Environments where the samplings were performed. Capture locations of *L. darwini*, *L. riojanus* and *A. longicauda*. Town of the Encón, Department of 25 de Mayo (A, B, C)



Fig. 3. Studied hosts. A = *Philodryas trilineata*, B = *Aurivela longicauda* (photo: Ignacio Hernandez), C = *Liolaemus darwini* (photo: Claudio Mendez), D = *Liolaemus riojanus*

RESULTS

Three taxa of nematode parasites were identified: *Physaloptera* sp. (larvae), *Physaloptera retusa* (adults) (Physalopteridae) and *Parapharyngodon riojensis* (adults) (Pharyngodonidae). Eight specimens of *A. longicauda* (4 males, 4 females) were examined, and 36 nematodes corresponding to two taxa were isolated in their stomachs: *Physaloptera retusa* (14 females, 2 males) and *Parapharyngodon riojensis* (15 females, 5 males). Fourty specimens of *L. darwinii* were examined (24 females, 16 males), and 16 nematodes were registered, corresponding to larval stages of *Physaloptera* sp. Only two specimens of *L. riojanus* and one of *P. trilineata* were analyzed, finding larvae of *Physaloptera* sp. (Table 2).

DISCUSSION

In the present study, *Physaloptera* (larvae and adults) infecting *L. riojanus*, *L. darwinii*, *P. trilineata* and *A. longicauda* from the Monte desert, represent new host records.

Currently in Argentina, 14 reptile species infected by *Physaloptera* spp. have been reported: *L. quilmes*, *L. ornatus*, *L. alticolor*, *L. koslowskyi*, *L. neuquensis*, *L. olongasta*, *Salvator rufescens*, *A. tergo-laevigata*, *H. underwoodi*, *T. etheridgei*, *L. catamarcensis*, *L. belli*, *P. scapulatus* and *X. merremi* (RAMALLO & DÍAZ 1998, O'GRADY & DEARING 2006, GOLDBERG *et al.* 2004, CRUZ *et al.* 1998, LAMAS *et al.* 2016, CASTILLO *et al.* 2019a,b,c,d, GALLARDO *et al.* 2019). *Physaloptera* sp. parasitizing the present hosts raises to 18 the number of reptile species infected by this genera of nematode in Argentina.

Table 2. Quantitative descriptors of nematodes infection in a sector from the Monte region, central-western Argentina. P = Prevalence, MI = Mean intensity; MA = Mean abundance; N = examined host; np = total number of parasite. Mean values are followed by ± 1 standard deviation (range).

Host	N	Nematode	np	P %	MI	MA	Site of infection
<i>Aurivela longicauda</i>	4	<i>Physaloptera retusa</i> (adults)	16	50	4 \pm 3.5 (1–8)	2 \pm 3.16	Stomach
		<i>Parapharyngodon riojensis</i> (adult)	20	12.5	20 \pm 0	2.5 \pm 7.07	Stomach*
<i>Liolaemus darwinii</i>	40	<i>Physaloptera</i> sp. (larvae)	16	30	1.3 \pm 0.4 (1–2)	0.4 \pm 0.6	Stomach
<i>Liolaemus riojanus</i>	2	<i>Physaloptera</i> sp. (larvae)	1	50	1	0.5 \pm 0.7 (0–1)	Intestine
<i>Philodryas trilineata</i>	1	<i>Physaloptera</i> sp. (larvae)	1	100	1	1	Stomach

* These nematodes migrate to the stomach due to the fixation method. Probable location = large intestine

Physaloptera has heteroxenous life cycle and its intermediate hosts are mainly from the Orders Orthoptera and Coleoptera (ANDERSON *et al.* 2000). This genus currently includes 105 species which are parasites in reptiles, amphibians, birds and mammals (PEREIRA *et al.* 2012, 2014). In the Neotropics, eight species that parasitize the stomach of reptiles are known: *P. bonnei* Ortlepp, 1922, *P. liophis* Vicente et Santos, 1974, *P. lutzii* Cristofaro, Guimaraes et Rodrigues, 1976, *P. monodens* Molin, 1860, *P. obtusissima* Molin, 1860, *P. retusa* Rudolphi, 1819, *P. tupinambae* PEREIRA, ALVES, ROCHA, LIMA et LUQUE, 2012 and *P. bainaie* PEREIRA, ALVES, ROCHA, LIMA et LUQUE, 2014 (CASTILLO *et al.* 2019c).

Reptiles have been mentioned as paratenic hosts (WIDMER 1970). *Liolaemus darwini* would have a role as a paratenic host in the life cycle of *Physaloptera* sp. A possible definitive host, *Salvator rufescens*, in which adult stages of *Physaloptera* were recorded (CASTILLO *et al.* 2019b). This was captured in the same place as *L. darwini*. *Aurivela longicauda* would have a definitive host role, probably by feeding on other lizards. We have records that *A. longicauda* has a tendency to feed on other lizards (BLANCO *et al.* 2012).

Parapharyngodon riojensis as a parasite of *A. longicauda* from the Monte desert, also represents a new host record. It is a generalist nematode, and in Argentina it parasitizes *Liolaemus* spp., *Phymaturus* spp. and *Tropidurus* spp. (RAMALLO *et al.* 2002, GOLDBERG *et al.* 2004, CASTILLO *et al.* 2017, RAMALLO *et al.* 2017, CASTILLO *et al.* 2018, CASTILLO *et al.* 2019d). *Parapharyngodon riojensis* is very common in reptiles of the San Juan province (CASTILLO & ACOSTA 2019, CASTILLO *et al.* 2019d).

According to our bibliographic review, in Argentina there are currently 15 lizard species parasitized by *Parapharyngodon* spp. (CRUZ *et al.* 1998, RAMALLO *et al.* 2002b, GOLDBERG *et al.* 2004, LAMAS & ZARACHO 2006, RAMALLO *et al.* 2016, RAMALLO *et al.* 2017, CASTILLO *et al.* 2017, CASTILLO *et al.* 2018, CASTILLO *et al.* 2019e, CASTILLO *et al.* 2019c, CASTILLO & ACOSTA 2019).

The probability that an uninfected host will become infected would depend on the frequency of movement (GETZ & PICKERING 1993). *Aurivela longicauda* is a more agile, faster and more moving species than *L. darwini*. Infection is more likely in *A. longicauda*. In addition, the relationship between lizard SVL (snout vent length) and the intensity of nematode infection suggest that body size is an important determinant of the infection rate (RIBAS *et al.* 1995, ANJOS *et al.* 2012). The larger digestive tract of the larger lizards (*A. longicauda*), provides more microhabitats suitable for nematode settlement compared to the relatively smaller one of *L. darwini*.

Considering the large number of lizard species registered in Argentina, current knowledge on their parasitic nematodes is incomplete and fragmented. The present report expands the number of host species and the geographi-

cal distribution of *P. riojensis* and *Physaloptera* spp. representing a valuable contribution to parasitological knowledge and adding to knowledge of this species in the Monte desert of Argentina.

We emphasize that the results obtained with respect to the parasites of *A. longicauda* are similar to those of other phylogenetically related species. *Parapharyngodon* spp. and *Physaloptera* spp. was reported in lizards of the family Teiidae (ÁVILA & SILVA 2010, GALLARDO *et al.* 2019, VIEIRA *et al.* 2019).

*

Acknowledgement – We thank the Sub-Secretary of the Environment for the permits granted (N° 1300-3097-16), and the rangers (Mariano Hidalgo, Jorge Cayuela, Jesús Quiroga and José Castro) for their help in field samplings. Sofia Nanni assisted us in drafting the English version. We thank two anonymous reviewers for improving the manuscript.

The authors declare no competing interests.

REFERENCES

- ABDALA, C. S., ACOSTA, J. C., ACOSTA, J. L., ÁLVAREZ, B. B., ARIAS, F., AVILA, L., BLANCO, M. G., BONINO, M. J., BORETTO, M., BRANCATELLI, G., BREITMAN, M. F., CABRERA, M., CAIRO, R. S., CORBALÁN, V., HERNANDO, A., IBARGUENGOYÍA, N. R., KACOLIRIS, F., LASPIUR, A., MONTERO, R., MORANDO, M., PELEGRÍN, N., PÉREZ, C. H. F., QUINTEROS, A., SEMHAN, S. R. V., TEDESCO, M. E., VEGA, L. & ZALBA, S. M. (2012): Categorización del estado de conservación de las lagartijas y anfisbenas de la República Argentina. – *Cuadernos de Herpetología* **26**: 215–248.
- AHO, J. M. (1990): Helminth communities of amphibian and reptiles: comparative approaches to understanding patterns and processes. Pp. 157–195. In: ESCH, G. W., BUSH, A. O. & AHO, J. M. (eds): *Parasite communities: Patterns and processes*. – Chapman and Hall, NY. https://doi.org/10.1007/978-94-009-0837-6_7
- ANDERSON, R. C. (2000): *Nematodes parasites of vertebrates: Their development and transmission*. 2nd ed. – CABI Publishing, Wallingford, UK, 650 pp.
- ANJOS, L. A., ROCHA, C. F. D., VRCIBRADIC, D. & VICENTE, J. J. (2005): Helminths of the exotic lizard *Hemidactylus mabouia* from a rock outcrop area in southeastern Brazil. – *Journal of Helminthology* **79**(4): 307–313. <https://doi.org/10.1079/JOH2005288>
- ANJOS, L. A., ÁVILA, R. W., RIBEIRO, S. C., ALMEIDA, W. O. & DA SILVA, R. J. (2012): Gastrointestinal nematodes of the lizard *Tropidurus hispidus* (Squamata: Tropiduridae) from a semi-arid region of north-eastern Brazil. – *Journal of Helminthology* **87**(4): 443–449. <https://doi.org/10.1017/S0022149X12000491>
- ÁVILA, R. W. & SILVA, R. J. (2010): Checklist of helminths from lizards and amphisbaenians (Reptilia, Squamata) of South America. – *Journal of Venomous Animals and Toxins including Tropical Diseases* **16**: 543–572. <https://doi.org/10.1590/S1678-91992010000400005>
- BLANCO, M. G., ACOSTA, J. C., PIAGGIO, L., NIEVA, R., VICTORICA, A. & CASTILLO, G. (2012): Saurofagia y canibalismo en dos especies de lagartos del centro-oeste de Argentina. – *Cuadernos de herpetología* **26**: 91–93.
- BRITO, S. V., CORSO, G., ALMEIDA, A. M., FERREIRA, F. S., ALMEIDA, W. O., ANJOS, L. A., MESQUITA, D. O. & VASCONCELLOS, A. (2014): Phylogeny and micro-habitats utilized by

- lizards determine the composition of their endoparasites in these miarid Caatinga of Northeast Brazil. – *Parasitology Research* **113**: 3963–3972. <https://doi.org/10.1007/s00436-014-4061-z>
- BUSH, A. O., LAFFERTY, K. D., LOTZ, J. M. & SHOSTAK, A. W. (1997): Parasitology meets ecology on its own terms. – *Journal of Parasitology* **83**: 575–583. <https://doi.org/10.2307/3284227>
- CABRERA, A. L. (1976): *Enciclopedia Argentina de agricultura y jardinería: regiones fitogeográficas Argentinas*. – Acme, Buenos Aires, 85 pp.
- CASTILLO, G. N., RAMALLO, G. & ACOSTA, J. C. (2017): *Liolaemus ruibali*. Endoparasites. – *Herpetological Review* **48**: 651–652.
- CASTILLO, G., ACOSTA, J. C., RAMALLO, G. & PIZARRO, J. (2018): Pattern of infection with *Parapharyngodon riojensis* Ramallo, Bursey, Goldberg 2002 (Nematoda: Pharyngodonidae) in the lizard *Phymaturus extrilidus* from Puna region, Argentina. – *Annals of Parasitology* **64**(2): 83–88.
- CASTILLO, G. N. & ACOSTA, J. C. (2019): Parasitism in two species of lizards of the genus *Liolaemus* (Wiegmann, 1834) from the puna Argentina. – *Neotropical Helminthology* **13**: 89–95.
- CASTILLO, G. N., GONZÁLEZ- RIVAS, C. J. & ACOSTA, J. C. (2019a): *Liolaemus olongasta* (Chelco Lizard). Endoparasites. – *Herpetological Review* **50**: 578–579.
- CASTILLO, G. N., GONZÁLEZ- RIVAS, C. J. & ACOSTA, J. C. (2019b): Nematode parasites in the lizards *Salvator rufescens*, *Teiusteyou* (Teiidae) and *Homonota underwoodi* (Phyllodactylidae) from the Monte Region in Central-Western Argentina. – *North-Western Journal of Zoology* **15**: 192–195.
- CASTILLO, G. N., ACOSTA, J. C. & ACOSTA, R. (2019c): *Liolaemus fitzgeraldi*. Endoparasites. – *Herpetological Review* **50**: 578–579.
- CASTILLO G. N., ACOSTA, J. C. & BLANCO, G. M. (2019d): Trophic analysis and parasitological aspects of *Liolaemus parvus* (Iguania: Liolaemidae) in the Central Andes of Argentina. – *Turkish Journal of Zoology* **43**: 277–286. <https://doi.org/10.3906/zoo-1812-33>
- CRUZ, F. B., SILVA, S. & SCROCCHI, G. J. (1998): Ecology of the lizard *Tropidurus etheridgei* (Squamata: Tropiduridae) from the dry Chaco of Salta, Argentina. – *Herpetological Natural History* **6**: 23–31. <https://doi.org/10.2307/1565679>
- GALLARDO, G. A., TULLI, M. J. & SCROCCHI, G. J. (2019): Dimorfismo sexual y ecología trófica de *Aurivelatergo laevigata* (Squamata, Teiidae). – *Revista del Museo Argentino de Ciencias Naturales nueva serie* **21**(1): 45–50. <https://doi.org/10.22179/REVMACN.21.606>
- GETZ, W. M. & PICKERING, J. (1983): Epidemic models thresholds and population regulation. – *American Naturalist* **121**: 892–898. <https://doi.org/10.1086/284112>
- GIRAUDO, A. R., ARZAMENDIA, V., BELLINI, G., BESSA, C. A., CALAMANTE, C. C., CARDOZO, G., CHIARAVIGLIO, M., COSTANZO, M. B., ETCHEPARE, E. G., COLA, V. & DI PIETRO, D. O. (2012): Categorización del estado de conservación de las Serpientes de la República Argentina. – *Cuadernos de herpetología* **26**: 303–374.
- GOLDBERG, S. R., BURSEY, C. R. & MORANDO, M. (2004): Metazoan endoparasites of 12 species of lizards from Argentina. – *Comparative Parasitology* **71**: 208–214. <https://doi.org/10.1654/4089>
- KARLIN, U. O., KARLIN, M. S., ZAPATA, R. M., COIRINI, R. O., CONTRERAS, A. M. & CARNERO, M. (2017): La provincia fitogeográfica del Monte: límites territoriales y su representación. – *Multequina* **26**: 63–75.
- LAMAS, M. & ZARACHO, V. (2006): *Tropidurus torquatus* (Brown Lizard). Endoparasites. – *Herpetological Review* **37**: 474–475.

- LAMAS, M. F., CÉSPEDEZ, J. A., RUIZ-GARCÍA, J. A. (2016): Primer registro de nematodos parásitos para la culebra *Xenodon merremi* (Squamata, Dipsadidae) en Argentina. – *Facena* **32**: 59–67.
- MÁRQUEZ, J., MARTÍNEZ CARRETERO, E. & DALMASSO, A. (2016): Provincias fitogeográficas de la Provincia de San Juan. Pp. 187–197. In: MARTÍNEZ CARRETERO, E. & GARCÍA A. (eds): San Juan Ambiental. 1^o edición. – Editorial INCA, Mendoza.
- O'GRADY, S. P. & DEARING, M. D. (2006): Isotopic insight into host–endosymbiont relationships in Liolaemidae lizards. – *Oecologia* **150**: 355–361. <https://doi.org/10.1007/s00442-006-0487-z>
- PEREIRA, F. B., SOUSA, B. M. & DE SOUZA LIMA, S. (2012): Helminth community structure of *Tropidurus torquatus* (Squamata: Tropiduridae) in a rocky outcrop area of Minas Gerais state, southeastern Brazil. – *Journal of Parasitology* **98**: 6–10. <https://doi.org/10.1645/GE-2689.1>
- PEREIRA, F. B., ALVES, P. V., ROCHA, B. M., DE SOUZA LIMA, S. & LUQUE, J. L. (2014): *Physaloptera baina* n. sp. (Nematoda: Physalopteridae) parasitic in *Salvator merianae* (Squamata: Teiidae), with a key to Physaloptera species parasitizing reptiles from Brazil. – *Journal of Parasitology* **100**: 221–228. <https://doi.org/10.1645/13-281.1>
- RAMALLO, G. R. & DÍAZ, F. (1998): *Physaloptera lutzi* (Nematoda, Physalopteridae) parasite de *Liolaemus* (Iguania, Tropiduridae) del noroeste Argentino. – *Boletín Chileno de Parasitología* **53**: 19–22.
- RAMALLO, G., BURSEY, C., CASTILLO, G. & ACOSTA, J. C. (2016): New species of Parapharyngodon (Nematoda: Pharyngodonidae) in *Phymaturus* spp. (Iguania: Liolaemidae) from Argentina. – *Acta Parasitologica* **61**: 461–465. <https://doi.org/10.1515/ap-2016-0062>
- RAMALLO, G., BURSEY, C. H., GOLDBERG, S., CASTILLO, G. & ACOSTA, J. C. (2017): *Phymaturus extrilidus* (Argentine Lizard). Endoparasites. – *Herpetological Review* **48**: 198.
- RAMALLO, G., BURSEY, C. R. & GOLDBERG, S. R. (2002): Parapharyngodon *riojensis* n. sp. (Nematoda: Pharyngodonidae) from the lizard *Phymaturus punae* (Squamata: Iguania: Liolaemidae) from northwestern Argentina. – *Journal of Parasitology* **88**: 979–982. <https://doi.org/10.2307/3285541>
- RIBAS, S. C., ROCHA, C. F. D., TEIXEIRA-FILHO, P. F. & VICENTE, J. J. (1995): Helminths (Nematoda) of the lizard *Cnemidophorus ocellifer* (Sauria: Teiidae): Assessing the effect of rainfall, body size and sex in the nematode infection rates. – *Ciencia e Cultura* **47**: 88–91.
- VAN SLUYS, M., ROCHA, C. F. D. & RIBAS, S. C. (1994): Nematodes infecting the lizard *Tropidurus itambere* in southeastern Brazil. – *Amphibia-Reptilia* **15**(4): 405–408. <https://doi.org/10.1163/156853894X00443>
- VIEIRA, F. M., DE SOUZA, T. T., NOVELLI, I. A., LIMA, S. S., MUNIZ-PEREIRA, L. C. & DE SOUSA, B. M. (2019): Nematode parasites of lizards (Squamata, Sauria) from the Cerrado biome in the State of Minas Gerais, Brazil. – *Herpetology Notes* **12**: 855–863.
- VRCIBRADIC, D., ROCHA, C. F. D., RIBAS, S. C. & VICENTE, J. J. (1999): Nematodes infecting the skink *Mabuya frenata* in Valinhos, São Paulo state, southeastern Brazil. – *Amphibia-Reptilia* **20**: 333–339. <https://doi.org/10.1163/156853899X00367>
- WIDMER, E. A. (1970): Development of third-stage Physaloptera larvae from *Crotalus viridis* Rafinesque, 1818 in cats with notes on pathology of the larvae in the reptile (Nematoda, Spiruroidea). – *Journal of Wildlife Diseases* **6**: 89–93. <https://doi.org/10.7589/0090-3558-6.2.89>

Received April 7, 2020, accepted August 7, 2020, published November 13, 2020

