

CONTRIBUTION OF CARBONATE PRECIPITATION TO THE PRESERVATION OF RIPPLES IN SILICICLASTIC SEDIMENTS COLONIZED BY MICROORGANISMS

Lucía Maisano^{1,2,*}, I. Emma Quijada³, Vanesa L. Perillo^{1,4}, Diana G. Cuadrado^{1,2},
L. Ariel Raniolo^{1,5}, Eduardo A. Gómez^{1,6}

¹ Instituto Argentino de Oceanografía (IADO), CONICET, Florida 5000, 8000 Bahía Blanca, Buenos Aires, Argentina.

² Departamento de Geología, Universidad Nacional del Sur, Av. Alem 1253, cuerpo B´ - 2º Piso, 8000 Bahía Blanca, Buenos Aires, Argentina.

³ Departamento de Geología, Universidad de Oviedo, C/ Jesús Arias de Velasco s/n, 33005 Oviedo, Spain.

⁴ Departamento de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur, San Juan 670, 8000 Bahía Blanca, Buenos Aires, Argentina.

⁵ Departamento de Ingeniería, Universidad Nacional del Sur Av. Alem 1253 .8000 Bahía Blanca, Buenos Aires, Argentina).

⁶ Universidad Tecnológica Nacional. Facultad Regional Bahía Blanca, 11 de Abril 378, 8000 Bahía Blanca, Buenos Aires, Argentina.

* Corresponding author: Lucia Maisano. Email: lmaisano@iado-conicet.gob.ar

ARTICLE INFO

Article history

Received July 16, 2021

Accepted December 2, 2021

Available online December 15, 2021

Handling Editor

Fernando Gómez

Keywords

Sedimentary structures

Ripples structures

Carbonate precipitation

Microbial mats-

Paso Seco, Argentina

ABSTRACT

The present study documents the biostabilization and early lithification of ripple marks created during high-energy hydrodynamic events in a coastal environment colonized by microbial mats. The presence of microorganisms and their characteristic secretion of exopolymeric substances (EPS) are biological factors that diminish the sediment permeability, a first step to maintain seawater over the sediment. Subsequently, seawater is evaporated causing an increase in salinity that leads to the precipitation of calcite and gypsum. The calcite precipitation covering the microbial mat and ripple marks is documented in detail. The study was conducted in Paso Seco (40°38'40"S; 62°12'22"W), a modern coastal flat within an elongated semi-closed basin colonized by microbial mats frequently inundated. Salinity was measured in different ponds in the flat with different degrees of evaporation, and in a tidal creek. Water samples were analysed to determine Ca²⁺, CO₃²⁻ and SO₄²⁻ concentrations. Seawater level fluctuations relative to the tidal flat were measured through a 10-month period in 2018 using a HOBO water level logger. Ripple field formation over the tidal flat was documented after the occurrence of a severe storm and was monitored throughout three field trips. Six months after the formation of the ripples, a sedimentary block containing their vertical section was taken to perform petrographic studies. A laterally continuous dense micritic calcite layer, 100–200 μm in thickness, was identified covering ripple structures, improving the immobilization of these bedforms, and promoting their early lithification. Our observations of modern sedimentary structures and the *in situ* study of their evolution and early lithification may be potentially useful to be taken into account for the interpretation of similar sedimentary structures in the geological record.

INTRODUCTION

Sedimentary dynamics have been shown to be biologically modified in environments where sediments are colonized by benthic microorganisms (Noffke and Paterson, 2008). However, one of the greatest challenges in the study of the geological record is to recognize biogenic influence in the sedimentary structures of ancient sediments. The study of modern microbial mats is especially useful for this purpose because it allows a precise analysis of present-day biogenic structures and the environmental conditions in which they are formed and can, then, be used as analogues for the fossil record (Cuadrado and Blasi, 2018; Cuadrado, 2020). On some occasions the general hydrodynamic regime in the modern sedimentary environment colonized by microbial mats may show differences between winter and summer due to water level fluctuations (Cuadrado, 2020). Such seasonal fluctuations can be recorded in the sediment through the precipitation of certain minerals. According to this, one of the most interesting aspects of microbial influence on sediments is their impact on carbonate precipitation, which has been proven to facilitate microbial mat preservation and lithification, especially in carbonate environments (Reid *et al.*, 2000, Dupraz *et al.*, 2004), enabling the identification of microbial mats in the fossil record. However, for precipitation to occur, it is essential to develop a delicate balance between physicochemical processes and microbial activities (Dupraz and Visser, 2005). Therefore, the study of mineral precipitation in these systems becomes a complex endeavour as it involves the interaction between biotic and abiotic factors (Maisano *et al.*, 2020).

Mineral precipitation in microbial ecosystems has been studied in detail in modern and past carbonate environments, where rapid mineralization favours preservation of microbialites (Reid *et al.*, 2000, Dupraz *et al.*, 2004). In contrast, in modern siliciclastic environments, microbial mats are generally unlithified and form planar soft organic layers that colonize depositional areas that are subject to erosive episodes by high-energy currents and to sediment deposition (Noffke, 2003). Nevertheless, the present investigation documents evidence of calcium carbonate precipitation in a siliciclastic coastal environment that enables the lithification of ripple marks and microbial mats.

Sedimentary structures such as ripple marks, formed by severe hydrodynamic events, are biostabilized by microbial activity and the presence of exopolymeric substances (EPS; Cuadrado, 2020), which may produce sinoidal structures. In addition, the presence of a carbonate lamination over these sand sedimentary structures possibly would enhance the lithification of the colonized ripple structure. The analysis of this carbonate precipitation is the focus of this paper, which coupled with the sinoidal structure, may provide potential for preservation in the fossil record. Paso Seco coastal area is an excellent case study to present the formation of carbonate lamination in microbial mats, which is a consequence of a combination of biogeochemical and physicochemical factors (Perillo *et al.*, 2019, Maisano *et al.*, 2020).

STUDY AREA

The Paso Seco coastal environment (40°38'40"S; 62°12'22"W; Fig. 1) is in the northern coast of Patagonia (Argentina), which is characterized by a mesotidal regime with mean and maximum tidal ranges of 1.62 m and 2.5 m, respectively. The climate of the region is semiarid, as stated by the recent climatic classification of the Argentinian Pampas proposed by Aliaga *et al.* (2017), characterized by a potential evapotranspiration greater than precipitation, which generates enough aridity to limit vegetation development. The elongated study area (3.5 km in length and 0.4 km in width) corresponds to an ancient tidal channel closed more than 100 years ago by the generation of a sand spit (1.8 km wide) due to longshore sediment transport. Thus, the study area is a coastal environment behind the sand spit, a semi-closed basin categorized as a supratidal zone because it is flooded by seawater under specific oceanographic conditions when storm surges provoke seawater overpass the sand spit (Stempels Bautista, 2019). The flooding might occur up to 43 times per year, but the area remains unaffected by daily flood tides (Perillo *et al.*, 2019; Maisano *et al.*, 2019). The high-energy flooding is followed by a slow receding current and the retention of a few cm-deep seawater layer over the sedimentary surface for several days to weeks. Furthermore, the extreme-energy water currents produce microbially-induced sedimentary structures of significant dimensions (e.g., large mat tears, between 2 and 8 m; large roll-



Figure 1. a) Location of the Paso Seco flat behind a sand spit (yellow square). b) Close-up view of the study area (yellow square in A). The blue mark indicates the location of the water level station. The yellow mark indicates the location of the tidal creek where a water sample was taken. c) Large microbially-induced sedimentary structures created by deformation processes over the tidal flat during extreme storm that occurred in June 2018, as seen in August 2018, 20 days after a moderate storm. I) microbial roll-ups, II) flipped-over edges.

ups \approx 25 cm in height; Fig. 1c) (Maisano *et al.*, 2019).

The area is characterized by \approx 1 cm-thick epibenthic mats that colonize the sediments (Cuadrado *et al.*, 2015). The main constructor member of this microbial consortium is the filamentous cyanobacteria in the order *Oscillatoria sp.* where the *Coleofasciculus (Microcoleus) chthonoplastes* is the most abundant cosmopolitan cyanobacteria species in the area (Cuadrado and Pan, 2018). Other important microorganisms of the microbial association are pennate diatoms that are the first colonizers (Pan *et al.*, 2017). Once the microbial mat is formed, the consortium of microorganisms, EPS and sediment produces a poorly permeable sediment surface, which causes slow water infiltration from the surface and allows gradual evaporation of seawater and evaporite precipitation (Perillo *et al.*, 2019). Thus, Paso Seco can be considered a saline basin conditioned by the hydrological conditions and biological processes (Perillo *et al.*, 2019).

METHODS

To perform hydrochemical analyses, six water samples were taken from the study area in September 2017: five samples from different ponds with different salinities and one sample from a tidal creek (Fig. 1b). Water Ca^{2+} , CO_3^{2-} and SO_4^{2-} concentrations were determined by titration (Ca^{2+} , CO_3^{2-}) and turbidimetric (SO_4^{2-}) methods following standard methods (APHA, 2005), whereas water salinity was determined gravimetrically as described by Strickland and Parsons (1972).

To know the frequency of seawater entering the study zone, water-level fluctuations were measured in the tidal flat using a HOBO water level logger (Onset-model U20; 2.5 cm diameter, 15 cm length). It was deployed into a vertically buried, perforated PVC pipe, 40 cm in depth from the sedimentary surface. The sensor recorded water level and temperature every 10 min. The water level data were corrected by atmospheric pressure by means of another logger placed in an upper level close to the tidal flat. The present study shows the water level and temperature recorded over a 10-month period in 2018.

Ripple field formation over the tidal flat was documented after the occurrence of a severe storm and was monitored throughout three field trips over a period of nearly six months since its creation, until December 2018. Firstly, ripples were examined

two days after their creation, then in a following field trip carried out 20 days after a moderate storm that caused deposition of a thin sand layer over the previous ripples, and finally, six months afterwards. During the third field trip (December 2018) a sedimentary block (12 x 12 x 5 cm) which included the monitored ripples was taken for petrographic analysis. In addition, sedimentary cores were obtained using PVC tubes (3 cm diameter and 7 cm length) to examine the sedimentary texture and microstructures. They were opened in the laboratory, and subsequently analysed under a SMZ Nikon 1500 stereoscopic zoom binocular microscope. In addition, thin sections (thicker than the 30- μm traditional thickness of petrographic thin sections to avoid losing fragments of sample) obtained from the sedimentary block were prepared for petrographic analysis. Thin sections were analysed by a Nikon Eclipse POL 50i transmitted-light microscope, coupled with a camera. The analysis was carried out under plane-polarized light and cross-polarized light to identify mineral species. Three sections were stained with a solution of Potassium Ferrocyanide Ke (CN) 6, Alizarin and Hydrochloric Acid for determining the nature of the carbonate (Dickson, 1966).

RESULTS

Hydrodynamic processes and water evaporation

The topography of the study area presents a 0.03% general gradient towards inland. The maximum elevation (1.8 m above mean sea level) is in the northeasternmost area of the basin (Fig. 2a). On the other hand, the minimum elevation (0.5 m above mean sea level) corresponds to a tidal creek located in the inland area of Paso Seco (Fig. 2a), which is connected to a major tidal channel at the west of the study area (see Fig. 1).

The water level records show the storm-related inundations of the tidal flat (Fig. 2b). From March to June 2018, the sediment flat remained exposed for most of the time. On 25 June 2018, an extreme storm produced a seawater inundation of 70 cm over the microbial flat (green arrow, Fig. 2b) and the formation of a ripple field. The first field trip was done two days after the inundation (red arrow 1, Fig. 2b). Subsequently, seawater remained near the sedimentary surface until a moderate inundation





Figure 2. a) Topographic map of the study area. b) Water level record from March to December 2018. Green arrow indicates a high-energy flooding, during which the ripple field was created. Pink arrow indicates a subsequent moderate inundation. Black dashed line indicates a period during which the flat was episodically inundated. Green dashed line indicates a period during which the flat was subaerially exposed. Orange dashed line indicates a period of several floodings lower than 25 cm interspersed with subaerial exposure periods. Red dashed arrows indicate the date of the field trips. During field trip 3 a sedimentary block was sampled.

on 22 July occurred (pink arrow, Fig. 2b). In this occasion, the water depth was 25 cm. Seawater remained near the microbial surface for 19 days including two subsequent inundations (black dashed line, Fig. 2b). A second field trip was conducted 20

days after the moderate inundation (red arrow 2, Fig. 2b). Then, the water level descended progressively below the tidal flat, causing subaerial exposure of the flat for 29 days (green dashed line, Fig. 2b). Later, during spring, the flat experienced eight floodings




Figure 3. a) Evaporites in the field site. Black arrows show evaporation rings around a shallow pond in the coastal flat. **b)** Comparison between Ca^{2+} , CO_3^{2-} and SO_4^{2-} concentrations from water samples obtained in different ponds and a tidal creek with different water salinities.

less than 20 cm water column, which alternated with flat exposures (orange dashed line, Fig. 2b), until the last field trip was conducted in December (red arrow 3, Fig. 2b).

The topography of Paso Seco is characterized by several shallow ponds (≈ 4 cm in depth) containing seawater after inundations, whose volume progressively diminishes as seawater evaporates. Hydrochemical analysis were carried out in September 2017 to determine the evolution of water composition as it evaporates after an inundation. Salinity increased in the different tidal ponds in agreement to the visual signs of volume water reduction and evaporite presence (e.g., visual gypsum and halite precipitation; Fig. 3a). The chemical analysis of the water composition in these ponds indicated that CO_3^{2-} concentration increased until salinity reached the value of 100, when an abrupt decrease in CO_3^{2-} concentration occurred (Fig. 3b). In contrast, Ca^{2+} concentration showed an increase as salinity reached a value close to 202, when the maximum Ca^{2+} concentration was attained, and subsequently dropped rapidly. Moreover, the SO_4^{2-} concentration raised gradually as salinity increased (Fig. 3b).

Ripple field characteristics

The extreme storm that occurred in June 2018 (green arrow in Fig. 2b) produced the modification of the microbial flat combined with the formation of sand ripple patches due to the sediment transport over the microbial mat. Ripples, made up of fine sand

sediment, were characterized by asymmetrical 2D and 3D morphologies with straight and sometimes sinuous crestlines (Fig. 4a). Wavelengths ranged from 5 to 10 cm and heights reached 1 cm, resting on a sharp and planar microbial mat (green dashed line, Fig. 4b). The visualization of ripples immediately after the storm (1 shown in Fig. 2b) showed a sand layer on the sediment surface with mud present only in the trough (Fig. 4a).

In July 2018 a moderate storm flooded Paso Seco (pink arrow in Fig. 2b). 20 days after this storm the study area was visited during a field trip (2 shown in Fig. 2b), allowing to observe that a thin biofilm colonized the whole ripple field (Fig. 4c). Moreover, cross-sections of the sediment permitted the recognition of the surficial biofilm (orange arrow in Fig. 4d, e), and below the biofilm the mud layer covering the ripple trough is identified (red dashed line in Fig. 4e).

Petrographic analysis

A sedimentary block containing the sand ripples deposited over a 1 mm-thick microbial mat (Fig. 5a) was petrographically analysed. The microbial mat (c frame in Fig. 5b) is composed of brownish organic matter laminae of $\approx 20 \mu\text{m}$ thickness (yellow arrows in Fig. 5c), interbedded with clay to fine-grained silt and micritic calcite laminae between 40 and $80 \mu\text{m}$ in thickness (orange arrows in Fig. 5c). The micritic layers sometimes are composed of peloids less than $100 \mu\text{m}$ in size (blue arrows in Fig. 5c-d), which may coalesce forming laterally continuous layers (pink




Figure 4. a) Asymmetrical 2D ripples observed in Paso Seco flat two days after the storm of June 25, 2018. Red arrows indicate fine sediment deposited in the troughs of the ripple. b) A vertical section of a ripple structure obtained two days after the storm of June 25 2018. The microbial mat is delineated by a green dashed line. c) Tidal flat appearance on August 2018, 20 days after the moderate storm in July 2018. Sediment surface shows a smooth appearance caused by the colonization of the rippled surface by a biofilm. d) A vertical section of a ripple colonized by a biofilm (orange arrow). e) Close view of the red square in (d). Red dashed line: clay and silt sediment layer in the ripple trough. Orange arrow: biofilm on the sediment surface.


arrows in Fig. 5c-d).

The sand ripple deposited over the microbial mat is 6 mm in height and its trough is covered by a layer made up of fine silt and clay (limited by red dashed line in Fig. 5b). This fine-grained layer presents variable thickness, being thicker (≈ 1 mm) in the trough, and thinner (few tens of micrometres) towards the crest of the ripple. This layer presents a brownish interference colour under cross-polarized light (red dashed line in Fig. 5b, e-g) which is covered by a thin biofilm along the upper red dashed line (Fig. 5b, e-g). Upwards, another tiny train of sand ripples (up to ≈ 1 mm in height) covers the previous fine sediment layer and the thin biofilm (green arrows in Fig. 5b- e, f). This second train of ripples had not been recognized during the field trips due to its small size, and its identification was only possible under

the microscope. Covering the previous laminae, a laterally continuous dense micritic calcite layer, 100-to-200 μm in thickness, is developed (m in Fig. 5e- f). The high-order interference colours of calcite under cross-polarized light allow to easily discern this lamina (Fig. 5h- i). The micritic layer is covered by an organic matter lamina forming a biofilm on the sediment surface (b in Fig. 5e- i).

DISCUSSION

The episodic storms control the water exchange between the study zone and the sea, while the geomorphology affects the sediment dynamics. Seawater enters the narrow mouth of the area forming fast currents that are essential for sand transport (Fig. 1b, 2a). During these severe storms,



water depths over 70 cm are quickly reached over the microbial mats (Maisano *et al.*, 2019). These quick currents produce large microbially-induced sedimentary structures created by deformation processes over the tidal flat, such as microbial roll-ups, flipped over structures, microbial folds, and erosional pockets (Fig. 1c). Moreover, the sediment transport during flooding forms ripple patches over the layered planar epibenthic microbial mat, and the presence of the microorganisms influences the stabilization of the ripples (Cuadrado, 2020). In that sense, once the ripples are created, when calm conditions are recovered, the filamentous microorganisms present in the underlying mat, such as *O. Oscillatoriales* (Cuadrado and Pan, 2018), can move quickly upward to the sand ripple due to their gliding motility (Shepard and Sumner, 2010). Consequently, the grains are entangled by microbial filaments and embedded by EPS, which causes the increase of the critical erosion threshold for the movement of particles (Paterson, 1994; Noffke *et al.*, 2001).

The analysis of the sedimentary block obtained from the flat after the formation and colonization of the ripples (Fig. 5a), coupled with the analysis of the water level record (Fig. 2b), allows the recognition of the successive hydrodynamic events that took place in Paso Seco along 2018 and the identification of all the processes involved in ripple preservation. The strong storm of June 2018 (green arrow, Fig. 2b) created the ripple resting on the microbial mat (yellow arrow, Fig. 5a-b). After that strong event and during the subsequent relative quiet conditions, when there was no sediment reworking, a clay and silt sediment layer of varying thickness was

deposited (red dashed line in Fig. 5b, e). Moreover, during this period, a reorganization of the microbial mat fabrics occurred, and binding and trapping processes took place. These processes would lead to the formation of a thin biofilm covering the sediment surface forming a sinoidal structure (see Noffke, 2010) (located along the upper red dashed line in Fig. 5b, e-g). The subsequent moderate storm in July (pink arrow in Fig. 2b) produced another ripple over that biofilm (green arrow in Fig. 5b), and finally the ripple was capped again by a thin biofilm formed during following subaerial exposure and several inundation events (green and orange dashed line in Fig. 2b). Although the genesis of the ripples was triggered by mechanical processes (thus abiogenic), that is, transport of non-cohesive detrital sediments (Kennedy, 1969), the presence of the filamentous microorganisms in the microbial mats over which the sand was deposited, produced a fast stabilization (Cuadrado, 2020).

The biofilm formed over the first ripple structure is not associated with a micritic lamina, probably because the evaporation rate between the storms in June and July 2018 was low. This is suggested by the fact that the sediment surface was not subaerially exposed during that period, as the water level remained approximately at the sediment surface (see period between green and pink arrow in Fig. 2b). On the other hand, the biofilm formed over the uppermost ripples covers a thin carbonate lamina (orange arrow in Fig. 5b). This carbonate was probably precipitated when the seawater retained in the saline basin reached CaCO_3 supersaturation before the surface exposure (green dashed line in Fig. 2b). Attainment of CaCO_3 supersaturation in Paso Seco is

Figure 5. **a)** Vertical sedimentary profile of the block containing the ripple formed in June 2018 (yellow arrow). The location of the analysed thin section is indicated by the orange square. **b)** Thin-section cross-polarised light photomicrograph of the orange square in (a). Yellow arrow indicates the body of the ripple formed in June 2018; red dashed line indicates a layer made up fine silt and clay; green arrow indicates another ripple formed because of a moderate storm; orange arrow indicates a micritic layer covered by an organic matter lamina forming a biofilm. **c)** Cross-polarised light photomicrograph of a close-up of the microbial mat below the ripple (square c in b). Yellow arrows indicate brownish organic matter laminae; orange arrows indicate micritic calcite laminae; blue arrows indicate peloids; pink arrows indicate coalescent peloids. **d)** Cross-polarised light photomicrograph of a close-up of the microbial mat of (c). Blue arrows indicate peloids; pink arrows indicate coalescent peloids. **e)** Plane-polarised light photomicrograph of a close-up of the uppermost sand ripple (square e-f in b). Green arrow indicates the body of the ripple; b indicates an organic matter lamina forming a biofilm at the surface; orange dashed lines (m) indicate a laterally continuous dense micritic layer. Red dashed line indicates a lamina composed of fine silt and clay. **f)** Cross-polarised light photomicrograph of (e). **g)** Thin-section cross-polarised light photomicrograph of the square g in (b). Red dashed line indicates a lamina made up fine silt and clay. **h)** Plane-polarised light photomicrograph of a close-up of yellow square in (e). b indicates the biofilm covering the micritic layer (m). **i)** Cross-polarised light photomicrograph of (h).

possible by a combination of factors, including a) the semiarid climatic conditions that characterize the area (Aliaga *et al.*, 2017); b) seawater retention over the sediment surface, influenced by the topographic gradient towards inland; c) the biologically-induced changes in sediment permeability by microbial mats (Perillo *et al.*, 2019), which helps seawater retention over the sediment; and d) the aid of fast blowing dry winds. The combination of all these factors induces the seawater evaporation that leads to precipitation firstly of calcite, and subsequently of gypsum when CO_3^{2-} ion is completely depleted due to the precipitation of calcite but the water still contains calcium and sulphate anions (Fig. 3b).

Moreover, Maisano *et al.* (2020) suggested that biotic processes also play an important role in carbonate precipitation in Paso Seco. On one hand, these authors described the presence of occasional whittings in the study area and suggested that microorganisms in suspension in the water column could act as nucleation sites and could even promote carbonate precipitation in relation to photosynthesis processes. A similar explanation, increased photosynthetic activity and cell metabolism of cyanobacterial picoplankton, was sustained by Thompson *et al.* (1997) for whittings in a lacustrine water system. On the other hand, Maisano *et al.* (2020) also suggested that the peloidal textures shown by some carbonate laminae in Paso Seco may be related with precipitation of calcite in the EPS of the microbial mats, similarly to what has been described in microbial mats from a salina environment (Gerdes *et al.*, 1994). Taking into account all these considerations, the formation of the carbonate lamina that covers the studied ripples resulted likely from the combination of CaCO_3 supersaturation attainment due to intense seawater evaporation retained over the sediment surface plus the existence of nucleation sites (suspended microorganisms and/or EPS in the mats) for carbonate precipitation.

The sedimentary processes identified in Paso Seco, in close relationship to the different hydrodynamic events that took place along 2018, allow us to interpret the main factors that contributed to the preservation of the studied ripples. First, the biostabilization of the ripples caused by the development of a thin biofilm over their surface could help to prevent these structures from erosion. Similarly, Cuadrado *et al.* (2021) demonstrated the important role of microbial mat development in the

preservation of a vertebrate footprint formed in Paso Seco, which agrees with the postulation of Seilacher (2008), who argued that cyanobacterial films are instrumental in the preservation of animal tracks and drag marks. Likewise, numerous authors have highlighted the role of microbial mats in enhancing exceptional fossil preservation (Varejão *et al.*, 2019). Secondly, the present study shows that carbonate precipitation within the microbial mats that colonize the ripples is crucial for the early lithification of these structures, which might significantly increase their preservation potential in the geological record.

Although this study on present-day microbial mats proves the importance that microbial colonization may have in the preservation of ripples in some environments the application of these results to the ancient successions is challenging due to the difficulties on recognizing microbial signatures in the fossil record. The observations on present-day ripples of Paso Seco show that a thin fine-grained sediment lamina over the ripple (upper red dashed line, Fig. 5b) is the result of the colonization of the sediment surface by a biofilm. However, this feature might be difficult to identify in ancient deposits, either because it is not preserved or because it is misinterpreted as a mud drape produced by purely physical processes (Cuadrado, 2020). Nevertheless, the recognition of a continuous fine-grained lamina and a continuous micritic carbonate lamina covering a ripple train is a good first step to consider that ripple preservation and early lithification may have been enhanced by the development of a biofilm or microbial mat.

CONCLUSIONS

This study documents ripple formation, stabilization, and the beginning of their lithification in a colonized coastal environment. Ripples are developed due to episodic high-energy events, and after that, subsequent calm conditions allow the accumulation of a clay and silt layer and the creation of a sinoidal structure by the development of a thin biofilm over the sedimentary sand structures. Furthermore, microbial mats change the permeability of the sediment, so the seawater that frequently enters the study area and provides Ca^{2+} and CO_3^{2-} ions can progressively evaporate forming a continuous lamina of micritic calcite, which promotes the early lithification of the sediment.

Overlying the micrite layer a biofilm is formed, which can enhance the biostabilization of the ripples by migration of microbial filaments and production of EPS. Therefore, the structures would acquire a first preservation that may protect them from following erosional processes. These observations on a present-day sedimentary environment may be particularly useful to better understand the formation, biostabilization and early lithification of ripple marks and may have a potential application for the interpretation of the stratigraphic record.

Acknowledgements

We thank the staff of the UNS/INGEOSUR Petronomy Laboratory for the skilful preparation of thin sections. This work was supported by Argentinian CONICET (grant number PIP 2013 N°4061), AGENCIA (PICT 2019-2182 PICT 2015 302) and SECYT-UNS (grant number PGI 24/H138), the International Association of Sedimentology (IAS, Post-Doctoral Research Grant) as well as by the Spanish Department of Science, Innovation and Universities (project PGC2018-094034-B-C21) and the Spanish Department of Education, Culture and Sports (José Castillejo grants CAS16/00124 and CAS17/00270). We also thanks to R. Agustin Mors, as well as Associate Editor Fernando Gomez, for their worthwhile suggestions to enhance a previous version of this paper.

REFERENCES

- Aliaga, V.S., Ferrelli, F., and Piccolo, M.C. (2017). Regionalization of climate over the Argentine Pampas. *International journal of climatology*, 37: 1237-1247.
- American Public Health Association, and American Public Health Association. (2005). Standard methods for the examination of water and wastewater. APHA. *American Water Works Association and Water Environment Federation*, 21st ed.; *American Public Health Association*: Washington, DC, USA.
- Cuadrado, D.G. (2020). Geobiological model of ripple genesis and preservation in a heterolithic sedimentary sequence for a supratidal area. *Sedimentology*, 67(5): 2747-2763.
- Cuadrado, D.G. and Blasi, A. (2018). Presencia de actividad microbiana en ambientes silicoclásticos actuales y en paleoambientes. Estudio comparativo para el establecimiento de análogos. *Latin American Journal of Sedimentology and Basin Analysis*, 24: 39-73.
- Cuadrado, D.G., and Pan, J. (2018). Field observations on the evolution of reticulate patterns in microbial mats in a modern siliciclastic coastal environment. *Journal of Sedimentary Research*, 88 (1): 24-37.
- Cuadrado, D.G., Pan, J., Gómez, E.A., and Maisano, L. (2015). Deformed microbial mat structures in a semiarid temperate coastal setting. *Sedimentary Geology*, 325: 106-118.
- Cuadrado, D.G., Maisano, L., and Quijada, I.E. (2021). Role of microbial mats and high sedimentation rates in the early burial and preservation of footprints in a siliciclastic tidal flat. *Journal of Sedimentary Research*, 91 (5): 479-494.
- Dickson, J.A.D. (1966). Carbonate identification and genesis as revealed by staining. *Journal of Sedimentary Research*, 36(2): 491-505.
- Dupraz, C., and Visscher, P.T. (2005). Microbial lithification in marine stromatolites and hypersaline mats. *Trends in microbiology*, 13 (9): 429-438.
- Dupraz, C., Visscher, P.T., Baumgartner, L.K., and Reid, R.P. (2004). Microbe-mineral interaction: Early carbonate precipitation in a hypersaline lake (Eleuthera Islands, Bahamas). *Sedimentology*, 51: 745-765.
- Gerdes, G., Dunajtschik-Piewak, K., Riege, H., Taher, A.G., Krumbein, W.E., and Reineck, H.E. (1994). Structural diversity of biogenic carbonate particles in microbial mats. *Sedimentology*, 41 (6): 1273-1294.
- Kennedy, J.F. (1969). The formation of sediment ripples, dunes, and antidunes. *Annual review of fluid mechanics*, 1 (1): 147-168.
- Maisano, L., Cuadrado, D.G., and Gómez, E.A. (2019). Processes of MISS-formation in a modern siliciclastic tidal flat, Patagonia (Argentina). *Sedimentary Geology*, 381: 1-12.
- Maisano, L., Quijada, I. E., Cuadrado, D. G., Perillo, V. L., Pan, J., and Martínez, A.M. (2020). Carbonate laminae recorded in a siliciclastic tidal flat colonized by microbial mats. *Sedimentary Geology*, 405: 105702.
- Noffke, N. (2003). Epibenthic cyanobacterial communities interacting with sedimentary processes in siliciclastic depositional systems (present and past). In *Fossil and recent biofilms* (pp. 265-280). Springer, Dordrecht.
- Noffke, N., and Paterson, D. (2008). Microbial interactions with physical sediment dynamics, and their significance for the interpretation of Earth's biological history. *Geobiology*, 6 (1): 1-4.
- Noffke, N. (2010). *Geobiology: Microbial mats in sandy deposits from the Archean Era to today*. Springer Science and Business Media.
- Noffke, N., Gerdes, G., Klenke, T., and Krumbein, W.E. (2001). Microbially induced sedimentary structures: a new category within the classification of primary sedimentary structures. *Journal of Sedimentary Research*, 71 (5): 649-656.
- Pan, J., Cuadrado, D.G., and Bournod, C.N. (2017). Diatom-driven recolonization of microbial mat-dominated siliciclastic tidal flat sediments. *FEMS Microbiology Ecology*, 93 (10).
- Paterson, D.M. (1994). Microbiological mediation of sediment structure and behaviour. In *Microbial mats* (pp. 97-109). Springer, Berlin, Heidelberg.
- Perillo, V. L., Maisano, L., Martínez, A. M., Quijada, I. E., and Cuadrado, D. G. (2019). Microbial mat contribution to the formation of an evaporitic environment in a temperate-latitude ecosystem. *Journal of Hydrology*, 575: 105-114.
- Reid, R.P., Visscher, P.T., Decho, A.W., Stolz, J.F., Bebout, B.M., Dupraz, C., and Steppe, T.F. (2000). The role of microbes in accretion, lamination and early lithification of modern marine stromatolites. *Nature*, 406: 989-992.
- Seilacher, A. (2008). Biomats, biofilms, and bioglyphs as preservational agents for arthropod trackways. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 270: 252-257.

- Shepard, R.N., and Sumner, D.Y. (2010). Undirected motility of filamentous cyanobacteria produces reticulate mats. *Geobiology*, 8 (3): 179-190.
- Stempels Bautista, C. (2019). *Ondas de tormenta en la zona costera de Paso Seco. Buenos Aires, Argentina*. (Dissertation, Tesis), Universidad Nacional del Sur, 66 pp.
- Strickland, J. D. H., and Parsons, T. R. (1972). *A practical handbook of seawater analysis*. Fisheries Research Board of Canada, 328 pp.
- Thompson, J.B., Schultze-Lam, S., Beveridge, T.J., and Des Marais, D.J. (1997) Whiting events: biogenic origin due to the photosynthetic activity of cyanobacterial picoplankton. *Limnology and Oceanography*, 42: 133-141
- Varejão, F. G., Warren, L. V., Simões, M. G., Fürsich, F. T., Matos, S. A., and Assine, M. L. (2019). Exceptional preservation of soft tissues by microbial entombment: insights into the taphonomy of the Crato Konservat-Lagerstätte. *Palaios*, 34(7): 331-348.