

DROUGHT STRESS

Leaf Photosynthesis During Grain Filling Under Mediterranean Environments: Are Barley or Traditional Wheat More Efficient Than Modern Wheats?

L. G. Abeledo¹, R. Savin² & G. A. Slafer^{2,3}

1 Departamento de Producción Vegetal, Facultad de Agronomía, Universidad de Buenos Aires, and CONICET, Buenos Aires, Argentina

2 Department of Crop and Forest Sciences and AGROTECNIO (Center for Research in Agrotechnology), University of Lleida, Lleida, Spain

3 ICREA (Catalonian Institution for Research and Advanced Studies), Lleida, Spain

Keywordsgrain-filling period; *Hordeum distichum* L.; instantaneous water-use efficiency; leaf photosynthetic rate; *Triticum aestivum* L.**Correspondence**

L.G. Abeledo
 Departamento de Producción Vegetal
 Facultad de Agronomía, Universidad de Buenos Aires
 Av. San Martín 4453
 (C1417DSE) Buenos Aires
 Argentina
 Tel. / Fax: +54 11 4524 8039 / 8053
 Email: abeledo@agro.uba.ar

Accepted February 14, 2014

doi:10.1111/jac.12054

Abstract

Barley is one of the most popular crops in dryland agricultural systems of Mediterranean areas, where it is assumed that barley, or traditional wheat cultivars, performs better than modern wheat under low-yielding conditions. It was tested whether variations in net leaf photosynthetic rate (P_N) during grain filling provide any basis for the potential better performance of barley and traditional wheat compared to modern wheats in Mediterranean areas. Two groups of field experiments were conducted in Agramunt (NE Spain) during 2005/06 (06) and 2006/07 (07) growing seasons combining low and high nitrogen (N) availabilities under rain-fed and irrigated conditions. Cultivars used in the first group of experiments were a traditional (Anza) and a modern (Soissons) wheat, whilst in a second group of experiments, a wheat (Soissons) and a barley (Sunrise) modern cultivars were used. Both wheat cultivars showed a similar P_N during grain filling but higher than that of the modern barley cultivar. Differences between species in P_N were maximized under high-yielding conditions. There were no differences between cultivars in instantaneous water-use efficiency. The barley cultivar showed a higher specific leaf area, but lower N content *per* unit of leaf area, than wheat. Photosynthetic nitrogen-use efficiency was similar between the traditional and the modern cultivar but lower than barley. Decreases in P_N after anthesis were not exactly observable in SPAD measurements. In conclusion, we found no consistent differences between cultivars in terms of post-anthesis photosynthetic activity to support the assumption of better performance under Mediterranean farm conditions of traditional wheat or barley against modern wheat.

Introduction

Wheat and barley are the most important cereal crops in Mediterranean areas, in which high temperatures and water scarcity impose a restriction to yield generation during grain filling (Araus et al. 2007, Francia et al. 2011, Jacobsen et al. 2012). Modern wheat cultivars have been normally selected for improved productivity under high-yielding conditions, whereas barley or traditional wheat cultivars have been usually defined as crops with better performance under poor environments than modern high-yielding wheats (Ceccarelli et al. 2007, Ryan et al. 2008). In fact, barley is the dominant cereal crop in the dryland systems

of the Mediterranean basin (as well as in many other low-yielding systems) as farmers assume that barley performs better than wheat under these conditions (see Gibbon 1981, Cossani et al. 2007, Ryan et al. 2008) and, when growing wheat, farmers in these systems tend to favour the use of traditional rather than high-yielding modern cultivars (assuming that high-yielding cultivars are more sensitive to stresses than their traditional counterparts; Byerlee 1996). Conflicting with these assumptions, the performance of a cultivar of high potential yield under stressful growing conditions is frequently equal or better than that of cultivars of lower yielding capacity (Richards 2000, Araus et al. 2002, Abeledo et al. 2003, Slafer and Araus

2007, Sadras and Lawson 2011). For instance, Calderini and Slafer (1999) showed for different countries that modern wheat cultivars outyielded their older counterparts not only in high-yielding conditions but also along a wide range of environments. We were interested in determining whether photosynthetic rates during grain filling (frequently the most stressed phase in Mediterranean dryland systems) provide any basis for the assumed improved performance of barley and traditional wheat compared to high-yielding wheats in the Mediterranean region of Catalonia.

Regarding breeding effects on leaf photosynthesis, there are no generalized conclusions. In some cases, it was increased (soybean, Morrison et al. 1999), whilst in others it remained virtually constant (maize, Tollenaar and Wu 1999) or even decreased (wheat, Koç et al. 2003). In the case of wheat and barley, increasing grain yield through improved partitioning (i.e. harvest index) has been far more important than through increasing photosynthesis. Consequently, despite that breeding increased yield, photosynthesis *per* unit leaf area did not seem to have been consistently improved so far (Richards 2000). Sadras et al. (2012) found, working with a set of old and modern Australian wheat cultivars, that radiation-use efficiency improved with year of cultivar release (due to changes in the leaf nitrogen concentration along the canopy profile) but without significant changes in flag leaf photosynthesis (measured at anthesis). Additional evidences at the crop level of organization showed that modern cultivars have higher radiation-use efficiency than old wheat cultivars (Calderini et al. 1995, Shearman et al. 2005), even in Mediterranean conditions with terminal stresses (Acreche et al. 2008a). This is in line with improvements in the post-anthesis radiation-use efficiency in modern cultivars (Acreche and Slafer 2009). Modern cultivars, in wheat (Shearman et al. 2005, Acreche et al. 2008b, Sadras and Lawson 2011) as well as in barley (Schittenhelm et al. 1996, Abeledo et al. 2003), sets up more grain number than old ones and, as a result, there was an increase on the demand for photoassimilates during grain filling (e.g. Reynolds et al. 2005). Accordingly, wheats with larger number of grains showed an increase in net photosynthetic rates during grain filling due to a feedback response associated with the increasing sink strength (Sharma-Natu and Ghildiyal 1994, Reynolds et al. 2005, Acreche and Slafer 2009). Although wheat and barley yields seem mostly limited by the sink size rather than by the source strength (Slafer and Savin 1994, Borrás et al. 2004, Cartelle et al. 2006, Bingham et al. 2007), modern cultivars could experience a certain degree of source–sink co-limitation (Shearman et al. 2005, Acreche et al. 2008b). Thus, it might be possible that modern cultivars had increased leaf photosynthetic rates during grain filling.

In this paper, we analysed comparatively variations in leaf photosynthetic rate during grain filling between a tra-

ditional and a modern wheat cultivar and between a modern barley and wheat cultivar. Experiments were always grown on a real farm located in a Spanish Mediterranean region. Within each experiment, genotypes were subjected to a wide range of water and nitrogen (N) conditions. Although results from this sort of field studies are more variable than those in which single factor effects are studied under controlled conditions, the approach provides a situation very close to reality of actual farmers.

Materials and Methods

General conditions and treatments

Two independent studies were conducted under field conditions in Agramunt (41° 47' 17''N, 1° 5' 59''E, altitude 337 m; Spain), a typical Mediterranean location, on soils classified as Xerofluvic typic (SSS 1999). Each study (one comparing a modern and a traditional wheat, and the other comparing modern wheat and barley, hereafter WW and WB experiments, respectively) was carried out in two consecutive growing seasons (2005/06 and 2006/07, from now on designated by their harvest year 06 and 07, respectively). These studies were not conducted in an experimental station but in a field at a real farm, that was rented to install the experiments, sown with the farmer machinery and in which weeds, insects and diseases were controlled with the practices used by the farmers. In each experiment, the treatments comprised a factorial combination of two genotypes, two N levels (unfertilized and fertilized) and two water regimes (rain-fed and irrigated). Cultivars used in experiments WW06 and WW07 were a traditional (Anza) and a modern (Soissons) bread wheat (*Triticum aestivum* L.), while in experiments WB06 and WB07, they were Soissons and a modern two-rowed barley (*Hordeum distichum* L. cv. Sunrise). Although we are aware that there is genotypic variability within each species, we had to choose only one cultivar for each genotypic treatment to manage different N × water availabilities and take several measurements during the growth cycle, particularly photosynthetic measurements in the field within few hours (see below). These cultivars represent the actual choices made by farmers in the Catalanian Mediterranean region in NE Spain (Cossani et al. 2007, Acreche et al. 2008b), and they were used as standard controls to test the performance of other cultivars for at least 10 years, including the period of the present experiments, by GENVCE (Group for the Evaluation of the New Cereals Varieties in Spain).

Previous to sowing, soil moisture and nitrate content were measured for the 0–100 cm depth. Initial N content (N₀) was 80 (WW06), 115 (WB06), 215 (WW07) and 150 kg N ha⁻¹ (WB07). N fertilization treatments consisted of 0 and 150 kg N ha⁻¹ applied as urea. The high

fertilizer application represents a common fertilization level in the region. Fertilizer applications were split in two identical quantities at seedling emergence and beginning of tillering (DC21, Zadoks *et al.* 1974). Thus, nitrogen availabilities for the fertilized treatments (N1) were 230 (WW06), 265 (WB06), 365 (WW07) and 300 kg N ha⁻¹ (WB07). Soil moisture content at sowing (0–100 cm) was, in both studies, 90 and 191 mm in 2005 and 2006, respectively. The rain-fed treatments (Rf) received 92 (06) and 333 mm (07) of water as rainfall during the crop cycle. The irrigated treatments (Ir) consisted of a weekly irrigation of ~10 mm each from beginning of stem elongation (DC31) to maturity (DC94) in both studies and both years. Throughout the crop cycle, ~121 mm and 220 mm were irrigated in WW06 and WW07 and 85 and 327 mm in WB06 and WB07, respectively, of which 25–30 % were irrigated from anthesis to maturity. A drip irrigation system was used for the distribution of water. The irrigation system was a set of hoses, parallel to the crop rows, derived from a central pipe connected to a bomb.

The experiments were arranged in a split block–split plot design with three replicates. Main plots were assigned to the combination of cultivars and irrigation treatments, and sub-plots to fertilization treatments. Each of the experimental units (sub-plots) was 3 m wide and 5 m long. The experiments of both studies were sown in the same dates, on 30 November 2005 and 06 November 2006. Planting density was, for both studies, 180 plants m⁻² in the first season and 245 plants m⁻² in the second season.

Measurements

Single leaf photosynthetic rate and related traits were determined weekly in each experimental unit from anthesis (DC65) to the end of grain filling using a portable photosynthesis system (Model LCI, ADC BioScientific, Great Amwell, UK). Measurements were performed in the leaf contributing most photoassimilates to the growing grains: the flag leaf in wheat (Cruz-Aguado *et al.* 1999, Yin *et al.* 2009) and the penultimate leaf in barley (Jenkyn and Anilkumar 1990). The flag leaf in wheat and the penultimate leaf in barley are the largest leaves in each case, and they are commonplace to measure post-anthesis photosynthesis in each species (see, for example, Pearman *et al.* 1979, Austin *et al.* 1982, Shearman *et al.* 2005, Zheng *et al.* 2011, for measurements in flag leaf in wheat, and Planchon *et al.* 1989, Sicher and Bunce 1997, Rybiński and Garczyński 2004, for measurements in the penultimate leaf in barley). Measurements were taken between middle morning and noon on cloudless days by holding the photosynthetic chamber perpendicularly to the direction of the sunrays (incident radiation always above 1000 µmol photons m⁻² s⁻¹). The traits measured were net CO₂ assimila-

tion (net photosynthetic rate, P_N) and leaf transpiration rate (E). To compare across time, transpiration rate was adjusted to the air vapour pressure deficit of each day (E_{VPD}) (Turner and Sinclair 1983). Instantaneous water-use efficiency (WUE_i ; Tambussi *et al.* 2007) for each treatment was calculated as the slope of the linear regression between P_N and E_{VPD} during the grain-filling period.

Leaf chlorophyll concentration was estimated *in situ* using a portable chlorophyll metre (SPAD-520; Minolta, Tokyo, Japan) from anthesis to maturity. Measurements were made in each leaf used to determine gas exchange plus in three-additional leaves (four leaves per sub-plot in total) taking three readings *per* leaf, and their average was recorded. At anthesis, immediately after the gas exchange measurements were taken, the four leaves were harvested and green area *per* leaf was recorded using a leaf area metre (AT Dias II, Delta-T Devices, Cambridge, UK); and individual leaf dry weight (after oven-drying for 72 h at 65 °C) was measured with an electronic precision balance (HR-200-AC; A&D Instruments Ltd., Tokyo, Japan). Specific leaf area (SLA) was estimated as the ratio of leaf area to dry weight. Leaves were then milled using a pestle and a mortar, and leaf N concentration was measured through the Dumas method (AACC 2000). Leaf N content was calculated by multiplying leaf dry matter and N concentration. Specific leaf nitrogen (SLN) was calculated as the ratio of leaf nitrogen content to leaf area. Photosynthetic nitrogen-use efficiency ($PNUE$) was assessed as the ratio of P_N to nitrogen content per unit leaf area.

Dynamics of leaf photosynthesis and SPAD measurements were related to time using thermal time units with a base temperature of 8.2 °C for wheat (Slafer and Savin 1991) and 7.1 °C for barley (Wallwork *et al.* 1998). For integrating the effects of treatments on P_N throughout grain filling, we calculated the integral of the area underneath the curve joining the data points for the relationship between P_N and thermal time, and by analogy with the term ‘leaf area duration’ (Welbank *et al.* 1966), this integral was termed post-anthesis P_N duration. P_N was regressed against accumulated thermal time using a linear equation as follows:

$$y = a + bx$$

where y is the net photosynthetic rate (P_N , µmol CO₂ m⁻² s⁻¹), a the intercept, b the rate of decrease of P_N throughout the grain-filling period (µmol CO₂ m⁻² s⁻¹ °Cd⁻¹) and x the thermal time from anthesis (°Cd). Parameters were iteratively estimated using an optimization model (Jandel Scientific 1991).

Environmental conditions

The main environmental difference between years was related to the amount and timing of rainfall. Total precipitation




Fig. 1 Maximum (closed circles) and minimum (open circles) temperatures (up panels), global solar radiation (closed triangles) and air vapour pressure deficit (open triangles, right Y axis) (middle panels), and rainfall (dashed columns) (down panels) during experiments carried out under field conditions comparing a traditional and modern wheat cultivar (WW experiments) and comparing a modern wheat and barley cultivar (WB experiments) during the 2005/06 (06) and 2006/07 (07) growing seasons in Agramunt, NE Spain. The arrows indicate sowing (solid arrow) and average anthesis date (open arrow).

throughout the 2005/06 growing season was 92 mm, but only 5 mm precipitated from anthesis to maturity. Precipitation in 2006/07, a wet season, was 333 mm, of which 74 mm fall from anthesis to maturity (Fig. 1). Temperature, solar radiation and air vapour pressure deficit regimes were quite similar between both growing seasons (Fig. 1). The period between anthesis and maturity had a mean temperature of 20.9 °C in 2005/06 and 20.5 °C in 2006/07, while the mean solar radiation was 25.8 and 25.3 MJ m⁻² d⁻¹, respectively. The daily air vapour pressure deficit increased during the grain-filling period from 0.85 (near anthesis) to 1.17 KPa (at the end of the period) (Fig. 1).

Statistical analyses

Data were subjected to analysis of variance (Statistix 2000) and differences among treatments established at a level of significance of $P < 0.05$. The degree of association between variables was estimated by linear regression models.

Results

Photosynthetic traits throughout grain filling

P_N in 2006, a dry season (Fig. 1), was significantly lower than P_N in 2007 when the plots remained rain-fed in both WW and WB experiments (Figs 2 and 3), and irrigation increased P_N far more in the dry than in the wet year (Figs 2 and 3). But, in any condition, no consistent differ-

ences were found in P_N (neither at anthesis nor in overall P_N during grain filling) between the traditional and the modern wheat cultivars (Fig. 2, WW experiments). Integrating post-anthesis P_N , measuring the area underneath the values, revealed no significant statistically differences between the traditional and the modern wheat cultivars (Table 1), although the tendency of the traditional wheat to show a higher integral photosynthesis due to an extended grain-filling period (mean of ~4 days). In experiment WB, Soissons showed a greater mean P_N at anthesis than Sunrise (Fig. 3, WB experiments). Although the rate of P_N reduction during grain filling was higher in the wheat (mean of $-0.633 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} 10 \text{ }^\circ\text{Cd}^{-1}$) than in the barley cultivar (mean of $-0.563 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} 10 \text{ }^\circ\text{Cd}^{-1}$), the differences between wheat and barley in P_N at anthesis overrode those in rate of P_N decrease during grain filling and, consequently, mean post-anthesis P_N duration in Soissons was either equal to, or higher than, Sunrise independently of the water or N level (Table 2). As the rate of decrease in P_N during grain filling was positively associated with P_N at anthesis ($R^2 = 0.75$, $P < 0.001$), differences between cultivars tended to be higher at the early phase of the post-anthesis period than at the end phase.

During grain filling, P_N was positively related to E_{VPD} in both experiments and both growing seasons (Fig. 4). WUE_i (the slope of the relationship between P_N and E_{VPD}) showed no consistent differences between the traditional and the modern wheat or between the modern barley and wheat (Fig. 4).




Fig. 2 Relationship between net leaf photosynthetic rate (P_N) and thermal time from anthesis for a traditional (Anza, TW, closed circles) and modern (Soissons, MW, open circles) wheat cultivar (WW experiments), grown under two nitrogen levels (unfertilized N0 and fertilized N1) and two water regimes (rain-fed Rf and irrigated Ir) in Agramunt, NE Spain, in 2005/06 (06) and 2006/07 (07) growing seasons. Each data point is the average of the experimental replicates, and vertical bars indicate 1 S.D. (when not seen it is because the length of the bar was smaller than the symbol). It is shown the overall rate of decrease in P_N during grain filling ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} 10 ^{\circ}\text{Cd}^{-1}$) (± 1 S.D.).

Leaf traits and its relationship with photosynthetic rate

In WW, there were relatively small variations in SLA between the traditional and the modern wheat cultivar (Table 1), and there was no clear association between P_N and SLA ($P > 0.10$). In WB, the SLA was higher




Fig. 3 Relationship between net leaf photosynthetic rate (P_N) and thermal time from anthesis for a modern barley (Sunrise, MB, closed squares) and wheat (Soissons, MW, open squares) cultivar (WB experiments), grown under two nitrogen levels (unfertilized N0 and fertilized N1) and two water regimes (rain-fed Rf and irrigated Ir) in Agramunt, NE Spain, in 2005/06 (06) and 2006/07 (07) growing seasons. Each data point is the average of the experimental replicates, and vertical bars indicate 1 S.D. (when not seen it is because the length of the bar was smaller than the symbol). It is shown the overall rate of decrease in P_N during grain filling ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} 10 ^{\circ}\text{Cd}^{-1}$) (± 1 S.D.).

($P < 0.005$) in Sunrise than in Soissons (mean SLA of 193 and 230 $\text{mm}^2 \text{ g}^{-1}$ for Sunrise, and of 126 and 171 $\text{mm}^2 \text{ g}^{-1}$ for Soissons in the 06 and 07 growing seasons, respectively). Differences between wheat and barley in P_N were negatively related to SLA, and this relationship was

Table 1 Post-anthesis P_N duration, specific leaf area (SLA), specific leaf nitrogen (SLN), and photosynthetic nitrogen use efficiency (PNUE) (\pm one standard deviation) for a traditional (Anza) and modern (Soissons) wheat cultivar (WW experiments), grown under two nitrogen levels (unfertilized N0 and fertilized N1) and two water regimes (rain-fed Rf and irrigated Ir) in Agramunt, NE Spain, in 2005/06 (06) and 2006/07 (07) growing seasons

Exp	N	Water	Post-anthesis P_N duration ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} \text{ }^\circ\text{Cd}$)		SLA ($\text{cm}^{-2} \text{ g}^{-1}$)		SLN (mg N cm^{-2})		PNUE ($\mu\text{mol CO}_2 \text{ g}^{-1} \text{ N s}^{-1}$)	
			Anza	Soissons	Anza	Soissons	Anza	Soissons	Anza	Soissons
WW06	N0	Rf	3714 \pm 227	3212 \pm 190	116.9 \pm 18.7	126.3 \pm 14.1	0.33 \pm 0.05	0.32 \pm 0.02	4.23 \pm 0.87	4.67 \pm 1.59
		Ir	6428 \pm 450	4717 \pm 611	135.0 \pm 11.1	148.8 \pm 13.7	0.31 \pm 0.08	0.26 \pm 0.05	6.37 \pm 1.94	7.29 \pm 0.54
	N1	Rf	3533 \pm 148	2326 \pm 157	126.7 \pm 10.9	130.4 \pm 32.6	0.34 \pm 0.02	0.28 \pm 0.03	2.93 \pm 0.71	3.67 \pm 2.43
		Ir	4821 \pm 643	4578 \pm 919	142.3 \pm 4.4	156.0 \pm 22.9	0.32 \pm 0.02	0.27 \pm 0.07	5.91 \pm 0.67	8.57 \pm 5.00
WW07	N0	Rf	6130 \pm 855	5285 \pm 218	143.8 \pm 17.3	117.7 \pm 35.0	0.24 \pm 0.04	0.38 \pm 0.06	9.65 \pm 2.26	6.92 \pm 3.47
		Ir	8086 \pm 133	6388 \pm 571	98.0 \pm 13.4	123.8 \pm 19.4	0.36 \pm 0.08	0.34 \pm 0.07	6.96 \pm 2.20	7.14 \pm 2.73
	N1	Rf	6091 \pm 172	5833 \pm 737	145.3 \pm 25.4	75.9 \pm 11.5	0.26 \pm 0.05	0.60 \pm 0.09	8.80 \pm 2.76	3.73 \pm 0.55
		Ir	6329 \pm 949	5752 \pm 62	104.6 \pm 14.7	85.5 \pm 17.0	0.37 \pm 0.08	0.49 \pm 0.04	6.18 \pm 1.94	4.33 \pm 0.72
LSD	Year		948*		23.9 ns		0.07 *		1.25 *	
	Cv		1063 ns		7.2*		0.07 *		1.62 ns	
	N		803*		14.8 ns		0.06 *		1.75 ns	
	Water		1173 ns		27.8 ns		0.09 ns		1.94 ns	
	Year \times Cv		1341 ns		10.2 *		0.08 *		2.30 *	
	Year \times N		1136 ns		21.0 ns		0.08 *		2.47 ns	
	Year \times Water		1660 ns		39.3 ns		0.11 ns		2.74 *	
	Cv \times N		570 *		17.5 *		0.06 *		2.05 ns	
	Cv \times Water		966 ns		15.2 *		0.06 *		2.42 ns	
	N \times Water		1136 ns		21.0 ns		0.09 ns		2.47 ns	

LSD values for main treatments and first degree interactions (ns $P > 0.05$, * $P \leq 0.05$).

particularly evident under wetter conditions, where differences in P_N between species were maximized. Differences in SLA between wheat and barley determined that wheat showed a higher N content *per* unit of leaf area (SLN) than barley (general mean of 0.28 mg N cm⁻² and 0.18 mg N cm⁻², respectively) (Table 2), which implied that the conditions with high P_N values were those with the greater SLN. SLN followed a linear and negative relationship with SLA ($R^2 = 0.83$, $P < 0.001$ for the whole dataset). For the global set of experiments, leaf N concentration explained just a 35 % of variations in P_N ($P < 0.05$). There were no consistent differences between the traditional and the modern wheat cultivar in *PNUE* (Table 1), but barley tended to show a higher *PNUE* than wheat (general mean of 10.7 $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ N s}^{-1}$ for barley and 8.1 $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ N s}^{-1}$ for wheat; Table 2). Variations in *PNUE* were independent of *WUEi* ($P > 0.10$).

Chlorophyll metre readings measured at anthesis were similar between the traditional and the modern wheat cultivars (Fig. 5) as well as between wheat and barley (Fig. 6). Chlorophyll metre readings maintained constant until 100–200 $^\circ\text{Cd}$ after anthesis and from then on decreased at a rate that was similar between cultivars in both experiments. Variations in P_N during post-anthesis between cultivars (particularly observed in WB) were not reflected in con-

comitant differences in SPAD measurements. Thus, decreases in P_N after anthesis were partially related to SPAD measurements ($R^2 = 0.64$, $P < 0.001$), being the major departure from linear fitness concentrated in the days immediately after anthesis (the stage in which photosynthesis began to decrease but SPAD measurements still remained constant).

Discussion

The present work was carried out with the aim of analysing differences in photosynthetic traits during grain filling between wheat and barley (experiments WB) and between a traditional and a modern wheat cultivars (experiments WW) in a Mediterranean environment, to ascertain whether the preference of growers to use barley or traditional wheat instead of modern and high-yielding wheats is supported in their photosynthetic performances. It was not part of this study the analysis of yield and yield generation, but agronomic data, collected by others in these experiments, showed that under the lowest yielding conditions explored, the modern wheat effectively yielded less than the traditional wheat (2.9 and 3.6 Mg ha⁻¹) and both wheats yielded less than barley (4.4 Mg ha⁻¹; Cossani et al. 2009). However, even when in the present study the assumption

Table 2 Post-anthesis P_N duration, specific leaf area (SLA), specific leaf nitrogen (SLN), and photosynthetic nitrogen use efficiency (PNUE) (\pm one standard deviation) for a modern barley (Sunrise) and wheat (Soissons) cultivar (WB experiments) grown under two nitrogen levels (unfertilized N0 and fertilized N1) and two water regimes (rain-fed Rf and irrigated Ir) in Agramunt, NE Spain, in 2005/06 (06) and 2006/07 (07) growing seasons

Exp	N	Water	Post-anthesis P_N duration ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} \text{ }^\circ\text{Cd}$)		SLA ($\text{cm}^{-2} \text{ g}^{-1}$)		SLN (mg N cm^{-2})		PNUE ($\mu\text{mol CO}_2 \text{ g}^{-1} \text{ N s}^{-1}$)	
			Sunrise	Soissons	Sunrise	Soissons	Sunrise	Soissons	Sunrise	Soissons
WB06	N0	Rf	2991 \pm 62	2311 \pm 62	184.0 \pm 9.1	126.2 \pm 11.4	0.17 \pm 0.01	0.31 \pm 0.03	6.69 \pm 1.67	5.23 \pm 1.65
		Ir	3280 \pm 441	3800 \pm 333	199.3 \pm 3.2	126.4 \pm 3.8	0.16 \pm 0.01	0.32 \pm 0.03	7.00 \pm 1.70	6.30 \pm 0.80
	N1	Rf	3043 \pm 19	2682 \pm 151	190.3 \pm 13.2	126.4 \pm 26.2	0.22 \pm 0.04	0.32 \pm 0.08	4.72 \pm 0.89	4.99 \pm 0.17
		Ir	3011 \pm 65	4015 \pm 192	198.2 \pm 12.9	124.2 \pm 12.2	0.19 \pm 0.03	0.34 \pm 0.09	4.66 \pm 0.00	6.42 \pm 2.83
WB07	N0	Rf	3530 \pm 142	4164 \pm 462	258.8 \pm 92.8	157.9 \pm 34.4	0.12 \pm 0.05	0.24 \pm 0.06	21.37 \pm 6.04	10.29 \pm 1.22
		Ir	4032 \pm 315	4385 \pm 144	190.2 \pm 80.8	189.3 \pm 33.7	0.27 \pm 0.08	0.20 \pm 0.05	12.50 \pm 10.5	12.00 \pm 2.71
	N1	Rf	2839 \pm 223	4385 \pm 144	239.5 \pm 88.2	170.4 \pm 44.8	0.16 \pm 0.07	0.26 \pm 0.08	15.02 \pm 5.96	9.32 \pm 1.44
		Ir	3460 \pm 192	4601 \pm 109	233.0 \pm 15.9	167.0 \pm 23.7	0.18 \pm 0.03	0.24 \pm 0.04	11.67 \pm 2.44	10.08 \pm 1.07
LSD	Year		312 *		53.2 *		0.08 ns		3.05 *	
	Cv		561 *		41.0 *		0.05 *		2.47 *	
	N		294 *		11.0 ns		0.04 ns		1.62 ns	
	Water		206 *		36.4 ns		0.03 ns		3.08 ns	
	Year \times Cv		793 ns		58.0 ns		0.07 *		3.49 ns	
	Year \times N		416 ns		15.6 ns		0.06 ns		2.29 ns	
	Year \times Water		291 *		51.4 ns		0.03 *		4.35 ns	
	Cv \times N		395 ns		35.2 ns		0.04 ns		2.42 ns	
	Cv \times Water		490 ns		96.6 ns		0.09 ns		6.34 ns	
	N \times Water		416 *		15.6 ns		0.06 ns		2.29 ns	

LSD values for main treatments and first degree interactions (ns $P > 0.05$, * $P \leq 0.05$).

that traditional wheat and barley would yield more than the modern wheat under low-yielding conditions was in line with observed, this cannot be considered a proof of concept for that assumptions as when in the same region the analysis is wider (considering more experiments and treatments), there was no support for a consistently better performance of barley than wheat (Cossani *et al.* 2009), or of traditional and modern cultivars (Acreche *et al.* 2008b). In our experiments, we found no consistent differences in terms of post-anthesis photosynthetic activity to support the assumption of better performance under farm conditions of traditional wheat or barley against modern wheat. In fact, and despite the range of environments explored, traditional wheat or barley did not show a superior overall P_N than modern wheat (quite the opposite in general for the WB experiments). This is in line with the report from Dias *et al.* (2010), comparing photosynthetic rates in bread and durum wheat cultivars under heat stress conditions, where the leaf photosynthesis was similar between both species under heat stress and superior in bread wheat under optimal conditions.

The lack of consistent relation between differences in grain yield and flag leaf photosynthesis reinforces the concept that measurements carried out at individual plant level do not always correlate with the performance of the crop. Photosynthetic capacity at the crop level is determined by

total leaf area and photosynthesis capacity of each individual leaf. Thus, inconsistencies in photosynthesis between the plant and the crop level can be related to (i) flag leaf is not the only green leaf during post-anthesis (specially at the beginning of the grain-filling period when the relative importance of flag leaf area is lower comparing with the area of the rest of the leaves in the whole crop), and it can be a counterbalance between leaf area at the crop level and photosynthesis per unit of leaf area, (ii) changes in canopy architecture alter the gradient of incident and absorbed irradiance through the canopy (de Pury and Farquhar 1997, Chenu *et al.* 2005) and, as a consequence, the canopy photosynthesis, (iii) changes in vertical leaf nitrogen distribution in canopy modified photosynthesis of the whole canopy (Werger and Hirose 1991, Dreccer *et al.* 2000) and (iv) the alternative importance of ear photosynthesis as source of photoassimilates (Blum 1985, Abbad *et al.* 2004), which is greater in barley than in wheat (Johnson *et al.* 1974).

Although there was in our work a contraposition between P_N at anthesis and its rate of decrease from then on, the rate of decrease did not compensate differences in P_N at anthesis (i.e. even though P_N decreased faster in wheat, it was always equal to or higher than P_N in barley during the whole grain-filling period). The relatively low value of P_N of the barley cultivar was related to its differen-




Fig. 4 Relationship between net leaf photosynthetic rate (P_N) and transpiration rate adjusted to VPD (E_{VPD}) during grain filling for a traditional (Anza, closed circles) and modern (Soissons, open circles) wheat cultivar (WW experiments), and a modern barley (Sunrise, closed squares) and wheat (Soissons, open squares) cultivar (WB experiments), grown under different nitrogen and water availabilities in Agramunt, NE Spain, in 2005/06 (06) and 2006/07 (07) growing seasons. It is shown the slope b of the linear regression (± 1 S.E.), which represents the instantaneous water-use efficiency ($\mu\text{mol CO}_2 \text{ mol H}_2\text{O}^{-1} \text{ kPa}$).

tial characteristics with respect to wheat in (i) stomatal behaviour (i.e. rate of transpiration) and (ii) leaf levels of organization (i.e. SLA, SLN).

At the leaf level, water-use efficiency can be estimated as the ratio of P_N to the water transpired E (Tambussi et al. 2007, Cabrera-Bosquet et al. 2009). In our work, WUE_i showed no consistent differences between wheats or between wheat and barley. This is in line with the evidences provided by Cossani et al. (2012), who found no consistent advantages in water-use efficiency at the crop level of organization ($\text{kg}_{\text{grain}} \text{ ha}^{-1} \text{ mm}^{-1}$) between barley and wheat under a wide range of Mediterranean conditions.

SPAD measurements remained constant for about 150–200 °Cd after anthesis, and from then on it decreased at a rate that was similar between the traditional and the modern wheat and between wheat and barley. Thus, variations in P_N between cultivars throughout the grain-filling period were not exactly observable in SPAD measurements. As photosynthesis decreased earlier than SPAD, SPAD measurements were a better predictor of P_N when advancing the grain-filling period. The maintenance of SPAD measurements from anthesis to mid-grain-filling was consistent with the maintenance of chlorophyll content of the flag leaf observed in previous works (Ommen et al. 1999, Prasad et al. 2011). Part of the failure of SPAD in copying P_N differences at the initial phase of the post-anthesis rate could




Fig. 5 Relationship between SPAD readings and thermal time from anthesis for a traditional (Anza, closed circles) and modern (Soissons, open circles) wheat cultivar (WW experiments), grown under two nitrogen levels (unfertilized N0 and fertilized N1) and two water regimes (rain-fed Rf and irrigated Ir) in Agramunt, NE Spain, in 2005/06 (06) and 2006/07 (07) growing seasons. Each data point is the average of the experimental replicates, and vertical bars indicate 1 S.D. (when not seen it is because the length of the bar was smaller than the symbol).

be related to the fact that there were no significant differences between cultivars in leaf N concentration, which is one of the determinants of SPAD measurements (Giunta et al. 2002, Lopez-Bellido et al. 2004).

In conclusion, the traditional and modern wheat showed a similar trend of leaf photosynthetic rate during grain filling, but these rates were higher than in barley. The modern wheat, in comparison with the barley culti-




Fig. 6 Relationship between SPAD readings and thermal time from anthesis for a modern barley (Sunrise, closed squares) and wheat (Soissons, open squares) cultivar (WB experiments), grown under two nitrogen levels (unfertilized NO and fertilized N1) and two water regimes (rain-fed Rf and irrigated Ir) in Agramunt, NE Spain, in 2005/06 (06) and 2006/07 (07) growing seasons. Each data point is the average of the experimental replicates, and vertical bars indicate 1 S.D. (when not seen it is because the length of the bar was smaller than the symbol).

var, showed higher P_N and higher degree of response with improving the environmental conditions associated with a greater transpiration rate (which did not result in differences between cultivars in water-use efficiency at the leaf level). Measurements of P_N under field conditions allow recognizing restrictions to biomass production at the crop level. No consistent evidences were found for an improved photosynthetic performance of

barley or traditional wheat over modern high-yielding wheat.

Acknowledgements

The authors acknowledge the invaluable assistance of Mariano Cossani and Julia Cartelle in installing and making the experiments. This work was supported partly by WatNit-MED INCO-Project of the European Union and partly by a grant from Ministry of Science and Technology of Spain (AGL2006-07814/AGR). LGA held a 'Juan de la Cierva' post-doctoral contract funded by the Spanish Ministry of Science during the execution of this research.

References

- AACC, 2000: Approved Methods of the American Association of Cereal Chemists, 10th edn. American Association of Cereal Chemists, Saint Paul, MN, USA.
- Abbad, H., S. El Jaafari, J. Bort, and J. L. Araus, 2004: Comparison of flag leaf and ear photosynthesis with biomass and grain yield of durum wheat under various water conditions and genotypes. *Agronomie* 24, 19–28.
- Abeledo, L. G., D. F. Calderini, and G. A. Slafer, 2003: Genetic improvement of yield responsiveness to nitrogen fertilization and its physiological determinants in barley. *Euphytica* 133, 291–298.
- Acreche, M., and G. A. Slafer, 2009: Grain weight, radiation interception and use efficiency as affected by sink-strength in Mediterranean wheats released from 1940 to 2005. *Field Crops Res.* 110, 98–105.
- Acreche, M., G. Briceño-Félix, J. A. Martín Sánchez, and G. A. Slafer, 2008a: Radiation interception and use efficiency as affected by breeding in Mediterranean wheat. *Field Crops Res.* 110, 91–97.
- Acreche, M., G. Briceño-Félix, J. A. Martín Sánchez, and G. A. Slafer, 2008b: Physiological bases of genetic gains in Mediterranean bread wheat yield in Spain. *Eur. J. Agron.* 28, 162–170.
- Araus, J. L., G. A. Slafer, M. P. Reynolds, and C. Royo, 2002: Plant breeding and water relations in C3 cereals: what to breed for? *Ann. Bot.* 89, 925–940.
- Araus, J. L., J. P. Ferrio, R. Buxo, and J. Voltas, 2007: The historical perspective of dryland agriculture: lessons learned from 10 000 years of wheat cultivation. *J. Exp. Bot.* 58, 131–145.
- Austin, R. B., C. L. Morgan, M. A. Ford, and S. G. Bhagwat, 1982: Flag leaf photosynthesis of *Triticum aestivum* and related diploid and tetraploid species. *Ann. Bot.* 49, 177–189.
- Bingham, I. J., J. Blake, M. J. Foulkes, and J. Spink, 2007: Is barley yield in the UK sink limited? I. Post-anthesis radiation interception, radiation use efficiency and source-sink balance. *Field Crops Res.* 101, 198–211.
- Blum, A., 1985: Photosynthesis and transpiration in leaves and ears of wheat and barley varieties. *J. Exp. Bot.* 36, 432–440.

- Borrás, L., G. A. Slafer, and M. E. Otegui, 2004: Seed dry weight response to source-sink manipulations in wheat, maize and soybean: a quantitative reappraisal. *Field Crops Res.* 86, 131–146.
- Byerlee, D., 1996: Modern varieties, productivity, and sustainability: recent experience and emerging challenges. *World Dev.* 24, 697–718.
- Cabrera-Bosquet, L., G. Molero, S. Nogues, and J. L. Araus, 2009: Water and nitrogen conditions affect the relationships of D13C and D18O to gas exchange and growth in durum wheat. *J. Exp. Bot.* 60, 1633–1644.
- Calderini, D. F., and G. A. Slafer, 1999: Has yield stability changed with genetic improvement of wheat yield? *Euphytica* 107, 51–59.
- Calderini, D. F., M. F. Dreccer, and G. A. Slafer, 1995: Genetic improvements in wheat yield and associated traits. A re-examination of previous results and latest trends. *Plant Breed.* 114, 108–112.
- Cartelle, J., A. Pedró, R. Savin, and G. A. Slafer, 2006: Grain weight responses to post-anthesis spikelet-trimming in an old and a modern wheat under Mediterranean conditions. *Eur. J. Agron.* 25, 365–371.
- Ceccarelli, S., S. Grando, and M. Baum, 2007: Participatory plant breeding in water-limited environments. *Exp. Agric.* 43, 411–435.
- Chenu, K., N. Franck, J. Dauzat, J. F. Barczi, H. Rey, and J. Lecoq, 2005: Integrated responses of rosette organogenesis, morphogenesis and architecture to reduced incident light in *Arabidopsis thaliana* results in higher efficiency of light interception. *Funct. Plant Biol.* 32, 1123–1134.
- Cossani, C. M., R. Savin, and G. A. Slafer, 2007: Contrasting performance of barley and wheat in a wide range of conditions in Mediterranean Catalonia (Spain). *Ann. Appl. Biol.* 151, 167–173.
- Cossani, C. M., G. A. Slafer, and R. Savin, 2009: Yield and biomass in wheat and barley under a range of conditions in a Mediterranean site. *Field Crops Res.* 112, 205–213.
- Cossani, C. M., G. A. Slafer, and R. Savin, 2012: Nitrogen and water use efficiencies of wheat and barley under a Mediterranean environment in Catalonia. *Field Crops Res.* 128, 109–118.
- Cruz-Aguado, J. A., F. Reyes, R. Rodes, I. Perez, and M. Dorado, 1999: Effect of source-to-sink ratio on partitioning of dry matter and ¹⁴C-photoassimilates in wheat during grain filling. *Ann. Bot.* 83, 655–665.
- Dias, A. S., J. Semedo, J. C. Ramalho, and F. C. Lidon, 2010: Bread and durum wheat under heat stress: a comparative study on the photosynthetic performance. *J. Agron. Crop Sci.* 197, 50–56.
- Dreccer, M. F., M. Van Oijen, A. H. C. M. Schapendonk, C. S. Pot, and R. Rabbinge, 2000: Dynamics of vertical leaf nitrogen distribution in a vegetative wheat canopy. Impact on Canopy Photosynthesis. *Ann. Bot.* 86, 821–831.
- Francia, E., A. Tondelli, F. Rizza, F. W. Badeck, O. L. D. Nicosia, T. Akar, S. Grando, A. Al-Yassing, A. Benbelkacem, W. T. B. Thomas, F. van Eeuwijk, I. Romagosa, A. M. Stanca, and N. Pecchioni, 2011: Determinants of barley grain yield in a wide range of Mediterranean environments. *Field Crops Res.* 120, 169–178.
- Gibbon, D., 1981: Rainfed farming systems in the mediterranean region. *Plant Soil* 58, 59–80.
- Giunta, F., R. Motzo, and M. Deidda, 2002: SPAD readings and associated leaf traits in durum wheat, barley and triticale cultivars. *Euphytica* 125, 197–205.
- Jacobsen, S. E., C. R. Jensen, and F. Liu, 2012: Improving crop production in the arid Mediterranean climate. *Field Crops Res.* 128, 34–47.
- Jandel Scientific, 1991: Table Curve V. 3.0. User's Manual. Version 3.0 AISN Software. Jandel Scientific, Corte Madera, CA, USA.
- Jenkyn, J. F., and T. B. Anilkumar, 1990: Effects of defoliation at different growth stages and in different grain-filling environments on the growth and yield of spring barley. *Ann. App. Biol.* 116, 591–599.
- Johnson, R. R., N. M. Frey, and D. N. Moss, 1974: Effect of water stress on photosynthesis and transpiration of flag leaves and spikes of barley and wheat. *Crop Sci.* 14, 728–731.
- Koç, M., C. Barutcular, and I. Genç, 2003: Photosynthesis and productivity of old and modern durum wheats in a Mediterranean environment. *Crop Sci.* 43, 2089–2098.
- Lopez-Bellido, R. J., C. E. Shepherd, and P. B. Barraclough, 2004: Predicting post-anthesis N requirements of bread wheat with a Minolta SPAD meter. *Eur. J. Agron.* 20, 313–320.
- Morrison, M. J., H. D. Voldeng, and E. R. Cober, 1999: Physiological changes from 58 years of genetic improvement of short-season soybean cultivars in Canada. *Agron. J.* 91, 685–689.
- Ommen, O. E., A. Donnelly, S. Vanhoutvin, M. van Oijen, and R. Manderscheid, 1999: Chlorophyll content of spring wheat flag leaves grown under elevated CO₂ concentrations and other environmental stresses within the 'ESPACE-wheat' project. *Eur. J. Agron.* 10, 197–203.
- Pearman, I., S. M. Thomas, and G. N. Thorne, 1979: Effect of nitrogen fertilizer on photosynthesis of several varieties of winter wheat. *Ann. Bot.* 43, 613–621.
- Planchon, C., A. Sarrafi, and R. Ecochard, 1989: Chlorophyll fluorescence transient as a genetic marker of productivity in barley. *Euphytica* 42, 269–273.
- Prasad, P. V. V., S. R. Pisipati, I. Momcilovic, and Z. Ristic, 2011: Independent and combined effects of high temperature and drought stress during grain filling on plant yield and Chloroplast EF-Tu expression in spring wheat. *J. Agron. Crop Sci.* 197, 931–2250.
- de Pury, D. G. G., and G. D. Farquhar, 1997: Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models. *Plant Cell Env.* 20, 537–557.
- Reynolds, M. P., A. Pellegrineschi, and B. Skovmand, 2005: Sink-limitation to yield and biomass: a summary of some investigations in spring wheat. *Ann. App. Biol.* 146, 39–49.

- Richards, R. A., 2000: Selectable traits to increase crop photosynthesis and yield of grain crops. *J. Exp. Bot.* 51, 447–458.
- Ryan, J., M. Singh, and M. Pala, 2008: Long-term cereal-based rotation trials in the Mediterranean region: implications for cropping sustainability. *Adv. Agron.* 97, 273–319.
- Rybiński, W., and S. Garczyński, 2004: Influence of laser light on leaf area and parameters of photosynthetic activity in DH lines of spring barley (*Hordeum vulgare* L.). *Int. Agrophysics* 18, 261–267.
- Sadras, V. O., and C. Lawson, 2011: Genetic gain in yield and associated changes in phenotype, trait plasticity and competitive ability of South Australian wheat varieties released between 1958 and 2007. *Crop and Pasture Sci.* 62, 533–549.
- Sadras, V. O., C. Lawson, and A. Montoro, 2012: Photosynthetic traits in Australian wheat varieties released between 1958 and 2007. *Field Crops Res.* 134, 19–29.
- Schittenhelm, S., J. A. Okeno, and W. Friedt, 1996: Prospects of agronomic improvement in spring barley based on a comparison of old and new germplasm. *J. Agron. Crop Sci.* 176, 295–303.
- Sharma-Natu, P., and M. C. Ghildiyal, 1994: Photosynthesis and dry matter production in *T. monococcum* and *T. aestivum* wheat in response to ear removal. *J. Agron. Crop Sci.* 173, 218–224.
- Shearman, V. J., R. Sylvester-Bradley, R. K. Scott, and M. J. Foulkes, 2005: Physiological processes associated with wheat yield progress in the UK. *Crop Sci.* 45, 175–185.
- Sicher, R. C., and J. A. Bunce, 1997: Relationship of photosynthetic acclimation to changes of Rubisco activity in field-grown winter wheat and barley during growth in elevated carbon dioxide. *Photosynth. Res.* 52, 27–38.
- Slafer, G. A., and J. L. Araus, 2007: Physiological traits for improving wheat yield under a wide range of conditions. In: J. H. J. Spiertz, P. C. Struik, and H. H. van Laar, eds. *Scale and Complexity in Plant Systems Research: Gene-Plant-Crop Relations*, pp. 147–156. Springer, Dordrecht, The Netherlands.
- Slafer, G. A., and R. Savin, 1991: Developmental base temperature in different phenological phases of wheat (*Triticum aestivum*). *J. Exp. Bot.* 42, 1077–1082.
- Slafer, G. A., and R. Savin, 1994: Grain mass change in a semi-dwarf and a standard-height wheat cultivar under different sink-source relationships. *Field Crops Res.* 37, 39–49.
- SSS, 1999: *Soil Taxonomy. A Basic System for Making and Interpreting Soil Surveys*. Agricultural Handbook No 436. Soil Survey Staff. US Government Print Office, Washington, DC, USA.
- Statistix, 2000: *Statistix 7 for Windows. User's Manual Analytical Software*. Tallahassee, FL, USA.
- Tambussi, E. A., J. Bort, and J. L. Araus, 2007: Water use efficiency in C3 cereals under Mediterranean conditions: a review of physiological aspects. *Ann. App. Biol.* 150, 307–321.
- Tollenaar, M., and J. Wu, 1999: Yield improvement in temperate maize is attributable to greater stress tolerance. *Crop Sci.* 39, 1597–1604.
- Turner, N. C., and T. R. Sinclair, 1983: Efficient water use in crop production research or re-search. In: H. Taylor, ed. *Limitations to Efficient Water Use in Crop Production*, pp. 1–28. ASA - SSA, Madison, WI, USA.
- Wallwork, M. A. B., S. J. Logue, L. C. MacLeod, and C. F. Jenner, 1998: Effect of high temperature during grain filling on starch synthesis in the developing barley grain. *Aust. J. Plant Physiol.* 25, 173–181.
- Welbank, P. J., S. A. W. French, and K. J. Witts, 1966: Dependence of yields of wheat varieties on their leaf area durations. *Ann. Bot.* 30, 291–299.
- Werger, M. J. A., and T. Hirose, 1991: Leaf nitrogen distribution and whole canopy photosynthetic carbon gain in herbaceous stands. *Vegetatio* 97, 11–20.
- Yin, X., W. Guo, and J. H. Spiertz, 2009: A quantitative approach to characterize sink-source relationships during grain filling in contrasting wheat genotypes. *Field Crops Res.* 114, 119–126.
- Zadoks, J. C., T. T. Chang, and C. F. Konzak, 1974: A decimal code for the growth stage of cereals. *Weed Res.* 14, 415–421.
- Zheng, T. C., X. K. Zhang, G. H. Yin, L. N. Wang, Y. L. Han, L. Chen, F. Huang, J. W. Tang, X. C. Xia, and Z. H. Hed, 2011: Genetic gains in grain yield, net photosynthesis and stomatal conductance achieved in Henan Province of China between 1981 and 2008. *Field Crops Res.* 122, 225–233.