


Doxycycline inhibits  $\alpha$ -synuclein-associated pathologies in vitro and in vivo

Antonio Dominguez-Meijide, Valeria Parrales, Eftychia Vasili, Florencia González-Lizárraga, Annekatrin König, Diana F. Lázaro, Annie Lannuzel, Stéphane Haik, Elaine Del Bel, Rosana Chehín, Rita Raisman-Vozari, Patrick P. Michel, Nicolas Bizat, Tiago Fleming Outeiro



PII: S0969-9961(21)00005-X

DOI: <https://doi.org/10.1016/j.nbd.2021.105256>

Reference: YNBDI 105256

To appear in: *Neurobiology of Disease*

Received date: 16 November 2020

Revised date: 29 December 2020

Accepted date: 6 January 2021

Please cite this article as: A. Dominguez-Meijide, V. Parrales, E. Vasili, et al., Doxycycline inhibits  $\alpha$ -synuclein-associated pathologies in vitro and in vivo, *Neurobiology of Disease* (2021), <https://doi.org/10.1016/j.nbd.2021.105256>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Doxycycline inhibits $\alpha$ -synuclein-associated pathologies *in vitro* and *in vivo*

Antonio Dominguez-Mejide<sup>1,2,3,§</sup>, Valeria Parrales<sup>4,§</sup>, Eftychia Vasili<sup>1</sup>, Florencia González-Lizárraga<sup>5</sup>, Annekatrin König<sup>1</sup>, Diana F. Lázaro<sup>1</sup>, Annie Lannuzel<sup>4,6</sup>, Stéphane Haik<sup>4,11</sup>, Elaine Del Bel<sup>7</sup>, Rosana Chehín<sup>5</sup>, Rita Raisman-Vozari<sup>4</sup>, Patrick P Michel<sup>4</sup>, Nicolas Bizat<sup>4,8,\*</sup> and Tiago Fleming Outeiro<sup>1,9,10,\*</sup>

<sup>1</sup>Department of Experimental Neurodegeneration, Center for Biostructural Imaging of Neurodegeneration, University Medical Center Goettingen, Goettingen, Germany

<sup>2</sup>Laboratory of Neuroanatomy and Experimental Neurology, Dept. of Morphological Sciences, CIMUS, IDIS, University of Santiago de Compostela, Santiago de Compostela, Spain

<sup>3</sup>Networking Research Center on Neurodegenerative Diseases (CIBERNED), Madrid, Spain

<sup>4</sup>Paris Brain Institute, Inserm U 1127, CNRS UMR 7225, Sorbonne Université, F-75013 Paris, France

<sup>5</sup>Instituto de Investigación en Medicina Molecular y Celular Aplicada (IMMCA) (CONICET-UNT-SIPROSA)

<sup>6</sup>University Hospital of Pointe-à-Pitre, Neurology Department, route de Chauvel, 97139 Abymes, Guadeloupe

<sup>7</sup>Department of Basic and Oral Biology, Faculty of Odontology of Ribeirão Preto, University of São Paulo (USP), Av do Café s/n, São Paulo, Brazil. eadelbel@usp.br.

<sup>8</sup>Faculté de Pharmacie de Paris, Paris University, 4 avenue de l'Observatoire, Paris, F-75006, France

<sup>9</sup>Max Planck Institute for Experimental Medicine, Goettingen, Germany

<sup>10</sup>Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Framlington Place, Newcastle Upon Tyne, NE2 4HH, UK

<sup>11</sup>AP-HP, Cellule Nationale de Référence des Maladies de Creutzfeldt-Jakob, University Hospital Pitié-Salpêtrière, Paris, F-75013, France

§ These authors contributed equally to this work

\*Corresponding authors:

**Email:** Prof. Dr. Tiago F. Outeiro

Department of Experimental Neurodegeneration

University Medical Center Goettingen

37073 Goettingen

Germany

Telephone: +495513913545

Email: tiago.outeiro@med.uni-goettingen.de

or

Dr. Nicolas Bizat

Institut du Cerveau et de la Moelle

Hôpital de la Pitié-Sapilêtrière

47, Bd de l'Hôpital

Paris F-75013

France

Telephone: +331 5727 4388

Email: nicolas.bizat@icm-institute.org

### Keywords

alpha-synuclein, Parkinson's disease, doxycycline, aggregation, *C. elegans*

### Highlights

- Repurposing of drugs may be a valuable strategy to help treat Parkinson's disease.
- Doxycycline decreases the number and size of alpha-synuclein inclusions in cells.
- Doxycycline lowers the production of mitochondrial-derived reactive oxygen species.
- Doxycycline induces redistribution of alpha-synuclein aggregates in *C.elegans*.

## Abstract

Parkinson's disease (PD) and dementia with Lewy bodies (DLB) are neurodegenerative disorders characterized by the misfolding and aggregation of alpha-synuclein (aSyn). Doxycycline, a tetracyclic antibiotic shows neuroprotective effects, initially proposed to be due to its anti-inflammatory properties. More recently, an additional mechanism by which doxycycline may exert its neuroprotective effects has been proposed as it has been shown that it inhibits amyloid aggregation. Here, we studied the effects of doxycycline on aSyn aggregation *in vivo*, *in vitro* and in a cell free system using real-time quaking induced conversion (RT-QuIC). Using H4, SH-SY5Y and HEK293 cells, we found that doxycycline decreases the number and size of aSyn aggregates in cells. In addition, doxycycline inhibits the aggregation and seeding of recombinant aSyn, and attenuates the production of mitochondrial-derived reactive oxygen species. Finally, we found that doxycycline induces a cellular redistribution of aggregates in a *C.elegans* animal model of PD, an effect that is associated with a recovery of dopaminergic function. In summary, we provide strong evidence that doxycycline treatment may be an effective strategy against synucleinopathies.

## Introduction

Parkinson's disease (PD) is the second most common neurodegenerative disease, affecting more than ten million people worldwide (Tysnes & Storstein, 2017). PD and some other synucleinopathies like dementia with Lewy bodies (DLB) are characterized by the presence of intraneuronal aggregates composed of the protein alpha-synuclein (aSyn; (Spillantini et al, 1998)) and the overexpression of this protein in neurons contributes to these pathologies (Mehra et al, 2019; Outeiro et al, 2019). Monomeric aSyn is an intrinsically disordered neuronal protein highly enriched in presynaptic nerve terminals, consistent with the hypothesis that aSyn is a regulator of dopaminergic (DA) neurotransmission (Abeliovich et al, 2000). Upon membrane binding, aSyn promotes SNARE complex formation and may protect against neurodegeneration (Burré et al., 2014). aSyn deficient mice exhibit a reduction in striatal dopamine and an attenuation of dopamine-dependent locomotor response. The molecular mechanisms involved in synucleinopathies, from aSyn misfolding and aggregation to the various cellular effects and pathologies associated remains unclear (Brás et al, 2020). This may happen most likely due to aggregated aSyn disrupting vesicular transport, impairing autophagy, causing mitochondrial deficits and affecting vesicle recycling, neuronal plasticity and synaptic integrity (Bonini & Giasson, 2005; Liang & Tamm, 2010; Lindström et al, 2017; Murphy et al, 2000; Winslow et al, 2010). Furthermore, aggregate forms of aSyn can self-propagate within neurons and to interconnected neural networks throughout the nervous system in a prion-like manner, resulting in the spread of pathology throughout the brain and the progression of PD (Angot et al, 2012). This spread may lead to the seeding of aggregation in receiving cells, and has been observed even when exogenous aSyn is used (Desplats et al., 2009; Luk et al., 2009). Doxycycline is a wide-spectrum antibiotic that belongs to the group of the tetracyclines reported to be neuroprotective (Reglodi et al, 2017; Santa-Cecília et al, 2019). The neuroprotective effect was proposed to be due to its anti-inflammatory properties similar to minocycline, another tetracycline antibiotic (Bortolanza et al, 2018; Cronin & Grealy, 2017). In fact, in animal models of PD, it has been seen that tetracycline derivatives can protect DA neurons in the substantia nigra (Cho et al, 2009; Du et al, 2001), possibly by restraining the glial response in the 6-OHDA animal model of PD (Lazzarini et al, 2013). In addition, doxycycline inhibits fibril formation of amyloid beta (A $\beta$ ), prion protein (PrP),  $\beta$ -microglobulin and aSyn (Costa et al, 2011; Giorgetti et al, 2011; González-Lizárraga et al, 2017; Hannaoui et al, 2014; Schmitz et al, 2016). More recently, it has been proposed that doxycycline may also act against amyloid

aggregation in neurodegenerative diseases (Socias et al, 2018; Stoilova et al, 2013). Even though the neuroprotective effects of doxycycline are clear in various model systems, the mechanism of its action remains to be elucidated.

The nematode *Caenorhabditis elegans* (*C. elegans*) possess a well-known neuronal system and has emerged as a valuable experimental *in vivo* model for understanding the neurobiological processes involved in neurodegenerative diseases like Parkinson, Alzheimer's, Huntington's and prion diseases (Alexander et al, 2014; Bizat et al, 2010; Dexter et al, 2012; Gaeta et al, 2019; Hannan et al, 2016; Rand & Nonet, 1997; White et al, 1986). To study *in vivo* the potential interactions that may exist between doxycycline and aSyn, we developed transgenic models expressing the human wild-type aSyn in the GFP-tagged DA neurons.

In the present study, we show that doxycycline reduces the formation of aSyn inclusions by inhibiting the seeding process and by disrupting the seeds, as observed by RT-QuIC and Thioflavin T (ThT) seeding assays. In addition, doxycycline inhibits the production of reactive oxygen species by mitochondria. Importantly, we found that, *in vivo*, doxycycline modifies the cellular distribution of aSyn inclusions in DA neurons and restores altered motor behaviours.

## Materials and methods

### Cell-free studies

#### Generation of pre-formed fibrils

Monomeric aSyn WT was produced by transforming *E. coli* BL21-DE3 competent cells with plasmids encoding the corresponding cDNA sequence (pET21-aSyn). Purification was performed as previously reported (Masaracchia et al, 2018; Volpicelli-Daley et al, 2014) and the protein stocks were frozen in single aliquots at -80°C. The stocks were subsequently diluted in a buffer containing 100 mM Tris, 500 mM NaCl and 10 mM EDTA to a final concentration of 5 mg/ml. Samples were incubated for seven days in an Eppendorf Thermomixer Comfort (Eppendorf, USA) at 600 rpm and 37°C. The transition from the monomeric form to aggregated aSyn was determined by thioflavin T fluorescence absorbance. In brief, thioflavin T was added to a final concentration of 20  $\mu$ M, samples were mixed by slowly inverting the tubes and the mix was added to a black 96-well plate (COSTAR, Corning Incorporated) in which 100  $\mu$ l of the reaction mixture was loaded. Fluorescence intensity measurements were performed in an Infinite M200 fluorescence plate reader (TECAN) at 480 nm. Three replicates of each sample were measured.

#### Seeding assay

Monomeric aSyn solutions (70  $\mu$ M) in 20 mM HEPES, 150 mM NaCl, pH 7.4, were incubated in a Thermomixer comfort (Eppendorf) at 37°C under orbital agitation at 600 rev./min to form aggregates. These species were diluted 1/10 when harvested and further incubated with 70  $\mu$ M of fresh monomers in the absence or presence of 100  $\mu$ M doxycycline. Aliquots were taken from the seeding aggregation at different times and mixed with ThT probe, according to LeVine (LeVine, 1993). Changes in the emission fluorescence spectra with the excitation wavelength set at 450 nm were monitored using FluoroMax-4 Spectrofluorometer. We employed similar proportions between oligomers, monomers and ThT as the ones used for the RT-QuIC.

#### Real-time quaking induced conversion (RT-QuIC)

RT-QuIC reactions were performed in black 96-well plates (COSTAR, Corning Incorporated) in which 100  $\mu$ l of the reaction mixtures were loaded. The final concentrations in the reaction mix were 150 mM NaCl, 1 mM EDTA, 10  $\mu$ M ThT, 70  $\mu$ M SDS, 7  $\mu$ M monomeric aSyn and 0.7  $\mu$ M of PFF in PBS buffer (pH= 7.1). Doxycycline was added at the indicated final concentrations. Plates were covered with sealing tape and incubated at 41°C in a plate reader (Infinite M200 fluorescence plate reader,

TECAN) with intermittent shaking cycles, consisting of one minute orbital shaking at 432 rpm followed by two minutes incubation and one minute pause to measure the ThT fluorescence intensity at 480 nm. Three replicates of each sample were measured for 250 amplification cycles. Pre-formed fibrils, generated by the method proposed by Volpicelli-Daley have been shown to have similar aggregation and spreading behavior to in vivo generated seeds (Patterson et al, 2019; Volpicelli-Daley et al, 2014), furthermore, as in vivo generated seeds need to be purified from patients there are several external factors affecting the homogeneity of the results including the baseline pathology (Candelise et al, 2019; Candelise et al, 2020). Thus, the use of in vitro generated seeds will lead to more comparable results. A sample lacking PFF and SDS in the reaction mix was used as a negative control as the lack of these components will lead to no aggregation after the process.

### **Dynamic light scattering**

Dynamic light scattering measurements were performed using a DynaPro Nanostar (Whyatt) operating at  $\lambda = 633$  nm. The measurement temperature was 25°C. Protein samples contained 7.7  $\mu$ M of aSyn in RT-QuIC aggregation buffer. Measurements were always performed in triplicate.

### **In vitro studies**

#### **Cell culture**

Human neuroglioma (H4) cells were maintained in OptiMEM 1x with Glutamax medium (Gibco, Carlsbad, Germany) supplemented with 10% FBS and 1% penicillin/streptomycin (PS) at 37°C in a 5% CO<sub>2</sub> atmosphere.

Human embryonic kidney (HEK293) cells were maintained in DMEM supplemented with 10% FBS and 1% PS at 37°C in a 5% CO<sub>2</sub> atmosphere.

Human neuroblastoma (SH-SY5Y) cells were used to measure reactive oxygen species produced by mitochondria. Cells were grown in DMEM-GlutaMAX supplemented with 10% FBS and 1% PS, at 37 °C and 5% CO<sub>2</sub>. After, the cells were seeded in 48 plates at 60,000 cell/well and differentiated 24 h after plating by treatment with 10  $\mu$ M retinoic acid (Sigma) in DMEM-GlutaMAX supplemented with 1% FBS and 1% PS, at 37 °C and 5% CO<sub>2</sub> for 5 days. Media were changed every 2 days.

#### **Cell transfection**

Twenty-four hours prior to transfection, approximately 100.000 H4 cells were plated per well in a 12-well plate (Corning, Corning, New York, USA). Six hours prior to transfection medium was replaced with fresh one. Five  $\mu$ l of a 2.5 M CaCl<sub>2</sub> solution was added to 45  $\mu$ l of a solution containing equal amounts of the plasmids encoding SynT and Synphilin-1-V5 (Lázaro et al, 2014). 2xHBS calcium phosphate buffer (50 mM N, N-bis (2-hydroxyethyl)-2-aminoethanesulfonic acid (BES) pH 7.05, 280 mM NaCl, 1.5 mM Na<sub>2</sub>PO<sub>4</sub> x 2 H<sub>2</sub>O), pH 7.05 was added drop wise to the solution and vigorously mixed. The mix was incubated for 20 minutes and added dropwise to the cells while the plate was gently rocked.

Additionally, HEK293 cells were transfected according to the following protocol: twenty-four hours prior to transfection, approximately 50.000 cells were plated per well in a 24-well live imaging plate (Zell-Kontakt, Nörtern-Hardenberg, Germany). Transfection was performed with metafectene according to the manufacturer's instructions. In brief, a solution containing equal amounts of the plasmids encoding aSyn-VC and VN-Synphilin-1 (Lázaro et al, 2014) was added to a solution containing metafectene leading to a final DNA:metafectene ratio of 1:3.

#### **Cell treatment**

Seventeen hours after transfection, H4 cells were fed with fresh medium and treated with doxycycline hyclate (Sigma-Aldrich, St. Louis, MO, USA) at the indicated final



concentrations. Additionally, some cells were treated with water as it was used as solvent for doxycycline.

Seventeen hours after transfection, HEK293 cells were fed with fresh medium and treated with doxycycline hyclate to a final concentration of 10  $\mu$ M.

Additionally, exogenous pre-formed fibrils (PFFs) (Volpicelli-Daley et al, 2014) were added to HEK293 cells stable for the expression of aSyn-EGFP at 0.1  $\mu$ M 48 hours after culture. Concomitantly, 1 and 10  $\mu$ M doxycycline were added to the cells. At the end of the treatment, cells were extensively washed with PBS to remove residual proteins still outside of the cells and immunocytochemistry was performed.

### **Mitochondrial superoxide levels**

Differentiated SH-SY5Y cells were treated with 25  $\mu$ l of 70  $\mu$ M aSyn aggregates formed in the absence or presence of 100  $\mu$ M doxycycline. ROS levels produced by cells were examined by determining mitochondrial superoxide using MitoSOX Red. This probe selectively targets mitochondria and undergoes oxidation by superoxide to form a stable fluorescent compound. Briefly, SH-SY5Y cells after the different treatments were incubated with 5  $\mu$ M MitoSOX Red reagent (Invitrogen) for 10 min, then washed and fixed with 4% formaldehyde in PBS before further analysis. Fluorescent images of randomly chosen fields were acquired with identical acquisition parameters using inverted fluorescence microscopy. The experiments were run in triplicate and fluorescence intensity values calculated by adding the intensities of each pixel in the image and then the average background was subtracted.

### **Immunoblotting**

For SDS-PAGE, after 250 amplification cycles, a volume of the RT-QuIC products containing 0.2  $\mu$ g of aSyn was boiled with Laemmli buffer at 95°C for 5 minutes and loaded into 12% Bis-Tris-polyacrylamide gels and transferred to nitrocellulose membranes. The membranes were incubated overnight with different anti-aSyn primary antibodies with different binding properties: either 1:1000 mouse anti-aSyn (BD Biosciences), 1:3000 mouse anti-aggregated-aSyn (clone 5G4, Millipore), 1:2000 rabbit anti C20-Synuclein (Santa Cruz) or 1:2000 rabbit anti- $\alpha$  $\beta$ syn (Santa Cruz). For the SDD-PAGE immunoblot, the method used was a modification of a previously proposed one (Olsson et al, 2018). In brief, a volume of the RT-QuIC products containing 0.2  $\mu$ g of aSyn was loaded into 10% tricine gels. The electrophoresis was performed with a running buffer containing 0.1 M Tris base, 0.1 M tricine and 0.1% SDS. Gels were then transferred to nitrocellulose membranes. The membranes were incubated overnight with 1:3000 mouse anti-aggregated-aSyn (clone 5G4, Millipore) at 4°C. Following three washes with TBS-Tween, membranes were incubated for two hours with the corresponding HRP conjugated secondary antibody (GE Healthcare). After incubation with the secondary antibody, membranes were washed three times with TBS-Tween and developed in a chemiluminescence system (Fusion FX Vilber Lourmat).

### **Immunocytochemistry**

Twenty-four hours after treatment with doxycycline, transfected H4 cells were washed with PBS and fixed with 4% PFA for 20 minutes at room temperature (RT), followed by a permeabilization step with 0.1% Triton X-100 (Sigma-Aldrich, St. Louis, MO, USA) for 20 minutes at RT. Following permeabilization, a blocking step with 1.5% BSA (NZYTech, Lisbon, Portugal) was performed for 1 hour. Primary antibodies used were: rabbit anti-aSyn (1:2000, Santa Cruz Biotechnology, Dallas, TX, USA) and anti V5 antibody (1:2000, Life Technologies-Invitrogen, Carlsbad, CA, USA). After overnight incubation with the primary antibody at 4°C, cells were washed with PBS and incubated for 2 hours at RT with Alexa Fluor 488 donkey anti-rabbit (1:2000, Life Technologies-Invitrogen, Carlsbad, CA, USA) and Alexa Fluor 555 donkey anti-mouse (1:2000, Life Technologies-Invitrogen, Carlsbad, CA, USA). Finally, cells were stained with DAPI (1:5000 in PBS) for 5 minutes and maintained in PBS for epifluorescence microscopy.

Additionally, 48 hours after treatment HEK293 cells stably expressing aSyn-EGFP were fixed with 4% PFA for 20 minutes at RT, followed by a permeabilization step with 0.1% Triton X-100 for 20 minutes at RT. Following permeabilization, a blocking step with 1.5% BSA was performed for 1 hour. Primary antibody used was mouse anti-Syn1 (1:1000, BD Biosciences, Heidelberg, Germany). After overnight incubation with the primary antibody at 4°C, cells were washed with PBS and incubated for 2 hours at RT with Alexa Fluor 555 donkey anti-mouse (1:1000, Life Technologies-Invitrogen). Cells were finally stained with DAPI (1:5000 in PBS) for 5 minutes and mounted in coverslips for epifluorescence microscopy.

### Microscopy

Pictures for quantification and determination of the size of aSyn inclusions were taken on a Leica DMI 6000B microscope, with an objective of 40x. The camera used was a Leica DFC 350 FX R2. The filter modes used were Y3 (666 nm), GFP (516 nm) and A4 (451 nm). The software used was Leica Application Suite AF.

### Determination of the size of aSyn inclusions

Average inclusion size was determined using *ImageJ*. For each image, individual inclusions were identified, the image was then duplicated and a Gaussian blur filter was applied to one of the images with a sigma radius value of 10. The blurred image was subtracted from the original and the image threshold of the image obtained was adjusted. The resulting image was used to analyse the particle size directly with the analyse particles plug-in.

### Quantification of aSyn inclusions

H4 cells transfected with aSyn were scored based on the aSyn inclusion pattern and classified into four different groups: cells without inclusions, cells with less than six inclusions, cells between six and ten inclusions and cells with more than ten inclusions. Results were expressed as percentage of the total number of transfected cells obtained from three independent experiments. For the PFF-treated cells, the presence or absence of inclusions was assessed and the data were expressed as percentage of cells with inclusions.

### Electron microscopy

Samples (100 µl) were adsorbed onto glow-discharged 200 mesh formvar carbon coated copper grids (Electron Microscopy Sciences). Counterstaining was done with 1% uranylacetate in methanol. The excess of liquid was removed and grids were allowed dry. Images were taken in a Philips CM120 electron microscope (Philips Inc.) using a TemCam F416 CMOS camera (TVIPS).

### In vivo studies

#### Growth and maintenance of *C. elegans* strains

Animals were generally maintained in Petri dishes on solid nematode growth medium (NGM) seeded with OP50 strain of *E. coli* as previously described (Brenner, 1974) and synchronized animals were prepared by the standard bleach method (Stiernagle, 2006). For liquid culture media, after hatching, larval stage L1 animals were incubated at 20°C in liquid S-medium enriched with heat-killed of *E. coli* OP50 strain as a food source as previously described (Couillault & Ewbank, 2002; Stiernagle, 2006).

#### Plasmids constructs and generation of transgenic lines

pNB30 (*dat-1p::gfp*) and pNB50 (*dat-1p::SNCA<sup>WT</sup>*) expression vectors were produced by a gene synthesis manufacturer (GenScript). For amplification, the vectors were transferred in competent bacteria (One Shot TOP 10, Thermo Fischer Scientific) in LB media overnight at 37°C under agitation at 200 rpm. For microinjection, the total genomic DNA from plasmids was extracted from fresh transformed bacteria using a



plasmid DNA Maxi Prep kit (Nucleobond XtraMaxi EF, Macherey-Nagel). Transgenic control GFP fluorescent marker expressing line (NB30; *nbls50[dat-1p::gfp]*; named *control<sup>GFP</sup>* in this study) and double transgenic line co-expressing the full length wild-type human gene of aSyn (417 bp) and the GFP proteins (NB50; *nbls70[dat-1p::gfp; dat-1p::SNCA<sup>wt</sup>]*; named *aSyn* in this study) were generated by microinjection following standard transformation techniques (Mello & Fire, 1995). For this, we injected in the gonads of the animals (i) for the *control<sup>GFP</sup>* line, GFP fluorescent marker plasmid (NB30) at the final concentration 40 ng/μl and (ii) for the *aSyn* line, the *SNCA* gene (NB50) in association with the GFP plasmid (NB30) at the final concentration of 60 ng/μl and 40 ng/μl respectively. To obtain stable transgenic lines, integration of extrachromosomal arrays were induced by gamma irradiation exposures (42 Gy; CellRad Faxitron, Edimax) of GFP-expressing preselected L4 animals. Lines showing stable and full transmission of the *gfp* transgene expression were cloned and backcrossed three times with the *N2* wild-type referenced background. The strains *N2* and *cat-2 (e1112)* (deficient mutant of ortholog of human tyrosine hydroxylase) (Lints & Emmons, 1999) were provided from the *Caenorhabditis Genetics Center* (University of Minnesota, St. Paul, MN).

### Drug treatment

Larval stage L1 after hatching were grown in liquid media in presence of doxycycline at the final concentration of 15 μM during 3 days at 20°C under agitation (70 rpm). To avoid small progeny larvae, young adult animals before eggging (approximately 72h of incubation) were put on solid NGM media and were individually picked to put back in culture liquid media with doxycycline treatment until day 6. L-Dopa (*Sigma-Aldrich*, St. Louis, MO, USA) was dissolved in water at 10 mM and young adult stage animals were incubated at a final concentration of 1 mM during 2h hours at 20°C under agitation (70 rpm).

### Behavioural analysis

To evaluate dopaminergic behaviours induced by aSyn we used a method described by Vidal-Gadea *et al* (Vidal-Gadea *et al*, 2011). Behavioural test of young adult animals was preceded by an incubation of 2 hour in M9 medium without food source at 20°C without doxycycline in presence or absence of L-Dopa (1 mM). To motivate the movement of the worms, assay plates containing bacteria were prepared by spreading a drop of OP50 *E. coli* in the center of the plates followed by incubation at 37°C overnight. A drop of 5 μl of about 10 worms was placed on the assay plate and the assays began 5 min after the drop was absorbed by the agar, i.e., at a time where the worms were able to escape and start crawling (recordings were done for 5 min on a dry zone). We performed movies during 1 min (7 fps, AZ100M microscope from Nikon) on a total of 30 worms per condition for each experiment. Crawling velocity was analysed by tracking the centroid of animals by video using the wrMTrck plugin of *ImageJ* software (Nussbaum-Krammer *et al*, 2015). Assays were performed with three independent experiments, n = 30 animals per experimental condition and per experiment.

### Immunostaining of *C. elegans*

Hermaphrodite worms were synchronized and young adults were prepared using a variation of the *freeze-crack* method from Albertson (Albertson, 1984). Worms were washed by centrifugation in M9 and placed on a poly-L-Lysine-coated slide (superfrost) previously incubated with a drop of poly-L-Lysine (*Sigma-Adrich*) for 15 min at 60°C. The samples were incubated with 50 μl of primary antibodies diluted in the blocking solution overnight at 4°C. Monoclonal anti-aSyn antibodies used for this study were Ab6176 (1:200, *Abcam*; recognizes the N-terminal part of the human aSyn protein, red signal in fig. 4B). The tyrosine hydroxylase enzyme (TH) was detected with the antibody #22941 (1:200, Immunostar, blue signal in fig. 4B). After washing, the worms

were incubated in a solution containing the secondary anti-mouse (for TH; Alexa Fluor, Red-548, Life Technologies) and anti-rabbit antibody (for aSyn; Alexa Fluor, Red-647, Life Technologies) for 2 hours at RT. Slides were mounted with 8  $\mu$ l of anti-fade gold (Life Technologies) and stored at 4°C.

#### **Quantification of dopaminergic neurons**

Quantification of GFP tagged DA neurons was performed in young adult worms raised at 20 °C. Animals fixed in PBS with 4 % PFA were then mounted on slides. The potential neuroprotective effect of doxycycline was evaluated with an epifluorescence microscope (Apotome.2 imaging system from Zeiss) using an x100 magnification lens. Quantifications were performed from three independent experiments,  $n = 25$  animals per experimental condition and per experiment. In a second analysis, images sections of PDE DA neurons were acquired, the surface area and the intensity of the GFP of each neuron was quantified using the *3D object counter* plugin of *ImageJ* software (90). Data were obtained from three independent experiments,  $n = 20$  animals per experimental condition and per experiment.

#### **aSyn distribution in DA neurons in *C. elegans***

To determine the localisation of the aSyn immunosignal, images of DA GFP tagged neurons were captured with an inverted confocal microscope (*Leica TCS SP8*) using an x63 oil immersion objective. For quantification of aSyn aggregates, stacks of 20 images sections of 0.228  $\mu$ m per neuron were acquired per animal. Analysis of the aSyn immunostaining distribution was assessed using the *3D object counter* plugin of *ImageJ* software. Assays were performed with three independent experiments,  $n = 20$  neurons per experimental condition and per experiment.

#### **Statistical analyses**

All data were obtained from at least three independent experiments and expressed as mean  $\pm$  SEM or mean  $\pm$  SD. Comparisons were made using ANOVA followed by *Tukey's* or *Dunnnett's* post-hoc test and Student's *t* test whenever required. Differences were considered as statistically significant at  $p < 0.05$ . Statistical analyses were carried out with GraphPad Prism 5 (San Diego, California, USA) and EXCEL (Microsoft, Seattle, WA, USA).

## Results

### **Doxycycline reduces aSyn aggregation in a cell free system**

In order to evaluate the effect of different doxycycline concentrations on aSyn aggregation we used the RT-QuIC aggregation assay (Wilham et al, 2010). The results obtained showed a dose dependent decrease in ThT fluorescence intensity. Basal values decreased with higher concentrations of doxycycline pointing not only to inhibition of aggregation but also to a possible change in the conformation of the aggregated aSyn (Fig. 1A). aSyn can also aggregate into polymorphs that are “invisible” to ThT (de Giorgi et al., 2020).


Samples collected before and after the RT-QuIC reaction were run on SDS-PAGE gels and, upon immunoblotting with the Syn-1 antibody, we observed a clear signal corresponding to monomeric aSyn in the samples prior to RT-QuIC amplification (Fig. 1B). On the other hand, after RT-QuIC amplification we generally observed no monomeric, neither multimeric aSyn signal. We then incubated the membranes with other anti-aSyn antibodies that bind to larger epitopes (Fernández et al, 2003) (pan  $\alpha\beta\gamma$ -Synuclein or C20-Synuclein), and detected a smear in samples incubated with 10  $\mu$ M doxycycline (Fig. 1C). As this suggested the presence of aggregated species, we incubated the membranes with the 5G4 anti-aggregated aSyn antibody, observing again a smear in the samples treated with 10  $\mu$ M doxycycline (Fig. 1D).

Interestingly, when the samples were run in a semi-denaturing polyacrylamide gel electrophoresis (SDS-PAGE) tricine gel, samples amplified in the presence of 0.1  $\mu$ M doxycycline showed smears ranging from approximately 35 kDa all the way up to the top of the gel while samples amplified in the presence of 10  $\mu$ M doxycycline showed only higher molecular weight species (Fig. 1E). The absence of the smear when we probed with the Syn-1 antibody might be due to the differences in the epitopes recognized, as pan  $\alpha\beta\gamma$ -Synuclein and C20-Synuclein bind to the C-terminus, and the 5G4 to a different region of the protein (Fig. 1F,G). Interestingly, this characteristic of the Syn-1 antibody has been used to selectively detect monomeric aSyn in disassembly assays (De Giorgi et al., 2020).

As it has been previously reported that, above equimolar concentrations, doxycycline, may change the conformation of the protein, leading to the formation of non-fibrillary species we then analysed the samples using electron microscopy (EM) (González-Lizárraga et al, 2017). Samples amplified without doxycycline or with 1  $\mu$ M doxycycline showed the presence of fibrils, while samples amplified in the presence of 10  $\mu$ M doxycycline showed some fibrils and large non-fibrillary structures. In the RT-QuIC assay, 1  $\mu$ M doxycycline showed an effect after more than 25 cycles (Fig. 1H).

Next, in order to further assess the size of the aggregated species formed, we used dynamic light scattering (DLS). We detected very high molecular weight (HMW) species in samples amplified in the presence of 10  $\mu$ M doxycycline (Fig. 1I, Fig. A.1). We also assessed the average molecular weight of the monomers used for amplification, obtaining a result of  $16.05 \pm 1.8$  kDa, consistent with the reported

molecular weight of 14.5 kDa..



**Figure 1. Effects of the presence of different doxycycline concentrations on *in vitro* amplification of aSyn by RT-QuIC.** **A.** Effect of doxycycline on aSyn aggregation by RT-QuIC. Increasing concentrations of doxycycline lead to lower ThT fluorescence intensity. **B.** SDS-PAGE membrane of the RT-QuIC products developed with the BD anti-aSyn antibody. (b: Sample before amplification. a: Sample after amplification). **C.** SDS-PAGE membrane of the RT-QuIC products developed with the anti-alpha beta gamma aSyn antibody. (b: Sample before amplification. a: Sample after amplification). **D.** SDS-PAGE membrane of the RT-QuIC products developed with the 5G4 anti-aggregated aSyn antibody. (b: Sample before amplification. a: Sample after 250 cycles of amplification). **E.** Semi-Denaturing gel membrane of the RT-QuIC products. (b: Sample before amplification. a: Sample after 250 cycles of amplification. n: n = 4)


*Sample without PFF and SDS after 250 cycles of amplification used as negative control for RT-QuIC*). Presence of doxycycline during RT-QuIC amplification lead to the formation of higher molecular weight species above equimolar concentrations. **F.** Epitopes for the anti-human aSyn (Syn-1 and C20) and anti- $\alpha\beta\gamma$ Syn antibodies. **G.** Epitope for the 5G4 anti-aggregated human aSyn antibody. **H.** Electron microscopy pictures of the samples obtained in the RT-QuIC. (a) In samples without docycycline, abundant fibrillar structures are observed. Scale bar 360 nm. (b) Below equimolar concentrations (1  $\mu$ M) abundant fibrillar structures are still observed. Scale bar 360 nm. (c) Above equimolar concentrations (10  $\mu$ M) few fibrils are seen. Scale bar 360 nm. (d) Above equimolar concentrations large non-fibrillar structures are observed. Scale bar 360 nm. **I.** Average molecular weight obtained by DLS. The presence of 10  $\mu$ M doxycycline leads to the formation of higher molecular weight species. Data presented as mean  $\pm$  SD. \* $p < 0.05$  in comparison with 0  $\mu$ M Doxycycline. \$ $p < 0.05$  in comparison with 1  $\mu$ M Doxycycline. \$ $p < 0.05$  in comparison with 10  $\mu$ M Doxycycline. Monomers displayed an average value of 16 kDa approximately.

**Doxycycline reduces seeded aSyn aggregation and the levels of reactive oxygen species generated by mitochondria**

Next, in order to further investigate the effect of doxycycline on aSyn aggregation, we tested whether it affected the seeded aggregation of monomeric aSyn, as this might relate to the spreading of aSyn pathology as disease progresses. In vitro, in cell free models, the process of aSyn aggregation follows a typical sigmoidal curve, with a latency time of 16 hours followed by an exponential phase that reached a stationary state around 44 hours (Fig. 2A), as previously described (Ávila et al, 2014). In contrast, the presence of aSyn aggregates triggered a rapid aggregation of monomeric aSyn, abolishing the lag phase. However, when doxycycline was added to the reaction, the increase in ThT fluorescence intensity was completely blocked. This suggests that doxycycline “poisons” the seeding of aSyn aggregates species.

Mitochondrial dysfunction leads to the production of reactive oxygen species and can directly interact with variety of specific proteins to be involved in neurodegenerative disorders (Lin & Beal, 2006). For example, in PD aSyn aggregates may induce neurodegeneration by causing a mitochondrial dysfunction and thereby, the increased production of reactive oxygen species (ROS) (Park et al, 2020; Santos & Outeiro, 2020; Szegő É et al, 2019; Wang et al, 2019). For this reason, we measured intracellular ROS production in differentiated SH-SY5Y cells treated with aggregates formed in the absence or presence of 100  $\mu$ M doxycycline. SH-SY5Y cells, as most neuronal cell lines, are more susceptible to oxidative stress than glial cells (Wang & Michaelis, 2010). Using MitoSOX Red, we observed that SH-SY5Y cells exhibited a significant increase in intracellular ROS when in the presence of aSyn aggregates (Fig. 2B). However, when the cells were treated with aggregated aSyn, formed in the presence of doxycycline, this effect was almost completely blocked.





**Figure 2. Doxycycline interferes with  $\alpha$ Syn seeding and mitochondrial ROS production.**

**A.** Doxycycline inhibits  $\alpha$ Syn seeding *in vitro*. Fresh solutions of monomeric  $\alpha$ Syn were seeded with aggregates and incubated in the absence (empty squares) or presence of doxycycline (empty diamonds) at 37 °C under orbital agitation. Aggregation was assayed by ThT fluorescence emission. Unseeded aggregation kinetics are also shown (empty circles).

**B.** Intracellular ROS production visualized with MitoSOX Red probe in differentiated SH-SY5Y cells treated with  $\alpha$ Syn aggregates produced in the presence or the absence of doxycycline.

Scale bar: 100  $\mu\text{m}$ . MitoSOX Red fluorescence intensities in the same culture conditions as in A  
 \* $p < 0.05$  vs NT.

### **Doxycycline decreases the number of aSyn inclusions in cells**


In order to assess the effect of doxycycline treatment on aSyn aggregation in cell cultures, we used the SynT/Synphilin-1 model of inclusion formation (Lázaro et al, 2014; Masaracchia et al, 2020). Synphilin-1 is an interacting partner of aSyn and SynT is a C-terminally modified variant of aSyn that is more prone to aggregation. H4 cells were transiently transfected with constructs encoding for Synphilin-1 and SynT, as previously described (Lázaro et al, 2014), in order to induce the formation of LB-like inclusions (Fig. A.2A), as characterized by our and other groups (Engelender et al, 1999; Masaracchia et al, 2020; McLean et al, 2001). Previous studies showed that some inclusions are positive to pS219aSyn (Smith et al., 2005; Büttner et al., 2010; Lázaro et al., 2014; Lázaro et al, 2016). Seventeen hours after transfection, cells were treated with doxycycline at the following concentrations: 0.01  $\mu\text{M}$ , 0.1  $\mu\text{M}$ , 1  $\mu\text{M}$  and 10  $\mu\text{M}$ . The presence and number of inclusions were assessed by immunocytochemistry (Fig. 3A) 24 hours after treatment. An average of 50 positive cells for aSyn expression per treatment and experiment were then arbitrarily divided into four different groups, depending on the number of inclusions present, i.e. cells without inclusions, cells with 1 to 5 inclusions, cells with 6 to 10 inclusions, and cells with more than 10 inclusions (Fig. 3 A-E). We observed no statistically significant differences in the percentages of transfected cells showing less than six inclusions with any of the doxycycline concentrations tested (Fig. 3 C). On the other hand, the percentage of cells containing more than 6 inclusions significantly decreased when cells were treated with 10  $\mu\text{M}$  doxycycline (Fig. 3 D, E). Interestingly, the percentage of cells showing more than 10 inclusions increased when cells were treated with 0.01, 0.1 and 1  $\mu\text{M}$ . The total number of inclusions per cell showed a tendency to increase in relation with the control cells at 0.1 and 1  $\mu\text{M}$ , with values of  $4.05 \pm 1.57$ ,  $3.97 \pm 1.41$ ,  $5.86 \pm 0.88$  and  $6.08 \pm 1.56$  inclusions per cell, respectively (Fig. A.2B). Treatment with 10  $\mu\text{M}$  resulted in a drastic reduction in the total number of inclusions per cell ( $1.28 \pm 0.39$  inclusions per cell).

### **Doxycycline reduces the size of aSyn inclusions at concentrations below 10 $\mu\text{M}$**

After observing a slight increase in the percentage of cells showing more than 10 inclusions (Fig. 3E), we wondered whether this was due to the formation of smaller inclusions. To assess this, we checked the average inclusions size. Untreated cells showed average values of  $7.85 \pm 0.82 \mu\text{m}^2$  and cells treated with 0.01  $\mu\text{M}$  of doxycycline showed average values of  $6.80 \pm 1.42 \mu\text{m}^2$ . Interestingly, we observed a significant decrease in average inclusion area when cells were treated with 0.1 and 1  $\mu\text{M}$ , values been  $5.12 \pm 0.56$  and  $3.91 \pm 0.79 \mu\text{m}^2$ , respectively ( $p = 0.03$  for 0.1  $\mu\text{M}$ ,  $n = 3$  and  $p = 0.004$  for 1  $\mu\text{M}$ ,  $n = 3$ ). This was reverted when cells were treated with 10  $\mu\text{M}$  doxycycline, as the mean area was  $10.50 \pm 3.72 \mu\text{m}^2$  (Fig. 3F). The greater average size of inclusions in cells treated with 10  $\mu\text{M}$  combined with the lower average number of inclusions may suggest that, at 10  $\mu\text{M}$  doxycycline, smaller inclusions are disrupted while a few larger ones still remain, suggesting a breakage phenomenon consistent with the in vitro and EM data. Interestingly, larger inclusions remain at higher concentrations, suggesting that doxycycline may reach a point where aSyn is present mainly as oligomers that organize into large cellular inclusions that are not toxic to cells. Furthermore, the total inclusion area in the cells was also smaller, which points to not only a disruption of smaller inclusions but to an overall lower presence of aggregated aSyn (Fig. A.2C).

To test whether doxycycline was disrupting the aggregates into smaller ones, we performed live cell imaging of cells transfected with aSyn-VC and VN-Synphilin-1 bimolecular fluorescence complementation constructs (Lázaro et al, 2014) and manuscript in preparation). As the inclusions were formed through the interaction

between aSyn and Synphilin-1, the VN- and VC- fragments of Venus fluorescent protein came into contact, thereby reconstituting the functional fluorophore and emitting fluorescence (Outeiro et al, 2008). Using this aggregation model we observed a decrease in particle size over time after treatment with 10  $\mu$ M doxycycline (Fig. A.2D), consistent with the disruption of aSyn inclusions




**Figure 3. Doxycycline reduces aSyn inclusion size and number in H4 cells. A.** Representative pictures of immunostained H4 cells without inclusions, with less than six inclusions, between six and ten inclusions and with more than 10 inclusions. Scale bar 25  $\mu$ m. **B.** Percentage of cells that showed no inclusions in the absence or in the presence of

doxycycline (0.01-10  $\mu$ M). **C.** Percentage of cells that showed less than six inclusions in the presence or not of doxycycline (0.01-10  $\mu$ M). **D.** Percentage of cells that showed between six and ten inclusions in the absence or presence of doxycycline (0.01-10  $\mu$ M). Treatment with 10  $\mu$ M doxycycline leads to a significant decrease in the percentage of transfected cells containing between 6 and 10 inclusions. **E.** Percentage of cells that showed more than ten inclusions over the total number of transfected cells in the absence or presence of doxycycline (0.01-10  $\mu$ M). There was a significant decrease in the percentage of cells treated with 10  $\mu$ M doxycycline. **F.** Average area of the inclusions in cells positive for aSyn expression at different doxycycline concentrations. Treatment with 0,1 and 1  $\mu$ M doxycycline leads to a significant decrease in the percentage of transfected cells that showed more than ten inclusions. Data presented as mean  $\pm$  SD. N=3 in all experiments \*  $p < 0.05$  relative to control. \*\*  $p < 0.005$  relative to control. ANOVA and subsequent t-test against control.

**Doxycycline prevents the seeding of aSyn inclusions in PFF-treated cells**

It has been reported that the addition of exogenous aSyn PFFs seeds the aggregation of endogenous aSyn (Volpicelli-Daley et al, 2014). Thus, based on our previous results, we hypothesized that doxycycline could inhibit this process. To assess this we treated, 48 hours after plating, cells stably expressing aSyn-EGFP with PFFs at a final concentration of 100 nM of PFFs with 1 or 10  $\mu$ M doxycycline. We observed a significant decrease in the number of inclusions in cells treated with 10  $\mu$ M doxycycline (Fig. 4A, B) when compared to vehicle-treated cells.




**Figure 4. Doxycycline prevents the formation of aSyn inclusions after treatment with exogenous PFFs.** Treatment with PFFs at a concentration of 0.1  $\mu\text{M}$  for 48hrs in absence or presence of doxycycline at concentrations of 1  $\mu\text{M}$  and 10  $\mu\text{M}$ . **A** Presence of aSyn aggregates in HEK293-aSyn:EGFP stable cells treated with PFF and visualized with anti-aSyn antibody (Scale bar 20  $\mu\text{m}$ ). DAPI was used for nuclear staining. **B**. Quantification of the inclusions formed in  $n>200$  cells per condition (mean  $\pm$  SD) reveals a significant decrease in the number of inclusions after treatment with 10  $\mu\text{M}$  doxycycline. In contrast, treatment with 1  $\mu\text{M}$  of doxycycline had no effect compared to non-treated cells.  $N = 3$  independent experiments \*\*  $p < 0.005$ .



**Expression of human aSyn in the dopaminergic neurons of *C. elegans***

We first characterised the presence of the aSyn in the tagged dopaminergic neurons. For this, we evaluated the specific targeting pattern of the *dat-1p* promoter using GFP as a reporter. We observed signals in all the eight DA neurons previously described (Sulston et al, 1975), including (i) the four CEP cell bodies, their associated neuritic extensions and the two ADEs neurons located in the anterior part of the animals, and (ii) the two PDEs neurons in a more posterior lateral position (Fig. 5A). Using an anti-human aSyn antibody, we confirmed that the aSyn transgenic line expressed aSyn in all GFP<sup>+</sup> DA neurons (Fig. 5B). In addition, GFP<sup>+</sup> neurons were also immuno-positive for TH. (Fig. 5B).



**Figure 5. Targeted expression of aSyn in *C. elegans* DA neurons.** **A.** Illustration describing a *C. elegans* control line expressing the GFP reporter gene (control<sup>GFP</sup>) specifically in DA neurons and another line co-expressing the wild-type human *SNCA* gene and the GFP reporter ( $\alpha$ S), in the same neurons. The observation was made on fixed samples using an epifluorescence microscope fitted with a high magnification (x100) objective. The micrographs depict the expression pattern of the GFP reporter in the two lines and confirm the presence of


six DA neurons in the head (CEPs and ADEs) and two DA neurons (PDEs) in the middle of the body. **B.** Panel showing the concomitant expression of TH (blue), aSyn (red) and GFP (green) in the CEP and PDE neurons. These observations confirm that aSyn is expressed in the soma and the dendrites of DA neurons.

### **Doxycycline alters the subcellular distribution of aSyn in dopaminergic neurons in *C. elegans***

The total number of DA neurons tagged with GFP was similar in animals expressing aSyn and controls suggesting that the expression of aSyn does not induce DA neuronal loss. However, the expression of aSyn induced both a significant increase in the surface area of the DA cell bodies of 60.58% ( $54.54 \pm 3.44 \mu\text{m}^2$  for control<sup>GFP</sup> versus  $87.58 \pm 7.73 \mu\text{m}^2$  for aSyn line;  $p = 0.0004$ ) and a significant decrease of GFP density of 23.87% ( $118.5 \pm 12.80$  A.U. for control<sup>GFP</sup> versus  $90.21 \pm 5.04$  A.U. for aSyn line;  $p = 0.03$ ) suggesting a deleterious effect of aSyn expression on DA neurons (Fig. 6A). Chronic treatment with 15  $\mu\text{M}$  doxycycline did not induce any change in the number of DA cell bodies or their neuritic processes in comparison to untreated controls. In contrast, this tetracycline concentration significantly decreased the swollen morphology of the soma induced by the aSyn transgene. In addition, doxycycline significantly improved the altered density of GFP fluorescent signal induced by aSyn when comparing to untreated animals (Fig. 6A). In addition, we performed immunohistochemistry of aSyn on whole animal observing a regular immunolabeling of the aSyn in the soma and along the axons of the PDEs GFP tagged neurons (Fig. 6B, left panel). The treatment with doxycycline altered the distribution of the protein in the soma compartment, presenting a punctuated expression pattern (white arrows) (Figure 6B, left panel). To evaluate these modifications, we quantified the ratio of the surface area signal between the aSyn associated immunofluorescence and the GFP signal in the soma of PDE neurons. In the presence of doxycycline, we found a significant reduction (55.20%) of the surface area of the aSyn distribution signal relative to the soma of the PDE neurons (Fig. 6B, right panel), suggesting, in combination with our previous results, that larger non-toxic aggregates are formed.

### **Doxycycline restores motor alteration in aSyn *C. elegans***

The functionality of DA neurons was evaluated using a specific dopaminergic behavioural test which measures the locomotor velocity of the animal in a plate preceded with a short swimming activity in a liquid media (Vidal-Gadea et al, 2011). As expected, we found that this parameter was significantly decreased in the *cat-2* (*e1112*) TH null mutant used as a reference line (Fig. 6C). Most interesting, this parameter was strongly reduced in the transgenic line expressing aSyn in DA neurons in comparison to the control line (Fig. 6C). L-Dopa treatment significantly restored dopaminergic function in both the *cat-2* and aSyn lines. These results suggest that aSyn overexpression in the DA neurons resulted in a DA deficit inducing behavioural dysfunction despite the absence of neuronal loss. Of interest, doxycycline treatment restored locomotor behaviour specifically on aSyn line (Fig. 6C) compared to the control strain expressing only GFP (not shown), suggesting that the effect of doxycycline on aSyn distribution in DA neurons improved motor behaviour.



**Figure 6. Cellular and behavioural consequences of doxycycline treatment in *C. elegans*.**

**A.** After synchronization, a homogeneous population of young adult animals treated or not with doxycycline (15 μM), were PFA-fixed (4%), mounted on slide and observed with an epifluorescence microscope at equipped with a x63 objective. (Upper panel) Images illustrate the most characteristic features of DA GFP-tagged neurons (PDE, arrows head). (Lower panel) Quantification of the number of GFP<sup>+</sup> cell bodies and dendrites, the surface area of GFP<sup>+</sup> cell

bodies and the somal intensity of the GFP fluorescent signal in control and aSyn-expressing animals. Bar graphs show that aSyn expression did not affect the number of DA neurons but induced significant morphological alterations, such as an increase in the somal surface area and a decrease of the GFP fluorescence signal. Such changes were prevented by a chronic exposure to doxycycline. Data are expressed as mean values  $\pm$  s.e.m. Three independent experiment;  $n = 25$ /condition/experiment. Statistical significance was calculated with various tests: (i) a one-way ANOVA followed by a *Dunnett's* multiple comparisons tests with all condition versus the untreated control line (control<sup>GFP</sup>), \* $p < 0,05$  and \*\*\* $p < 0,005$ ; (ii) a two-tailed unpaired *t*-test for aSyn animals treated or not with doxycycline, # $p < 0,05$ , and ## $p < 0,01$ ; n.s : not significant. **B. Left panel:** Illustrations showing the impact of doxycycline (15  $\mu$ M) on the distribution of human aSyn in the soma of DA neurons. Representative images of the wild-type human aSyn immunosignal (red signal) in the PDE-DA GFP tagged neurons (green). In untreated animals the immunosignal for aSyn is distributed over a large part of the cell body in contrast with worms treated with doxycycline where the signal is punctuated and restricted to a small intracellular compartment (white arrows). **Right panel:** Measurement of the impact of the doxycycline treatment on the surface area occupied by the aSyn immunosignal in the soma of PDE DA neurons. The surface area of the aSyn immunosignal expressed as a ratio to the total somal surface area of GFP positive neurons, appears significantly decreased. Data are mean values  $\pm$  s.e.m. Three independent experiment,  $n = 20$ /condition/experiment. Statistical significance was calculated with two-tailed unpaired *t*-test for the two experimental conditions; \*\*\* $p < 0,005$ . **C.** Impact of the doxycycline treatment on behavioural crawling velocity after swim-to-crawl transition. Note that the loss of velocity in animals expressing wild-type aSyn is significantly prevented by a chronic doxycycline treatment in comparison with the untreated line. The null mutant *cat-2* (*e1112*) /TH was used as positive control for the assays. A treatment of the animals with 1 mM of L-DOPA 2h before the test prevented the behavioural defect observed in the line expressing aSyn. These results suggest that DA neurons are affected by the presence of the aSyn. Data are mean values  $\pm$  s.e.m. Three independent experiment,  $n = 30$ /condition/experiment. Statistical significance was calculated with various tests: (i) a one-way ANOVA followed by a *Dunnett's* multiple comparisons test versus the GFP control line (Control<sup>GFP</sup>); \*\* $p < 0,01$ , n.s is not significant; (ii) a two-tailed unpaired *t*-test for *cat-2* (*e1112*) animals treated or not with L-Dopa, #### $p < 0,0001$ ; (iii) a two-tailed unpaired *t*-test for aSyn animals treated or not with L-Dopa, ## $p < 0,01$ ; (iv) a two-tailed unpaired *t*-test for aSyn animals treated or not with doxycycline, \$\$ $p < 0,01$ .

## Discussion

The aggregation of aSyn is one of the hallmarks of synucleinopathies. In recent years, several aggregation inhibitors have been developed and tested, and the search for new ones continues (Lamirgue-Meijide et al, 2020; Hengartner & Fernandez, 2019). Different drugs have been developed and tested, such as pyrazole derivatives and hydrazones (Cukierman et al, 2020; Cukierman et al, 2017; Wagner et al, 2013). Nonetheless, most of the drugs developed are either toxic or show low bioavailability in the brain due to low water solubility or other factors. To circumvent this, new therapeutic strategies like pharmacokinetic modulation (like complexation with cyclodextrins) or repurposing of existing drugs have been developed (Barros et al, 2018; Heemskerk et al, 2002). Examples of recent repurposing of existing drugs include inhibitors of the renin-angiotensin system, anti-asthmatic drugs, COX-2 inhibitors and other NSAIDs (Hengartner & Fernandez, 2019; Jang et al, 2017; Moghaddam et al, 2007; Rodriguez-Perez et al, 2018). Following this trend, the anti-neuroinflammatory properties of tetracyclines such as minocycline and doxycycline have been studied (Bortolanza et al, 2018; Lazzarini et al, 2013). More recently, it has been proposed that tetracyclines and, specifically, doxycycline, have not only anti-neuroinflammatory properties, but can also inhibit the aggregation of misfolded aSyn (González-Lizárraga et al, 2017).

Our results in cell free systems showed that increasing concentrations of doxycycline lead to a decrease in ThT fluorescence intensity in an RT-QuIC assay. This is probably



due not only to a decrease in the aggregation rate of aSyn, but also to a possible change in the conformation of the aSyn aggregates formed. As ThT becomes fluorescent upon binding to beta-sheet structures, a possible change in aSyn aggregation may lead to lower ThT binding and, thereby, to lower ThT fluorescence (Wu et al, 2009). Indeed, it has been proposed that treatment with doxycycline induces a conformational change, as infrared studies demonstrated the presence of parallel beta-sheet instead of the typical antiparallel beta-sheet found in on-pathway toxic oligomeric species (Celej et al, 2012). Furthermore, our WB, EM and DLS results confirm previous observations where it was reported that the presence of doxycycline induces a remodelling of aSyn oligomers into off-pathway non-toxic, non-seeding, high molecular weight species (González-Lizárraga et al, 2017). In fact, it has been shown that different aSyn fibril polymorphs can exhibit differential ThT binding, consistent with some of the high molecular weight species we detected (De Giorgi et al., 2020). Our results and previous ones also prove that this effect is dose-dependent (González-Lizárraga et al, 2017). In addition, our RT-QuIC results show a decrease in ThT fluorescence intensity at high doxycycline concentrations. As these reactions were seeded, this suggests additional effects on the fibrils. Furthermore, we also demonstrate that while aSyn aggregates are very efficient at seeding native aSyn, as revealed by ThT assays, they lose all seeding ability when incubated with doxycycline. Our results in cells also show the ability of doxycycline to break intracellular aSyn-positive inclusions, as we observed an increase in the number inclusions per cell that was accompanied by a decrease in their size. This was further confirmed by live cell imaging. The lower percentage of cells with more than ten inclusions when cells were treated with 10  $\mu$ M, and the higher average inclusion size may be either due to more compact inclusions that cannot be decomposed by doxycycline or to a possible reorganization of aSyn inside individual cells that leads to fewer but larger inclusions. In addition, it has been proposed that exogenous aSyn fibrils can be internalized and act as templates for the formation of aggregates by addition of endogenous monomeric aSyn (Grassi et al, 2018; Masaracchia et al, 2018; Volpicelli-Daley et al, 2014). These results, in combination with those observed in cell free systems, further support the idea of doxycycline being able to exert inhibition of seeding. Thus, doxycycline not only degrades already formed aggregates, but it can also inhibit the formation of new ones. Interestingly, our results also suggest that the formation of certain aggregated species may be higher in the aSyn-VC and VN-Synphilin-1 BiFC models than those in the SynT/Synph-1 model, as higher doxycycline concentrations in this model still render smaller inclusions. It has been reported that an equimolar concentration of doxycycline to aSyn would be suitable to reach a protective effect (González-Lizárraga et al, 2017). The concentration of 10  $\mu$ M may be already above equimolar levels as the concentration of aSyn in cell synapse is ~50  $\mu$ M (Perni et al, 2017; Wilhelm et al, 2014). Furthermore, the neuroprotective and anti-inflammatory doses of doxycycline are below the ones needed to exert its antibiotic activity (González-Lizárraga et al, 2017; Santa-Cecília et al, 2019).

An additional pathological effect of aSyn does not come from its aggregation and propagation. It may also cause mitochondrial dysfunction and enhanced production of ROS, which may, in turn, further modulate aSyn oligomerization and cytotoxicity, leading to a vicious circle. The generation of ROS by aSyn aggregates can affect macromolecules or activate signalling pathways that culminate in central nervous system neurodegeneration (Park et al, 2020; Wang et al, 2019). For this reason, the inhibition of oxidative stress could offer an effective therapeutic approach to mitigate the progression of PD. In concordance with this concept, we show that aSyn aggregates increase the production of ROS in differentiated SH-SY5Y cells. However, aSyn aggregates formed in presence of doxycycline are less prone to induce oxidative stress in cell cultures. In this regard, minocycline and doxycycline have been proposed to protect against mitochondrial dysfunction. Furthermore, the proposed mechanism of

inhibition of aSyn aggregation may also reduce its detrimental effects on mitochondrial function (Garcia-Martinez et al, 2010; Santa-Cecília et al, 2016).

In recent years, several aSyn-inducible cell models have been developed, including tet-on and tet-off inducible systems, which are based on the induction or blocking of aSyn expression in the presence of doxycycline, respectively (Vekrellis et al, 2009; Zhu et al, 2002). These induction systems have been developed in several cell lines and animal models (Harding et al, 1997; Takahashi et al, 2007; Vasquez et al, 2018; Vekrellis et al, 2009). In tet-on systems, addition of doxycycline leads to the expression of aSyn at levels that enable the formation of inclusions (Vasquez et al, 2018). The concentrations of doxycycline used in these publications range between 2.25 and 6.7  $\mu$ M (Harding et al, 1997; Vasquez et al, 2018). Our results show that a concentration of 10  $\mu$ M efficiently modulates aSyn aggregation in cellular models. Therefore, the concentration of doxycycline required to modulate protein aggregation in our culture setting appears higher than that required for inducing aSyn expression in aSyn-inducible cell models.

To further test the biochemical properties of doxycycline in this context, we generated a new *in vivo* transgenic model of PD in *C. elegans* expressing wild-type aSyn in the DA neurons. Unlike previously established *C. elegans* models designed to model different features of PD (Cooper & Van Raamsdonk, 2018), and target tissues like muscle (Hamamichi et al, 2008; van Ham et al, 2008), the entire neuronal system (Kuwahara et al, 2006; Kuwahara et al, 2008; Lakso et al, 2003), or the DA system (Cao et al, 2005; Lakso et al, 2003), our model uses the *dat-1* promoter. *Dat-1* is the ortholog of the human dopamine transporter in the nematode and is selectively expressed in the DA system. Thus, our model enables the *in vivo* study of the specific vulnerability of DA neurons expressing aSyn. We verified the correct targeting of the transgene in the cell soma using GFP as a reporter. Human wild-type aSyn, co-expressed with the GFP reporter under the same genetic regulatory system, is present in the soma of the DA neurons. In young-adult animals, aSyn did not induce significant DA neuronal loss until the worms reached 6 days of age. In another *C. elegans* model expressing aSyn in the DA system, the authors described the presence of aSyn-dependent neuronal injuries, probably resulting from differences in the expression levels of the transgene (Cao et al, 2005). However, in our model aSyn induced abnormal neuronal morphologies consistent with a dysfunctional DA system, as indicated by a noticeable alteration of DA-associated behaviour, suggesting that overexpression of aSyn impacts DA homeostasis. These irregular neuronal morphologies and pathological behaviour are restored after chronic treatment with doxycycline, consistent with the effects observed *in vitro* model, and reinforcing the hypothesis that doxycycline may act as a neuroprotective agent.

It has been proposed that *C. elegans* movement depends not only on DA, but also on cholinergic neuron-dependent muscle contraction and GABAergic neuron-dependent relaxation. However, it has been shown that blockade of DA D2 receptors does not fully limit paralysis triggered by changes in cholinergic and GABAergic signaling (Refai & Blakely, 2019). The opposite results have also been shown, i.e. changes in cholinergic and GABAergic signaling due to D2 antagonism (Krum et al, 2020). Furthermore, it was also shown that changes in DA may affect GABAergic egg release in the worms (Cermak et al, 2020). These results demonstrate that further research into these pathways is necessary, but this is beyond the scope of our study.

In our study, we employed a classic transgenesis technique (Mello & Fire, 1995) to produce our line, resulting in the production of transgenic animals carrying multi-copy arrays of the *SNCA* transgene, which may induce non-specific neurobiological process. Additionally, some studies showed that expression of aSyn in DA motor neurons of *C. elegans* recapitulates features of aSyn neuronal propagation. Using the BiFC assay (Outeiro et al, 2008), we showed the protein could disseminate to the neighbouring serotonergic neurons and interact with another host targeted aSyn (Tyson et al, 2017). These processes have been extensively described in mammals for synucleinopathies and other proteinopathies (Peng et al, 2020). Here, we did not

observe a significant neuronal propagation of aSyn, which might be possibly explained by the fact that we used a different promoter gene to induce aSyn expression. In addition, we have not investigated the tissue expression pattern of aSyn in old animals, as this was beyond the scope of the present study.

In *C. elegans* the treatment with doxycycline induces a significant effect on development by reducing the growth rate (Letizia et al, 2018), and it appears to promote longevity via inhibition of *mrps-5* ribosomal protein and *atf-4* transcription factor mitochondrial pathways (Molenaars et al, 2020). Doxycycline has been described to play role in the inhibition of the translation process effect by suppressing the binding of tRNAs to the ribosomal-machinery system and inducing a global decrease in protein production (Nelson & Levy, 2011). Thus, we cannot rule out that the neuroprotective effect of doxycycline in our experimental conditions may also be explained *via* an indirect activation of the aging-metabolic cellular pathways and the observed effect on mitochondria and ROS levels, as decreased ROS may lead to a lower level of neuroinflammation and neuroprotective effects (Picca et al, 2020). Since aging is the major risk factor for PD, it will be interesting to assess the potential interaction between aging-associated pathways and aSyn in the novel transgenic lines we generated.

Strikingly, we observed modifications in aSyn distribution in the DA neurons in aSyn-transgenic animals chronically incubated with doxycycline. In particular, we observed a reduction of the sub-cellular distribution of the protein in the soma. This occurred together with the effects on aSyn aggregation, further strengthening the anti-aggregation properties of doxycycline *in vivo* and suggesting additional complementary effects on aSyn to those observed *in vitro* and in cell-free models. The punctate pattern observed in DA neurons *in vivo* are consistent with a combined effect of changes in cellular distribution and the formation of fewer and larger inclusions inside the cells, as observed *in vitro*. These changes lead to an overall decrease in aSyn area in relation to the total surface of the cell soma of the PDE neurons. This is not an independent effect, but the combination of different effects, i.e. inhibition of aggregation, disruption of pre-formed inclusions, blocking of seeding, and changes in inclusion area leading, ultimately, to the distinct dotted pattern observed *in vivo*. Additionally, aggregated species of aSyn produced in the presence of doxycycline lost their capacity to stimulate mitochondrial ROS production, which may also account for the restoration of DA neuron function. In fact, our results indicate that a restoration of motor behavior by doxycycline can be explained by the combination of the aforementioned decrease in ROS, combined with changes in aSyn processing, seeding and distribution.

As mentioned above, doxycycline has neuroprotective effects in several models of neurodegenerative disease and, in particular, has anti-aggregation properties (Costa et al, 2011; Paldino et al, 2020; Santa-Cecília et al, 2019). In a *C. elegans* amyloidogenesis model induced by the expression of a human pathogenic variant of  $\beta$ 2-microglobulin (D76N of  $\beta$ <sub>2</sub>-m) in the muscle (Faravelli et al, 2019), doxycycline decreases the proportion of oligomeric fraction produced by the  $\beta$ <sub>2</sub>-m aggregation, resulting in an improvement of the motor symptoms (Raimondi et al, 2017). These findings are consistent with those we report herein.

## Conclusions

In summary, our data provides novel insight into the mechanisms by which doxycycline may inhibit the aggregation of aSyn by remodelling aSyn oligomers into off-pathway non-toxic, non-seeding, high molecular weight species. We showed this effect not only *in vitro* but also in cell free models. The extreme need for alternative strategies against PD calls for attempts for repurposing existing drugs. In this context, our data suggests that doxycycline may act at different levels, modulating the aggregation process and

the effects of aSyn aggregation, promoting somatic repair, and leading to motor recovery in *C. elegans*.

## Acknowledgments

ADM is supported by a postdoctoral fellowship from the Galician Government (Programa de axuda á etapa posdoutoral, XUGA, GAIN, ED481B 2017/053).

The authors want to thank Dr. Filippo Favretto for his help and advice with the DLS measurements, Patrícia Santos for her help with the SDD western blots.

TFO is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC 2067/1- 390729940, and by SFB1286 (project B8). VP is the recipient of a Thesis scholarship from Association France Parkinson. PPM is supported by the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement No 821522 (PL MitoQUANT; this Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation program and EFPIA and Parkinson's UK). This work also benefited from support by Investissements d'Avenir (ANR-10-IAI14U-06) and the Translational Research Infrastructure for Biotherapies in Neurosciences (ANR-11-INBS-0011-NeurATRIS). This work also was supported by PICS 2015 N° 3201. The study was supported by the São Paulo State Foundation for the Support of Research (FAPESP, Brazil; Grant 2014/25029-4). EDB is a recipient of grants from the National Council for Scientific and Technological Development (CNPq, Brazil), CAPES, and the French Committee for the Evaluation of Academic and Scientific Cooperation with Brazil, Grant #848/15). EDB is a CNPq senior research fellow.

## Author Contributions

EDB, RC, RRV, PPM, NB and TFO designed research. ADM, VP, EV and FGL performed research. AK and DFL contributed new reagents or analytic tools. ADM, VP, EF, FGL, AL, SH, EDB, RC, RRV, PPM, NB and TFO analyzed data. ADM, VP, EF, FGL, AL, SH, EDB, RC, RRV, PPM, NB and TFO wrote the paper.

## Conflict of interest

The authors declare no conflict of interest.

## References

Abeliovich A, Schmitz Y, Fariñas I, Choi-Lundberg D, Ho WH, Castillo PE, Shinsky N, Verdugo JM, Armanini M, Ryan A, Hynes M, Phillips H, Sulzer D, Rosenthal A (2000) Mice lacking alpha-synuclein display functional deficits in the nigrostriatal dopamine system. *Neuron* **25**: 239-252

Albertson DG (1984) Localization of the ribosomal genes in *Caenorhabditis elegans* chromosomes by in situ hybridization using biotin-labeled probes. *The EMBO journal* **3**: 1227-1234

Alexander AG, Marfil V, Li C (2014) Use of *Caenorhabditis elegans* as a model to study Alzheimer's disease and other neurodegenerative diseases. *Frontiers in genetics* **5**: 279

Angot E, Steiner JA, Lema Tomé CM, Ekström P, Mattsson B, Björklund A, Brundin P (2012) Alpha-synuclein cell-to-cell transfer and seeding in grafted dopamineergic neurons in vivo. *PloS one* **7**: e39465

Ávila CL, Torres-Bugeau CM, Barbosa LR, Sales EM, Oliveira MO, Socías SB, Celej MS, Raisman-Vozari R, Papy-Garcia D, Itri R, Chehín RN (2014) Structural characterization of heparin-induced glyceraldehyde-3-phosphate dehydrogenase protofibrils preventing  $\alpha$ -synuclein oligomeric species toxicity. *The Journal of biological chemistry* **289**: 13838-13850

Barros MCF, Ribeiro ACF, Estes MA (2018) Cyclodextrins in Parkinson's Disease. *Biomolecules* **9**

Bizat N, Peyrin JM, Haïk S, Cochois V, Beaudry P, Laplanche JL, Néri C (2010) Neuron dysfunction is induced by prior protein with an insertional mutation via a Fyn kinase and reversed by sirtuin activation in *Caenorhabditis elegans*. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **30**: 5394-5403

Bonini NM, Giasson BI (2005) Snaring the function of alpha-synuclein. *Cell* **123**: 359-361

Bortolanza M, Nascimento GC, Socias SB, Ploper D, Chehín RN, Raisman-Vozari R, Del-Bel E (2018) Tetracycline repurposing in neurodegeneration: focus on Parkinson's disease. *Journal of neural transmission (Vienna, Austria : 1996)* **125**: 1403-1415

Brás IC, Xylaki M, Outeiro TF (2020) Mechanisms of alpha-synuclein toxicity: An update and outlook. *Progress in brain research* **252**: 91-129

Brenner S (1974) The genetics of *Caenorhabditis elegans*. *Genetics* **77**: 71-94



Burré J, Sharma M, Südhof TC (2014)  $\alpha$ -Synuclein assembles into higher-order multimers upon membrane binding to promote SNARE complex formation. *Proceedings of the National Academy of Sciences of the United States of America* **111**:E4274-E4283

Büttner S, Delay C, Franssens V, Bammens T, Ruli D, Zaunschirm S, de Oliveira RM, Outeiro TF, Madeo F, Buée L, Galas MC, Winderickx (2010) Synphilin-1 enhances alpha-synuclein aggregation in yeast and contributes to cellular stress and cell death in a Sir2-dependent manner. *PLoS One* **5**:e13700.

Candelise N, Schmitz M, Llorens F, Villar-Piqué A, Cramm M, Thom T, da Silva Correia SM, da Cunha JEG, Möbius W, Outeiro TF, Álvarez VG, Banchelli M, D'Andrea C, de Angelis M, Zafar S, Rabano A, Matteini P, Zerr I (2019) Seeding variability of different alpha synuclein strains in synucleinopathies. *Annals of neurology* **85**: 691-703

Candelise N, Schmitz M, Thüne K, Cramm M, Rabano A, Zafar S, Stoops E, Vanderstichele H, Villar-Piqué A, Llorens F, Zerr I (2020) Effect of the micro environment on  $\alpha$ -synuclein conversion and implication in seeded conversion assays. *Translational neurodegeneration* **9**: 5

Cao S, Gelwix CC, Caldwell KA, Caldwell GA (2005) Torin-mediated protection from cellular stress in the dopaminergic neurons of *Caenorhabditis elegans*. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **25**: 3801-3812

Celej MS, Sarroukh R, Goormaghtigh E, Fidelio GD, Ruyschaert JM, Raussens V (2012) Toxic prefibrillar  $\alpha$ -synuclein amyloid oligomers adopt a distinctive antiparallel  $\beta$ -sheet structure. *The Biochemical journal* **443**: 719-726

Cermak N, Yu SK, Clark R, Huang YC, Baskoylu SN, Flavell SW (2020) Whole-organism behavioral profiling reveals a role for dopamine in state-dependent motor program coupling in *C. elegans*. *eLife* **9**

Cooper JF, Van Raamsdonk JM (2018) Modeling Parkinson's Disease in *C. elegans*. *Journal of Parkinson's disease* **8**: 17-32

Costa R, Speretta E, Crowther DC, Cardoso I (2011) Testing the therapeutic potential of doxycycline in a *Drosophila melanogaster* model of Alzheimer disease. *The Journal of biological chemistry* **286**: 41647-41655

Couillault C, Ewbank JJ (2002) Diverse bacteria are pathogens of *Caenorhabditis elegans*. *Infection and immunity* **70**: 4705-4707

Cronin A, Grealy M (2017) Neuroprotective and Neuro-restorative Effects of Minocycline and Rasagiline in a Zebrafish 6-Hydroxydopamine Model of Parkinson's Disease. *Neuroscience* **367**: 34-46



Cukierman DS, Lázaro DF, Sacco P, Ferreira PR, Diniz R, Fernandez CO, Outeiro TF, Rey NA (2020) X1INH, an improved next-generation affinity-optimized hydrazonic ligand, attenuates abnormal copper(i)/copper(ii)- $\alpha$ -Syn interactions and affects protein aggregation in a cellular model of synucleinopathy. *Dalton transactions (Cambridge, England : 2003)*

Cukierman DS, Pinheiro AB, Castiñeiras-Filho SL, da Silva AS, Miotto MC, De Falco A, de PRT, Maisonette S, da Cunha AL, Hauser-Davis RA, Landeira-Fernandez J, Aucélio RQ, Outeiro TF, Pereira MD, Fernández CO, Rey NA (2017) A moderate metal-binding hydrazone meets the criteria for a bioinorganic approach towards Parkinson's disease: Therapeutic potential, blood-brain barrier crossing evaluation and preliminary toxicological studies. *Journal of inorganic biochemistry* **170**: 160-168

Cho Y, Son HJ, Kim EM, Choi JH, Kim ST, Ji IJ, Choi DH, Joh TH, Kim YS, Hwang O (2009) Doxycycline is neuroprotective against nigral dopaminergic degeneration by a dual mechanism involving MMP-3. *Neurotoxicity research* **16**: 361-371

De Giorgi F, Laferrière F, Zinghirino F, Faggiani E, Lendé A, Bertoni M, Yu X, Grélard A, Morvan E, Morvan E, Habenstein B, Dutheil N, Doudnikoff E, Danilov J, Claverol S, Quin C, Loquet A, Bezard E, Ichas F (2020) Novel self-replicating  $\alpha$ -synuclein polymorphs that escape ThT monitoring can spontaneously emerge and acutely spread in neurons. *Science Advances* **6**:eabc4364

Desplats P, Lee HJ, Bae EJ, Patrick C, Rockenstein E, Crew L, Spencer B, Masliah E, Lee SJ (2009) Inclusion formation and neuronal cell death through neuron-to-neuron transmission of alpha-synuclein. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 13010-13015

Dexter PM, Caldwell KA, Caldwell GA (2012) A predictable worm: application of *Caenorhabditis elegans* for mechanistic investigation of movement disorders. *Neurotherapeutics : the journal of the American Society for Experimental Neurotherapeutics* **9**: 393-404

Dominguez-Meijide A, Vasili E, König A, Cima-Omori MS, Ibáñez de Opakua A, Leonov A, Ryazanov S, Zweckstetter M, Griesinger C, Outeiro TF (2020) Effects of pharmacological modulators of  $\alpha$ -synuclein and tau aggregation and internalization. *Scientific reports* **10**: 12827

Du Y, Ma Z, Lin S, Dodel RC, Gao F, Bales KR, Triarhou LC, Chernet E, Perry KW, Nelson DL, Luecke S, Phebus LA, Bymaster FP, Paul SM (2001) Minocycline prevents nigrostriatal dopaminergic neurodegeneration in the MPTP model of Parkinson's disease. *Proceedings of the National Academy of Sciences of the United States of America* **98**: 14669-14674

Engelender S, Kaminsky Z, Guo X, Sharp AH, Amaravi RK, Kleiderlein JJ, Margolis RL, Troncoso JC, Lanahan AA, Worley PF, Dawson VL, Dawson TM, Ross CA (1999) Synphilin-1 associates

with alpha-synuclein and promotes the formation of cytosolic inclusions. *Nature genetics* **22**: 110-114

Faravelli G, Raimondi S, Marchese L, Partridge FA, Soria C, Mangione P, Canetti D, Perni M, Aprile FA, Zorzoli I, di Schiavi E, Lomas DA, Bellotti V, Sattelle DB, Giorgetti S (2019) C. elegans expressing D76N  $\beta$ (2)-microglobulin: a model for in vivo screening of drug candidates targeting amyloidosis. *Scientific reports* **9**: 19960

Gaeta AL, Caldwell KA, Caldwell GA (2019) Found in Translation: The Utility of C. elegans Alpha-Synuclein Models of Parkinson's Disease. *Brain sciences* **9**

Garcia-Martinez EM, Sanz-Blasco S, Karachitos A, Bandez MJ, Fernandez-Gomez FJ, Perez-Alvarez S, de Mera RM, Jordan MJ, Aguirre N, Galindo MF, Villalobos C, Navarro A, Kmita H, Jordán J (2010) Mitochondria and calcium flux as targets of neuroprotection caused by minocycline in cerebellar granule cells. *Biochemical pharmacology* **79**: 239-250

Giorgetti S, Raimondi S, Pagano K, Relini A, Bucciantini M, Corazza A, Fogolari F, Codutti L, Salmona M, Mangione P, Colombo L, De Luigi A, Porcari R, Gliozzi A, Stefani M, Esposito G, Bellotti V, Stoppini M (2011) Effect of tetracyclines on the dynamics of formation and destructure of beta2-microglobulin amyloid fibrils. *The Journal of biological chemistry* **286**: 2121-2131

González-Lizárraga F, Socías SB, Ávila CL, Torres-Bugeau CM, Barbosa LR, Binolfi A, Sepúlveda-Díaz JE, Del-Bel E, Fernandez CO, Papy-Garcia D, Itri R, Raisman-Vozari R, Chehín RN (2017) Repurposing doxycycline for synucleinopathies: remodelling of  $\alpha$ -synuclein oligomers towards non-toxic parallel beta-sheet structured species. *Scientific reports* **7**: 41755

Grassi D, Howard S, Zhou M, Diaz Perez N, Urban NT, Guerrero-Given D, Kamasawa N, Volpicelli-Daley LA, LoGrasso P, Lasmezas CI (2018) Identification of a highly neurotoxic  $\alpha$ -synuclein species inducing mitochondrial damage and mitophagy in Parkinson's disease. *Proceedings of the National Academy of Sciences of the United States of America* **115**: E2634-e2643

Hamamichi S, Rivas RN, Knight AL, Cao S, Caldwell KA, Caldwell GA (2008) Hypothesis-based RNAi screening identifies neuroprotective genes in a Parkinson's disease model. *Proceedings of the National Academy of Sciences of the United States of America* **105**: 728-733

Hannan SB, Dräger NM, Rasse TM, Voigt A, Jahn TR (2016) Cellular and molecular modifier pathways in tauopathies: the big picture from screening invertebrate models. *Journal of neurochemistry* **137**: 12-25

Hannaoui S, Gougerot A, Privat N, Levavasseur E, Bizat N, Hauw JJ, Brandel JP, Haïk S (2014) Cyclin efficacy on the propagation of human prions in primary cultured neurons is strain-specific. *The Journal of infectious diseases* **209**: 1144-1148

Harding TC, Geddes BJ, Noel JD, Murphy D, Uney JB (1997) Tetracycline-regulated transgene expression in hippocampal neurones following transfection with adenoviral vectors. *Journal of neurochemistry* **69**: 2620-2623

Heemskerk J, Tobin AJ, Bain LJ (2002) Teaching old drugs new tricks. Meeting of the Neurodegeneration Drug Screening Consortium, 7-8 April 2002, Washington, DC, USA. *Trends in neurosciences* **25**: 494-496

Hengartner D, Fernandez HH (2019) The next chapter in symptomatic Parkinson disease treatments. *Parkinsonism & related disorders* **59**: 39-48

Jang H, Kim S, Lee JM, Oh YS, Park SM, Kim SR (2017) Montelukast treatment protects nigral dopaminergic neurons against microglial activation in the 6-hydroxydopamine mouse model of Parkinson's disease. *Neuroreport* **28**: 242-249

Krum BN, Martins AC, Jr., Queirós L, Ferrer B, Milne GL, Soares FAA, Fachinetto R, Aschner M (2020) Haloperidol Interactions with the dop-3 Receptor in *Caenorhabditis elegans*. *Molecular neurobiology*

Kuwahara T, Koyama A, Gengyo-Ando K, Masuda M, Kowa H, Tsunoda M, Mitani S, Iwatsubo T (2006) Familial Parkinson mutant alpha-synuclein causes dopamine neuron dysfunction in transgenic *Caenorhabditis elegans*. *The Journal of biological chemistry* **281**: 334-340

Kuwahara T, Koyama A, Koyama S, Yoshida S, Ren CH, Kato T, Mitani S, Iwatsubo T (2008) A systematic RNAi screen reveals involvement of endocytic pathway in neuronal dysfunction in alpha-synuclein transgenic *C. elegans*. *Human molecular genetics* **17**: 2997-3009

Lakso M, Vartiainen S, Miettinen AM, Sirviö J, Thomas JH, Nass R, Blakely RD, Wong G (2003) Dopaminergic neuronal loss and motor deficits in *Caenorhabditis elegans* overexpressing human alpha-synuclein. *Journal of neurochemistry* **86**: 165-172

Lázaro DF, Dias MC, Carija A, Navarro S, Madaleno CS, Tenreiro S, Ventura S, Outeiro TF (2016) The effects of the novel A53E alpha-synuclein mutation on its oligomerization and aggregation. *Acta Neuropathologica Communications* **4**:128

Lázaro DF, Rodrigues EF, Langohr R, Shahpasandzadeh H, Ribeiro T, Guerreiro P, Gerhardt E, Kröhnert K, Klucken J, Pereira MD, Popova B, Kruse N, Mollenhauer B, Rizzoli SO, Braus GH, Danzer KM, Outeiro TF (2014) Systematic comparison of the effects of alpha-synuclein mutations on its oligomerization and aggregation. *PLoS genetics* **10**: e1004741

Lazzarini M, Martin S, Mitkovski M, Vozari RR, Stühmer W, Bel ED (2013) Doxycycline restrains glia and confers neuroprotection in a 6-OHDA Parkinson model. *Glia* **61**: 1084-1100

Letizia MC, Cornaglia M, Trouillon R, Sorrentino V, Mouchiroud L, Bou Sleiman MS, Auwerx J, Gijs MAM (2018) Microfluidics-enabled phenotyping of a whole population of *C. elegans* worms over their embryonic and post-embryonic development at single-organism resolution. *Microsystems & nanoengineering* **4**: 6

LeVine H, 3rd (1993) Thioflavine T interaction with synthetic Alzheimer's disease beta-amyloid peptides: detection of amyloid aggregation in solution. *Protein science : a publication of the Protein Society* **2**: 404-410

Liang B, Tamm LK (2018) Solution NMR of SNAREs, complexin and  $\alpha$ -synuclein in association with membrane-mimetics. *Progress in nuclear magnetic resonance spectroscopy* **105**: 41-53

Lin MT, Beal MF (2006) Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* **443**: 787-795

Lindström V, Gustafsson G, Sanders LH, Howlett EH, Sigvardsson J, Kasrayan A, Ingelsson M, Bergström J, Erlandsson A (2017) Extensive uptake of  $\alpha$ -synuclein oligomers in astrocytes results in sustained intracellular deposits and mitochondrial damage. *Molecular and cellular neurosciences* **82**: 143-156

Lints R, Emmons SW (1999) Patterning of dopaminergic neurotransmitter identity among *Caenorhabditis elegans* ray sensory neurons by a TGFbeta family signaling pathway and a Hox gene. *Development (Cambridge, England)* **127**: 5819-5831

Luk KC, Song C, O'Brien P, Stieber A, Brundin JR, Brunden KR, Trojanowski JQ, Lee VMY (2009) Exogenous alpha-synuclein fibrils seed the formation of Lewy body-like intracellular inclusions in cultured cells. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 20051-20056

Masaracchia C, Hnida M, Bernardt E, Lopes da Fonseca T, Villar-Pique A, Branco T, Stahlberg MA, Dean C, Fernández CO, Milosevic I, Outeiro TF (2018) Membrane binding, internalization, and sorting of alpha-synuclein in the cell. *Acta neuropathologica communications* **6**: 79

Masaracchia C, König A, Valiente-Gabioud AA, Peralta P, Favretto F, Strohäker T, Lázaro DF, Zweckstetter M, Fernandez CO, Outeiro TF (2020) Molecular characterization of an aggregation-prone variant of alpha-synuclein used to model synucleinopathies. *Biochimica et biophysica acta Proteins and proteomics* **1868**: 140298

McLean PJ, Kawamata H, Hyman BT (2001) Alpha-synuclein-enhanced green fluorescent protein fusion proteins form proteasome sensitive inclusions in primary neurons. *Neuroscience* **104**: 901-912

Mehra S, Sahay S, Maji SK (2019)  $\alpha$ -Synuclein misfolding and aggregation: Implications in Parkinson's disease pathogenesis. *Biochimica et biophysica acta Proteins and proteomics* **1867**: 890-908

Mello C, Fire A (1995) DNA transformation. *Methods in cell biology* **48**: 451-482

Moghaddam HF, Hemmati A, Nazari Z, Mehrab H, Abid KM, Ardestani MS (2007) Effects of aspirin and celecoxib on rigidity in a rat model of Parkinson's disease. *Pakistan journal of biological sciences : PJBS* **10**: 3853-3858

Molenaars M, Janssens GE, Williams EG, Jongejan A, Lan J, Rabot S, Joly F, Moerland PD, Schomakers BV, Lezzerini M, Liu YJ, McCormick MA, Kennedy BK, van Weeghel M, van Kampen AHC, Aebersold R, MacInnes AW, Houtkooper RH (2020) A Conserved Mito-Cytosolic Translational Balance Links Two Longevity Pathways. *Cell metabolism* **31**: 549-563.e547

Murphy DD, Rueter SM, Trojanowski JQ, Lee VM (2000) Synucleins are developmentally expressed, and alpha-synuclein regulates the size of the presynaptic vesicular pool in primary hippocampal neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **20**: 3214-3220

Nelson ML, Levy SB (2011) The history of the tetracyclines. *Annals of the New York Academy of Sciences* **1241**: 17-32

Nussbaum-Krammer CI, Neto MF, Brielmann RM, Pedersen JS, Morimoto RI (2015) Investigating the spreading and toxicity of prion-like proteins using the metazoan model organism *C. elegans*. *Journal of visualized experiments : JoVE*: 52321

Olsson TT, Klementieva O, Gouras GK (2018) Prion-like seeding and nucleation of intracellular amyloid- $\beta$ . *Neurobiology of disease* **113**: 1-10

Outeiro TF, Koss DJ, Erskine D, Walker L, Kurzawa-Akanbi M, Burn D, Donaghy P, Morris C, Taylor JP, Thomas A, Attwood J, McKeith I (2019) Dementia with Lewy bodies: an update and outlook. *Molecular neurodegeneration* **14**: 5

Outeiro TF, Putcha P, Tetzlaff JE, Spoelgen R, Koker M, Carvalho F, Hyman BT, McLean PJ (2008) Formation of toxic oligomeric alpha-synuclein species in living cells. *PloS one* **3**: e1867

Paldino E, Balducci C, La Vitola P, Artioli L, D'Angelo V, Giampà C, Artuso V, Forloni G, Fusco FR (2020) Neuroprotective Effects of Doxycycline in the R6/2 Mouse Model of Huntington's Disease. *Molecular neurobiology* **57**: 1889-1903

Park JH, Burgess JD, Farooqi AH, DeMeo NN, Fiesel FC, Springer W, Delenclos M, McLean PJ (2020) Alpha-synuclein-induced mitochondrial dysfunction is mediated via a sirtuin 3-dependent pathway. *Molecular neurodegeneration* **15**: 5

Patterson JR, Polinski NK, Duffy MF, Kemp CJ, Luk KC, Volpicelli-Daley LA, Kanaan NM, Sortwell CE (2019) Generation of Alpha-Synuclein Preformed Fibrils from Monomers and Use In Vivo. *Journal of visualized experiments : JoVE*

Peng C, Trojanowski JQ, Lee VM (2020) Protein transmission in neurodegenerative disease. *Nature reviews Neurology* **16**: 199-212

Perni M, Galvagnion C, Maltsev A, Meisl G, Müller MB, Challa PK, Kirkegaard JB, Flagmeier P, Cohen SI, Cascella R, Chen SW, Limbocker R, Sormanni P, Heller GT, Aprile FA, Cremades N, Cecchi C, Chiti F, Nollen EA, Knowles TP, Vendruscolo M, Bax A, Zaslhoff M, Dobson CM (2017) A natural product inhibits the initiation of  $\alpha$ -synuclein aggregation and suppresses its toxicity. *Proceedings of the National Academy of Sciences of the United States of America* **114**: E1009-e1017

Perrin RJ, Payton JE, Barnett DH, Wraight CL, Woods WS, Ye J, George JM (2003) Epitope mapping and specificity of the anti-alpha-synuclein monoclonal antibody Syn-1 in mouse brain and cultured cell lines. *Neuroscience letters* **349**: 133-135

Picca A, Calvani R, Coelho-Junior HJ, Landi F, Bernabei R, Marzetti E (2020) Mitochondrial Dysfunction, Oxidative Stress, and Neuroinflammation: Intertwined Roads to Neurodegeneration. *Antioxidants (Basel, Switzerland)* **9**

Raimondi S, Porcari R, Mangione PP, Verona G, Marcoux J, Giorgetti S, Taylor GW, Ellmerich S, Ballico M, Zanini S, Pardon E, Al-Shawi R, Simons JP, Corazza A, Fogolari F, Leri M, Stefani M, Bucciantini M, Gillmore JD, Hawkins PR, Valli M, Stoppini M, Robinson CV, Steyaert J, Esposito G, Bellotti V (2017) A specific nanobody prevents amyloidogenesis of D76N  $\beta$ (2)-microglobulin in vitro and modifies its tissue distribution in vivo. *Scientific reports* **7**: 46711

Rand JB, Nonet ML (1997) Synaptic transmission. In *C. elegans II*, Riddle DL, Blumenthal T, Meyer BJ, Priess JR (eds). Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press

Copyright © 1997, Cold Spring Harbor Laboratory Press.

Refai O, Blakely RD (2019) Blockade and reversal of swimming-induced paralysis in *C. elegans* by the antipsychotic and D2-type dopamine receptor antagonist azaperone. *Neurochemistry international* **123**: 59-68

Reglodi D, Renaud J, Tamas A, Tizabi Y, Socías SB, Del-Bel E, Raisman-Vozari R (2017) Novel tactics for neuroprotection in Parkinson's disease: Role of antibiotics, polyphenols and neuropeptides. *Progress in neurobiology* **155**: 120-148

Rodriguez-Perez AI, Sucunza D, Pedrosa MA, Garrido-Gil P, Kulisevsky J, Lanciego JL, Labandeira-Garcia JL (2018) Angiotensin Type 1 Receptor Antagonists Protect Against Alpha-Synuclein-Induced Neuroinflammation and Dopaminergic Neuron Death. *Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics* **15**: 1063-1081



Santa-Cecília FV, Leite CA, Del-Bel E, Raisman-Vozari R (2019) The Neuroprotective Effect of Doxycycline on Neurodegenerative Diseases. *Neurotoxicity research* **35**: 981-986

Santa-Cecília FV, Socias B, Ouidja MO, Sepulveda-Diaz JE, Acuña L, Silva RL, Michel PP, Del-Bel E, Cunha TM, Raisman-Vozari R (2016) Doxycycline Suppresses Microglial Activation by Inhibiting the p38 MAPK and NF- $\kappa$ B Signaling Pathways. *Neurotoxicity research* **29**: 447-459

Santos PI, Outeiro TF (2020) Protein trapping leads to altered synaptic proteostasis in synucleinopathies. *The FEBS journal*

Schmitz M, Cramm M, Llorens F, Candelise N, Müller-Cramm D, Vargas D, Schulz-Schaeffer WJ, Zafar S, Zerr I (2016) Application of an in vitro-amplification assay as a novel pre-screening test for compounds inhibiting the aggregation of prion protein scrapie. *Scientific reports* **6**: 28711

Smith WW, Margolis RL, Li X, Troncoso JC, Lee MK, Dawson VL, Dawson TM, Iwatsubo T, Ross CA (2005) Alpha-synuclein phosphorylation enhances eosinophilic cytoplasmic inclusion formation in SH-SY5Y *Journal of neuroscience* **25**:5544-5552

Socias SB, González-Lizárraga F, Avila CL, Vera C, Acuña L, Sepulveda-Diaz JE, Del-Bel E, Raisman-Vozari R, Chehin RN (2018) Exploring the therapeutic potential of ready-to-use drugs: Repurposing antibiotics against amyloid aggregation in neurodegenerative diseases. *Progress in neurobiology* **162**: 17-36

Spillantini MG, Crowther RA, Jakes R, Hasegawa M, Goedert M (1998) alpha-Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with lewy bodies. *Proceedings of the National Academy of Sciences of the United States of America* **95**: 6469-6473

Stiernagle T (2006) Maintenance of *C. elegans*. *WormBook : the online review of C elegans biology*: 1-11

Stoilova T, Colombo L, Forloni G, Tagliavini F, Salmona M (2013) A new face for old antibiotics: tetracyclines in treatment of amyloidoses. *Journal of medicinal chemistry* **56**: 5987-6006

Sulston J, Dew M, Brenner S (1975) Dopaminergic neurons in the nematode *Caenorhabditis elegans*. *The Journal of comparative neurology* **163**: 215-226

Szegő É M, Dominguez-Mejide A, Gerhardt E, König A, Koss DJ, Li W, Pinho R, Fahlbusch C, Johnson M, Santos P, Villar-Piqué A, Thom T, Rizzoli S, Schmitz M, Li J, Zerr I, Attems J, Jahn O, Outeiro TF (2019) Cytosolic Trapping of a Mitochondrial Heat Shock Protein Is an Early Pathological Event in Synucleinopathies. *Cell reports* **28**: 65-77.e66

Takahashi M, Ko LW, Kulathingal J, Jiang P, Sevlever D, Yen SH (2007) Oxidative stress-induced phosphorylation, degradation and aggregation of alpha-synuclein are linked to upregulated CK2 and cathepsin D. *The European journal of neuroscience* **26**: 863-874

Tysnes OB, Storstein A (2017) Epidemiology of Parkinson's disease. *Journal of neural transmission (Vienna, Austria : 1996)* **124**: 901-905

Tyson T, Senchuk M, Cooper JF, George S, Van Raamsdonk JM, Brundin P (2017) Novel animal model defines genetic contributions for neuron-to-neuron transfer of alpha-synuclein. *Sci Rep* **7**: 7506

van Ham TJ, Thijssen KL, Breitling R, Hofstra RM, Plasterk RH, Nollen EA (2008) C. elegans model identifies genetic modifiers of alpha-synuclein inclusion formation during aging. *PLoS genetics* **4**: e1000027

Vasquez V, Mitra J, Perry G, Rao KS, Hegde ML (2018) An Inducible Alpha-Synuclein Expressing Neuronal Cell Line Model for Parkinson's Disease1. *Journal of Alzheimer's disease : JAD* **66**: 453-460

Vekrellis K, Xilouri M, Emmanouilidou E, Stefanis L (2009) Inducible over-expression of wild type alpha-synuclein in human neuronal cells leads to caspase-dependent non-apoptotic death. *Journal of neurochemistry* **109**: 1318-1362

Vidal-Gadea A, Topper S, Young L, Crisp A, Kressin L, Elbel E, Maples T, Brauner M, Erbguth K, Axelrod A, Gottschalk A, Siegel D, Pierre-Shimomura JT (2011) Caenorhabditis elegans selects distinct crawling and swimming traits via dopamine and serotonin. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 17504-17509

Volpicelli-Daley LA, Luk KC, Lee VM (2014) Addition of exogenous  $\alpha$ -synuclein preformed fibrils to primary neuronal cultures to seed recruitment of endogenous  $\alpha$ -synuclein to Lewy body and Lewy neurite-like aggregates. *Nature protocols* **9**: 2135-2146

Wagner J, Ryazanov S, Leonov A, Levin J, Shi S, Schmidt F, Prix C, Pan-Montojo F, Bertsch U, Mitteregger-Kretschmar G, Geissen M, Eiden M, Leidel F, Hirschberger T, Deeg AA, Krauth JJ, Zinth W, Tavan P, Pilger J, Zweckstetter M, Frank T, Bähr M, Weishaupt JH, Uhr M, Urlaub H, Teichmann U, Samwer M, Bötzel K, Groschup M, Kretschmar H, Griesinger C, Giese A (2013) Anle138b: a novel oligomer modulator for disease-modifying therapy of neurodegenerative diseases such as prion and Parkinson's disease. *Acta neuropathologica* **125**: 795-813

Wang X, Becker K, Levine N, Zhang M, Lieberman AP, Moore DJ, Ma J (2019) Pathogenic alpha-synuclein aggregates preferentially bind to mitochondria and affect cellular respiration. *Acta neuropathologica communications* **7**: 41

Wang X, Michaelis EK (2010) Selective neuronal vulnerability to oxidative stress in the brain. *Frontiers in aging neuroscience* **2**: 12

White JG, Southgate E, Thomson JN, Brenner S (1986) The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philosophical transactions of the Royal Society of London Series B, Biological sciences* **314**: 1-340

Wilham JM, Orrú CD, Bessen RA, Atarashi R, Sano K, Race B, Meade-White KD, Taubner LM, Timmes A, Caughey B (2010) Rapid end-point quantitation of prion seeding activity with sensitivity comparable to bioassays. *PLoS pathogens* **6**: e1001217


Wilhelm BG, Mandad S, Truckenbrodt S, Kröhnert K, Schäfer C, Rimmner B, Koo SJ, Claßen GA, Krauss M, Haucke V, Urlaub H, Rizzoli SO (2014) Composition of isolated synaptic boutons reveals the amounts of vesicle trafficking proteins. *Science (New York, NY)* **344**: 1023-1028

Winslow AR, Chen CW, Corrochano S, Acevedo-Arozena A, Gordon DE, Peden AA, Lichtenberg M, Menzies FM, Ravikumar B, Imarisio S, Brown S, O'Kane CJ, Rubinsztein DC (2010)  $\alpha$ -Synuclein impairs macroautophagy: implications for Parkinson's disease. *The Journal of cell biology* **190**: 1023-1037

Wu C, Biancalana M, Koide S, Shea JE (2005) Binding modes of thioflavin-T to the single-layer beta-sheet of the peptide self-assembly mimics. *Journal of molecular biology* **394**: 627-633


Zhu Z, Zheng T, Lee CG, Homer RJ, Elia JA (2002) Tetracycline-controlled transcriptional regulation systems: advances and application in transgenic animal modeling. *Seminars in cell & developmental biology* **13**: 121-128

## Appendices



**Figure A.1. The presence of doxycycline leads to changes in average molecular weight and average particle size of RT-QuIC products.** **A.** Average molecular weight of the lower weight species measured by DLS. There are no statistically significant differences amongst groups. **B.** Average particle size of the RT-QuIC products measured by DLS. Treatment with 10  $\mu$ M Doxycycline leads to a significantly higher average particle size. **C.** Average particle size of the lower weight species measured by DLS. The presence of 1  $\mu$ M doxycycline during RT-QuIC leads to a statistically significant decrease in average particle size in comparison with untreated RT-QuIC products before and after a posterior sonication. **D.** DLS measurements of the samples obtained in the RT-QuIC. Above equimolar concentrations higher molecular weight species lead to higher times for the decrease of the intensity correlation. N = 3 in all

experiments \* $p < 0.05$  in comparison with 0  $\mu\text{M}$  Doxycycline.  $\$p < 0.05$  in comparison with 1  $\mu\text{M}$  Doxycycline.  $\$p < 0.05$  in comparison with 10  $\mu\text{M}$  Doxycycline.



**Figure A.2. Treatment with doxycycline leads to an increase in the average number of inclusions per cell.** **A.** Immunostaining for aSyn and Synphilin-1-V5 shows presence of both proteins in inclusions. **B.** Average number of inclusions per cell. Treatment with different concentrations of doxycycline leads to a significantly different average number of inclusions per cell. Data presented as mean  $\pm$  SD.  $N = 3$  in all experiments \*  $p < 0.05$  **C.** Total inclusion area (in  $\mu\text{m}^2$ ) of control cells and cells treated with 10  $\mu\text{M}$  doxycycline. Treatment with 10  $\mu\text{M}$  of doxycycline lead to a significantly different overall inclusion area in cells. Data presented as mean  $\pm$  SD.  $N = 3$  in all experiments \*  $p < 0.05$ . **D.** Images of a cell transfected with

VNSynph/aSynVC and treated with 10  $\mu$ M doxycycline at different time points. An apparent increase in the number of inclusions accompanied by a decrease in their size can be seen.

Journal Pre-proof



**Credit author statement**

**Antonio Dominguez-Meijide:** Investigation, visualization, formal analysis, writing – original draft. **Valeria Parrales:** Investigation, visualization, methodology, formal analysis, writing – original draft. **Eftychia Vasili:** Investigation, visualization, formal analysis, writing – original draft. **Florencia González-Lizárraga:** Investigation, visualization, formal analysis, writing – original draft. **Annekatriin König:** Resources. **Diana F. Lázaro:** Resources, methodology. **Annie Lannuzel:** Formal analysis, writing – review and editing. **Stéphane Haik:** Writing – review and editing. **Elaine Del Bel:** Conceptualization, formal analysis, funding acquisition, writing – review and editing. **Rosana Chehín:** Conceptualization, funding acquisition, writing – review and editing. **Rita Raisman-Vozari:** Conceptualization, funding acquisition, writing – review and editing. **Patrick P. Michel:** Conceptualization, funding acquisition, writing – review and editing. **Nicolas Bizat:** Conceptualization, funding acquisition, writing – review and editing. **Tiago F. Outeiro:** Conceptualization, funding acquisition, writing – review and editing.

### Highlights

- Repurposing of drugs may be a valuable strategy to help treat Parkinson's disease.
- Doxycycline decreases the number and size of alpha-synuclein inclusions in cells.
- Doxycycline lowers the production of mitochondrial-derived reactive oxygen species.
- Doxycycline induces redistribution of alpha-synuclein aggregates in *C.elegans*.