

SOCIOECONOMIC INEQUALITIES IN SELF-REPORTED HEALTH AND PHYSICAL FUNCTIONING IN ARGENTINA: FINDINGS FROM THE NATIONAL SURVEY ON QUALITY OF LIFE OF OLDER ADULTS 2012 (ENCaViAM)

SANTIAGO RODRÍGUEZ LÓPEZ*¹, SONIA E. COLANTONIO*†
AND DORA E. CELTON*†

**Research and Study Centre on Culture and Society, National Scientific and
Technical Research Council (CIECS-CONICET-UNC), Córdoba, Argentina and
†National University of Córdoba (UNC), Argentina*

Summary. This study aimed to evaluate educational and income inequalities in self-reported health (SRH), and physical functioning (limitations in Activities of Daily Living (ADL)/Instrumental Activities of Daily Living (IADL)), among 60-year-old and older adults in Argentina. Using cross-sectional data from the Argentinian National Survey on Quality of Life of Older Adults 2012 (Encuesta Nacional sobre Calidad de Vida de Adultos Mayores, ENCaViAM), gender-specific socioeconomic inequalities in SRH and ADL and IADL limitations were studied in relation to educational level and household *per capita* income. The Relative Index of Inequality (RII) – an index of the relative size of socioeconomic inequalities in health – was used. Socioeconomic inequalities in the studied health indicators were found – except for limitations in ADL among women – favouring socially advantaged groups. The results remained largely significant after full adjustment, suggesting that educational and income inequalities, mainly in SRH and IADL, were robust and somehow independent of age, marital status, physical activity, the use of several medications, depression and the occurrence of falls. The findings add to the existing knowledge on the relative size of the socioeconomic inequalities in subjective health indicators among Argentinian older adults, which are to the detriment of lower socioeconomic groups. The results could be used to inform planning interventions aimed at decreasing socioeconomic inequalities in health, to the benefit of socially disadvantaged adults.

¹ Corresponding author. Email: santiago.rodriguez@conicet.gov.ar

Introduction

The persistence of socioeconomic inequalities in health, even in the highly developed welfare states of Western Europe, is one of the greatest failures of public health (Mackenbach, 2012). A growing body of evidence provides a clear indication of the adverse effects of socioeconomic inequality on individuals' morbidity and mortality (Brandt *et al.*, 2012). It has been suggested that high levels of social inequality have a direct and negative causal effect on the health of populations (Wagstaff & van Doorslaer, 2000; Lynch *et al.*, 2004). Although this is highly supported by the literature, it has been argued that some other factors, such as the time of exposure to those inequalities, may play a key role, particularly among older adults (Vries *et al.*, 2014).

A recent review suggested that wide income differences play a causal role leading to worse health, especially among individuals of lower socioeconomic status (Pickett & Wilkinson, 2015). Nevertheless, high levels of inequality negatively affect not only the health of the most disadvantaged groups but also the health of the affluent. This is so, mainly because inequality reduces social cohesion or social capital, which leads to more stress, fear and insecurity for everyone (Kawachi & Kennedy, 1999). Therefore, health at the individual level may not respond simply to, for example, absolute income, but also to relativities in society (Wagstaff & van Doorslaer, 2000). Thus, it has been shown that in states with high levels of income inequality, low-income women are 80% more likely to report fair or poor health than low-income women in low-income-inequality states (Kahn *et al.*, 2000).

As in high-income Western Europe countries, the wide socioeconomic inequalities in Latin American middle-income countries may represent a determinant social factor of health (Reygadas, 2006; Cardona *et al.*, 2013). Actually, socioeconomic inequalities are more prominent in most middle- and low-income countries, and life expectancy has increased much faster in these countries than in richer ones (Palloni *et al.*, 2002; Lima-Costa *et al.*, 2012). In particular, Argentina has had one of the most disappointing social performances in Latin America in the last three decades (Gasparini, 2007). The year 2001, which was characterized by serious economic and social crisis in the country, displayed the highest educational inequalities in mortality in comparison to either 1991 or 2010 (Manzelli, 2014). However, the high economic growth experienced by the region during the past decade has had a positive impact on Argentinian social indicators (Maurizio, 2014). Thus, and in spite of the fact that the magnitude of educational differences has been much higher in Argentina than in other Latin American countries (Manzelli, 2014), educational inequalities have declined in almost all metropolitan areas in the past years (Morales & Paz Terán, 2010). In a recent ecological study, Argentina showed the second most favourable health status in the region (Cardona *et al.*, 2013). Therefore, while the social situation has substantially improved in recent years, poverty and inequality are still important (Gasparini, 2007). Furthermore, the prevalence of different cardiovascular risk factors has increased, mainly in individuals of lower socioeconomic status (Ferrante *et al.*, 2011).

In this situation, addressing current differences in health among individuals of different socioeconomic status has become especially important. Health inequalities have been estimated in Argentina in terms of both education and income (De Maio, 2007, 2008; Alzraqui *et al.*, 2009; Manzelli, 2014). These studies have shown patterns of

socioeconomic inequalities in morbidity favouring socially advantaged groups (Hoffman & Centeno, 2003; Cardona *et al.*, 2013). Nevertheless, the demographic transition in most Latin American countries is generating populations with unprecedented numbers of older adults exposed to significant social inequalities (Lima-Costa *et al.*, 2012). However, few studies have addressed health inequalities among Argentinian older adults, considering their social and biological vulnerability.

A gender perspective is important when studying socioeconomic inequalities in health. It has been argued that i) gender is an influential factor that contributes to the prevalence of poor self-reported health during the ageing process (WHO, 2002), ii) differences in health status and perception are linked to gender inequalities in the workplace and in economic and personal autonomy (Montero López *et al.*, 2011). Gender differences in subjective health have been reported previously for Argentina (Alazraqui 2005), and it has been suggested that multi-morbidity explains part of these disparities (Assari & Moghani Lankarani, 2015). Additionally, ways in which social determinants contribute to a poorer health status in women compared with men vary among countries (Hosseinpoor *et al.*, 2012). Thus, it is relevant to examine whether the patterns of association between education/income and subjective health indicators vary for older men and women in Argentina.

Within this context, the present study aimed to analyse educational and income inequalities – by means of an inequality index – in three subjective health indicators based on a representative sample of Argentinian older adults. Addressing health inequalities in Argentina is important due to the high proportion of older, particularly socially disadvantaged adults. These are, largely, more likely to benefit from both social policies leading to improvements in the welfare state (Qin & Liu, 2013) and reduction in socioeconomic inequalities (Stuckler *et al.*, 2009).

Methods

Sample

Data were obtained from the Argentinian National Survey on Quality of Life of Older Adults 2012 (Encuesta Nacional sobre Calidad de Vida de Adultos Mayores, ENCaViAM) (INDEC, 2014). ENCaViAM is the first cross-sectional national survey on adults' quality of life to include a representative subsample of older adults living in households located in urban areas of Argentina. Data collection was carried out during 2012 by the National Institute of Statistics and Census (INDEC) together with the Regional Directions of Statistics from Argentina.

ENCaViAM was mainly developed to evaluate the most urgent needs of public policy application among Argentinian older adults. The survey gathered valuable information on access to health services and medicines, self-reported health, life satisfaction and limitations in physical activities of daily living, among other things (INDEC, 2014). The studied sample included 4654 60-year-old and older individuals (mean age 70.5 years; 57.4% women) with complete data. Since ENCaViAM is a probabilistic sample, every primary sampling unit (household) has an expansion factor (calibrated weight), which is the inverse of the selection probability. Consequently, all analyses were performed using calibrated weights. Finally, the non-response rate was

relatively low (13.4% of selected individuals). Further methodological aspects of ENCaViAM are described elsewhere (INDEC, 2014).

Variables

Socioeconomic status. Two individual indicators of socioeconomic status were considered to summarize adults' socioeconomic status: educational level and family *per capita* income.

Educational level was categorized as: no formal education, primary school, high school and university/higher education. These categories were equivalent and comparable with the ones proposed by the International Standard Classification of Education (ISCED) (UNESCO, 2006) as follows: 1 = no formal education; 2 = primary school (Level 1 ISCED); 3 = high school (Level 2 and 3 ISCED); and 4 = university/higher education (Levels 5 and 6 ISCED).

For family *per capita* income decile groups were created and transformed into quintiles for analytical purposes: 1st (poorest), 2nd, 3rd, 4th and 5th (richest).

Health indicators. Subjective health predicts mortality and a variety of other health outcomes beyond objective health (Franz *et al.*, 2016). Two different subjective health indicators were evaluated: general self-reported health status (SRH) and self-assessments of physical functioning. Both are valid and useful indicators for measuring general physical well-being and population health (Lundberg & Manderbacka, 1996; Miilunpalo *et al.*, 1997; Lantz *et al.*, 2001). While SRH is a static indicator capturing deviations from a norm perceived as healthy (Ziebarth, 2010), limitations in physical activities of daily living is a functional one, capturing the inability to perform certain tasks. Overall, the use of dichotomized self-reported health measures is a standard approach in the literature on health inequalities (Ziebarth & Frick, 2010) given its simplicity and wide use of logistic regression, particularly in epidemiology (Manor *et al.*, 2000). Additionally, the use of self-reports can provide information about individuals' own perceptions regarding their health that cannot be measured using an objective assessment tool (CEPAL, 2008).

Self-reported health (SRH) is a subjective indicator that has proved to be highly correlated with objective measures of physical health, as well as a good predictor of functional decline (Lee, 2000) and mortality (Idler & Benyamini, 1997; Gasparini, 2007; Haas, 2007). Originally, it was assessed using a 5-point scale: excellent, very good, good, fair and poor. However, for analytical purposes and following conventional practice, it was recoded as a binary outcome with 1 = fair/poor, and 0 = excellent/very good/good (De Maio *et al.*, 2012). Although dichotomization of the self-rated health variable involves a loss of information and has other drawbacks (Ziebarth, 2010), only small reductions in power and statistical efficiency have been reported (Manor *et al.*, 2000).

Physical functioning. Activities of Daily Living (ADL) are defined as those activities essential for independent living, while Instrumental Activities of Daily Living (IADL) are more complex and require a higher level of personal autonomy. IADL refer to tasks implying greater interaction with the environment and enough capacity to make

decisions (WHO, 2001). Based on these differences, deficits in IADL normally precede deficits in ADL (Judge *et al.*, 1996). Limitations in ADL (or IADL) are defined as having difficulty with and/or receiving help from another person or being unable to do one or more of the following tasks: dressing, walking across the home, bathing or showering, combing oneself/brushing teeth, eating/cutting up food, getting in or out of bed, using the toilet (including getting up/down) and climbing/climbing down one flight of stairs. Similarly, IADL include seven activities: difficulties in using public transport, preparing a hot meal, shopping for groceries, making telephone calls, taking medications, doing work around the house/garden and managing money/paying bills/keeping track of expenses. Consequently, disabilities in ADL and in IADL were defined as: 1 = one or more limitations in ADL/IADL, and 0 = without any limitations.

Other variables. In addition to age, other factors such as marital status, the manifestation of depression within the past year (Lorant *et al.*, 2003), the occurrence of falls within the past two years, level of physical activity (Vagetti *et al.*, 2014) and the use of eight or more medications per day might be influencing socioeconomic inequalities in health. Including these variables in the analysis is important in order to evaluate the robustness of the associations of interest.

Statistical analysis

The prevalence of adults with fair/poor SRH and ADL and IADL limitations was analysed by gender. The Relative Index of Inequality (RII) was estimated in order to measure the magnitude of educational and income inequalities in the three health outcomes (SRH, ADL, IADL). The RII is a regression-based measure that takes the complete socioeconomic distribution into account, rather than comparing only the two extreme groups (Mackenbach & Kunst, 1997). The RII resembles relative risk in that it compares the health of the extremes of the social distribution, but it is estimated using data from all social categories (Singh-Manoux *et al.*, 2005). To estimate RII, first both indicators of socioeconomic status were transformed into summary measures (ridit scores). Then educational level and family *per capita* income were scaled from 0 (highest level of education/income) to 1 (lowest level of education/income) and weighted to reflect the share of the sample at each educational level/income category (Ernstsen *et al.*, 2012). Then, the population in each education/income category was assigned a modified ridit score based on the mid-point of the range in the cumulative distribution of the population of participants in the given categories (Khang *et al.*, 2008; Ernstsen *et al.*, 2012; Kroll, 2013). For example, if the most educated women comprised 10% of the population, the range of women in this category was assigned a value of 0.05 (0.1/2), and if the second category comprised 40% of the population, then every individual in this category was assigned a value of 0.3 (0.1 + [0.4/2]), and so forth. Then, gender-specific generalized linear models for the binomial family were used to calculate age-adjusted RII as follows (Kiadaliri *et al.*, 2015):

$$g(Y) = \beta_0 + \beta_1 \text{ridit} + \beta_2 \text{age} + \text{error}$$

where the error term has a binomial distribution, $Y = 1$ for exposure to the risk factor under study (e.g. fair/poor SRH, 1 + ADL/1 + IADL limitations) and $Y = 0$ for

no exposure. The coefficient β_1 is the coefficient of interest and expresses RII when the link function is log, and ridit is the ridit score (replaces educational level/family *per capita* income). Thus, RII can be interpreted as the rate ratio between the least and the most educated people/the lowest and highest income quintile (i.e. an $RII > 1$ implies a negative relationship between fair/poor SRH, ADL and IADL limitations and education/income, and an $RII < 1$ implies a positive relationship between these variables). In other words, the RII represents the predicted value of the health outcome in the least advantaged divided by the predicted value in the most advantaged (Kunst *et al.*, 1995). Finally, gender-specific RII for education and income were calculated adjusted by i) age and ii) other variables potentially associated with both exposures and outcomes (marital status, depression, falls, physical activity and use of medications).

Data analysis was conducted with the STATA 13 statistical software package (StataCorp 2013).

Results

The sample was characterized by a high proportion of younger men and very old women (Table 1). More than half of the participants had primary level education, and most were in the middle-income distribution with the lowest income quintile being least represented. Compared with men, women showed a higher prevalence of both limitations in ADL and IADL, while the percentage of individuals with fair/poor SRH was relatively high and similar in both genders.

Table 2 shows the prevalence of fair/poor SRH and ADL and IADL limitations by level of education and RII for each outcome. The frequency of fair/poor SRH decreased from the lowest to the highest level of education in both men and women (Table 2). Moreover, there were educational inequalities (age-adjusted RII) in fair/poor SRH for both genders: e.g. women without formal education had 86% (RII 1.859, 95% CI 1.264, 2.455) increased odds of experiencing fair/poor SRH than those with university education.

Additionally, the prevalence of ADL and IADL limitations was markedly higher in women compared with men, with the highest frequency among individuals with no formal education. However, educational inequalities in ADL and IADL were higher in men compared with women (Table 2): e.g. men with no formal education had 36% (age-adjusted RII 1.363, 95% CI 0.071, 2.655) and 97% (age-adjusted RII 1.970, 95% CI 1.052, 2.889) higher odds of having limitations in ADL and IADL, respectively, than those with higher education.

After adjusting for other variables such as age, marital status, occurrence of falls within the past two years, depression within the past year, physical activity and the use of several medications (Table 2, multivariate adjusted RII), educational inequalities decreased slightly although they remained statistically significant for most of the outcomes except for inequalities in ADL in men. Similarly, Table 3 shows the prevalence of fair/poor SRH, and limitations in ADL and IADL by income quintiles and RII for each outcome. The frequency of fair/poor SRH, and limitations in one or more ADL/IADL, decreased from the first (poorest) to the fifth (richest) income quintiles in both men and women (Table 3). Compared with men, the prevalence of ADL and IADL limitations was higher in women throughout the income distribution. Participants within the lowest/poorest income quintile had 66% higher odds of experiencing fair/poor SRH

Table 1. Unadjusted sociodemographic characteristics and prevalence of subjective health indicators in Argentinian men and women aged 60+ years

Variable	Men	Women
	(<i>n</i> = 1984) (%)	(<i>n</i> = 2670) (%)
Age group		
60–64	29.9	26.3
65–69	22.3	23.9
70–74	22.5	17.8
75–89	11.4	13.0
80–84	10.1	11.4
85+	3.8	7.6
Educational level		
No formal education	3.1	3.1
Primary school	55.1	58.5
High school	25.6	23.7
University/higher	16.2	14.7
Income quintile		
1st (poorest)	11.6	9.4
2nd	16.9	19.1
3rd	26.2	25.1
4th	22.2	24.4
5th (richest)	23.1	22.0
Health outcome		
Fair/poor SRH	40.5	40.7
1 + ADL limitations	5.9	12.2
1 + IADL limitations	15.7	26.5

SRH, self-reported health status; ADL, activities of daily living; IADL, instrumental activities of daily living.
Data source: Argentinian National Survey on Quality of Life of Older Adults 2012 (ENCaViAM).

than those within the highest/richest quintile (statistically significant age-adjusted RII was around 1.66 in both genders). Similar income inequalities in IADL limitations existed in both genders while inequalities in ADL limitations were observed only in men. Similarly to what was observed for education in the multivariate adjusted analysis (Table 3), while income inequalities decreased marginally, they remained statistically significant for most of the outcomes.

Discussion

This study evaluated, using the relative index of inequality (RII), educational and income inequalities in SRH and ADL and IADL limitations, among 60-year-old and older Argentinians. Overall, the results showed education- and income-related

Table 2. Prevalence of fair/poor SRH and ADL and IADL limitations by level of education and RII among Argentinian men and women aged 60+ years

Educational level	Fair/poor SRH	Limitations to physical functioning	
		(1 + ADL)	(1 + IADL)
Men (n = 1984)			
No formal education	64.1	8.4	39.3
Primary school	45.5	7.3	18.9
High school	40.4	5.3	12.5
University+	19.2	1.9	5.5
Age-adjusted RII (95% CI)	1.446 (0.756, 2.137)	1.363 (0.071, 2.655)	1.970 (1.052, 2.889)
Multivariate adjusted RII (95% CI) ^a	1.584 (0.872, 2.296)	1.187 (-0.092, 2.466)	1.791 (0.815, 2.767)
Women (n = 2670)			
No formal education	64.9	30.0	63.5
Primary school	48.3	13.9	30.2
High school	30.7	6.2	18.9
University+	21.9	11.1	16.4
Age-adjusted RII (95% CI)	1.859 (1.264, 2.455)	0.806 (-0.311, 1.922)	1.316 (0.593, 2.040)
Multivariate adjusted RII (95% CI) ^a	1.784 (1.142, 2.426)	0.247 (-0.882, 1.375)	1.097 (0.338, 1.856)

^aMultivariate adjusted RII includes: age, marital status, occurrence of falls within the past two years, depression within the past year, physical activity and the use of several medications.

RII, Relative Index of Inequality; CI, confidence interval; 1 + ADL, one or more limitations in activities of daily living; 1 + IADL, one or more limitations in instrumental activities of daily living. Data source: Argentinian National Survey on Quality of Life of Older Adults 2012 (ENCaViAM).

inequalities in the studied subjective health indicators favouring socially advantaged groups. The results remained largely significant after controlling for other associated variables suggesting that education- and income-related inequalities in health indicators – mainly in SRH and IADL – were robust and somehow independent of marital status, physical activity, the use of several medications, depression and the occurrence of falls.

By and large the study findings were in the expected direction for the effects of educational and income inequalities on the studied indicators, to the detriment of socially disadvantaged groups. Nevertheless, it has previously been shown that education and income did not have a similar effect in relation to self-reported health in Argentina (De Maio, 2007, 2008); that is, while poorer income quintiles were associated with higher probabilities of poor health, lower level of education displayed lower probabilities of poor health. The present findings agreed with the described income-related inequalities against disadvantaged groups. After full adjustment, the results on educational inequalities suggested that men and women without formal education had, respectively, 58% and 78% increased odds of experiencing fair/poor SRH than those with university education. Additionally, the prevalence of fair/poor SRH was similar in men and women. This contrasted with evidence from most other countries

Table 3. Prevalence of fair/poor SRH and ADL and IADL limitations by income quintiles and RII among Argentinian men and women aged 60+ years

Income quintile	Fair/poor SRH	Limitations in physical functioning	
		(1 + ADL)	(1 + IADL)
Men (n = 1984)			
1st (poorest)	57.3	11.4	30.7
2nd	51.3	8.5	22.4
3rd	47.0	7.4	17.1
4th	31.9	8.6	18.3
5th (richest)	30.5	3.2	9.2
Age-adjusted RII (95% CI)	1.662 (1.034, 2.291)	1.405 (0.565, 2.245)	1.856 (1.601, 2.551)
Multivariate adjusted RII (95% CI) ^a	1.638 (0.996, 2.280)	1.235 (0.307, 2.163)	1.786 (1.061, 2.511)
Women (n = 2670)			
1st (poorest)	51.8	15.3	43.9
2nd	54.9	14.4	32.9
3rd	44.8	13.0	28.3
4th	33.1	7.8	18.7
5th (richest)	25.8	10.2	16.8
Age-adjusted RII (95% CI)	1.665 (1.134, 2.197)	0.862 (-0.007, 1.731)	1.752 (1.146, 2.358)
Multivariate adjusted RII (95% CI) ^a	1.484 (0.914, 2.053)	0.614 (-0.271, 1.499)	1.624 (0.974, 2.274)

^aMultivariate adjusted RII includes: age, marital status, occurrence of falls within the past two years, depression within the past year, physical activity, and the use of several medications. RII, Relative Index of Inequality; CI, confidence interval; 1 + ADL, one or more limitations in activities of daily living; 1 + IADL, one or more limitations in instrumental activities of daily living.

Data source: from the Argentinian National Survey on Quality of Life of Older Adults 2012 (ENCaViAM).

(Bardage *et al.*, 2005; Hosseinpoor *et al.*, 2012), in which the incidence of poor SRH was higher among adult women.

This study reported a higher prevalence of limitations in physical activities – both ADL and IADL – in women compared with men. This might be due to i) inequalities in the physical and psychological burden associated with activities mainly conducted by women, such as care of elderly/disabled people or family members (Verbrugge, 1985), and/or ii) a greater awareness of health issues in relation to men, which might lead to a higher consciousness of health problems. Additionally, income inequalities in ADL were present in men but not in women after full adjustment. This might be explained by i) the above described tasks being mainly carried out by women, somehow independently of their socioeconomic status; and/or ii) the differences for men and women possibly being a consequence of the gender-stratified analysis: insufficient sample sizes in some subgroups, or the confounding effect of other variables in the regressions; and/or iii) RII might not be sensitive enough to detect inequalities in ADL limitations, which are smaller than, for example, limitations in IADL. In this sense, the lack of correlations

between SRH and activity limitations with different inequality indexes has been previously highlighted (De Maio, 2008).

Finally, gender differences in educational and income inequalities in IADL limitations were not observed in the present study. Compared with participants with university education, men and women in the lowest educational group showed increased odds (97% and 32%, respectively) of experiencing limitations in IADL. The results for income inequalities in IADL showed a similar pattern, favouring higher-income groups. The evidence is inconsistent as to whether socioeconomic inequalities in health are different in women compared with men, across various health measures and life stages (Moss, 2002). The findings of the present study suggest that the patterns of the associations between education/income and subjective health indicators – at least for SRH and IADL – do not vary substantially for older men and women in Argentina.

The study has several limitations that need to be highlighted. Probably the most important one is the use of subjective health indicators. Most studies on social inequalities have been especially focused on subjective health indicators. This is so since national health interviews and level of living surveys usually cover various aspects of subjective health such as perceived general health, physical complaints and quality of life, among others (Kunst *et al.*, 1995). Although there is a large amount of longitudinal evidence showing that perceived or subjective health is a significant predictor of both mortality and morbidity (Idler & Benyamini, 1997; Menec *et al.*, 1999; Maurizio, 2014), some studies have reported a lack of agreement between objective and subjective indicators (De Maio, 2007; Cramm *et al.*, 2015). This calls for the inclusion of both subjective and objective health indicators since they seem to catch ‘different parts’ of a person’s overall health. As such, reliance solely on self-reported health measures seems to provide information that may not reflect actual health performance (Cramm *et al.*, 2015).

Furthermore, it has been shown that the magnitude of health inequality measures depends on the underlying health measure (De Maio, 2008). Some studies reported the highest degree of inequality when subjective health measures like self-assessed health are dichotomized (Ziebarth, 2010). Additionally, it has been suggested that the choice of the welfare indicator determines the size of welfare-related health inequality (Ziebarth & Frick, 2010): welfare measures have a particularly large impact when dichotomized subjective health measures are used to calculate welfare-related health inequalities. The effect is much less pronounced when using quasi-objective generic health measures or objective health measures (Ziebarth & Frick, 2010).

Another limitation is that ENCaViAM does not provide within-country comparisons, which somewhat limits the scope of the present study. Previous evidence from social inequities in Argentina has shown that respondents from deprived regions were most likely to report being in good health, despite living in areas with low life expectancy and high infant mortality (De Maio, 2007); in other words, respondents from areas of poor population health reported good individual health. Thereby, it was suggested that under-reporting of poor health in deprived regions might be because people living in these areas are usually less aware of treatable conditions (Sen, 2002). If true, this will ultimately lead to an underestimate of the social gradient in health (De Maio *et al.*, 2012) and needs to be taken into account when interpreting the findings of the present study. Additionally, the survey only covered urban populations

(close to 85% of the total population of the country; Dachs *et al.*, 2002), avoiding comparisons between urban and rural adults. Finally, the cross-sectional design of ENCaViAM did not allow for making any assumptions on causality.

Despite these limitations, one interesting aspect of the study is that more than half of the studied population had primary school or lower level of education (near 10% had never received any formal education) and around 30% of the sample was represented in the two lowest income quintiles. Thus, the high prevalence of individuals in the lower social strata has strong implications, since the results showed that both low education and income were associated to an increased likelihood of poor health and more physical impairments.

Argentina is undergoing an advanced demographic transition process, being one of the most aged countries in Latin America (CEPAL, 2008). The population aged 60 years and older represents 14.3% of the whole population of the country (INDEC, 2010). Considering the process of population ageing is set to continue, information is needed to successfully face the social challenge of living longer with the best possible quality of life. Efforts to tackle health inequality will contribute to decreasing the inequalities between men and women and different socioeconomic subgroups. Despite the recent implementation of diverse mechanisms of universal access to health care aimed at decreasing the inequitable socioeconomic pattern in older Argentinian older adults, much work still needs to be done (Minoldo *et al.*, 2015).

Acknowledgments

The authors are grateful for comments made on the draft version of this paper by Estela López and the anonymous reviewers.

References

- Alazraqui, M., Diez Roux, A. V., Fleischer, N. & Spinelli, H. (2009) Self-rated health and social inequalities, Buenos Aires, Argentina, 2005 [in Spanish]. *Cadernos de Saúde Pública* **25**(9), 1990–2000.
- Assari, S. & Moghani Lankarani, M. (2015) Does multi-morbidity mediate the effect of socioeconomic on self-rated health? Cross-country differences. *International Journal of Preventive Medicine* **6**, 85.
- Bardage, C., Pluijm, S. M. F., Pedersen, N. L., Deeg, D. J. H., Jylhä, M., Noale, M. *et al.* (2005) Self-rated health among older adults: a cross-national comparison. *European Journal of Ageing* **2**, 149–158.
- Brandt, M., Deindl, C. & Hank, K. (2012) Tracing the origins of successful aging: the role of childhood conditions and social inequality in explaining later life health. *Social Science & Medicine* **74**(9), 1418–1425.
- Cardona, D., Acosta, L. D. & Bertone, C. L. (2013) Inequities in health among Latin American and Caribbean countries (2005–2010) [in Spanish]. *Gaceta Sanitaria* **27**(4), 292–297.
- CEPAL (2008) *Transformaciones demográficas y su influencia en el desarrollo de América Latina y el Caribe*. Comisión Económica, Santo Domingo.
- Cramm, J. M., Borscheuer, L., Selivanova, A. & Lee, J. (2015) The health of India's elderly population: a comparative assessment using subjective and objective health outcomes. *Population Ageing* **8**, 245–259.

- Dachs, J. N. W., Ferrer, M., Florez, C. E., Barros, A. J. D., Narváez, R. & Valdivia, M.** (2002) Inequalities in health in Latin America and the Caribbean: descriptive and exploratory results for self-reported health problems and health care in twelve countries. *Revista Panamericana de Salud Pública* **11**, 335–355.
- De Maio, F. G.** (2007) Health inequalities in Argentina: patterns, contradictions, and implications. *Health Sociology Review* **16**, 279–291.
- De Maio, F. G.** (2008) Ecological analysis of the health effects of income inequality in Argentina. *Public Health* **122**(5), 487–496.
- De Maio, F. G., Linetzky, B., Ferrante, D. & Fleischer, N. L.** (2012) Extending the income inequality hypothesis: ecological results from the 2005 and 2009 Argentine National Risk Factor Surveys. *Global Public Health* **7**(6), 635–647.
- Ernstsen, L., Heine Strand, B., Nilsen, S. M., Arild Espnes, G. & Krokstad, S.** (2012) Trends in absolute and relative educational inequalities in four modifiable ischaemic heart disease risk factors: repeated cross-sectional surveys from the Nord-Trøndelag Health Study (HUNT) 1984–2008. *BMC Public Health* **12**, 266.
- Ferrante, D., Linetzky, B., Konfino, J., King, A., Virgolini, M. & Laspiur, S.** (2011) 2009 National Risk Factors Survey: evolution of the epidemic of chronic non communicable diseases in Argentina. Cross sectional study [in Spanish]. *Revista Argentina de Salud Pública* **2**(6), 34–41.
- Franz, C. E., Finkel, D., Panizzon, M. S., Spoon, K., Christensen, K., Gatz, M. et al.** (2016) Facets of subjective health from early adulthood to old age. *Journal of Aging and Health*, doi: 10.1177/0898264315625488.
- Gasparini, L.** (2007) *Monitoring the socio-economic conditions in Argentina 1992–2006*. Working Paper, Center for Distributional, Labor and Social Studies (CEDLAS), Universidad Nacional de La Plata. URL: www.depeco.econo.unlp.edu.ar/cedlas/monitoreo/pdfs/argentina.pdf (accessed 21st June 2016).
- Haas, S. A.** (2007) The long-term effects of poor childhood health: an assessment and application of retrospective reports. *Demography* **44**(1), 113–135.
- Hoffman, K. & Centeno, M.** (2003) The lopsided continent: inequality in Latin America. *Annual Review of Sociology* **29**, 363–390.
- Hosseinpoor, A. R., Stewart Williams, J., Amin, A., Araujo de Carvalho, I., Beard, J., Boerma, T. et al.** (2012) Social determinants of self-reported health in women and men: understanding the role of gender in population health. *PLoS One* **7**(4), 1–9.
- Idler, E. L. & Benyamini, Y.** (1997) Self-rated health and mortality: a review of twenty-seven community studies. *Journal of Health and Social Behavior* **38**(1), 21–37.
- INDEC** (2010) *Censo Nacional de Población, Hogares y Viviendas 2010. Censo del Bicentenario. Resultados definitivos*. Series B, No. 2. Instituto Nacional de Estadística y Censos, Ciudad Autónoma de Buenos Aires.
- INDEC** (2014) *Encuesta Nacional sobre Calidad de Vida de Adultos Mayores 2012*, 1st edition. Instituto Nacional de Estadística y Censos, Ciudad Autónoma de Buenos Aires.
- Judge, J., Schechtman, K. & Cress, E.** (1996) The relationship between physical performance measures and independence in instrumental activities of daily living. *Journal of the American Geriatrics Society* **44**(11), 1332–1341.
- Kahn, R. S., Wise, P. H., Kennedy, B. P. & Kawachi, I.** (2000) State income inequality, household income, and maternal mental and physical health: cross sectional national survey. *British Medical Journal* **321**(7272), 1311–1315.
- Kawachi, I. & Kennedy, B.** (1999) Income inequality and health: pathways and mechanisms. *Health Services Research* **34**(1, part 2), 215–227.
- Kheng, Y.-H., Yun, S.-C. & Lynch, J. W.** (2008) Monitoring trends in socioeconomic health inequalities: it matters how you measure. *BMC Public Health* **8**, 66.

- Kiadaliri, A. A., Asadi-Lari, M., Kalantari, N., Jafari, M., Mahdavi, M. R. V. & Faghihzadeh, S. (2015) Absolute and relative educational inequalities in obesity among adults in Tehran: findings from the Urban HEART Study-2. *Obesity Research & Clinical Practice*. doi: 10.1016/j.orcp.2015.05.002.
- Kroll, L. E. (2013) *RIIGEN: Stata Module to Generate Variables to Compute the Relative Index of Inequality*. Boston College Department of Economics. URL: <http://EconPapers.repec.org/RePEc:boc:bocode:s457736> (accessed 28th March 2016).
- Kunst, A. E., Geurts, J. J. M. & van den Berg, J. (1995) International variation in socioeconomic inequalities in self reported health. *Journal of Epidemiology and Community Health* **49**, 117–123.
- Lantz, P. M., Lynch, J. W., House, J. S., Lepkowski, J. M., Mero, R. P., Musick, M. A. & Williams, D. R. (2001) Socioeconomic disparities in health change in a longitudinal study of US adults: the role of health-risk behaviors. *Social Science & Medicine* **53**(1), 29–40.
- Lee, Y. (2000) The predictive value of self assessed general, physical, and mental health on functional decline and mortality in older adults. *Journal of Epidemiology and Community Health* **54**(2), 123–129.
- Lima-Costa, M. F., De Oliveira, C., Macinko, J. & Marmot, M. (2012) Socioeconomic inequalities in health in older adults in Brazil and England. *American Journal of Public Health* **102**(8), 1535–1541.
- Lorant, V., Deliège, D., Eaton, W., Robert, A., Philippot, P. & Anseau, M. (2003) Socioeconomic inequalities in depression: a meta-analysis. *American Journal of Epidemiology* **157**(2), 98–112.
- Lundberg, O. & Manderbacka, K. (1996) Assessing reliability of a measure of self-rated health. *Scandinavian Journal of Social Medicine* **24**(3), 218–224.
- Lynch, J., Davey Smith, G., Harper, S., Hillemeier, M., Ross, N., Kaplan, G. A. & Wolfson, M. (2004) Is income inequality a determinant of population health? Part 1. A systematic review. *Milbank Quarterly* **82**(1), 5–99.
- Mackenbach, J. P. (2012) The persistence of health inequalities in modern welfare states: the explanation of a paradox. *Social Science & Medicine* **75**(4), 761–769.
- Mackenbach, J. P. & Kunst, A. E. (1997) Measuring the magnitude of socio-economic inequalities in health: an overview of available measures illustrated with two examples from Europe. *Social Science & Medicine* **44**(6), 757–771.
- Manor, O., Matthews, S. & Power, C. (2000) Dichotomous or categorical response? Analysing self-rated health and lifetime social class. *International Journal of Epidemiology* **29**(1), 149–157.
- Manzelli, H. M. (2014) The relationship between adult mortality and educational attainment in Argentina. PhD thesis, University of Texas at Austin. URL: <https://repositories.lib.utexas.edu/handle/2152/26037>.
- Maurizio, R. (2014) Labour formalization and declining inequality in Argentina and Brazil in 2000s: a dynamic approach. International Labour Force. *ILO Research Paper No.9*.
- Menec, V., Chipperfield, J. & Perry, R. (1999) Self-perceptions of health: a prospective analysis of mortality, control, and health. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences* **54**(2), 85–93.
- Miilunpalo, S., Vuori, I., Oja, P., Pasanen, M. & Urponen, H. (1997) Self-rated health status as a health measure: the predictive value of self-reported health status on the use of physician services and on mortality in the working-age population. *Journal of Clinical Epidemiology* **50**(5), 517–528.
- Minoldo, M. S. T., Zavattiero, C., Peláez, E. & Féliz, J. (2015) Contributory social security systems: equity as a pendent task. Discussions through the cases of Argentina, Paraguay and Dominican Republic [in Spanish]. *Revista Latinoamericana de Población* **9**(16), 75–108.
- Montero López, P., Fernández-Ballesteros, R., Zamarrón, M. D. & Rodríguez López, S. (2011) Anthropometric, body composition and health determinants of active ageing: a gender approach. *Journal of Biosocial Science* **43**(5), 597–610.

- Morales, M. & Paz Terán, C.** (2010) *Educational inequality in Argentina: the best and worst*. Working Paper, Instituto de Estudios Laborales y del Desarrollo Económico (ielde), Universidad Nacional de Salta (UNSa), Argentina.
- Moss, N. E.** (2002) Gender equity and socioeconomic inequality: a framework for the patterning of women's health. *Social Science & Medicine* **54**, 649–661.
- Palloni, A., Pinto-Aguirre, G. & Pelaez, M.** (2002) Demographic and health conditions of ageing in Latin America. *International Journal of Epidemiology* **31**(4), 762–771.
- Pickett, K. E. & Wilkinson, R. G.** (2015) Income inequality and health: a causal review. *Social Science & Medicine* **128**, 316–326.
- Qin, X. & Liu, G.** (2013) Does the US health care safety net discourage private insurance coverage? *European Journal of Health Economics* **14**(3), 457–469.
- Reygadas, L.** (2006) Latin America: persistent inequality and recent transformations. In Hershberg, E. & Rosen, F. (eds) *Latin America after Neoliberalism: Turning the Tide in the 21st Century?* The New Press, pp. 120–143.
- Sen, A.** (2002) Health: perception versus observation. *British Medical Journal* **324**(7342), 860–861.
- Singh-Manoux, A., Ferrie, J. E., Lynch, J. W. & Marmot, M.** (2005) The role of cognitive ability (intelligence) in explaining the association between socioeconomic position and health: evidence from the Whitehall II prospective cohort study. *American Journal of Epidemiology* **161**(9), 831–839.
- Stuckler, D., Basu, S., Suhrcke, M. & McKee, M.** (2009) The health implications of financial crisis: a review of the evidence. *Ulster Medical Journal* **78**(3), 142–145.
- UNESCO** (2006) *International Standard Classification of Education: ISCED 1997*. Volume 5, re-edition. UNESCO, Canada.
- Vagetti, G. C., Barbosa Filho, V. C., Moreira, N. B., Oliveira, V. d., Mazzardo, O. & Campos, W. d.** (2014) Association between physical activity and quality of life in the elderly: a systematic review, 2000–2012. *Revista Brasileira de Psiquiatria* **36**, 76–88.
- Verbrugge, L.** (1985) Gender and health: an update on hypotheses and evidence. *Journal of Health and Social Behavior* **26**(3), 156–182.
- Vries, R., Blane, D. & Netuveli, G.** (2014) Long-term exposure to income inequality: implications for physical functioning at older ages. *European Journal of Ageing* **11**, 19–29.
- Wagstaff, A. & van Doorslaer, E.** (2000) Income inequality and health: what does the literature tell us? *Annual Review of Public Health* **21**, 543–567.
- WHO** (2001) *International Classification of Functioning, Disability and Health: ICF*. WHO Library Cataloguing-in-Publication Data, Geneva.
- WHO** (2002) *Active Ageing: A Policy Framework*. WHO, Geneva.
- Ziebarth, N.** (2010) Measurement of health, health inequality, and reporting heterogeneity. *Social Science & Medicine* **71**(1), 116–124.
- Ziebarth, N. & Frick, J.** (2010) *Revisiting the Income–Health Nexus: The Importance of Choosing the 'Right' Indicator*. Technical Report, SOEP papers 274/2010, DIW, Berlin.