



Abstract Book

SETAC Latin America 14th Biennial Meeting
Latin America, Diversity of Knowledge for a Sustainable Future

26-29 SEPTEMBER 2021 • VIRTUAL

Abstract Book

SETAC Latin America 14th Biennial Meeting

Table of Contents

Abstracts by Session

01A. Effects and Risks of Pesticide Used in Agriculture I	4
01B. Effects and Risks of Pesticide Used in Agriculture II	9
02. Micro- and Macroplastics Pollution	13
03. Chemistry and Exposure Assessment	19
04. Contaminants of Emerging Concern	27
05. Traditional Knowledge, Policy, Management and Communication	36
06A. Aquatic Toxicology, Ecology and Stress Response I	40
06B. Aquatic Toxicology, Ecology and Stress Response II	52
07. Ecotoxicology of Birds and Mammals: Novelty and Challenges	63
08. Endocrine Disruptors Compounds: Environmental Presence and Effects	67
09. Behavioral Ecotoxicology: Where Organism and Population-level Effects Meet	71
10. Engineering, Remediation and Restoration	75
11. Sources, Fate and Effects of Pollutants in Marine Ecosystems	79
12. Environmental Toxicology and Stress Response	82
13. Terrestrial Ecotoxicology and Ecosystem Services	87
14. Environmental Impact of Mining and Other Industrial Activities	94
15. Alternative Approaches to Animal Testing in Ecotoxicity and Risk Assessment	97
16. Occurrence and Risks of Pollutants in Latin America	102

Indices

Keyword Index	109
Author Index	113

This book comprises the abstracts of the presentations for the platform and poster sessions of the Society of Environmental Toxicology and Chemistry (SETAC) Latin America 14th Biennial Meeting, conducted virtually from 26–29 September 2021. The abstracts are reproduced as accepted by the Scientific Program Committee and appear in numerical order. In each abstract, the presenting author’s name is underlined. The author index cross-references the corresponding abstract numbers.

No part of this publication may be reproduced, distributed, stored, or transmitted in any form or by any means, including photocopying, recording or other electronic or mechanical methods, without permission in writing from the copyright holder.

All rights reserved. Authorization to photocopy items for internal or personal use, or for the purpose or internal use of specific clients, may be granted by the Society of Environmental Toxicology and Chemistry (SETAC), provided that the appropriate fee is paid directly to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923 USA (+1 978 750 8400) or to SETAC. Before photocopying items for educational classroom use, please contact the Copyright Clearance Center (www.copyright.com) or the SETAC Office in North America (+1 850 469 1500, setac@setac.org).

SETAC's consent does not extend to copying for general distribution, promotion, creating new works or resale. Specific permission must be obtained in writing from SETAC for such copying. Direct inquiries to SETAC, PO Box 12804 Pensacola, Florida, 32591-2804 USA.

© 2021 Society of Environmental Toxicology and Chemistry (SETAC)

previous studies show that the structural family of citronellol may have interesting antimicrobial activity. In order to tackle antibacterial resistance world-wide issue, the discovery of new bactericide molecules is utterly important. Despite such products from natural origin are credited with safety, citronellol shows some dermal and oral toxicity when tested at high concentrations in laboratory animals. Nevertheless, reports of its ecotoxicity in non-target organisms can hardly be found when citronellol is used as an active ingredient in biopesticides. This study assesses the antimicrobial activity of citronellol on a selection of gram-positive and gram-negative pathogenic bacteria, as well as its ecotoxicological effects on non-target soil organisms (*Eisenia fetida* and *Allium cepa* L.). *E. fetida* was exposed to different concentrations of citronellol ranged from 0.1 to 200 mg/L. *A. cepa* root grow was monitored after exposure to the following concentrations of citronellol: 300, 30, 3, 0.3 and 0.03 mg/L, measuring, therefore, the phytotoxicity on this vascular culture plant. The dose-response analysis allowed us to calculate the ecotoxicity of citronellol on *E. fetida*: $LC_{50} = 12.07$ mg /L (8.47-17.28) and $LC_{10} = 3.36$ (1.54-5.21); and over *A. cepa*: $LC_{50} = 172.40$ mg /L (122.91-252.50) and $LC_{10} = 0.033$ (0.018-0.054). These results show that citronellol is not harmless on non-target soil organisms and it may produce both ecotoxicity and phytotoxicity. These systematic studies are relevant since they provide valuable information about the effects of biopesticide compounds on the soil environment and especially in non-target organisms, when taking decisions on their commercialization. *The authors thank the financial support of Gobierno de Aragón-FSE-FEDER "Construyendo Europa desde Aragón" (Grupo E39_17R y RIS3 LMP28_18) and Catedra NOVALTIA.*

13.17 Bioaccumulation and Translocation of Cu, Cd, Zn, and As in Four Native Tree and Shrub Species Growing on Soils Contaminated by an Abandoned Gold Mine

B. Heredia, Instituto Nacional de Tecnología Agropecuaria / Estación Experimental Agropecuaria San Juan; R. Tapia, Consejo Nacional de Investigaciones Científicas y Técnicas-Instituto Nacional de Tecnología Agropecuaria / Estación Experimental Agropecuaria San Juan; B.J. Young, Instituto Nacional de Tecnología Agropecuaria / Instituto de Microbiología y Zoología Agrícola; G. Roqueiro, Instituto Nacional de Tecnología Agropecuaria / Estación Experimental Agropecuaria San Juan

Plants are known to have the ability to bioaccumulate metal(loid)s in tissues and translocate them from roots to aerial biomass. Native plants grow in soil contaminated with mining waste in the town of La Planta (Caucete, San Juan, Argentina). The objective of this study was to evaluate the bioaccumulation and translocation capacity of Cu, Cd, Zn and As in *Prosopis flexuosa* (Pf), *Larrea cuneifolia* (Lc), *Bulnesia retama* (Br), and *Plectrocarpa tetraantha* (Pt) that grow in contaminated soil. Organ samples (leaf, branches, stem, bark and root) of 3 plants per species (n=12) were collected. Rhizospheric soil samples around each tree and shrub were taken from the first 20 cm of depth (n=12). Concentrations of metal(loid)s in organs and soil were determined. Samples were digested with a combination of HNO₃, H₂O₂ and HF, and quantifications were made using ICP-MS. Bioaccumulation Factor (BAF) and Translocation Factor (TF) were calculated. Results showed that the most concentrated metal(loid)s in Lc, Pf and Br were Cu and Zn, and in Pt were Zn, Cu and As (p< 0.001). BAF and TF values greater than 1 were obtained for the four species, indicating that these species are bioaccumulators. The highest BAF values was obtained for Cu in the four species (p< 0.001). Regarding TF, higher values of Zn were only observed in leaf of Lc (p< 0.001). The high concentrations of

metal(loid)s measured in plant organs of the four native species plus the values of BAF and TF indicate a high potential for phytoextraction. Comparing bioaccumulation capacities, $Pt > Pf > Lc$ for As, $Pf > Lc > Pt > Br$ for Cu, $Pf > Pt > Lc > Br$ for Zn, and only Pt was effective for Cd. Bioaccumulation capacity of Cu, Cd, Zn and As found in the native plants studied in this work, generates baseline information for the implementation of phytoremediation strategies. These species present anatomical, morphological and structural adaptations that allow them to survive adverse climatic conditions that characterize arid and semi-arid environments. This would avoid the use of exotic species that could generate disturbances in the polluted environment and a higher economic cost.

13.18 Acute Toxicity of Soils Contaminated With Waste From an Abandoned Gold Mine on Seeds of Prosopis Flexuosa and Larrea cuneifolia

P. Diaz, Universidad Nacional de San Juan / Facultad de Ingeniería; B.J. Young, Instituto Nacional de Tecnología Agropecuaria / Instituto de Microbiología y Zoología Agrícola; R. Tapia, Consejo Nacional de Investigaciones Científicas y Técnicas-Instituto Nacional de Tecnología Agropecuaria / Estación Experimental Agropecuaria San Juan; G. Roqueiro, B. Heredia, Instituto Nacional de Tecnología Agropecuaria / Estación Experimental Agropecuaria San Juan

The presence of metal(loid)s in soil causes toxicity in plants such as physiological disorders, and alterations in growth and development. The objective of this study was to evaluate the acute toxicity of soils contaminated with mining waste on seeds of *Prosopis flexuosa* (Pf) and *Larrea cuneifolia* (Lc), two native species from the Monte region (La Planta, San Juan, Argentina). Seeds were collected in the surroundings of La Planta and soil samples were taken in two sites: contaminated (S1) and reference sites (S2). Seeds were exposed to 6 increasing concentrations of soil mixtures from Site 1 and 2: 0, 10, 25, 50, 70, and 100% (v/v), where 0 corresponds to S2 soil and 100% to S1 soil. Experimental unit consisted of 20 seeds placed on wet soil into a Petri dish, reaching a total of 5 repetitions per treatment. A pre-germination treatment was applied to the seeds to ensure the seedling emergence. Toxicity test was carried out in a germination chamber in darkness at 25±2 °C. Due to differences in germination time of each species, exposure of Pf and Lc lasted 7 and 15 d, respectively. Mean germination time (MGT), germination index (GI), relative growth index (RGI) and IC₅₀ of root (r) and hypocotyl (h) length were estimated. The concentration of the main metal(loid)s in soils were: S1) As=6608 mg kg⁻¹; Zn=10892 mg kg⁻¹; Cu=260 mg kg⁻¹; Cd=90 mg kg⁻¹, and S2): Zn=46 mg kg⁻¹; As, Cu and Cd were not detected. For Pf a significant increase in MGT was observed from treatment 70%, whereas a significant decrease in RGI_r and RGI_h, and GI were observed from 50% S1 concentration (p< 0.001). IC_{50r} and IC_{50h} were estimated in 21.1 and 40.3%, respectively. For Lc, seed germination inhibition of 100% was observed from 70% S1 concentration. Statistical differences between the values of RGI_r, RGI_h, and GI for remaining treatments were obtained (p< 0.001), while MGT showed no significant differences. The strongest inhibition was observed in 50% S1 concentration. IC_{50r} and IC_{50h} were estimated in 27.7% and 15.1%, respectively. The values of the toxicity endpoints and the phytotoxicity indexes showed that these native species are more tolerant to contaminated soil of La Planta than horticultural species evaluated in previous studies (IC_{50r}< 1% S1 concentration). In future studies, chronic effects should be evaluated to determine the underlying physiological mechanisms of the metal(loid) tolerance in Pf and Lc.