

A new formulation of oxalic acid for *Varroa destructor* control applied in *Apis mellifera* colonies in the presence of brood

Matías MAGGI^{1,2}, Elian TOURN^{3,4,5}, Pedro NEGRI^{1,2}, Nicolás SZAWARSKI¹,
Alfredo MARCONI^{3,4,5}, Liliana GALLEZ⁶, Sandra MEDICI^{1,2}, Sergio RUFFINENGO⁷,
Constanza BRASESCO¹, Leonardo De FEUDIS¹, Silvina QUINTANA⁸, Diana SAMMATARO⁹,
Martin EGUARAS^{1,2}

¹CIAS, Centro de Investigación en Abejas Sociales (ex Laboratorio de Artrópodos), Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Funes 3350, 7600, Mar del Plata, Argentina

²CONICET, Consejo Nacional de Investigaciones Científicas y Técnicas, Rivadavia 1917, C1033AJ Buenos Aires, Argentina

³Departamento de Agronomía, Universidad Nacional del Sur, Bahía Blanca, Buenos Aires, Argentina

⁴Ministerio de Agricultura, Ganadería y Pesca de la Nación, EEA INTA Bordenave, Bahía Blanca, Buenos Aires, Argentina

⁵Cooperativa de Trabajo Apícola Pampero Ltda., Bahía Blanca, Buenos Aires, Argentina

⁶LabEA, Depto. Agronomía, Universidad Nacional del Sur, 8000, Bahía Blanca, Argentina

⁷Cátedra de Apicultura, Facultad de Ciencias Agrarias, Universidad Nacional de Mar del Plata, 7620, Balcarce, Argentina

⁸Laboratorio de Biología Molecular, Fares Taie Instituto de Análisis, Mar del Plata, Argentina

⁹USDA ARS Carl Hayden Honey Bee Research Center, Tucson, AZ, USA

Received 14 July 2014 – Revised 4 August 2015 – Accepted 26 October 2015

Abstract – An organic product based on oxalic acid was evaluated for use in *Varroa* control under spring/summer climatic conditions in Argentina. The formulation consists of four strips made of cellulose impregnated with a solution based on oxalic acid. Forty-eight beehives were used to assess the product efficacy. Residues of the product were also tested in honey, bees, and wax. Each trial had respective control groups without oxalic treatment. At the beginning of the experiment, four strips of the formulation were applied to the colonies belonging to the treated group. Falling mites were counted after 7, 14, 21, 28, 35, and 42 days. After the last count, the strips were removed and colonies received two flumethrin strips for 45 days. Falling mites were counted throughout this period. Average efficacy of the organic product was 93.1 % with low variability. This product is an organic treatment designed for *Varroa* control during brood presence and represents a good alternative to the synthetic treatments.

Varroa destructor / control / oxalic acid / bee brood / Argentina

1. INTRODUCTION

Varroa destructor (Acari: Varroidae), an obligate ectoparasitic mite of the honey bee, *Apis*

mellifera (Hymenoptera: Apidae), feeds on the hemolymph of adult bees, larvae, and pupae in capped brood cells. In Argentina, severe *V. destructor* infestation of *A. mellifera* results in honey bee colony death within 1 to 2 years of initial infestation (Eguaras and Ruffinengo 2006). In recent years, resistance to acaricides has become a major problem in the control of mite populations. Increased tolerance to the most widely used synthetic active ingredients has been

Corresponding author: M. Maggi,
biomaggi@gmail.com

Manuscript editor: Yves Le Conte

observed (Maggi et al. 2010a, 2011). Because of the resistance to acaricides, there is a renewed interest in Argentina for substances of natural origin, such as essential oils and their components or organic acids, especially formic acid and oxalic acid (Eguaras et al. 2001, 2003; Maggi et al. 2010b; Ruffinengo et al. 2014). This situation related to *Varroa* resistance has been observed around the world (Sammataro et al. 2005).

Oxalic acid (OA) is widely used for controlling *V. destructor* because of its high efficacy (>90 %) and low risk of hive contamination (Gregorc and Planinc 2001, 2002; Nanetti et al. 2003; Marinelli et al. 2006; Rademacher and Harz 2006; Bacandritsos et al. 2007). OA is applied to colonies by spraying or trickling a solution of OA and sugar-water over the bees or by evaporating crystals with heat (Rademacher and Harz 2006). However, most tests have been carried out during a broodless period, and have reported more than 95 % efficacy (Higes et al. 1999; Charriere and Imdorf 2002). So, mite mortality after treatment with oxalic acid seems to be directly influenced by the presence of brood because oxalic acid does not kill the mites in sealed brood cells (Imdorf et al. 2003). Several reports agree on the use of oxalic acid when no brood is present in winter (Barbero et al. 1997). Its efficacy when brood is present is around 60 % (Charrière 1997; Rademacher and Harz 2006). Thus, the usefulness of the acid seems limited in warm climates with a long brood rearing period.

Taking into account that oxalic acid is a suitable compound for the control of *V. destructor* in broodfree colonies during the autumn and winter period within the concept of Integrated *Varroa* Control, the major challenge to scientists is to develop a new formulation that can be used during permanent brood presence in colonies of *A. mellifera*. The main goal of this article was to evaluate the efficacy of a new formulation based on oxalic acid for when brood was present.

2. MATERIAL AND METHODS

Location of study. Summer field trials were carried out during March and April 2010 in the experimental apiary located near Bahía Blanca city (Buenos Aires, Argentina) and during January and February 2011 in

the experimental apiary of The Charrúas Government (Downtown of Charrúas, Entre Ríos, Argentina). The average temperature recorded during this period in Bahía Blanca city was 22 °C (range 9–33 °C), and in Charrúas, the average was 31 °C (range 14–42 °C). Autumn field trials were carried out during May and June 2013, in the experimental apiary of Arthropods Laboratory (National University of Mar del Plata), near Mar del Plata city. The average temperature recorded during this period was 14 °C (range –2–20 °C).

Field trials. Forty-eight beehives were used to assess product efficacy. Three experiments were conducted to test the efficacy of a new formulation with oxalic acid (Aluen CAP) made by Cooperativa de Trabajo Apícola Pampero Ltda. In the first two trials, the treatment efficacy was assessed during the summer season: one trial was performed in Bahía Blanca (15 colonies divided in three groups of five colonies each one: a control group “A” without treatment, a second control group “B” with cellulose strips embebed with glycerin and a treatment group “C” with Aluen CAP) and the other in Charrúas (19 colonies divided in two groups: one of nine (control group) and the other of ten colonies (treatment group, Aluen CAP)). In a third trial, the honey and wax was tested for any OA residues (apiary located at Mar del Plata, 14 colonies divided in two groups of 7 colonies: control and treatment groups). All colonies used during trials were previously equalized for bee population, brood area, and honey and pollen stores. Hive bottoms specially adapted for the collection of dead mites were placed in each colony. In each trial, control groups without oxalic treatment were established. Apiaries were selected based on geographic locations at least 5 km away to avoid reinfestation phenomena.

At the beginning of the experiment, the new formulation were applied to the treatment group. Aluen Cap treatment consists of four strips U-shaped. The matrix of these strips is composed of cellulose (45 cm×3 cm×1.5 mm); each one contains 10 g of OA mixed with 20 mL of glycerin. Each strip was placed astride on frames 2, 4, 6, and 8 of the brood chamber (Figure 1). Falling mites were counted after 7, 14, 21, 28, 35, and 42 days using the hive bottoms specially adapted for the collection of dead mites. After the last count, the strips were removed and at the same day, colonies received four flumethrin strips according to the instructions of the manufacturer (registered trade name: Flumevar[®],



Figure 1. The application of the new formulation Aluen CAP, an organic product containing oxalic acid in strips of cellulose and glicerine to a hive.

supplied by APILAB SA, Argentina, <http://www.apilab.com/flumevarcostarica.pdf>). No resistance phenomena to flumethrin had been reported previously in these apiaries or in the region where assays were performed. This treatment was left in the colonies for 45 days. Falling mites also were counted throughout this period.

The efficacy of oxalic acid treatment was calculated as a percentage: $((\text{number of dead mites during oxalic acid treatment})/(\text{number of dead mites collected during the treatment with OA and flumethrin})) \times 100$. The cumulative mite fall after oxalic acid and flumethrin treatment was assumed to be 100 %. Data on percent efficacy were analyzed by analysis of variance (ANOVA) after arcsine transformation in the case of percentages, to reduce the heterogeneity of the variance.

Colony population development. The progress of the hives treated with the oxalic acid was monitored prior to each treatment application and also 1 week after the acaricide application. As an example, at T_0 , the colonies were inspected and the first treatment application was applied. Any change was compared with the untreated control hives. All other conditions (weather, nourishment, and supervision) were identical. The parameters

to quantify the general state of the colonies during the evaluation were as follows: number of combs fully covered with bees and open and sealed brood areas. These procedures were satisfactory to assess colony development in previous research (Maggi et al. 2013; Negri et al. 2015). All colonies were checked by visual observations for dead brood bees and queens after treatment. Data were analyzed by ANOVA to evaluate if OA treatment produces changes in the variables measured.

Oxalic acid extraction from bees, honey, and beeswax. During the autumn assay, samples of honey, bees, and wax were collected from colonies to detect possible residues generated by oxalic acid treatment. The bees were analyzed as a whole. Samples of each colony were taken prior the oxalic treatment application (control, timepoint T_0) and 42 days after it (timepoint T_2). Samples were collected according to the protocol of the European Working Group (2001) and were stored at -80°C until analysis. Ten grams of sample was then diluted with 80 % ethanol. Beeswax sampled was heated in a water bath $62 \pm 2^\circ\text{C}$ for 15 min; the samples had been previously acidified with 100 μL of pure HCl (Merck®). Tubes were shaken vigorously for 1 min. An aliquot of 50 μL was placed in a 2-mL vial and

evaporated until dried, using a nitrogen stream. The extract was resuspended in 1 mL of ethanol: acid solution. Vials were placed in incubator at 50 ± 2 °C for 5 h.

GF-FID analysis. A gas chromatograph HP 6890 with a FID detector and an autosampler for 100 samples was used. For GF-FID analysis, an Autosampler Agilent 7683 was used. The injector conditions were: temperature, 100 °C; pressure, 12.45 p.s.i.; split ratio, 10:1; and gas saver, 20 mL/min. An Agilent Column (30 m \times 0.25 m \times 0.25 μ m HP-5MS Ultra Inerte (p/n 19091S-433UI)) was employed using Helium 5.0 (Praxair) in constant pressure mode (flow, 1 mL/min; oven temperature, 50 °C; injection volume, 1 μ L; detector: FID: temperature, 250 °C) as gas carrier. The average recovery of the oxalic acid ranged from 87 to 110 %, and detection limits were honey: detection limit, 0.5 mg/kg–quantification limit, 1 mg/kg; wax: detection limit, 1.5 mg/kg–quantification limit, 12 mg/kg; bees: detection limit, 0.8 mg/kg–quantification limit, 2 mg/kg. Data were analyzed by ANOVA to evaluate if OA treatment produces changes in the variables measured.

3. RESULTS

Efficacy. The efficacy of the treatment in *Varroa* control was determined by comparing the number of falling mites recorded during the oxalic acid treatment period with the total number of falling mites recorded during the whole trial (including the flumethrin treatment). Results are presented in Table I. The average final efficacy in colonies treated with the new oxalic acid formulation was 94 % (in the experimental apiary located near to Charrúas downtown, summer trial), 92.7 % (in experimental apiary placed near to Bahia Blanca city, summer trial) and 92.8 % (in experimental apiary placed near to Mar del Plata city, autumn trial). Final efficacy of the OA treatment had significant differences to the control ($P < 0.05$). All trials had a low variability in the final efficacy (range between 85.9 ± 98.8 %). The highest mortality for the three assays was recorded during the first 22 days, with an average partial efficacy of 74.4 %.

Effects on colony population Colony population parameters before and after treatment are shown

in Table II. For summer trials, treated colonies (oxalic acid treatment) finished with an average of 4.5 brood combs and 8.2 frames of bees. In the autumn trial, colonies finished with 3.2 combs covered with brood (open+sealed brood) and 8 frames covered with adult bees. Adverse effects on the colonies (dead bee brood or queens) were not detected during and after treatment. Adverse effects on the bee populations were not detected in the treated colonies in all three trials ($P > 0.05$).

Oxalic acid detection from bees, honey and beeswax. Table III summarizes the measurements of oxalic acid in all samples taken before and after treatment. The natural oxalic acid content varied between 2.5 and 33.8 mg/kg. There was no increase in oxalic acid content of honey, wax, and bees after treatments, in all three trials ($P > 0.05$). All samples of bees and beeswax were negative before and after the OA treatment.

4. DISCUSSION

The new formulation with oxalic acid as the active ingredient, presents some advantages over the current methods of organic acid treatments. Polymer-cellulose matrices not only have all the benefits of the oxalic acid liquid treatments but also delay the release of OA so that it can remain longer in the colony (Eguaras et al. 2003). This reduces the number of visits to apiaries to reapply OA treatments. In this study, we report a new formulation based on oxalic acid with high acaricide activity against *V. destructor*. However, taking into account that mite fall was registered with hive bottom specially adapted, the estimated efficacy results not exclusively from the application of the test medication but also from a combinatory effect (natural mite mortality). Nonetheless, strong differences were detected with the control group, demonstrating an excellent acaricide activity of the new treatment.

Other research has demonstrated that good efficacy has been obtained using at least three applications of liquid oxalic acid per colony. In other cases, three or more applications were not enough for effective *Varroa* control (Rademacher and

Table 1. Number of falling mites with oxalic acid treatment and flumethrin treatment for each of the three field assays.

Trial	Treatment	Colony	No. of falling mites with treatment at day						Total	No. of falling mites with shock treatment	Efficacy of formulation (%)	Mean efficacy of formulation (%)
			7	14	21	28	35	42				
Summer trial (Bahia Blanca 2010)	Oxalic acid formulation	1	32	0	56	8	11	18	125	15	89.3	92.7
		2	395	852	872	212	77	52	2460	40	98.4	
		3	55	97	123	89	94	80	538	44	92.4	
		4	211	228	221	44	68	54	826	65	92.7	
		5	107	30	97	139	86	10	469	61	88.5	
	Control I (strips with glycerine)	0	0	4	5	3	0	0	12	42	22.2	20.6
		2	2	6	8	11	13	4	44	49	47.3	
		1	1	3	1	0	0	0	5	164	2.9	
		12	12	5	3	2	0	16	38	106	26.4	
		1	1	2	1	0	0	3	7	169	3.9	
		11	0	0	4	7	7	5	23	77	23	28.9
		12	0	0	2	0	4	6	12	84	12.5	
Summer trial (Charrúas, Bahia Blanca 2011)	Oxalic acid formulation	13	2	0	1	0	0	10	13	60	17.8	
		14	2	5	8	6	12	18	51	142	26.4	
		15	14	8	11	9	17	74	133	72	64.9	
		16	54	185	144	4	15	1	403	21	95	94
		17	0	282	46	14	0	1	343	14	96.1	
		18	8	695	595	15	3	3	1,319	16	98.8	
		19	19	186	112	18	3	4	342	1	99.7	
		20	367	0	1	0	5	3	376	15	96.1	
		21	0	915	348	41	1	2	1,307	0	100	
		22	88	620	340	815	253	221	2,337	70	97	
		23	42	42	93	2	3	1	183	20	90.1	
		24	26	10	33	1	3	0	73	23	76	
25	46	67	49	22	5	0	189	18	91.3			
Untreated	26	8	14	8	1	10	0	41	2032	1.9	5.2	
	27	20	6	9	1	13	2	51	1295	3.8		

Table 1 (continued)

Trial	Treatment	Colony	No. of falling mites with treatment at day							Total	No. of falling mites with shock treatment	Efficacy of formulation (%)	Mean efficacy of formulation (%)
			7	14	21	28	35	42					
Autumn trial (Mar del Plata 2013)	Oxalic acid formulation	28	16	3	1	0	0	0	2	22	592	3.6	
		29	62	5	6	2	0	0	1	76	1920	3.8	
		31	1	9	2	1	1	3	17	121	121	12.3	
		32	20	11	4	5	16	1	57	897	897	5.97	
		33	54	9	20	0	8	1	92	1120	1120	7.5	
		34	8	1	6	9	2	0	26	796	796	3.1	
		35	28	6	8	11	13	0	66	1234	1234	5.1	
		36	34	48	27	1	0	0	110	9	92.4	92.8	
		37	120	106	24	1	0	0	251	21	92.2		
		38	56	214	56	0	2	0	328	13	96.1		
		39	12	47	22	2	0	0	83	1	98.8		
		40	87	119	17	1	1	2	227	42	84.3		
		41	20	42	2	1	4	2	71	3	95.9		
		42	41	12	9	0	0	0	62	7	89.8		
		43	1	3	0	1	2	0	7	246	2.7	13.7	
		44	0	1	0	0	4	1	6	326	1.8		
		45	0	0	1	9	0	0	10	128	7.2		
		40	0	0	7	2	0	2	11	147	6.9		
446	2	8	3	12	2	0	27	56	32.5				
47	3	1	4	0	0	0	8	16	33.3				
48	1	3	4	0	1	3	12	91	11.6				

Table II. Colony population parameters before and after treatment for the three trials.

Colony population parameter	Treatment	Apiary I (Bahía Blanca)		Apiary II (Charrías, Entre Ríos)		Apiary III (Mar del Plata)	
		Before treatment	After treatment	Before treatment	After treatment	Before treatment	After treatment
Number of combs covered with adult bees	OA	9.8a±0.4 (5)	8.6a±0.4 (5)	8.4a±2.2 (10)	7.9a±2.1 (10)	8.7a±1.2 (7)	8a±0.5 (7)
	Control	9.8a±0.4 (10)	8.8a±0.4 (10)	7.4b±1.6 (9)	7.4b±1.7 (9)	9a±0.4 (7)	8.6a±0.2 (7)
Number of combs covered with brood	OA	6.4c±2 (5)	2.9d±.2 (5)	4.6d±1.4 (10)	4.7d±1.8 (10)	4.6d±0.8 (7)	3.2d±0.4 (7)
	Control	5.8c±2.4 (10)	4.1d±1.7 (10)	4.8d±1.3 (9)	5d±1 (9)	4.7d±0.3 (7)	3.1d±0.3 (7)

Mean±standard deviations. The number of colony sampled are enclosed in parentheses. Uppercase (a–d) letters indicate statistical differences among treatments (*t* test, *P* < 0.05)

Harz 2006). Also, during the summer season, the efficacy obtained by liquid treatments is reduced as oxalic acid does not kill the mites in sealed brood cells. In addition, Hatjina and Haristos (2005) have shown a detrimental effect on brood development plus low effectiveness using the trickling method of OA when open brood is present. They suggest that this method is not as safe as has been reported in the past.

Marinelli et al. (2006) evaluated the efficacy of cellulose strips with oxalic acid in comparison with other oxalic acid treatments (trickling and vaporization methods). In spring trials, the effectiveness of the cellulose strips was not statistically different from the natural fall of *Varroa*. Similarly, the autumn trials had the same results. However, when spraying, trickling, and vaporizing oxalic acid, good control of *Varroa* was seen in central Italy. They conclude that the unsatisfactory efficacy of cellulose strips versus the good control results of trickling and vaporization could be explained by the high acidity of the oxalic acid water solutions. A pH around 1 may be responsible for the best oxalic acid activity against the mites. Here, we show that one application of the Aluen Cap formulations is enough to have good mite control even with the presence of bee brood. In this study, the oxalic acid formulation had a high efficacy and it was possible to keep low *Varroa* prevalence indices in the colonies. The matrix of the OA strip may help to keep an adequate acid concentration inside the colonies for a long time (42 days). In addition, the combination of glycerin with AO could help to maintain the drug for a longer time inside the beehive and consequently with mites (including those that are emerging from brood cells). As it was reported by Segur and Oberstar (1951), glycerin possesses a high viscosity. This chemical property would help to disseminate the acid among bees during a longer time in comparison with other strip formulations without glycerin, increasing its efficacy (these hypotheses should be examined in later studies). During the season when this work was done, the temperature range was -2–42 °C, and in the three trials, there were no problems with the higher temperatures in the colonies.

This new product offers a good alternative for *Varroa* control as it does not have the two most

Table III. Oxalic acid residues in wax, bees and honey after treatment.

Treatment	Matrix	Before treatment	After treatment
Control	Wax	0 (7)	0 (7)
	Bees	0 (7)	0 (7)
	Honey	14.2 ^b ±11.6 (7)	25.6 ^c ±12.5 (7)
OA	Wax	0 (7)	0 (7)
	Bees	0 (7)	0 (7)
	Honey	7.5 ^a ±5.2 (7)	17.3 ^b ±15 (7)

Mean±standard deviations (SD) and number of samples (*n*). Honey: detection limit, 0.5 mg/kg–quantification limit, 1 mg/kg; wax detection limit, 1.5 mg/kg–quantification limit, 12 mg/kg; bees: detection limit, 0.8 mg/kg–quantification limit, 2 mg/kg. Uppercase letters (a–c) indicate statistical differences among treatments (*t* test, *P*<0.05)

frequent disadvantages of organic products used for mite control. First, it is easy to use, safe for beekeepers, and presents low variability between colonies in its efficacy. Second, it does not present a risk to colony development and so it can be used to safely reduce the damage done by *Varroa* mites even during the summer season.

In Argentina (as in many other parts of the world), the use of synthetic acaricides is being restricted due to the emergence of resistant mite populations (Maggi et al. 2009, 2010a, 2011; Sammataro et al. 2005), as well as honey and wax contamination (Bogdanov et al. 2003; Wallner 1999; Medici et al. 2015). Oxalic acid is a natural constituent of honey; values between 8 and 17,000 mg/kg have been found in different honeys (Mutinelli et al. 1997; Bernardini and Gardi 2001; Bogdanov et al. 2002).

Most vegetables contain much higher amounts of oxalic acid than honey so the total daily intake is negligible. Thus, from a nutritional point of view, oxalic acid should, like formic acid, also have a generally recognized as safe (GRAS) status. Moreover, no significant residues are expected after oxalic acid treatments as demonstrated in our research. Indeed, there is no risk of honey residues after all types of oxalic acid treatments (Radetzki 1994; Mutinelli et al. 1997; Del Nozal et al. 2000; Bernardini and Gardi 2001; Radetzki and Barmann 2001; Bogdanov et al. 2002).

Spraying and trickling of oxalic acid are accepted for use against *V. destructor* in most European countries and are widely used by beekeepers throughout Europe (Charriere and Imdorf 2002).

This new formulation is not dangerous for the user or for the bee colony. Thus, Aluen CAP could be a valid alternative for *Varroa* mite control.

ACKNOWLEDGMENTS

The authors would like to thank the Universidad Nacional de Mar del Plata (UNMDP) and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). This research was supported by a grant of ANPCyT, PICT 2011 to M.M.

Une nouvelle formulation de l'acide oxalique appliquée dans les colonies d'*Apis mellifera* pour lutter contre *Varroa destructor*, en présence du couvain

Acari / lutte antiparasitaire / couvain d'abeille / Argentine

Eine neue Oxalsäure-Formulierung zur Bekämpfung von *Varroa destructor* in Völkern von *Apis mellifera* mit Brut

***Varroa destructor* / Bekämpfung / Oxalsäure / Bienenbrut / Argentinien**

REFERENCES

- Bacandritsos, N., Papanastasiou, I., Saitanis, C., Nanetti, A., Roiniotti, E. (2007) Efficacy of repeated trickle applications of oxalic acid in syrup for varroosis control in *Apis mellifera*: Influence of meteorological conditions and presence of brood. *Vet. Parasitol.* **148**, 174–178

- Barbero, R., Panella, F., Bonizzoni, L. (1997) Ácido oxálico y el tratamiento de limpieza radical de otoño-invierno. *Vida Apícola* **85**, 8–13
- Bernardini, M., Gardi, T. (2001) Influence of acaricide treatments for *Varroa* control on the quality of honey and beeswax. *Apitalia* **28** (7–8), 21–24
- Bogdanov, S., Charrière, J., Imdorf, A., Kilcherman, V., Fluri, P. (2002) Determination of residues in honey after treatments under field conditions with formic and oxalic acid. *Apidologie* **33** (4), 399–409
- Bogdanov, S., Kilchenmann, V., Butikofer, U. (2003) Determination of acaricide residues in beeswax: Collaborative study. *Apiacta* **38** (3), 235–245
- Charrière, J.D. (1997) Potentiel et limites de l'emploi des acides organiques. *Sanitaire* **161**, 219–227
- Charrière, J.D., Imdorf, A. (2002) Oxalic acid treatment by trickling against *Varroa destructor*: recommendations for use in central Europe and under temperate climate conditions. *Bee World* **83** (2), 51–60
- Del Nozal, M., Bernal, J., Diego, J., Gomez, L., Ruiz, J., Higes, M. (2000) Determination of oxalate, sulfate and nitrate in honey and honeydew by ion-chromatography. *J. Chromat. A* **881**, 629–638
- Eguaras, M., Ruffinengo, S. (2006) Estrategias para el control de *Varroa*. Ed. Martin, Mar del Plata.
- Eguaras, M., del Hoyo, M., Palacio, A., Ruffinengo, S., Bedascarrasbure, E. (2001) A new product with formic acid for *Varroa jacobsoni* control. I Efficacy. *J. Vet. Med., (Series b)*, **48**, 11–14
- Eguaras, M., Palacio, A., Faverin, C., Basualdo, M., Del Hoyo, M., Velis, G., Bedascarrasbure, E. (2003) Efficacy of formic acid in gel for *Varroa* control in *Apis mellifera*: importance of the dispenser position inside the hive. *Vet. Parasitol.* **111**, 241–245
- Gregorc, A., Planinc, I. (2001) Acaricidal effect of oxalic acid in honeybee (*Apis mellifera*) colonies. *Apidologie* **32**, 333–340
- Gregorc, A., Planinc, I. (2002) The control of *Varroa destructor* using oxalic acid. *Vet. J* **163**, 306–310
- Hatjina, F., Haristos, L. (2005) Indirect effects of oxalic acid administered by trickling method on honey bee brood. *J. Apic. Res.* **44** (4), 172–174
- Higes, M., Meana, A., Suárez, M., Llorente, J. (1999) Negative long-term effects on bee colonies treated with oxalic acid against *Varroa jacobsoni* Oud. *Apidologie* **30**, 289–292
- Imdorf, A., Charrière, J.D., Kilchenmann, V., Bogdanov, S., Fluri, P. (2003) Alternative strategy in central Europe for the control of *Varroa destructor* in honey bee colonies. *Apiacta* **38**, 258–278
- Maggi, M., Ruffinengo, S., Damiani, N., Sardella, N., Eguaras, M. (2009) A first detection of *Varroa destructor* resistance to coumaphos in Argentina. *Exp. App. Acarol.* **47** (4), 317–320
- Maggi, M., Ruffinengo, S., Negri, P., Eguaras, M. (2010a) Resistance phenomena to amitraz from populations of the ectoparasitic mite *Varroa destructor* of Argentina. *Parasitol. Res.* **107** (5), 1189–1192
- Maggi, M., Ruffinengo, S., Gende, L., Sarlo, G., Bailac, P., Ponzi, M., Eguaras, M. (2010b) Laboratory evaluations of *Syzygium aromaticum* (L.) Merr. et Perry essential oil against *Varroa destructor*. *J. Ess. Oil Res* **22**, 119–122
- Maggi, M., Ruffinengo, S., Mendoza, Y., Ojeda, P., Ramallo, G., Floris, I., Eguaras, M. (2011) Susceptibility of *Varroa destructor* (Acari: Varroidae) to synthetic acaricides in Uruguay: *Varroa* mites' potential to develop acaricide resistance. *Parasitol. Res.* **108**, 815–821
- Maggi, M., Negri, P., Plischuk, S., Szawarski, N., De Piano, F., De Feudis, L., Eguaras, M., Audisio, C. (2013) Effects of the organic acids produced by a lactic acid bacterium in *Apis mellifera* colony development, *Nosema ceranae* control and fumagillin efficiency. *Vet. Microbiol.* . doi:10.1016/j.vetmic.2013.07.030
- Marinelli, E., Formato, G., Vari, G., De Pace, F.M. (2006) *Varroa* control using cellulose strips soaked in oxalic acid water solution. *Apiacta* **41**, 54–59
- Medici, S., Maggi, M., Sarlo, E., Ruffinengo, S., Marioli, J., Eguaras, M. (2015) Presence of synthetic acaricides in beeswax and its relationship with the development of resistance in *Varroa destructor*. *J. Apic. Res.* *in press*
- Mutinelli, F., Baggio, A., Capolongo, F., Piro, R., Prandin, L., Biaison, L. (1997) A scientific note on oxalic acid by topical application for the control of varroosis. *Apidologie* **28** (6), 461–462
- Nanetti, A., Büchler, R., Charriere, J.-D., Fries, I., Helland, S., Imdorf, A., Korpela, S., Kristiansen, P. (2003) Oxalic acid treatments for *Varroa* control (review). *Apiacta* **38**, 81–87
- Negri, P., Maggi, M., Ramirez, L., De Feudis, L., Szawarski, N., Eguaras, M., Lamattina, L. (2015) Abscisic acid enhances the immune response in *Apis mellifera* and contributes to the colony fitness. *Apidologie* **46** (4), 542–557
- Rademacher, E., Harz, M. (2006) Oxalic acid for the control of varroosis in honey bee colonies – a review. *Apidologie* **37**, 98–120
- Radetzki, T. (1994) Oxalsäure, eine weitere organische Säure zur Varroabehandlung. *Allgemeine Deutsche Imkerzeitung* **28** (12), 11–15
- Radetzki, T., Bärmann, M. (2001) Verdampfungsverfahren mit Oxalsäure. Feldversuch mit 1509 Völkern im Jahr 2000. *Allgemeine Deutsche Imkerzeitung* **35** (9), 20–23
- Ruffinengo, S., Maggi, M., Fuselli, S., De Piano, F., Negri, P., Brasesco, C., Satta, A., Floris, I., Eguaras, E. (2014) Bioactivity of microencapsulated essential oils and perspectives of their use in the control of *Varroa destructor*. *Bull. Insectol.* **67** (1), 81–86

- Sammataro, D., Untalan, P., Guerro, F., Finley, J. (2005) The resistance of *Varroa* mites (Acari: Varroidae) to acaricides and the presence of esterase. *Int. J. Acarol.* **31** (1), 67–74
- Segur, J., Obrstar, H. (1951) Viscosity of glycerol and its aqueous solutions. *Ind. Eng. Chem.* **43** (9), 2117–2120
- Wallner, K. (1999) Varroacids and their residues in bee products. *Apidologie* **30**, 235–248