

RESEARCH

Open Access



Trypanosoma cruzi infection in *Triatoma infestans* and high levels of human–vector contact across a rural-to-urban gradient in the Argentine Chaco

Alejandra Alvedro^{1,2}, María Sol Gaspe^{1,2}, Hannah Milbourn³, Natalia Paula Macchiaverna^{1,2}, Mariano Alberto Laiño^{1,2}, Gustavo Fabián Enriquez^{1,2}, Ricardo Esteban Gürtler^{1,2} and Marta Victoria Cardinal^{1,2*}

Abstract

Background: Peri-urban and urban settings have recently gained more prominence in studies on vector-borne transmission of *Trypanosoma cruzi* due to sustained rural-to-urban migrations and reports of urban infestations with triatomines. Prompted by the finding of *Triatoma infestans* across the rural-to-urban gradient in Avia Terai, an endemic municipality of the Argentine Chaco, we assessed selected components of domestic transmission risk in order to determine its variation across the gradient.

Methods: A baseline vector survey was conducted between October 2015 and March 2016, following which we used multistage random sampling to select a representative sample of *T. infestans* at the municipal level. We assessed *T. cruzi* infection and blood-feeding sources of 561 insects collected from 109 houses using kinetoplast DNA-PCR assays and direct enzyme-linked immunosorbent assays, respectively. We stratified triatomines according to their collection site (domestic or peridomestic), and we further categorized peridomestic sites in ecotopes of low- or high-risk for *T. cruzi* infection.

Results: The overall adjusted prevalence of *T. cruzi*-infected *T. infestans* was 1.8% (95% confidence interval [CI] 1.3–2.3) and did not differ between peri-urban (1.7%) and rural (2.2%) environments. No infection was detected in bugs captured in the urban setting; rather, infected triatomines were mainly collected in rural and peri-urban domiciles, occurring in 8% of *T. infestans*-infested houses. The main blood-feeding sources of domestic and peridomestic triatomines across the gradient were humans and chickens, respectively. The proportion of triatomines that had fed on humans did not differ between peri-urban (62.5%) and rural (65.7%) domiciles, peaking in the few domestic triatomines collected in urban houses and decreasing significantly with an increasing proportion of chicken- and dog- or cat-fed bugs. The relative odds ratio (OR) of having a *T. cruzi* infection was nearly threefold higher in bugs having a blood meal on humans (OR 3.15), dogs (OR 2.80) or cats (OR: 4.02) in a Firth-penalized multiple logistic model.

Conclusions: *Trypanosoma cruzi* transmission was likely occurring both in peri-urban and rural houses of Avia Terai. Widespread infestation in a third of urban blocks combined with high levels of human–triatomine contact in the few infested domiciles implies a threat to urban inhabitants. Vector control strategies and surveillance originally conceived for rural areas should be tailored to peri-urban and urban settings in order to achieve sustainable interruption of domestic transmission in the Chaco region.

*Correspondence: mvcardinal@ege.fcen.uba.ar; mvictoriacardinal@gmail.com

¹ Laboratorio de Eco-Epidemiología, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina
Full list of author information is available at the end of the article



© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords: *Trypanosoma cruzi*, *Triatoma infestans*, Urbanization, Blood-feeding sources

Introduction

Chagas disease, a neglected tropical disease endemic to the Americas, affects about 6–7 million people [1]. Its etiological agent, the protozoan *Trypanosoma cruzi*, is associated with cardiac and digestive syndromes in a third of infected patients and causes nearly 10,000 premature deaths each year [2]. Vector-borne transmission mediated by triatomine bugs has been traditionally the main route of transmission in endemic areas and still occurs in 21 countries [1].

Originally linked to impoverished rural environments where mud-and-thatch houses predominated and harbored large bug colonies, the landscape of Chagas disease has changed over the years. Migration has driven originated the globalization and urbanization of Chagas disease, with big cities worldwide now witnessing both the occurrence of congenital transmission [3] and of infected people [4, 5]. This phenomenon is not unique to Chagas disease; other diseases, such as visceral leishmaniasis and schistosomiasis, are experiencing the same shifting epidemiology [6–8].

Urban infestations and vector-borne transmission of *T. cruzi* have increasingly been reported in endemic regions throughout Central and South America [9–16]. In Cochabamba (Bolivia) and Arequipa (Peru), peri-urban infestations with *Triatoma infestans*, the main vector in the region, have been linked to the immigration of farmers from rural areas [13, 17]. In these peri-urban areas, *T. cruzi* infection in *T. infestans* ranged from 14 to 19% and was associated with the presence of guinea pigs [10, 17].

Domestic *T. cruzi* transmission to humans depends on the contact rates between infected triatomines and susceptible hosts. Ecological, environmental, socio-economic and cultural factors favor or hinder these encounters [18]. Housing type, building materials, husbandry practices and the number and density of domestic hosts may boost triatomine population size and hence facilitate transmission [19, 20]. The presence and increased number of *T. cruzi*-infected dogs and cats (the main domestic reservoir hosts) increase the infection risk of householders [21]. On the contrary, appropriate domestic animal management, regular cleaning activities, domestic insecticide use, window or door screening and plastering are protective factors [22–24].

As part of an intervention program aimed at suppressing house infestation with *T. infestans* in a sustainable manner and reducing disease burden in Avia Terai, Argentine Chaco, we conducted a survey that found a decreasing prevalence of house infestation (i.e. domestic

or/and peridomestic infestation) with *T. infestans*, ranging from 42.4% in rural houses to 22.5 and 4.5% in peri-urban and urban areas, respectively [25]. Domestic infestation, defined as the occurrence of *T. infestans* in human sleeping quarters or domiciles, also decreased across the rural-to-urban gradient, from 9.9 to 1.1%, respectively, as did the household number of domestic animals and peridomestic structures. *Triatoma infestans* seemed to be long established in Avia Terai, unlike in Arequipa and Cochabamba [17, 25, 26].

In the study reported here, we evaluated three selected components of *T. cruzi* transmission risk across a rural-to-urban gradient: (i) parasite infection prevalence in domestic and peridomestic *T. infestans*; (ii) triatomine bloodmeal sources; and (iii) degree of human-vector contact via the human blood index (HBI), defined as the proportion of reactive triatomines that fed on humans [27, 28]. Based on a decreasing prevalence of house and domestic infestation with *T. infestans* across the rural-to-urban gradient, we hypothesized that the prevalence of *T. cruzi* infection in *T. infestans* and the HBI in domestic bug populations would decrease across the rural-to-urban gradient. Secondly, we hypothesized that the presence of dogs, cats and chickens would decrease the HBI throughout the three environments.

Methods

Study area

Fieldwork was conducted in the municipality of Avia Terai (26°42' S, 60°44' W), Chaco Province, Argentina, which has been described elsewhere [25]. The last house spraying campaign with pyrethroid insecticides conducted by Chagas vector control personnel was carried out in 2011–2013 (2–4 years prior to our survey) and did not achieve full coverage. For the purpose of this study, we divided the municipality into three environments based on house arrangement patterns: urban (blocks), peri-urban (blocks and dispersed houses) and rural (dispersed houses) [25]. The peri-urban environment was highly heterogeneous and was further subclassified into established and recent neighborhoods [25].

Study design

A cross-sectional survey was conducted in all environments to evaluate house and domestic infestation with triatomines at baseline between October 2015 and March 2016, as described elsewhere [25]. In total, 2296 inhabited houses were surveyed. Domestic and peridomestic ecotopes were searched for triatomines if dwellers had

reported having seen triatomines or had peridomestic structures housing animals, and in one of every three houses that did not fulfil both conditions. Timed manual searches for bugs were performed for 15 min in each domestic and peridomestic structure by technicians of the Chagas control program with the aid of a flushing-out agent (0.2% tetramethrin; Espacial, Buenos Aires, Argentina). All rural houses and infested houses and their adjacent ones in peri-urban and urban environments were immediately sprayed with insecticide after the search for triatomines. A standard dose of beta-cypermethrin (50 mg/m²; Sipertrin, Chemotecnica, Argentina) in domiciles and a double dose in peridomestic were used to afford prolonged residual efficacy. Recently built peri-urban neighborhoods were excluded from the present analysis since no triatomines were found there [25].

Additional bug collections were attempted by technicians during and after insecticide applications or after the timed manual searches, and separately by householders; these insects were kept separately and labeled accordingly. These additional bug collections were performed to increase sample sizes for other purposes or to enhance triatomine detection; they were only used for abundance estimation in five houses (see text below). All triatomines were identified to species, sex and stage, and immediately frozen at -20 °C. Two other triatomine species captured in the study area, *Triatoma sordida* and *Panstrongylus geniculatus*, were excluded from the current analysis [25].

Bug sampling design

The bugs collected were classified according to where they had been collected (domestic or peridomestic ecotopes). Peridomestic ecotopes were further subdivided into two categories according to the probability of bug infection with *T. cruzi*: (i) those with a high risk of bug infection due to the presence of dogs, cats and rodents, including kitchens, granaries, mud ovens, storerooms and kennels; and (ii) those with a low risk of bug infection, mainly associated with poultry, which are refractory to *T. cruzi* (i.e. chicken nests, trees where chickens roosted, chicken coops) and goat and pig corrals.

A multistage sampling plan was used to select bugs for dissection. Only third-instar nymphs to adult bugs were examined. In the urban environment, we examined all triatomines collected in domiciles and high-risk peridomestic ecotopes, whereas in the peri-urban and rural environments we randomly selected up to ten and five triatomines per site, respectively. Additionally, in the three environments we randomly selected 25% of infested houses with low-risk peridomestic ecotopes; up to 5 bugs per site were examined for infection in each of these houses. If more than one low-risk ecotope was infested at a given house, we randomly examined one ecotope. We

separated the rectal ampoule, blood meal contents and the rest of each individual triatomine into labeled microtubes, as described previously [29].

Diagnosis of *T. cruzi* infection

Rectal ampoules were preserved in 50 µl of sterile water for parasite DNA extraction, boiled for 10 min and the DNA extracted using DNAzol® (Invitrogen, Carlsbad, CA, USA), following a previously described protocol [29]. A hot-start PCR targeting a 330-bp amplicon of the kinetoplast minicircle was employed [30]. PCR products were run in a 3% agarose gel and visualized under UV employing Gel Red® (Biotium, Hayward, CA USA).

Blood meal identification

Blood meal contents were stored in microtubes containing phosphate buffered saline with crystal violet. Identification of the blood meal was carried out using a direct enzyme-linked immunosorbent assay (ELISA) as described by Gürtler et al. [31]. The antisera employed in the ELISA assays included the most frequent (peri-)domestic blood meal sources of *T. infestans*: human, dog, cat, chicken, mice, goat and pig. We considered a bug as reactive when it was positive against any of the antisera tested in the ELISA assay and as a mixed blood meal when it was reactive to at least two different host species. The infective blood meal index was estimated as the percentage of *T. cruzi*-infected bugs with an identified blood meal on host X relative to the number of bugs examined for *T. cruzi* infection that had an identified blood meal on host X [32].

Data analysis

Agresti-Coull binomial 95% confidence intervals (95% CI) were calculated for the percentage of houses harboring infected triatomines, bug infection prevalence and HBI. Wilson 95% CI were calculated when the number of examined bugs was < 40 or the proportions were near 0 or 1 [33]. Proportions were compared using Fisher's exact or Pearson's χ^2 tests. Based on the observed prevalence of infection by bug collection ecotope (domicile, low-risk and high-risk peridomestic ecotope), we estimated the number of infected bugs that would have been detected if all collected third-instar nymphs and later stages had been examined. With these figures we estimated the adjusted prevalence of infection for each environment (i.e. urban, peri-urban and rural) and for the entire study area.

The relative abundance was calculated as the number of live triatomines collected during the timed manual searches per unit of search effort (i.e. 15 person-min per site). As an exception, in three rural and two peri-urban houses with examined bugs that had not been collected

during the timed manual searches, we estimated bug abundance considering these additional collections.

We examined the relationship between human blood-feeding or bug infection with *T. cruzi* (binary response variables) and *a priori* selected predictors (see text below) by means of a random-intercept and a Firth-penalized logistic regression, respectively. Odds ratios (OR) and their 95% CI were calculated using the 'sjPlot' package [34]. Numerical variables were categorized according to their quartiles or medians. All predictors were established *a priori* based on existing empirical evidence [28, 31, 35]. Host numbers were obtained from house-to-house surveys [25].

For the HBI model, we performed the regression analyses clustered by house using the 'lmer' function implemented in the 'lme4' package for R (R Foundation for Statistical Computing, Vienna, Austria). We included as predictors: the relative abundance of domestic bugs, bug stage (with four levels: third- or fourth-instar nymphs, fifth instars, adult males and adult females), the number of human residents, the number of dogs and/or cats in the household, dog or cat blood index and the chicken blood index (expressed as proportions), and only considered reactive domestic bugs (total: 157 bugs from 35 houses).

In order to identify risk factors associated with *T. cruzi* infection in triatomines we employed univariate and multiple Firth-penalized logistic regressions implemented in the 'logistf' package in R (version 2.15.1) [36, 37] given that few infested bugs were detected and complete separation of our dataset was observed for the urban environment. We included as predictors: relative bug abundance, type of environment (urban, peri-urban or rural), bug stage (with four levels: third- or fourth-instar nymphs, fifth instars, adult males and adult females), the number of human residents, the number of dogs and/or cats in the household and whether or not the bug had an identified blood meal on humans, dogs, cats or chickens (binary variables). This model only included examined bugs from domiciles and high-risk peridomestic ecotopes; thus, it comprised 408 bugs from 76 houses. Low-risk peridomestic ecotopes were excluded from this model because no infested bug was found in these ecotopes. We evaluated and found no significant effects of the interaction between type of environment and each explanatory variable on bug infection. No multicollinearity was detected (variance inflation factor < 3.9 for every variable). The global fit of the random-effects logistic model to the data was analyzed using the Hosmer–Lemeshow test with the 'Resource Selection' package [38].

The spatial distribution of bug infection was assessed through global and local point pattern analyses at the house level (rural environment) or using a combination

of block and house levels for peri-urban neighborhoods. Global spatial analyses of house (rural) and house or block (peri-urban) infection were performed using the K-function implemented in Programita [39]. Random labeling was used to test the null hypothesis of random events among the fixed spatial distribution of infested houses (or blocks). We set cell sizes of 300 m for rural houses and 50 m for urban or peri-urban houses or blocks, and maximum distances of search of 15 and 0.6 km, respectively. We computed the 95% confidence envelopes using 999 simulations, as reported in Gaspe et al. [25]. Local spatial aggregation of bug infection was tested using the Getis statistic (G^*) [40] implemented in PPA [41], with the same parameters as for the global analysis.

Results

Infection prevalence

In total, 561 *T. infestans* were selected and examined for infection from among the 4408 bugs collected (Table 1). The dissected bugs were collected from 66.7, 100 and 96.2% of the infested urban, peri-urban and rural domiciles, respectively and from 87.5, 85.7 and 89.7% of the urban, peri-urban and rural houses with infested high-risk peridomestic ecotopes (Table 1). Houses not examined for bug infection only harbored first or second-instar nymphs or bugs that were preserved for other studies (including 1 urban, 6 peri-urban and 5 rural houses).

The overall prevalence of infection was 5.7%. No infested bug was detected in the urban environment whereas no significant difference in the prevalence of infection was found between the peri-urban (8.1%) and rural (6.7%) environments (Fisher's exact test: $df=1$, $P=0.7$) (Table 1). Infection prevalence peaked in domiciles, followed by high-risk peridomestic ecotopes and was non-existent in low-risk peridomestic ecotopes in both peri-urban and rural environments. The adjusted prevalence of bug infection (estimated as if all collected bugs had been examined) was 1.8% (95% CI 1.3–2.3), and increased from 0% in the urban environment to 1.7% (95% CI 1.1–2.6) and 2.2% (95% CI 1.6–2.8) in the peri-urban and rural environments, respectively.

Bug infection was found in 11.9% of 109 infested houses examined for infection. The percentage of infested houses with at least one infested bug in the peri-urban (16.1%) and rural (13.6%) environments was not significantly different (Fisher's exact test: $df=1$, $P=1.0$) (Table 1); if all collected bugs had been examined, the adjusted prevalence of infection decreased to 8.2% (95% CI 6.7–24.4) and 7.9% (95% CI 3.2–12.8), respectively. Nearly half (53.8%) of the houses where infection was detected harbored only one infested triatomine. In houses with infested domiciles or high-risk peridomestic

ecotopes, the mean number of *T. cruzi*-infected triatomines per house ranged from 0 (urban) to 0.5 (peri-urban) to 0.4 (rural). Only two rural houses harbored infected triatomines in more than one site within the house compound. No significant global or local spatial clustering of *T. cruzi* infection was detected in the peri-urban and rural environments (Fig. 1).

The number of triatomines examined for infection increased with higher bug stage across the rural-to-urban gradient, with adult bugs accounting for 44.9% of all examined triatomines (252/561 insects) (Fig. 2). There was no significant statistical difference in infection prevalence among bug stages for all types of environments combined ($\chi^2 = 2.3$, $df = 3$, $P = 0.5$) (Fig. 2).

Blood-feeding sources

Most triatomines examined for blood meal sources were reactive (85.4%) to at least one of the antisera tested (Additional file 1: Table S1). Reactivity (i.e. the percentage of ELISA-reactive bugs) differed significantly among environments ($\chi^2 = 18.2$, $df = 2$, $P < 0.001$) and was significantly higher in peri-urban (93.3%) than in rural (85.6%) and urban environments (74.6%). Reactivity peaked in low-risk peridomestic ecotopes (95.2%) (Additional file 1: Table S1). Among reactive *T. infestans* with mixed blood meals (18.0%), 93.0% had fed on two different host

species, and the remaining (7.0%) bugs were positive for three host species.

The most frequent blood meal source of *T. infestans* across environments was chickens, followed by humans and dogs (Fig. 3a). The rural environment was the only one that showed a full range of host types, having reactive samples for all the seven species tested (Fig. 3a). In domiciles across all environments, humans were the main blood meal source (urban: 100.0%, peri-urban: 62.5%, rural: 65.7%) and cats were the secondary source (20.0, 22.9, 11.1%, respectively) (Fig. 3b). Dogs and chickens were also secondary blood meal sources of domestic bugs in the peri-urban and rural environments: rural houses exhibited twice as many dog-fed domestic bugs (21.2%) than peri-urban ones (10.4%) (Fig. 3b). The percentage of chicken-fed domestic bugs was threefold higher in rural houses (22.2%) than in peri-urban ones (8.3%). The main blood meal source in peridomestic ecotopes was chickens, ranging from 67.4 to 98.6% of the reactive bugs across environments (Fig. 3c, d). Dogs were the secondary blood meal source in peridomestic ecotopes.

Cats, pigs, goats and mice contributed little as blood meal sources of peridomestic bugs. In the urban environment, pig (15.2%) and goat (2.2%) blood meals were detected in peridomestic ecotopes (Fig. 3). Pig blood meals were detected in four adult triatomines and three nymphs collected in a mud oven, a kennel and a woodpile

Table 1 Distribution of *Trypanosoma cruzi* infection in *Triatoma infestans* by environment and type of ecotope, Avia Terai, Chaco, 2015–2016

Environment and ecotope (no. of houses inspected) ^a	Bug level			House level		
	% of bugs examined (no. collected)	% of bugs infected (no. examined)	95% CI	% examined for infection (no. infested)	% harboring infected bugs (no. examined)	95% CI
Urban (416)						
Domiciles	85.7 (14)	0.0 (12)	0.0–24.3	66.7 (3)	0.0 (2)	0.0–65.8
Peridomestic HR	100.0 (67)	0.0 (67)	0.0–5.4	87.5 (8)	0.0 (7)	0.0–32.2
Peridomestic LR	7.2 (485)	0.0 (35)	0.0–9.9	25.6 (39)	0.0 (10)	0.0–27.8
Total	20.1 (566)	0.0 (114)	0.0–3.2	38.0 (50)	0.0 (19)	0.0–16.8
Peri-urban (210)						
Domiciles	98.1 (53)	17.3 (52)	9.2–30.0	100.0 (14)	21.4 (14)	6.8–48.3
Peridomestic HR	24.3 (226)	5.5 (55)	1.9–14.9	85.7 (14)	18.2 (12)	4.0–48.9
Peridomestic LR	4.3 (974)	0.0 (42)	0.0–8.4	24.3 (37)	0.0 (9)	0.0–29.9
Total	11.9 (1253)	8.1 (149)	4.7–13.6	51.7 (60)	16.1 (31)	7.1–32.6
Rural (276)						
Domiciles	50.9 (228)	9.5 (116)	5.2–16.3	96.2 (26)	24.0 (25)	11.2–43.2
Peridomestic HR	29.6 (358)	8.5 (106)	4.4–15.6	89.7 (29)	15.4 (26)	5.8–35.2
Peridomestic LR	3.8 (2003)	0.0 (76)	0.0–4.8	26.6 (64)	0.0 (17)	0.0–18.4
Total	11.5 (2589)	6.7 (298)	4.4–10.1	57.3 (103)	13.6 (59)	7.0–25.5
Total (902)	12.7 (4408)	5.7 (561)	4.1–8.0	51.2 (213)	11.9 (109)	7.7–20.5

CI, Confidence interval


^a Peridomestic HR, Peridomestic ecotopes with high risk of bug infection; peridomestic LR, peridomestic ecotopes with low risk of bug infection



from three urban houses having no pigs, although pigs were recorded in neighboring uninfested houses. The only goat-fed adult triatomine detected in the urban environment was collected in a house with no goats. However, a neighboring family had two infested houses, one in the urban environment and another in the rural one. In the rural house goat-fed bugs were detected.

Human blood index

The HBI of domestic *T. infestans* peaked in the urban environment, where the few bugs examined contained human blood. No significant difference in HBI was found between rural (65.7%) and peri-urban domiciles (62.5%) (Fisher's exact test: $df=1$, $P=0.7$). The relative abundance of domestic bugs was not significantly



associated with the HBI in the model clustered by house and adjusted for bug stage. The HBI significantly decreased with increasing proportions of dog- or cat- and chicken-fed bugs in a logistic regression (OR 0.004, 95% CI 0.00–0.13 and OR 0.03; 95% CI 0.00–0.86, respectively) (Additional file 2: Table S2; Fig. 4). The number of human residents exerted a significant positive effect on the HBI although with large uncertainty on its magnitude (OR 8.93, 95% CI 1.12–71.17 for domiciles with 4 or 5 humans). No significant effect of the household number of dogs or cats was detected (Hosmer–Lemeshow χ^2 test = 8.88, $df = 8$, $P = 0.35$). The type of environment was not included as a predictor in this model due to perfect separation of urban triatomines.

Relationship between *T. cruzi* infection and blood meal sources

The blood meal source was identified in 87.5% of 32 *T. cruzi*-positive bugs. Mixed blood meals were two-fold more frequent in infected bugs (31.3%; 95% CI 17.6–48.6%) than in uninfected ones (16.9%; 95% CI 13.7–20.6%). In peri-urban houses, the main blood meal sources of infected *T. infestans* were humans and dogs (infective blood meal indices: 29.0 and 12.1%, respectively), whereas in rural houses, cat-fed bugs displayed peak infective blood meal indices (27.3%) (Table 2). No infection was detected among bugs fed on goats, pigs or mice (i.e. “other”).


Firth-penalized multiple logistic regression showed that the relative odds ratio of *T. cruzi* infection was 16.6- and 13.2-fold greater for bugs in the peri-urban and rural environments, respectively, with respect to urban bugs, although confidence intervals were large. *Trypanosoma*

cruzi infection in *T. infestans* collected in domiciles and high-risk peridomestic ecotopes was significantly associated with having a blood meal on humans, dogs or cats (Table 3). The relative odds ratio of having a *T. cruzi* infection was nearly three- to fourfold higher in bugs having a blood meal on humans (OR 3.15, 95% CI 1.27–8.17), dogs (OR 2.80, 95% CI 1.03–7.31) or cats (OR 4.02, 95% CI 1.46–10.39) than in bugs not fed on these hosts. In univariate analyses, the relative odds ratio of infection was significantly associated with the household number of dogs and cats and was ninefold higher in houses harboring > 8 dogs and cats than in houses with ≤ 3 cats and/or dogs (OR 9.49, 95% CI 3.49–31.66).

Discussion

Based on the prevalence of *T. cruzi* infection in triatomines and on human–vector contact rates measured by HBI (the risk components examined in this study), infested peri-urban and rural houses would be under a similar risk of transmission if other risk components (e.g. prevalence of human, cat and dog infection) were equal among environments. These patterns partially contradict the expected decrease in the prevalence of bug infection across the rural-to-urban gradient given the decreasing trends in house infestation and bug abundance. Our study reveals (peri)domestic *T. cruzi* transmission was likely occurring in peri-urban and rural houses from Avia Terai municipality 2–4 years after community-wide residual spraying with pyrethroid insecticides. Unexpectedly, infested urban houses had no infected *T. infestans* despite frequent occurrence of human infection and human–vector contact in the few domestic bugs collected (see below).

Trypanosoma cruzi infection was detected in 2% of all triatomines and in 8% of the infested houses. Lack of spatial aggregation of *T. cruzi*-infected bugs suggests independent transmission foci. These results evince a low prevalence but spatially spread distribution of *T. cruzi* infection in *T. infestans* in the context of rather recent insecticide applications by vector control personnel. For example, widespread bug infection in peri-urban scenarios were also reported for San Juan Province, western Argentina [42], in Chaco province in the 1940s [43] and in the outskirts of Cochabamba [17]. In contrast, *T. cruzi* infection in *T. infestans* in peri-urban Arequipa was spatially aggregated and restricted to certain locations [10]. Spatially spread house infestation and *T. cruzi* infection encompassing peri-urban areas, as in Avia Terai, pose additional challenges to vector elimination efforts originally conceived for rural areas. In addition to the much higher number of target houses and insensitivity of triatomine detection methods, the occurrence of closed or



nonparticipating households may hinder access to premises in peri-urban and urban areas.

In our study, peri-urban and rural environments had a similar prevalence of *T. cruzi* infection and similar percentages of infested domiciles (21–24%) and high-risk peridomestic ecotopes (15–18%) harboring infected bugs. These patterns remained despite rural houses having higher house infestation and higher bug abundance. This can be related to the high proportional contribution of low-risk peridomestic ecotopes to rural infestation and

bug abundance [25] (Table 1), but in which no infected bug was detected.

The main blood meal source detected in domestic bugs was humans, whereas chickens and dogs were the main blood meal sources in the peridomestic ecotopes, as in other settings [27, 28, 44–47]. Domestic *T. infestans* collected in the Paraguayan Chaco nearly exclusively fed on humans [48], in a context of strong vector control activities. In Avia Terai, chickens were relevant blood meal sources in all ecotopes across the rural-to-urban gradient, with the exception of urban domiciles where we




Table 2 Distribution of *T. cruzi* infection in *T. infestans* according to blood meal source, Avia Terai, Chaco, 2015–2016

Blood meal source	Infective blood meal index (no. reactive bugs)		
	Peri-urban environment	Rural environment	Total
Humans	29.0 (31)	9.0 (67)	15.3 (98)
Dogs	12.1 (33)	8.2 (49)	9.8 (82)
Cats	6.7 (15)	27.3 (22)	18.9 (37)
Chickens	1.3 (79) ^a	5.2 (153) ^b	3.9 (232)
Others	0.0 (6)	0.0 (24)	0.0 (30)

The infective blood meal index was estimated as the percentage of *T. cruzi*-infected bugs with an identified blood meal on host X relative to all bugs examined for *T. cruzi* infection that had an identified blood meal on host X

^a All infected triatomines exhibited a mixed blood meal with dogs, cats and humans

^b All infected triatomines exhibited a mixed blood meal with cats, with two exceptions

found bugs only fed on humans and cats. Chicken-fed bugs in domiciles are intimately linked to the widespread rural practice in the Chaco region of letting hens nest indoors for protection, especially in the spring and summer [19, 49], which corresponds to the timing of the current survey.

Cats apparently played a different epidemiological role across the gradient, serving as a highly infectious and frequent blood meal source in rural houses and urban

domiciles, respectively, while contributing little in peri-urban settings. These results are probably related to variations in domestic host abundance [25] and domestic animal management practices across the gradient.

The host-feeding patterns of *T. infestans* and other major triatomine vectors usually correlate closely with the main local resident host(s) in the collection ecotope [18]. The main blood meal sources identified in this study matched closely the *a priori* classification of ecotopes based on the expected occurrence of infected hosts. A high frequency of chicken-fed bugs was recorded in high-risk peridomestic ecotopes, and 31 (22%) insects from low-risk peridomestic ecotopes in rural and peri-urban houses had mixed blood meals on chickens and dogs or cats. These results point to the high mobility of triatomines and unconfined hosts. We also registered a wide diversity of blood meal sources in urban and peri-urban environments, including pigs and goats. Long-standing rural practices explain the presence of domestic ungulates in peri-urban and urban settings [25]. Additionally, active bug dispersal and passive bug transport from the rural environment may explain the few blood meals on host species not recorded at the bug collection house. Although most urban infestations consisted of triatomine colonies (88%) rather than invading adult triatomines [25], short distances between neighboring houses favor triatomine dispersal, as in Arequipa [50]. Frequent shifts from the refractory chickens or other

Table 3 Risk factors for *Trypanosoma cruzi* infection in *Triatoma infestans* collected in domiciles and high-risk peridomestic ecotopes, Avia Terai, Chaco

Predictor	Univariate analyses			Firth-penalized multiple logistic regression (n = 408)		
	Odds ratio	95% CI	P	Odds ratio	95% CI	P
Environment						
Urban	1.00	–	–	1.00	–	–
Peri-urban	20.81	2.67–2681.96	< 0.01*	16.60	2.07–2150.73	< 0.01*
Rural	16.10	2.17–2055.80	< 0.01*	13.23	1.73–1699.53	< 0.01*
Stage						
Third-fourth nymph	1.00	–	–	1.00	–	–
Fifth-instar nymph	0.67	0.26–1.65	0.38	0.63	0.24–1.65	0.35
Male	0.64	0.23–1.65	0.36	0.64	0.22–1.79	0.40
Female	0.45	0.15–1.26	0.13	0.39	0.12–1.12	0.08
Blood meal on						
Dog	1.85	0.77–4.10	0.16	2.80	1.03–7.31	0.04*
Cat	3.89	1.49–9.29	< 0.01*	4.02	1.46–10.39	< 0.01*
Human	2.69	1.29–5.54	< 0.01*	3.15	1.27–8.17	0.01*
Chicken	0.64	0.28–1.35	0.25	1.13	0.43–2.88	0.80
Site bug abundance (triatomines captured per one-person h)						
≤ 8	1.00	–	–	–	–	–
9–20	1.34	0.44–4.71	0.62	–	–	–
21–52	0.83	0.27–2.99	0.77	–	–	–
> 52	0.49	0.15–1.76	0.25	–	–	–
Number of dogs and cats per house ^a						
≤ 3	1.00	–	–	–	–	–
4–5	0.76	0.13–3.50	0.73	–	–	–
6–8	2.21	0.71–7.86	0.17	–	–	–
> 8	9.49	3.49–31.66	< 0.01*	–	–	–
Number of human residents per house ^b						
≤ 2	1.00	–	–	–	–	–
3	0.57	0.14–1.71	0.34	–	–	–
4–7	0.17	0.03–0.55	< 0.01*	–	–	–
> 7	0.60	0.23–1.44	0.26	–	–	–

*Significant at $P \leq 0.05$ ^a The number of dogs and cats was unknown for 5 examined bugs^b The number of residents was unknown for 98 examined bugs

uninfected hosts to dogs or cats, which frequently are infected and highly infectious, increase the risk of transmission [27, 51].

In our study, the urban environment had significantly fewer reactive triatomines than did rural and peri-urban ones. Bugs may have less access to blood-feeding in the urban environment due to a lower abundance of hosts, fewer peridomestic structures and/or better housing quality and animal management practices compared to rural and peri-urban environments [25, 52]. These specific factors observed in urban houses may also explain the unexpected lack of bug infection. Further investigations on *T. infestans* feeding frequency

across the rural-to-urban gradient may shed light on this issue.

Domestic triatomines in urban settings displayed the highest HBI (100%), the lowest prevalence of mixed blood meals (6%) and the absence of dog-fed bugs. These results may be related to urban householders restraining domestic animals, particularly dogs, from entering domiciles more frequently than peri-urban and rural dwellers. None of the two urban households with infested domiciles reported allowing dogs inside, whereas this occurred in 14% of infested peri-urban domiciles and in 27% of rural ones. Specifically designed studies are needed to confirm these findings since data on animal resting sites were missing for many houses.

Peri-urban and rural houses had similar frequencies of domestic bugs with mixed blood meals (19 and 22%) and HBI (62 and 65%), respectively. As expected, the proportions of dog- or cat-fed bugs (dog-cat blood index) and of chicken-fed bugs (chicken blood index) significantly decreased the HBI of domestic *T. infestans* [27, 28, 45]. These results further support the hypothesis that the occurrence of domestic animals indoors strongly affects human–vector contact rates and triatomine infection.

In contrast to our initial hypothesis, the HBI was not associated with the number of dogs or cats in a household in the multivariate analyses, which considered the absolute number of chickens, dogs and cats in each house instead of those actually resting or nesting indoors. Another limitation of our study was the lack of data on human overcrowding and other direct measures of host density more directly related to local host availability. These limitations hamper the estimation of the number of available hosts indoors and its relationship with transmission components. One strength of the current study is the molecular diagnosis of *T. cruzi* infection in a large sample of triatomines selected carefully across an environmental gradient. Molecular methods enhance the detection of light infections, a fraction of which are usually missed by optical microscopy. The false negative rate reached 10–13% in xenodiagnosis bugs examined by optical microscopy [53]. An additional advantage of using molecular diagnostic methods was the immediate freezing of collected triatomines upon completion of daily fieldwork. This at least partially explains the high reactivity of blood meal identification tests regardless of where the insects had been collected.

Triatomines that fed on humans in peri-urban houses and on cats in rural ones displayed high infective blood meal indices. Humans, dogs and cats contributed significantly to *T. infestans* infection in Avia Terai. Nearly 22–24% of urban households in Avia Terai reportedly harbored at least one *T. cruzi*-infected resident in 2012 [54]. A human serosurvey involving > 1000 people conducted in 2017 across the gradient showed that nearly 15% were seropositive for *T. cruzi* (Cardinal et al., unpublished). Hence, humans are also a potential source of *T. cruzi*. Although no data on the seroprevalence of *T. cruzi* infection in dogs and cats are currently available, the infective blood meal indices in our study suggest that infected dogs and cats were likely present.

Conclusions

The results of our study reveal that active transmission was likely occurring in peri-urban and rural settings of Avia Terai. Despite no infected bug being found in urban houses, the high human–vector contact detected,

the occurrence of low-density domestic colonies, widespread human *T. cruzi* infection and high density of households and hosts set a favorable scenario for vector-borne transmission if domestic infestations are allowed to thrive. In endemic countries, densely populated urban and peri-urban areas represent an unexpected challenge for vector control programs, mainly conceived for rural environments. Tailored vector control activities adapted to peri-urban and urban environments, strong community mobilization and sustained vector surveillance are required to effectively suppress the risk of domestic vector-borne transmission of *T. cruzi* and eliminate *T. infestans*.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13071-020-04534-z>.

Additional file 1: Table S1. Distribution of bloodmeal reactivity in *T. infestans* according to type of ecotope and environment.

Additional file 2: Table S2. Relationships between human blood index and selected factors in domestic *T. infestans*, Avia Terai, 2015–2016.

Additional file 3: Table S3. Dataset including infection status, bloodmeal source identification, environmental and demographic variables, Avia Terai, Chaco, 2015–2016.

Abbreviations

HR: High risk; LR: Low risk.

Acknowledgements

We thank the local communities for their warm welcome and commitment to the intervention program; municipal authorities and head of Avia Terai Hospital for providing field technicians and laboratory facilities for sample processing; and the National and Chaco Chagas disease control programs for field support and provision of dislodging aerosols and pyrethroid insecticides. We are grateful to Nicasio Vargas and Remigio Vargas for their valuable contribution to the field work over the last decade; to Roberto González and Martín Zárate, local vector control technicians, for their close collaboration and continuous support. HM thanks Tapan Bhattacharyya and Michael A. Miles for their supervision and training at LSHTM, Amanda Brunetti for support and encouragement in Argentina and the Bayer Fund for financial support.

Authors' contributions

AA, MVC, MSG, GFE and REG conceived and designed the experiments. AA, HM, MVC, NPM and MAL performed the experiments. AA, MVC, MSG and REG analysed the data. AA, HM, MVC, MSG, GFE, NPM, MAL and REG wrote the paper. All authors read and approved the final manuscript.

Funding

This study received financial support from Fundación Bunge y Born (2015–2018), and additional support from Agencia Nacional de Promoción Científica y Tecnológica (PICT 2015-2921), and Universidad de Buenos Aires (UBACYT 200201701007798A, UBACYT 200201301008438A).

Availability of data and materials

The dataset supporting the conclusions of this article is included within the article and in the Additional files. The dataset generated during and/or analyzed during the present study are available in Additional file 3: Table S3.

Ethics approval and consent to participate

The study protocol was approved by the "Comité de Ética en Investigación Clínica" (Ethics Committee in Clinical Research) of Buenos Aires, Argentina (IRB No. 00001678; Protocol No. TW-01-004).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Laboratorio de Eco-Epidemiología, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina. ² Instituto de Ecología, Genética y Evolución de Buenos Aires (IEGEB), Consejo Nacional de Investigaciones Científicas y Técnicas–Universidad de Buenos Aires, Buenos Aires, Argentina. ³ London School of Hygiene and Tropical Medicine, London, UK.

Received: 21 September 2020 Accepted: 10 December 2020

Published online: 09 January 2021

References

- Pan American Health Organization. Guidelines for the diagnosis and treatment of Chagas disease. Washington, DC: Pan American Health Organization; 2019.
- Stanaway JD, Roth G. The burden of Chagas disease estimates and challenges. *Glob Heart*. 2015;10(3):139–44.
- Bisio M, Seidenstein ME, Burgos JM, Ballering G, Rizzo M, Pontoriero R, et al. Urbanization of congenital transmission of *Trypanosoma cruzi*: prospective polymerase chain reaction study in pregnancy. *Trans R Soc Trop Med Hyg*. 2011;105(10):543–9. <https://doi.org/10.1016/j.trstmh.2011.07.003>.
- Klein N, Hurwitz I, Durvasula R. Globalization of Chagas disease: a growing concern in nonendemic countries. *Epidemiol Res Int*. 2012;2012:1–13.
- Requena-Méndez A, Aldasoro E, de Lazzari E, Sicuri E, Brown M, Moore DAJ, et al. Prevalence of Chagas disease in Latin-American migrants living in Europe: a systematic review and meta-analysis. *PLoS Negl Trop Dis*. 2015;9(2):e0003540.
- Ximenes R, Southgate B, Smith PG, Guimarães NL. Socioeconomic determinants of schistosomiasis in an urban area in the Northeast of Brazil. *Pan Am J Public Health*. 2003;14(6):409–21.
- Alinol E, Getaz L, Stoll B, Chappuis F, Loutan L. Urbanisation and infectious diseases in a globalised world. *Lancet Infect Dis*. 2011;11(2):131–41. [https://doi.org/10.1016/S1473-3099\(10\)70223-1](https://doi.org/10.1016/S1473-3099(10)70223-1).
- Harhay MO, Olliaro PL, Costa DL, Costa CHN. Urban parasitology: visceral leishmaniasis in Brazil. *Trends Parasitol*. 2011;27(9):403–9.
- Vallvé SL, Rojo H, Wisnivesky-Colli C. Urban ecology of *Triatoma infestans* in San Juan, Argentina. *Mem Inst Oswaldo Cruz*. 1996;91(4):405–8.
- Levy MZ, Bowman NM, Kawai V, Waller LA, Cornejo del Carpio JG, Cordova Benzaquen E, et al. Periurban *Trypanosoma cruzi*-infected *Triatoma infestans*, Arequipa, Peru. *Emerg Infect Dis*. 2006;12(9):1345.
- Guzman-Tapia Y, Ramirez-Sierra MJ, Dumonteil E. Urban infestation by *Triatoma dimidiata* in the city of Mérida, Yucatán, México. *Vector-Borne Zoonotic Dis*. 2007;7(4):597–606.
- Medrano-Mercado N, Ugarte-Fernandez R, Butrón V, Uber-Busek S, Guerra HL, De Araújo-Jorge TC, et al. Urban transmission of Chagas disease in Cochabamba, Bolivia. *Mem Inst Oswaldo Cruz*. 2008;103(5):423–30.
- Bayer AM, Hunter GC, Gilman RH, Del Carpio JGC, Naquira C, Bern C, et al. Chagas disease, migration and community settlement patterns in Arequipa, Peru. *PLoS Negl Trop Dis*. 2009;3:e567.
- Carvalho DB, Almeida CE, Rocha CS, Gardim S, Mendonça VJ, Ribeiro AR, et al. A novel association between *Rhodnius neglectus* and the *Livistona australis* palm tree in an urban center foreshadowing the risk of Chagas disease transmission by vectorial invasions in Monte Alto City, São Paulo, Brazil. *Acta Trop*. 2014;130:35–8.
- Provecho YM, Gaspe MS, del Pilar FM, Enriquez GF, Weinberg D, Gürtler RE. The peri-urban interface and house infestation with *Triatoma infestans* in the Argentine Chaco: an underreported process? *Mem Inst Oswaldo Cruz*. 2014;109(7):923–34.
- Carvalho Parente C, Bezerra FSM, Parente PI, Dias-Neto RV, Xavier SCC, Ramos AN, et al. Community-based entomological surveillance reveals urban foci of chagas disease vectors in Sobral, State of Ceara, Northeastern Brazil. *PLoS ONE*. 2017;12(1):1–11.
- Albarracín-Veizaga H, De Carvalho ME, Do Nascimento EMM, Rodrigues VLCC, Casanova C, Barata JMS. Chagas disease in an area of recent occupation in Cochabamba, Bolivia. *Rev Saude Publica*. 1999;33(3):230–6.
- Gürtler R, Fernández M, Cardinal M. Eco-epidemiology of the domestic vector-borne transmission of *Trypanosoma cruzi*. In: Guarneri A, Lorenzo M, editors. *Triatominae: the biology of Chagas disease vectors*. New York: Springer; 2021 (in press).
- Cecere MC, Gürtler RE, Chuit R, Cohen JE. Factors limiting the domestic density of *Triatoma infestans* in north-west Argentina: a longitudinal study. *Bull World Health Organ*. 1998;76:373–84.
- Brenière SF, Villacis A, Aznar C. Vector transmission: how it works, what transmits, where it occurs. In: Telleria J, Tibayrenc M, editors. *American trypanosomiasis Chagas disease: one hundred years of research*. 2nd ed. Amsterdam: Elsevier; 2017. p. 497–516.
- Gürtler RE, Cardinal MV. Reservoir host competence and the role of domestic and commensal hosts in the transmission of *Trypanosoma cruzi*. *Acta Trop*. 2015;151:32–50.
- Campbell-Lendrum DH, Angulo VM, Esteban L, Tarazona Z, Parra GJ, Restrepo M, et al. House-level risk factors for triatomine infestation in Colombia. *Int J Epidemiol*. 2007;36(4):866–72.
- Waleckx E, Camara-Mejia J, Ramirez-Sierra MJ, Cruz-Chan V, Rosado-Valdado M, Vazquez-Narvaez S, et al. An innovative ecohealth intervention for Chagas disease vector control in Yucatan, Mexico. *Trans R Soc Trop Med Hyg*. 2014;109(2):143–9.
- Lardeux F, Depickère S, Aliaga C, Chavez T, Zambrana L. Experimental control of *Triatoma infestans* in poor rural villages of Bolivia through community participation. *Trans R Soc Trop Med Hyg*. 2015;109:150–8.
- Gaspe MS, Fernández MP, Cardinal MV, Enriquez GF, Rodríguez-Planes L, Macchiaverna NP, et al. Urbanisation, risk stratification and house infestation with a major vector of Chagas disease in an endemic municipality of the Argentine Chaco. *Parasites Vectors*. 2020;13(1):1–4.
- Cornejo Donayre A. Enfermedad de Chagas: Estado actual en el Perú. *An Fac Med Lima*. 1958;39:428–53.
- Gürtler RE, Cohen JE, Cecere MC, Chuit R. Shifting host choices of the vector of Chagas disease, *Triatoma infestans*, in relation to the availability of host in houses in North-West Argentina. *J Appl Ecol*. 1997;34(3):699–715.
- Gürtler RE, Cecere MC, Vázquez-Prokopec GM, Ceballos LA, Gurevitz JM, del Pilar FM, et al. Domestic animal hosts strongly influence human-feeding rates of the Chagas disease vector *Triatoma infestans* in Argentina. *PLoS Negl Trop Dis*. 2014;8(5):e2894.
- Maffey L, Cardinal MV, Ordóñez-Krasnowski PC, Lanati LA, Lauricella MA, Schijman AG, et al. Direct molecular identification of *Trypanosoma cruzi* discrete typing units in domestic and peridomestic *Triatoma infestans* and *Triatoma sordida* from the Argentine Chaco. *Parasitology*. 2012;139(12):1570–9.
- Schijman AG, Altcheh J, Burgos JM, Biancardi M, Bisio M, Levin MJ, et al. Aetiological treatment of congenital Chagas' disease diagnosed and monitored by the polymerase chain reaction. *J Antimicrob Chemother*. 2003;52(3):441–9.
- Gürtler RE, Ceballos LA, Ordóñez-Krasnowski P, Lanati LA, Stariolo R, Kitron U. Strong host-feeding preferences of the vector *Triatoma infestans* modified by vector density: implications for the epidemiology of Chagas disease. *PLoS Negl Trop Dis*. 2009;3(5):e447.
- Zarate LG, Zarate RJ, Tempelis CH, Goldsmith RS. The biology and behavior of *Triatoma barberi* (Hemiptera: Reduviidae) in Mexico. I. Blood meal sources and infection with *Trypanosoma cruzi*. *J Med Entomol*. 1980;17:103–16.
- Brown LD, Cai TT, Das GA. Interval estimation for a binomial proportion. *Stat Sci*. 2001;16(2):101–17.
- Lüdecke D. sjPlot: data visualization for statistics in social science. R package version 2.8.6; 2020. <https://CRAN.R-project.org/package=sjPlot>. Accessed 1 Sept 2020.
- Cardinal MV, Orozco MM, Enriquez GF, Ceballos LA, Gaspe MS, Alvarado-Otegui JA, et al. Heterogeneities in the ecoepidemiology of *Trypanosoma cruzi* infection in rural communities of the Argentinean Chaco. *Am J Trop Med Hyg*. 2014;90(6):1063–73.
- Heinze G, Ploner M. logistf: Firth's bias-reduced logistic regression. 2018. <https://cemsiis.meduniwien.ac.at/en/kb/science-research/software/statistical-software/flogistf/>. Accessed 20 May 2020.

37. R Development Core Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2012.
38. Lele S, Keim J, Solymos P. ResourceSelection: resource selection (probability) functions for use-availability data. 2017. <https://cran.r-project.org/web/packages/ResourceSelection/index.html>. Accessed 15 May 2020.
39. Wiegand T, Moloney KA. Rings, circles, and null-models for point pattern analysis in ecology. *Oikos*. 2004;104:209–29.
40. Getis A, Ord JK. Spatial analysis and modeling in a GIS environment. In: McMaster RB, Userly EL, editors. A research agenda for geographic information science. Boca Raton: CRC Press; 1996. p. 157–96.
41. Chen D, Getis A. Point pattern analysis. San Diego: San Diego State Univ; 1998.
42. Carrizo Páez R, Pickenhayn J, Carrizo PM. Chagas urbano en San Juan: Diagnóstico, revisión y propuesta para un sistema integrado de ataque. *Rev Argent Cardiol*. 2008;76:480–7.
43. Nájera L, Ruiz G. El índice de infección de la vinchuca doméstica (*Triatoma infestans*) en la zona periurbana de Resistencia. In: Sesión Científica de la Misión de Estudios de Patología Regional Distrito Chaco. Actas y Trabajos, Argentina. Resistencia: Universidad Nacional de Tucumán; 1949. p. 10–2.
44. Gürtler RE, Cecere MC, Vazquez DP, Chuit R, Cohen JE. Host-feeding patterns of domiciliary *Triatoma infestans* (Hemiptera: Reduviidae) in Northwest Argentina: seasonal and instar variation. *J Med Entomol*. 1996;33(1):15–26.
45. Ordóñez-Krasnowski PC, Lanati LA, Gaspe MS, Cardinal MV, Ceballos LA, Gürtler RE. Domestic host availability modifies human-triatomine contact and host shifts of the Chagas disease vector *Triatoma infestans* in the humid Argentine Chaco. *Med Vet Entomol*. 2020;34(4):459–69.
46. Rabinovich JE, Kitron UD, Obed Y, Yoshioka M, Gottdenker N, Chaves LF. Ecological patterns of blood-feeding by kissing-bugs (Hemiptera: Reduviidae: Triatominae). *Mem Inst Oswaldo Cruz*. 2011;106(4):479–94.
47. Wisnivesky-Colli C, Paulone I, Pérez A, Chuit R, Gualtieri J, Solarz N, et al. A new tool for continuous detection of the presence of triatomine bugs, vectors of Chagas disease, in rural households. *Medicina (B Aires)*. 1987;47(1):45–50.
48. Fraenkel S, Salvioni OD, de Arias AR, Arze VP, Rolón M, Ramirez N, et al. Identification of bloodmeal sources of triatomines captured in the Paraguayan Chaco region of South America by means of molecular biology analysis. *Pathog Glob Health*. 2020;114(1):30–9.
49. Gurevitz JM, Ceballos LA, Gaspe MS, Alvarado-Otegui JA, Enriquez GF, Kitron U, et al. Factors affecting infestation by *Triatoma infestans* in a rural area of the humid Chaco in Argentina: a multi-model inference approach. *PLoS Negl Trop Dis*. 2011;5(10):e1349.
50. Levy MZ, Barbu CM, Castillo-Neyra R, Quispe-Machaca VR, Ancca-Juarez J, Escalante-Mejia P, et al. Urbanization, land tenure security and vector-borne Chagas disease. *Proc R Soc B*. 2014;281:20141003.
51. Cohen JE, Gürtler RE. Modeling household transmission of American trypanosomiasis. *Science*. 2001;293(5530):694–8.
52. Hassell JM, Begon M, Ward MJ, Fèvre EM. Urbanization and disease emergence: dynamics at the wildlife–livestock–human interface. *Trends Ecol Evol*. 2017;32(1):55–67.
53. Enriquez GF, Bua J, Orozco MM, Wirth S, Schijman AG, Gürtler RE, et al. High levels of *Trypanosoma cruzi* DNA determined by qPCR and infectiousness to *Triatoma infestans* support dogs and cats are major sources of parasites for domestic transmission. *Infect Genet Evol*. 2014;25:36–43.
54. Genero S, Zorzo LR, Chaparro RM. Impact of an educational program related to basic knowledge of Chagas disease in a population of the Argentine Northeast. *Rev Fac Cien Med Univ Nac Cordoba*. 2018;75(3):168–75.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

