

# Dynamics of quantum information scrambling under decoherence effects

Federico D. Domínguez<sup>1,\*</sup> and Gonzalo A. Álvarez<sup>1,2,3,†</sup>

<sup>1</sup>*Centro Atómico Bariloche, CONICET, CNEA, S. C. de Bariloche, Argentina.*

<sup>2</sup>*Instituto Balseiro, CNEA, Universidad Nacional de Cuyo, S. C. de Bariloche, Argentina.*

<sup>3</sup>*Instituto de Nanociencia y Nanotecnología, CNEA, CONICET, S. C. de Bariloche, 8400, Argentina*

Developing quantum technologies requires the control and understanding of the non-equilibrium dynamics of quantum information in many-body systems. Local information propagates in the system by creating complex correlations known as information scrambling, as this process prevents extracting the information from local measurements. In this work, we develop a model adapted from solid-state NMR methods, to quantify the information scrambling. The scrambling is measured via time reversal Loschmidt Echoes (LE) and Multiple Quantum Coherences experiments that intrinsically contain imperfections. Considering these imperfections, we derive expressions for Out-of-Time-Order Correlations (OTOC) to quantify the observable information scrambling based on measuring the number of active spins where the information was spread. Based on the OTOC expressions, decoherence effects arise naturally by the effects of the non-reverted terms in the LE experiment inducing localization of the measurable degree of information scrambling. These effects define a localization cluster size for the observable number of active spins that determines a dynamical equilibrium. We contrast the model's predictions with quantum simulations performed with solid-state NMR experiments, that measure the information scrambling with time-reversal echoes with controlled imperfections. An excellent quantitative agreement is found with the dynamics of quantum information scrambling and its localization effects determined from the experimental data. The presented model and derived OTOCs set tools for quantifying the quantum information dynamics of large quantum systems (more than  $10^4$  spins) consistent with experimental implementations that intrinsically contain imperfections.

## I. INTRODUCTION

In a quantum many-body system, local information can propagate into many degrees of freedom, creating complex correlations as entanglement that prevents extracting the information from local measurements. This propagation process of information is known as *scrambling* [1, 2], and characterizing and understanding its dynamics is an outstanding problem that connects different fields of physics such as quantum statistical mechanics, cosmology and quantum information processing [3–6]. The complexity of information scrambling limits our ability to study many-body quantum systems and employ them in technological developments [7, 8]. As many-body systems generate high-order quantum correlations that spread over the system's degrees of freedom [4, 9–13], a high degree of scrambling is produced that make these large quantum states more sensitive to perturbations [9, 14–18].

An accepted measure of scrambling is the tripartite information, which is near maximally negative for quantum channels that scramble the information [19]. This measure proves that information scrambling is manifested through the decay of Out-of-Time Order Correlators (OTOC). OTOCs are special correlators

$$F_{AB}(t) = \frac{1}{|A| \cdot |B|} \langle A^\dagger(t) B^\dagger A(t) B \rangle \quad (1)$$

that quantifies the degree of non-commutativity between two local, initially commuting operators  $A$  and  $B$ , where  $A(t) = e^{-i\mathcal{H}t} A e^{i\mathcal{H}t}$  and  $\mathcal{H}$  is the Hamiltonian of the interacting system [5, 20, 21]. The decay of the correlator  $F_{AB}$  captures the essence of the quantum butterfly effect: a time-evolved local operator fails to commute with almost all the other local operators [22]. The dynamics of  $F_{AB}$  is a probe for quantum chaos [23–26].

The OTOC functions are more accessible experimentally than the tripartite information, therefore these correlators are employed for observing information scrambling in a wide variety of quantum systems using time reversal of quantum evolutions with Loschmidt Echo (LE) experiments [9, 15, 17, 18, 20, 21, 27–32]. A LE experiment is performed by first evolving forward in time with a Hamiltonian  $\mathcal{H}$  and then evolving backward in time with the Hamiltonian  $-\mathcal{H}$ . An ubiquitous problem of LE experiments is that the forward and backward Hamiltonian are not identical, therefore leading to the existence of non-reverted interactions, such as imperfections or environmental interactions, that introduce non-unitary decays [33–38]. These decoherence effects add up an additional decay to the OTOC functions that should not be confused with the information scrambling generated by the system interactions  $\mathcal{H}$ . Therefore, it is necessary to model the information scrambling in open quantum systems to correctly interpret scrambling measurements obtained from LE experiments [12, 15, 18, 39–43].

An experimental framework for observing the information scrambling dynamics in large interacting quantum systems is the Multiple Quantum Coherences (MQC) technique [9, 15, 17, 18, 20, 27–29], originally devel-

\* federico.dominguez@cab.cnea.gov.ar

† gonzalo.alvarez@cab.cnea.gov.ar

oped in solid-state Nuclear Magnetic Resonance (NMR) [44, 45]. Under specific dynamical models, the MQC experiments provide the number of correlated spins  $K$  [46–48], which is an alternative way of interpreting the information scrambling within the spin system [17, 18, 27]. Since solving exactly the dynamics of general and large spin-systems is not possible with present technologies, a variety of models were developed to describe the dynamics of  $K$  during the quantum evolution with MQC experiments [44, 45, 49–55]. Levy and Gleason [50] propose a method to simplify the complex dynamics of a multi-spin system, by describing the quantum state as a mixture of average operators of  $L$  correlated spins. An initial excitation on local spin magnetization operators flows between these average operators with rates determined by the Liouville-von Neumann equation. Then the number of correlated spins is determined by a weighted average of the average operators of  $L$  spins that contributes to the system state. This model successfully explains the growth of the cluster size of correlated spins in different crystalline samples [50, 56–58].

In this work, we revisit the Levy and Gleason model and adapt it to quantify the information scrambling of a local excitation using LE and MQC experiments. We derive expressions for OTOC functions for determining the information scrambling considering imperfect LE experiments. A more general OTOC than the one described in Eq. (1) is obtained, that allows modeling decoherence effects induced by the non-reverted terms that affects the outcome of information scrambling measurements. The decoherence effects arise naturally as a leakage of the unitary dynamics of the ideal echo experiment with a rate that depends on the number of correlated spins  $L$  on a system state,  $\Gamma(L)$ . We show that the scrambling degree can be interpreted as the average number of active spins  $K$  –equivalent to a mean Hamming distance– where the information was spread by the evolution that survived the perturbation effects. Using these general OTOC expressions, we introduce the effects of non-reversed interactions into MQC experiments. This allows to model the dynamics of the effective cluster size of correlated spins  $K$  that provides a measurable degree of scrambling of information into system under decoherence effects.

We contrast the model’s predictions with quantum simulations that measure the information scrambling with imperfect time-reversal echoes performed with solid-state NMR experiments. The presented experiments are based on previous methods and results, where a spin system is quenched by a control Hamiltonian that induces the scrambling of a magnetization that play the role of a localized initial information [9, 15, 18, 59]. By adding tunable perturbations to the control Hamiltonian that produces the scrambling of information, we observe localization effects on the scrambling dynamics determined by the effective number of correlated spins [15, 59]. It was experimentally demonstrated that this localization effects are described by a dynamical equilibrium [15, 59, 60]. It was also observed that there ex-

ists a critical perturbation strength that, once surpassed, corresponds to a localized dynamics that, however, below the critical threshold the scrambling dynamics is delocalized [9]. This localized-delocalized transition as a function of the perturbation strength in the information scrambling dynamics evidence a sudden change on the scaling law of the decoherence effects induced by the perturbation [18]. The model developed in this article provides predictions that are consistent with the experimental observations and the findings in previous works that evidence a scaling transition on the scrambling dynamics behavior. In particular, our model predicts a localization cluster size of the scrambling dynamics as a dynamical equilibrium state. The model also manifests a transition from a localized to a delocalized dynamics as a function of the perturbation strength in finite systems. By extracting the decoherence leakage rates induced by the perturbations from the experiments, and introducing them into our model, we accurately predict the growth of the cluster size of correlated spins of the experimental data. The results show excellent quantitative agreement for the estimation of the localization sizes. Therefore the presented model combined with the derivation of OTOCs can be useful tools to predict the information scrambling dynamics of large quantum systems, and address the effect of imperfections on the control Hamiltonian that drives the quantum evolutions.

Our article is organized as follows, in Sec. II, we introduce the considered spin system, the OTOCs used and how they can be determined with LE and MQC experiments. We also provide a measure for information scrambling based on a cluster size of active –correlated– spins. We introduce the effects of imperfect echo experiments to quantify the scrambling dynamics. In Sec. III, we first introduce the original Levy and Gleason model, and then we adapt it to describe quantum information scrambling dynamics. Based on the derived OTOCs, we introduce the decoherence effects induced by imperfection on the time-reversal procedure. In Sec. IV, we analyze our model to show how scrambling is modified in the presence of decoherence effects. In Sec. V, we contrast our model with experimental results, and show the consistency on the predictions of the observable information scrambling bounds measured with NMR quantum simulations. Finally, in Sec. VI we give the conclusions.

## II. QUANTUM INFORMATION SCRAMBLING IN SPIN SYSTEMS

### A. The system out of equilibrium

We consider a system of  $N$  interacting spins  $I = 1/2$  in presence of a strong magnetic field along the  $z$  direction. We assume the Larmor frequency  $\omega_z \gg d_{ij}$ , with  $d_{ij}$  the dipole-dipole interaction strength between the  $i$  and  $j$  spins. In a frame of reference rotating at the Larmor frequency, the Hamiltonian of the system is given by the

truncated dipolar interaction

$$\begin{aligned} \mathcal{H} &= \mathcal{H}_{dd} \\ &= \sum_{i < j} d_{ij} [2I_z^i I_z^j - (I_x^i I_x^j + I_y^i I_y^j)], \end{aligned} \quad (2)$$

where we have neglected the non-secular terms of the dipolar Hamiltonian [61]. Therefore the Hamiltonian  $\mathcal{H}_{dd}$  conserves the total magnetization of the system as  $[\mathcal{H}_{dd}, I_z] = 0$ , with  $I_z = \sum_j I_z^j$  the total spin operator on the  $z$  direction. The operators  $I_v^i$  are the angular momentum of the  $i$ -th spin in the  $v$  direction. We assume the initial state of the system at a thermal equilibrium with the Zeeman interaction. Its density matrix is  $\rho(0) = e^{-\omega_z \hbar I_z \beta} / \text{Tr}(e^{-\omega_z \hbar I_z \beta})$ , where we have neglected the dipolar interaction due to  $\omega_z \gg d_{ij}$ . In the high-temperature limit  $\beta^{-1} = k_B T \gg \hbar \omega_z$ , the thermal state is approximated to [61]

$$\rho(0) \sim (\mathbb{I} - \beta \omega_z \hbar I_z) / \text{Tr}(\mathbb{I}). \quad (3)$$

The identity operator  $\mathbb{I}$  does not evolve over time and does not contribute to the expectation value of an observable magnetization, since  $\text{Tr}(I_v \cdot \mathbb{I}) = 0$  for any direction  $v$ . This initial state thus represents an ensemble of local operators  $I_z^j$ , that plays the role of an initial local information that will be scrambled into the system degrees of freedom. To induce the spreading of the quantum information from the initial state operator  $\rho(0)$ , we drive the system away from equilibrium by quenching the Hamiltonian [9, 15, 18]. In NMR, this is typically done by applying rotations to the spins with electromagnetic field pulses that engineer an average Hamiltonian that does not commute with the initial state [62]. In particular, we consider the effective interaction given by a double-quantum Hamiltonian

$$\mathcal{H}_0 = \frac{1}{2} \sum_{i,j} d_{ij} (I_+^i I_+^j + I_-^i I_-^j), \quad (4)$$

that allows to probe the growth of the number of correlated spins as a function of time [9, 15, 44] as a measure of the degree of scrambling of information into the system [17, 18, 20, 27–29, 63]. Here,  $I_\pm^i$  are the ladder operators of the  $i$ -th spin. In the Zeeman basis, the quantum states  $|\vec{m}\rangle = |m_1, \dots, m_N\rangle$  are characterized by the magnetization numbers  $m_i = \pm \frac{1}{2}$  associated with the local spin operators  $I_z^i$ . The double-quantum Hamiltonian flips simultaneously two spins with the same orientation  $m_i$ , therefore inducing transitions from a state  $|\vec{m}\rangle$  to a state  $|\vec{n}\rangle$  that change the coherence order  $M = h(\vec{m}, \vec{n}) = \sum_i m_i - n_i$  by  $\Delta M = \pm 2$ . Here  $h(\vec{m}, \vec{n})$  is the Hamming distance between the states  $|\vec{m}\rangle$  and  $|\vec{n}\rangle$ . Since the initial state  $\rho(0)$  is diagonal, it only has non-vanishing elements for  $M = 0$  and the double-quantum Hamiltonian only creates even coherence orders.

We consider as our observable the magnetization operator  $I_z$ . The observable signal is then  $\text{Tr}[I_z \rho(t)] \propto \text{Tr}[I_z I_z(t)]$ , which is proportional to the time evolution of the  $I_z$  operator as the identity term of the initial state gives null trace. Therefore, we consider the evolution of the  $I_z$  operator in the following. At the evolution time  $t$ , the scrambled state  $I_z^0(t) = U_0(t) I_z U_0^\dagger(t)$  can be expanded in coherence orders as

$$\begin{aligned} I_z^0(t) &= \sum_M \sum_{h(\vec{m}, \vec{n})=M} I_{z, \vec{m}\vec{n}}^0(t) |\vec{m}\rangle \langle \vec{n}| \\ &= \sum_M I_{z, M}^0(t), \end{aligned} \quad (5)$$

where the operator  $I_{z, M}^0 = \sum_{h(\vec{m}, \vec{n})=M} I_{z, \vec{m}\vec{n}}^0(t) |\vec{m}\rangle \langle \vec{n}|$  contains all the elements of the density operator involving coherences of order  $M$ . The superindex 0 indicates evolution under the double-quantum Hamiltonian  $\mathcal{H}_0$ , i.e.  $U_0(t) = e^{-i\mathcal{H}_0 t}$ . The amplitude of each coherence order is  $f_M^0 = \text{Tr}(I_z^2)^{-1} \text{Tr}[I_{z, M}^0(t) I_{z, M}^{0\dagger}(t)]$ , that defines the Multiple Quantum Coherence (MQC) spectrum [44].

## B. Measuring information scrambling with Multiple Quantum Coherences

The many-body Hamiltonian  $\mathcal{H}_0$  scrambles the initial local state into the degrees of freedom of the system. The information spreading dynamics is contained in the evolved state  $I_z^0(t) = U_0(t) I_z U_0^\dagger(t)$ . Usually, the degree of information scrambling is quantified between distant local operators  $A$  and  $B$  in Eq. (1) [19]. In our case, we consider the observation of scrambling based on monitoring the spreading of an ensemble of local operators  $I_z$  by its own time evolution  $I_z^0(t)$  via the commutator  $[I_z^0(t), I_z]$ , as it is accessible by time-reversal echoes common in NMR experiments [18, 27–29, 63]. Based on Eq. (1), the OTO commutator

$$\begin{aligned} &\frac{1}{\text{Tr}(I_z^2)} \langle [I_z^0(t), I_z] [I_z^0(t), I_z]^\dagger \rangle = \\ &= \frac{2}{\text{Tr}(I_z^2)} \{ \langle I_z I_z^0(t) I_z^{0\dagger}(t) I_z^\dagger \rangle - \text{Re} [\langle I_z^{0\dagger}(t) I_z^\dagger I_z^0(t) I_z \rangle] \} \\ &= \frac{2}{\text{Tr}(I_z^2)} \langle I_z I_z^0(t) I_z^{0\dagger}(t) I_z^\dagger \rangle - 2\text{Re}(F_{I_z, I_z}) \end{aligned} \quad (6)$$

is related to the OTOC function  $F_{I_z, I_z} = \text{Tr}(I_z^2)^{-1} \langle I_z^{0\dagger}(t) I_z^\dagger I_z^0(t) I_z \rangle$ , where  $\langle \mathcal{O} \rangle = \text{Tr}(\rho_\beta \mathcal{O})$  is the expectation value of an operator  $\mathcal{O}$  considering the system on the density matrix state  $\rho_\beta$ . If the system state is in the infinite temperature limit  $\rho_\beta = \mathbb{I}$ , then  $\langle \mathcal{O} \rangle_{\beta=0} = \text{Tr}(\mathcal{O})$ . Therefore

$$\begin{aligned} &\lim_{\beta \rightarrow 0} \left[ \frac{2}{\text{Tr}(I_z^2)} \langle I_z I_z^0(t) I_z^{0\dagger}(t) I_z^\dagger \rangle - 2\text{Re}(F_{I_z, I_z}) \right] = \\ &= \frac{1}{\text{Tr}(I_z^2)} \text{Tr} \left\{ [I_z^0(t), I_z] [I_z^0(t), I_z]^\dagger \right\} \end{aligned} \quad (7)$$




Figure 1. Protocol for measuring Multiple Quantum Coherence and monitoring the quantum information scrambling. (a) The MQC protocol consists of a sequence of quantum evolutions. First, a forward in time evolution driven by the double quantum Hamiltonian  $\mathcal{H}_F = \mathcal{H}_0$  that pumps coherences of different orders  $M$ . The rotation  $\phi_z = e^{iI_z\phi}$  encodes the coherence order with an order-dependent phase  $\phi M$ . Then a backward in time evolution driven by the Hamiltonian  $\mathcal{H}_B = -\mathcal{H}_0$  reverses the time evolution to allow the detection of the MQC spectrum  $f_M^0$  from the signal  $f_0(t, \phi) = \text{Tr}(I_z^2)^{-1} \sum_M e^{i\phi M} f_M^0$ . The signal results from measuring the magnetization  $I_z$ , starting from the initial state  $\rho(0) = (\mathbb{I} - \beta\omega_z \hbar I_z) / \text{Tr}(\mathbb{I})$ . (b) Modified MQC protocol to assess the sensitivity of a quantum dynamics to perturbations in the Hamiltonian. The forward Hamiltonian  $\mathcal{H}_F = (1-p)\mathcal{H}_0 + p\Sigma$  differs from the backward Hamiltonian  $\mathcal{H}_B = -\mathcal{H}_0$ . The resulting MQC spectrum  $f_M(t, p) = \text{Tr}(I_z^2)^{-1} \text{Tr}[I_{z,M}^0(t) I_{z,M}^\dagger(t)]$  quantifies the inner-product of the coherence orders between the ideal  $I_z^0(t)$  and the perturbed evolution  $I_z(t)$ . (c) Typical form of the MQC spectrum. Its second moment is determined from the squared width of the MQC distribution, and provides a measure for the average number of active spins  $K(t)$  where the information was scrambled shared by the perturbed and unperturbed dynamics.

quantifies the scrambling of local information on  $I_z$  due to the evolution driven by  $\mathcal{H}_0$  in the system at infinite temperature [18, 29, 63].

Determining the Multiple Quantum Coherence (MQC) spectrum allows to measure OTOC functions [20]. A scheme of an experimental implementation of a concatenation of quantum evolutions used to extract the MQC spectrum is shown in Fig. 1(a) [44]. First, the system is at its initial state, then it is quenched by suddenly turning on the double-quantum Hamiltonian  $\mathcal{H}_0$ . The system evolves forward in time into the scrambled state  $I_z^0(t) = U_0(t) I_z U_0^\dagger(t)$ . A rotation along the  $z$ -direction  $\phi_z = e^{-i\phi I_z}$  is applied to the state  $I_z^0(t)$  to label each coherence term  $I_{z,M}^0(t)$  of Eq. (5), as they acquire a phase  $M\phi$ . The  $I_z^0(t)$  operator becomes

$$\begin{aligned} I_z^0(\phi, t) &= \phi_z I_z^0(t) \phi_z^{-1} \\ &= \sum_M e^{iM\phi} I_{z,M}^0(t). \end{aligned} \quad (8)$$

Finally, the dynamics driven by the double-quantum Hamiltonian  $\mathcal{H}_0$  is reversed in time by changing the sign

of  $\mathcal{H}_0$  by a pulse sequence design [44]. This backward evolution in time is driven by the evolution operator  $U_0^{-1}(t) = U_0^\dagger(t)$ . This leads to a many-body Loschmidt Echo [33, 34, 37] on the resulting magnetization along  $I_z$ , where we obtain for the normalized signal at the end of the evolution

$$\begin{aligned} f_0(\phi, t) &= \frac{1}{\text{Tr}(I_z^2)} \text{Tr}[I_z \cdot U_0^\dagger(t) \phi_z U_0(t) I_z U_0^\dagger(t) \phi_z^\dagger U_0(t)] \\ &= \frac{1}{\text{Tr}(I_z^2)} \text{Tr}[I_z^0(t) I_z^{\text{ot}}(\phi, t)] \\ &= \frac{1}{\text{Tr}(I_z^2)} \sum_{MM'} e^{iM\phi} \text{Tr}[I_{z,M}^0(t) I_{z,M'}^{\text{ot}}(t)] \delta_{MM'}. \end{aligned} \quad (9)$$

We have used the cyclic property of the trace, and the normalization factor  $\text{Tr}(I_z^2)^{-1}$  to ensure the normalization of the signal  $f_0(\phi, t=0) = 1$ . This time reversal of the quantum dynamics combined with the rotation  $\phi_z$  allows quantifying the contribution of the different coherence orders  $f_{0,M} = \text{Tr}(I_z^2)^{-1} \text{Tr}[I_{z,M}^0(t) I_{z,M}^{\text{ot}}(t)]$  to the global spin state, and therefore the scrambling of information into multi-spin states by performing a Fourier transform on  $\phi$  [18, 44].

As the relevant observable term in the initial state  $\rho(0) = (\mathbb{I} - \beta\omega_z \hbar I_z) / \text{Tr}(\mathbb{I})$  is proportional to the observable magnetization  $I_z$ , the Loschmidt Echo  $f_0(\phi, t)$  can be considered as a fidelity function quantified by the inner product  $\text{Tr}[I_z^0(t) I_z^{\text{ot}}(\phi, t)]$  between the rotated state  $I_z^0(\phi, t)$  and the original one  $I_z^0(t)$ . Moreover,  $f_0(\phi, t)$  also defines an OTOC function  $F_{\phi_z, I_z}$  according to Eq. (7)

$$\begin{aligned} f_0(t, \phi) &= \frac{1}{\text{Tr}(I_z^2)} \text{Tr}[I_z^0(t) \phi_z I_z^0(t) \phi_z^\dagger] \\ &= F_{\phi_z, I_z} \\ &= 1 - \frac{1}{2\text{Tr}(I_z^2)} \left\langle |[I_z^0(t), \phi_z]|^2 \right\rangle_{\beta=0}, \end{aligned} \quad (10)$$

where again  $\text{Tr}(\mathcal{O}) = \langle \mathcal{O} \rangle_{\beta=0}$  is the expectation value of  $\mathcal{O}$  with the system state at infinite temperature  $\rho_\beta = \mathbb{I}$  [18, 29, 63, 64] and we have used that  $\langle \phi_z I_z^0(t) I_z^{\text{ot}}(t) \phi_z^\dagger \rangle_{\beta=0} = \text{Tr}(I_z^2)$ . Using a Taylor expansion as a function of  $\phi$  in Eq. (10), the second moment of the MQC spectrum  $K_0$  is determined by [18, 28, 63, 65]

$$\begin{aligned} K_0 &= \sum_M M^2 f_{0,M}(t) \\ &= \frac{1}{\text{Tr}(I_z^2)} \text{Tr}\left([I_z, I_z^0(t)] [I_z, I_z^0(t)]^\dagger\right), \end{aligned} \quad (11)$$

which is related to an OTOC function following Eq. (7). The second moment  $K_0$  quantifies therefore the information scrambling into the system driven by the double-quantum Hamiltonian  $\mathcal{H}_0$  starting from the localized information at  $I_z$ .

### C. Cluster size of correlated spins created by the information scrambling

In the NMR community, the second moment  $K_0$  is typically used for spin counting of the number of correlated spins by the dynamics induced by a many-body Hamiltonian [44, 46, 48]. The second moment quantifies the cluster size of spins where an initial local state was spread into the system [9, 15]. To relate the second moment of the MQC spectrum  $K_0$  to a number of correlated spins by the information scrambling, it is necessary to know the propagation model of excitations within the system. Baum *et. al.* proposed a simple model by assuming that all coherence orders are equally excited for a given system size [44, 45]. The corresponding MQC spectrum has a Gaussian distribution, whose second moment determines the system size  $K_0$  which corresponds to the instantaneous number of correlated spins. While this simple assumption works reasonable well in several solid-state systems [46, 50, 66], the MQC distribution is not always Gaussian [47], and other models are required to give a quantitative number of correlated spins [27, 47, 65]. There is no general method to define the number of correlated spins independently of the system dynamics.

We here derive a formal definition for the *cluster size of correlated spins* from the second moment  $K_0$  created by the information scrambling, that does not require assumptions regarding the dynamics of the system. This definition for the cluster size is used then in Sec. III to describe the information scrambling by OTOC functions determined from an imperfect echo experiment.

We consider the basis  $\{P_{\vec{u}} = (\sqrt{2})^N \otimes_{i=1}^N I_{u_i}^{(1)}\}$  of the composite Hilbert space for  $N$  1/2-spins, where the index  $i$  labels the spins,  $u_i \in \{0, x, y, z\}$  and  $\{I_{u_i}^{(1)}\}$  is the set of spin operators of a single 1/2-spin, including the identity operator  $I_0^{(1)} \equiv \mathbb{I}$ . As these single spin operators satisfy the orthogonal relation  $\text{Tr}(I_{u_i}^{(1)} \cdot I_{v_j}^{(1)}) = \frac{1}{2} \delta_{u_i, v_j}$ , the operators set  $\{P_{\vec{u}}\}$  is an orthonormal basis of the complete Hilbert space. The evolved quantum state  $I_z^0(t)$  can then be expanded on this basis as

$$I_z^0(t) = \sum_{\vec{u}} C_{\vec{u}}^0(t) P_{\vec{u}}, \quad (12)$$

where  $C_{\vec{u}}^0(t)$  are time-dependent complex coefficients. Using this expansion, the cluster size of correlated spins results

$$K_0(t) = \sum_{\vec{u}, \vec{v}} C_{\vec{u}}^0(t) C_{\vec{v}}^{0*}(t) \mathcal{L}(\vec{u}, \vec{v}), \quad (13)$$

where  $*$  is the complex conjugate. The function  $\mathcal{L}(\vec{u}, \vec{v})$  has a functional dependence on  $\vec{u}$  and  $\vec{v}$  defined by (see Appendix A for a demonstration)

$$\mathcal{L}(\vec{u}, \vec{v}) = \begin{cases} L(\vec{u}) & \vec{u} = \vec{v} \\ 2 & h_{z0}(\vec{u}, \vec{v}) = 0, h_{xy}(\vec{u}, \vec{v}) = 2 \\ & \text{and } \vec{v} = \Pi(\vec{u}) \\ -2 & h_{z0}(\vec{u}, \vec{v}) = 0, h_{xy}(\vec{u}, \vec{v}) = 2, \\ & \text{and } \vec{v} \neq \Pi(\vec{u}) \\ 0 & \text{Other cases} \end{cases} \quad (14)$$

The diagonal terms when  $\vec{u} = \vec{v}$  in this expression gives the number  $L(\vec{u})$  of elements in  $\vec{u}$  that are equal to  $x$  or  $y$ . For the non-diagonal terms  $\vec{u} \neq \vec{v}$ , the conditions on the right hand side have to be satisfied simultaneously. The function  $h_{z0}(\vec{u}, \vec{v})$  is the Hamming distance between  $\vec{u}$  and  $\vec{v}$  only considering the elements that are equal to 0 and  $z$ . In other words, the condition  $h_{z0}(\vec{u}, \vec{v}) = 0$  implies that  $u_i = v_i$  for every  $u_i = 0, z$ . Similarly,  $h_{xy}(\vec{u}, \vec{v})$  is the Hamming distance between  $\vec{u}$  and  $\vec{v}$  only considering the elements that are equal to  $x$  and  $y$ . The condition  $\vec{v} = \Pi(\vec{u})$  indicates that the vector  $\vec{v}$  is a permutation of vector  $\vec{u}$ . The cross terms  $\vec{u} \neq \vec{v}$  in Eq. (13) have complex numbers with phases that interfere destructively when they are summed, and therefore their contribution goes to zero as the cluster size  $K_0(t)$  increases [67]. Therefore the cross terms contribution is negligible for large  $K_0(t)$ , obtaining the cluster size of correlated spins

$$K_0(t) \sim \sum_{\vec{u}} |C_{\vec{u}}^0(t)|^2 L(\vec{u}). \quad (15)$$

Here the magnitude  $L(\vec{u})$  quantifies the number of *active spins* associated to a coherence transfer process  $\langle \vec{n} | P_{\vec{u}} | \vec{m} \rangle$  [68, 69]. The transition element  $\langle \vec{n} | P_{\vec{u}} | \vec{m} \rangle \neq 0$  if and only if there are  $L(\vec{u})$  spins that flip their state during the transition  $|\vec{m}\rangle \rightarrow |\vec{n}\rangle$  or, equivalently  $L = h(\vec{m}, \vec{n})$  with  $h$  the Hamming distance. This implies that  $K_0$  is the average number of active spins in the state  $I_z^0(t)$  weighted by the coefficients  $C_{\vec{u}}^0(t)$  that depends on the quantum dynamics, thus providing an interpretation for the information scrambling in spin systems. The cluster size  $K_0$  provides the average number of spins correlated by quantum superpositions generated by the information scrambling dynamics. This expression for  $K_0$  is similar to the “average correlation length” introduced by Wei *et. al.* in Ref. [27], but  $K_0$  is determined by the average number of spins that does not contain identity operators on the system state. A formal connection between OTOC functions and “average correlation length” is also provided in Ref. [27], but only for a particular non-interacting system defined by a spin-chain network topology. In our case,  $L(\vec{u})$  quantifies the number of non-identity operators and non- $I_z$  operators in  $P_{\vec{u}}$ . Based on Eqs. (13) and (15), we provide an average Hamming distance as a way of quantifying a correlation length which is directly connected with the quantum information spreading derived from the OTOC function in Eq. (11). Moreover, our expression to determine  $K_0$  and the corresponding OTOC is independent of the spin-network topology, and it does not require assumptions on the MQC dynamics.

### D. Imperfect echo effects on estimating the information scrambling dynamics

An implementation of an echo experiment for measuring the information scrambling from the MQC spectrum, always contains imperfections that lead to a time reversion not fully performed. Different sources of imperfections might occur induced by i) non-idealities of the control operations and ii) the existence of external degrees of freedom considered as an environment [38]. Both imperfection terms in the Hamiltonian cannot be typically reversed. As a paradigmatic model of these imperfections, we consider a perturbation term  $p\Sigma$  in the forward Hamiltonian as generally considered within the Loschmidt Echo formalism [35–37] [see Fig. 1(b)]. The perturbation strength  $p$  is a dimensionless parameter and  $\Sigma$  is a perturbation Hamiltonian. This perturbation spoils the time reversion process and thus induces decoherence effects [9, 15, 18, 38]. As shown in Fig. 1(b), the forward evolution is driven by the evolution operator  $U_P(t) = e^{-i\mathcal{H}(p)t}$ , where the forward Hamiltonian is  $\mathcal{H}_F = \mathcal{H}(p) = (1-p)\mathcal{H}_0 + p\Sigma$ . After the forward evolution, again the global rotation  $\phi_z = e^{-i\phi I_z}$  encodes the coherence orders as  $\phi_z I_z(t) \phi_z^\dagger = \sum_M e^{-iM\phi} I_{z,M}(t)$ , and then the system is evolved backward in time with an ideal evolution operator  $U_0^{-1} = e^{i\mathcal{H}_0 t}$ . The resulting state is then projected on  $I_z$ , which is the observable magnetization, leading to the signal

$$\begin{aligned} f(\phi, t, p) &= \\ &= \frac{1}{\text{Tr}(I_z^2)} \text{Tr} [I_z \cdot U_0(t) \phi_z U_p(t) I_z U_p^\dagger(t) \phi_z^\dagger U_0^{-1}(t)] \\ &= \frac{1}{\text{Tr}(I_z^2)} \sum_M e^{-iM\phi} \text{Tr} [I_{z,M}^0(t) I_{z,M}^\dagger(t)]. \end{aligned} \quad (16)$$

The MQC spectrum is now defined by the inner-product between the coherence orders of the forward and backward density matrix terms  $f_M(t, p) = \text{Tr}(I_z^2)^{-1} \text{Tr} [I_{z,M}^0(t) I_{z,M}^\dagger(t)]$ . The non-ideal time reversion evolution therefore compares the two different information scrambling dynamics by comparing the terms  $I_{z,M}^0(t)$  and  $I_{z,M}(t)$  based on their overlap  $f_M$ . As derived in Eq. (10), the Loschmidt Echo of Eq. (16) can be recast as

$$\begin{aligned} f(\phi, t) &= \frac{1}{\text{Tr}(I_z^2)} \text{Tr} [I_z^0(t) I_z^\dagger(t)] - \\ &= \frac{1}{2\text{Tr}(I_z^2)} \text{Tr} \left( [\phi_z, I_z^0(t)] [\phi_z, I_z(t)]^\dagger \right) \\ &= f(\phi = 0, t) - \\ &= \frac{1}{2\text{Tr}(I_z^2)} \text{Tr} \left( [\phi_z, I_z^0(t)] [\phi_z, I_z(t)]^\dagger \right). \end{aligned} \quad (17)$$

Here, the Loschmidt Echo is related to a more general OTO commutator  $\text{Tr}(I_z^2)^{-1} \text{Tr} \left( [\phi_z, I_z^0(t)] [\phi_z, I_z(t)]^\dagger \right)$  than the one described in Eq. (1). This OTOC quantifies

the overlap of the commutators  $[\phi_z, I_z^0(t)]$  and  $[\phi_z, I_z(t)]$  via the inner-product [18]. The second moment of the MQC spectrum  $f_M(t, p)$  is now

$$\sum_M M^2 f_M(t) = \frac{1}{\text{Tr}(I_z^2)} \text{Tr} \left( [I_z, I_z^0(t)] [I_z, I_z(t)]^\dagger \right). \quad (18)$$

In this imperfect echo experiment, the cluster size is then

$$K(t) = \frac{\sum_M M^2 f_M(t)}{f(\phi=0, t, p)}, \quad (19)$$

where the second moment of the MQC spectrum must be normalized to the fidelity  $f(\phi = 0, t, p) = \text{Tr}(I_z^2)^{-1} \text{Tr} [I_z^0(t) I_z^\dagger(t)]$ . This normalization is required to extract the MQC distribution width, as the fidelity decays as a function of time, and therefore the overall amplitude of the MQC distribution. The second moment  $K(t)$  is an *effective cluster size* that represents the common number of correlated spins between the ideal and the perturbed information scrambling dynamics, quantified by the commutators  $[I_z, I_z^0(t)]$  and  $[I_z, I_z(t)]$  respectively.

By expanding  $I_z(t)$  with the multi-spin operators  $\{P_{\vec{u}}\}$ ,

$$I_z(t) = U_p(t) I_z U_p^\dagger(t) = \sum_{\vec{u}} C_{\vec{u}}^p(t) P_{\vec{u}}, \quad (20)$$

the perturbed version of Eq. (13) results

$$K(t) = \frac{1}{f(\phi = 0, t, p)} \sum_{\vec{u}, \vec{v}} C_{\vec{u}}^0(t) C_{\vec{v}}^{p*}(t) \mathcal{L}(\vec{u}, \vec{v}), \quad (21)$$

where  $\mathcal{L}(\vec{u}, \vec{v})$  is defined as in Eq. (14). Again, the cross-terms in Eq. (21) are negligible when  $K(t)$  is large [67], and we obtain

$$\begin{aligned} K(t) &\simeq \frac{1}{f(\phi = 0, t, p)} \sum_{\vec{u}} C_{\vec{u}}^0(t) C_{\vec{u}}^{p*}(t) L(\vec{u}) \\ &= \frac{1}{\sum_{\vec{u}} C_{\vec{u}}^0(t) C_{\vec{u}}^{p*}(t)} \sum_{\vec{u}} C_{\vec{u}}^0(t) C_{\vec{u}}^{p*}(t) L(\vec{u}). \end{aligned} \quad (22)$$

This expression demonstrates that  $K(t)$  is an average Hamming distance based on the active spins, weighted by the product between the forward and backward coefficients  $C_{\vec{u}}^0(t) C_{\vec{u}}^{p*}(t)$  that determine the respective dynamics evolutions. Therefore the expression we derived here for  $K(t)$  gives the average number of active spins shared by the forward and backward dynamics. As experimental implementations of time reversions have always a non-reverted interaction, our expression for  $K(t)$  provides a definition of what is actually measured as information scrambling with echo experiments.

### III. MODEL FOR THE DYNAMICS OF QUANTUM INFORMATION SCRAMBLING UNDER DECOHERENCE

#### A. Revisiting the Levy-Gleason model: Quantum information scrambling dynamics

We develop a phenomenological model to describe the time evolution of the information scrambling observed via the second moment of the MQC distribution  $K(t)$ . We consider that decoherence effects or perturbations to the ideal Hamiltonian affect the time reversion of the protocol described in Sec. II B. Levy and Gleason developed a model to describe the MQC dynamics in solid state systems [50]. We here adapt this model for determining the information scrambling dynamics in a spin system by using the expressions of Eqs. (15) and (22).

The complexity of many-spin system dynamics is an outstanding problem in physics that only allows to obtain exact solutions for  $I_z(t)$  for very special cases [70–74]. In general, approximations or phenomenological models need to be implemented [44, 45, 49–52, 67, 75, 76]. The Levy-Gleason model describes the growth of the cluster size of correlated spins  $K_0(t)$  as a function of time, driven by the double-quantum Hamiltonian  $\mathcal{H}_0$  in an  $N$  spin system. The model introduces an average operator  $P_k$  that contains all the operators  $P_{\vec{u}}$  with the same number of non-identity operators  $k$ . This average operator is used to simplify Eq. (12) to

$$I_z^0(t) = \sum_{\vec{u}} C_{\vec{u}}^0(t) P_{\vec{u}} \approx \sum_k C_k^0(t) P_k, \quad (23)$$

where the coefficient  $C_k^0(t)$  describes an averaged evolution of the operators with the same number  $k$ . Applying the Liouville-von Neumann equation to Eq. (23), the dynamic of  $C_k^0(t)$  is determined by the set of  $N$  coupled linear differential equations

$$\frac{d}{dt} C_k^0 = -\frac{i}{4} W_{k-1} C_{k-1}^0 - \frac{i}{4} W_k C_{k+1}^0, \quad (24)$$

where  $W_{k-1}$  and  $W_k$  are transition probabilities.

Based on the result of Eq. (15), which shows that the active spins are the relevant ones for quantifying the information scrambling dynamics, we adapt the Levy-Gleason model by recasting Eq. (23) to the form

$$I_z^0(t) = \sum_{\vec{u}} C_{\vec{u}}^0(t) P_{\vec{u}} \approx \sum_L C_L^0(t) P_L, \quad (25)$$

where now  $L$  is the number of active spins, defined by the number of non-identity and non- $I_z$  operators in  $P_{\vec{u}}$  as described in Sec. II C. Similarly to Eq. (24), the dynamics of the coefficients  $C_L^0(t)$  is now determined by the set of equations

$$\frac{d}{dt} C_L^0 = -\frac{i}{4} W_{L-1} C_{L-1}^0 - \frac{i}{4} W_L C_{L+1}^0, \quad (26)$$

where  $W_L$  and  $W_{L-1}$  are the transition probabilities of increasing or reducing the number of active spins  $L(\vec{u})$  by one spin respectively. Levy and Gleason assume that the operator  $P_k$  represents a contiguous group of  $k$  spins in real space, and that the only relevant interactions are between near neighbors. Therefore, they conclude that the transition probability  $W_k$  is determined by an effective dipolar coupling strength  $d$  multiplied by the number of spins at the cluster edge  $k^\delta$ , where the exponent  $\delta$  can be estimated from the the spatial dimension  $D$  of the system as  $\delta \sim 1 - 1/D$ . The effective coupling  $d$  is of the same order of magnitude than the width of the spins' resonance line and can be determined from a fit to the experimental data [50]. Based on these assumptions, we extend this result to the transition probability

$$W_L = dL^\delta, \quad (27)$$

as a function of the number of active spins  $L$ .

In the original Levy and Gleason model, the link between the coefficients  $C_k^0(t)$  and the second moment of the MQC spectrum is assumed by imposing  $K_0(t) = \sum_k k |C_k^0(t)|^2$ . The consistency of this assumption relies on the fact that the model reproduces well several experimental results in solid-state systems where power-law growth of the cluster size evolution is seen [50]. The predicted growth rates and power law exponents were also consistent with the spin-spin coupling network topologies. Its time evolution can then be obtained by solving a set of  $N$  differential equations rather than  $2^N$  as in the exact Liouville-von Neumann equation for an  $N$  spin system. Our results of Sec. II C that derive the Eq. (15), show that the second moment  $K_0(t)$  and hence the information scrambling of Eq. (11), are determined by the average number of active spins weighted by the dynamics coefficients  $|C_L(t)|^2$

$$K_0(t) = \sum_L L |C_L^0(t)|^2. \quad (28)$$

Therefore, the connection between the cluster size of correlated spins and the second moment of the MQC spectrum arise naturally based on the average number of active spins involved in the information scrambling correlations.

#### B. Model for the decoherent dynamics of quantum information scrambling

The product of the two complex coefficients  $C_{\vec{u}}^p$  and  $C_{\vec{u}}^0$  of the forward and backward dynamics respectively in Eq. (22), produces a reduction of the effective cluster size  $K(t)$  compared to the one determined by the ideal echo experiment case  $K_0(t)$ . For weak perturbations, we consider that the coefficients  $C_{\vec{u}}^0$  and  $C_{\vec{u}}^p$  mainly differ by a phase, where  $C_{\vec{u}}^p \sim |C_{\vec{u}}^0| e^{i\phi_{\vec{u}}}$ . Therefore the cluster size of Eq. (22) is




Figure 2. Schematic representation of the model for the decoherent dynamics of quantum information scrambling. It models the evolution of the effective cluster size  $K(t)$  in the presence of decoherence effects induced by a perturbation on the echo experiment described in Fig. 1. The rate  $W_L$  is the transition probability between  $P_{L-1}$  and  $P_L$ , and it is determined by the strength of the dipolar interaction  $d$ , the size of the cluster of active spins  $L$  and the lattice dimension  $D$ . The spin clusters are affected by a decoherence process, characterized by the decoherence rate  $\Gamma_L$ .

$$K(t) \sim \frac{1}{f(\phi=0, t, p)} \sum_{\vec{u}} |C_{\vec{u}}^0(t)|^2 \cos(\phi_{\vec{u}}) L(\vec{u}). \quad (29)$$

We here only consider the real part  $\cos(\phi_{\vec{u}})$  of the phase term, since  $K(t)$  is real and the imaginary terms cancel out. We then recast Eq. (29) in terms of the number of active spins  $L$

$$K(t) \sim \frac{1}{f(\phi=0, t, p)} \sum_L L \sum_{\vec{u}_L} |C_{\vec{u}_L}^0(t)|^2 \cos(\phi_{\vec{u}_L}). \quad (30)$$

The coefficients  $C_L^0(t) \equiv \sum_{\vec{u}_L} |C_{\vec{u}_L}^0(t)|^2$  of the average operator  $P_L$ , in Eq. (25), need now to be modified as the phase  $\cos(\phi_{\vec{u}_L})$  introduces an attenuation factor that depends on  $L$ . The effective cluster size  $K$  is now described by the attenuated coefficients  $C_L(t) = \sum_{\vec{u}_L} |C_{\vec{u}_L}^0(t)|^2 \cos(\phi_{\vec{u}_L})$ , with  $|C_L(t)| < |C_L^0(t)|$ . The perturbation term in the echo experiment, is therefore modeled as a source of decoherence in an open quantum system. We model this attenuation by adding to Eq. (26) a leakage term with a rate  $\Gamma_L$  that destroys the quantum superpositions given by the operator product  $P_L$ ,

$$\frac{d}{dt} C_L = -\frac{i}{4} W_{L-1} C_{L-1} - \frac{i}{4} W_L C_{L+1} - \Gamma_L C_L. \quad (31)$$

The rate  $\Gamma_L$  is the average decoherence rate for the product of  $L$  active spin operators. An schematic representation of the model is shown in Fig. 2.

The matrix representation of Eq. (31) is

$$\frac{d}{dt} \vec{C} = -\Upsilon \cdot \vec{C},$$

$$\begin{bmatrix} \frac{d}{dt} C_1 \\ \frac{d}{dt} C_2 \\ \vdots \\ \frac{d}{dt} C_N \end{bmatrix} = -i \begin{bmatrix} -i\Gamma_1 & \frac{1}{4}W_1 & 0 & \cdots & 0 \\ \frac{1}{4}W_1 & -i\Gamma_2 & \frac{1}{4}W_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{1}{4}W_{N-1} & -i\Gamma_N \end{bmatrix} \begin{bmatrix} C_1 \\ C_2 \\ \vdots \\ C_N \end{bmatrix}, \quad (32)$$

and the effective cluster size  $K$  in terms of  $C_L$ , results

$$K(t) = \left[ \sum_L |C_L(t)|^2 \right]^{-1} \sum_L L |C_L(t)|^2. \quad (33)$$

If the rate  $\Gamma_L$  has no dependence on  $L$ , then all the coefficients  $C_L$  are affected by a global attenuation factor. The effective cluster size  $K(t)$  evolves then equally to the case without perturbation, consistently with the predictions derived in Ref. [63]. However, typically the decoherence effects induced by perturbations increase with the number of active spins that are correlated, and therefore  $\Gamma_L$  generally depends on  $L$  [14, 18, 59, 77–80].

#### IV. EFFECTIVE CLUSTER SIZE EVOLUTION: DECOHERENT DYNAMICS OF QUANTUM INFORMATION SCRAMBLING

To analyze the effective cluster size evolution predicted by our model, we consider the diagonal base for  $\Upsilon$  to solve Eq. (32)

$$\frac{d}{dt} \tilde{C}_i = -\lambda_i \tilde{C}_i, \quad (34)$$

where  $\tilde{C}_i$  is the  $i$ -th component of the populations vector  $\vec{C}$  in the eigenbasis of  $\Upsilon$ , and  $\lambda_i$  is the  $i$ -th eigenvalue. The solution is then

$$\tilde{C}_i(t) = \tilde{C}_i(0) e^{-\lambda_i t}, \quad (35)$$

where  $\tilde{C}_i(0)$  gives the initial condition  $C(0)$  expressed in the  $\Upsilon$ -eigenbasis. We consider that the decoherence rate increases as a power law with the cluster size of active spins  $\Gamma_L = \Gamma_1 L^\alpha$ , where  $\alpha$  is the scaling exponent, as reported in quantum simulations in solid-state spin systems [14, 18] and this dependence is also expected for spin-boson models [77–79].

We determine the evolution of the cluster size of correlated spins for the case without perturbation. Figure 3(a) shows the cluster size evolution  $K_0(t)$  for  $\Gamma_1 = 0$  and different system sizes  $N$ . The cluster size grows with a power law as  $K_0(t) = K_1 t^a$ , until it reaches a maximum value due to finite-size effects determined by the system size  $N$ . The power-law exponent  $a$  and the constant  $K_1$  are determined by  $D$  and  $d$  respectively, from Eq. (27). After  $K_0(t)$  reaches its maximum value, it begins to oscillate indefinitely, again due to finite-size effects. Therefore, this solution is only useful before finite-size effects dominate the dynamical behavior.






Figure 3. Effective cluster size of correlated spins  $K(t)$  that defines the information scrambling predicted by our model. (a) Cluster size dynamics  $K_0(t)$  in the ideal case, i.e. without considering decoherence effects ( $\Gamma_1 = 0$ ). Three different system sizes  $N$  are shown in the legend. The cluster size grows as  $K_0(t) \propto t^a$  until finite size effects distort the evolution. All solutions are equivalent before the finite size effects appear independently of the value of  $N$ . (b) Effective cluster size evolution  $K(t)$ , now including the decoherence effects. When  $\Gamma_1 \ll \Gamma_{\text{crit}}$ , the growth of  $K(t)$  is delocalized until finite-size effects appear as in the case of  $\Gamma_1 = 0$ . Near the critical decoherence strength  $\Gamma_{\text{crit}}$ ,  $K(t)$  tends to localize although it exhibits low-frequency oscillations. For  $\Gamma_1 > \Gamma_{\text{crit}}$ ,  $K(t)$  localizes indefinitely. (c) Finite-size effects on the observation of the localization dynamics. If  $\Gamma_1 \neq 0$ ,  $K(t)$  will localize only if  $N$  is large enough. If not, finite-size effects avoid the observation of the localization effects.

We then introduce the decoherence effects. For  $\Gamma_1 > 0$ , an initial power-law growth for the effective cluster size  $K(t)$  is observed similarly to the case of  $\Gamma_1 = 0$  [Fig. 3(b)]. However the exponent  $a$  decreases slightly with increasing  $\Gamma_1$ . Again, at long times when finite-size effects are reached, oscillations are seen on the dynamics of  $K(t)$ . These oscillations decrease in frequency and amplitude as  $\Gamma_1$  increases. When the decoherence strength  $\Gamma_1$  increases above a critical value  $\Gamma_{\text{crit}}$ , the effective cluster size  $K(t)$  reaches a plateau which is maintained indefinitely in time. This plateau defines a localization size for the observed information scrambling determined by the effective cluster size. The predicted localization size is consistent with the experimental observations of Refs. [9, 15, 18]. If the system-size of the model  $N \rightarrow \infty$ , we observe that  $\Gamma_{\text{crit}} \rightarrow 0$ . This implies that considering power laws scalings for the leakage rates  $\Gamma_L = \Gamma_1 L^\alpha$ , our model predicts that the effective cluster size dynamics localizes, provided that the system size  $N$  is large enough if  $\Gamma_1 \neq 0$ . We have also observed that localization effects are manifested even for slow-growing scalings as for  $\Gamma_L \propto \log(L)$  or  $\Gamma_L \propto L^\alpha$  with  $\alpha < 1$ . Notice that, if the decoherence rate is independent of  $L$ , localization effects are not observed.

Figure 3(c) shows the dynamics of  $K(t)$  for different values of  $N$  for a given  $\Gamma_1$ , manifesting that localization effect are observed once  $N$  is large enough. All the curves behave equally before finite size effects are significant. Therefore if  $N$  is large enough to manifest localization for a fixed rate  $\Gamma_1 > \Gamma_{\text{crit}}$ , the predicted curve for  $K(t)$  is independent of  $N$ .

The effective cluster size  $K(t)$  of Eq. (33) can be written in terms of the eigenbasis of  $\Upsilon$  as

$$K(t) = \frac{\sum_L L |\sum_i c_{Li} \tilde{C}_i(t)|^2}{\sum_L |\sum_i c_{Li} \tilde{C}_i(t)|^2}, \quad (36)$$

where the populations  $C_L(t)$  are in term of a linear

combination of the evolution of  $\tilde{C}_i(t)$ , i.e.  $C_L(t) = \sum_i c_{Li} \tilde{C}_i(t)$  with  $c_{Li}$  the eigenvector coefficients. Using the solution for  $\tilde{C}_i(t)$  determined from Eq. (35), we get

$$K(t) = \frac{\sum_L L \left| \sum_i c_{Li} \tilde{C}_i(0) e^{-t\lambda_i} \right|^2}{\sum_L \left| \sum_i c_{Li} \tilde{C}_i(0) e^{-t\lambda_i} \right|^2}. \quad (37)$$

The transition from a delocalized to a localized scrambling dynamics is evidenced in the behavior of the eigenvalues of  $\Upsilon$  as a function of  $\Gamma_1$  [Fig. 4(a) and (b)]. The eigenvalues  $\{\lambda_i\}$  are purely imaginary for  $\Gamma_1 = 0$ , which implies the conservation of  $\sum_L |C_L|^2$ . However, if  $\Gamma_1 \neq 0$ , the eigenvalues  $\{\lambda_i\}$  are in general complex numbers  $\lambda_i = \gamma_i + i\omega_i$ , with real  $\gamma_i$  and imaginary  $\omega_i$  components. We consider the set  $\{\lambda_i\}$  sorted by its real value, so as  $\gamma_i \leq \gamma_{i+1}$ . If  $\Gamma_1 < \Gamma_{\text{crit}}$ , then  $\lambda_1 = \lambda_2^*$ , which implies that two different frequencies  $\omega_1$  and  $\omega_2 = -\omega_1$  has the same decay constant  $\gamma_1 = \gamma_2$ . Hence, in the long time limit, terms with decay constants  $\gamma_i > \gamma_1$  become negligible and we obtain

$$K(t \rightarrow \infty) = \frac{\sum_L L \left| e^{-t\lambda_1} \sum_i c_{Li} \tilde{C}_i(0) e^{-t(\lambda_i - \lambda_1)} \right|^2}{\sum_L \left| e^{-t\lambda_1} \sum_i c_{Li} \tilde{C}_i(0) e^{-t(\lambda_i - \lambda_1)} \right|^2} \\ \sim \frac{\sum_L L \left| (c_{L1} \tilde{C}_1(0) + c_{L2} \tilde{C}_2(0) e^{-2it\omega_1}) \right|^2}{\sum_L \left| (c_{L1} \tilde{C}_1(0) + c_{L2} \tilde{C}_2(0) e^{-2it\omega_1}) \right|^2}. \quad (38)$$

This solution provides the oscillations observed for  $K(t)$  due to the finite-size effects in Fig. 3.

If  $\Gamma_1 > \Gamma_{\text{crit}}$ , then  $\lambda_1$  becomes a non-degenerate real eigenvalue ( $\omega_1 = 0$  and  $\gamma_2 > \gamma_1$ ), implying that the effective cluster size  $K(t)$  attain a localization value at long times




Figure 4. Eigenvalues of the transition rate operator  $\Upsilon$  derived from our model. (a) Imaginary and (b) real parts of the eigenvalues  $\lambda_i$  as a function of the decoherence strength  $\Gamma_1$ . We use a low number of  $N = 10$  to show clearer the functional behavior of the eigenfrequencies, but the qualitative behavior is analogous for larger  $N$ . For  $\Gamma_1 = 0$ , all the frequencies are pure imaginary numbers, therefore the global amplitude  $\sum_L |C_L|^2$  of the system is strictly conserved. For  $\Gamma_1 \neq 0$ , the solutions  $\tilde{C}(t)$  become damped oscillations as  $\lambda_i$  are complex numbers. There is a critical decoherence strength  $\Gamma_{\text{crit}}$  from which the smallest eigenvalue  $\lambda_1$  become a real number. Localization effects can be deduced from the eigenspectrum: the existence of a non-oscillating, long-living solution  $\tilde{C}_1(t)$  for  $\Gamma_1 > \Gamma_{\text{crit}}$  implies that observed scrambling dynamics by  $K(t)$  will eventually localize for long times.

$$\lim_{t \rightarrow \infty} K(t) = K_{\text{loc}} = \frac{\sum_L L |c_{L1}|^2}{\sum_L |c_{L1}|^2}. \quad (39)$$

This demonstrate the existence of a localized regime for the observable information scrambling determined by  $K(t)$  when  $\Gamma_1 > \Gamma_{\text{crit}}$ . Moreover,  $K(t)$  converges always to the same stationary value, providing that the initial condition  $\vec{C}(0)$  has a non-zero contribution of the  $\tilde{C}_1$  eigenvector. This is because the coefficients  $c_{L1}$  are independent of the initial condition  $\vec{C}(0)$  in Eq. (39). Therefore, independently of the initial cluster size of correlated spins, the effective cluster size in the long time limit will converge to the same localization size consistently with experimental observations [15, 59, 60].

The delocalization-localization transition on the dynamical behavior of the information scrambling manifested by the evolution of  $K(t)$ , resembles the quantum dynamical phase transitions induced by decoherence effects [81, 82] that are connected with exceptional points ubiquitous of non-hermitian Hamiltonian [83–86].

## V. MODEL VS. EXPERIMENTS: EVALUATION OF THE DECOHERENT DYNAMICS OF INFORMATION SCRAMBLING

### A. Information scrambling determined with NMR quantum simulations

We evaluate here the presented model as a framework to describe quantum information scrambling dynamics




Figure 5. Quantum information scrambling determined from MQC experiments with imperfect time reversals. (a) Effective cluster size  $K(t)$  evolution determined from quantum simulations with solid-state NMR experiments on powdered adamantane for several perturbations strengths  $p$  (symbols) with the perturbation Hamiltonian  $\Sigma = \mathcal{H}_{dd}$ . The corresponding predictions of our model are shown in dashed lines. (b) Experimentally determined effective cluster size  $K(t)$  evolution starting from different initial conditions  $K_i$  for  $p = 0.108$  (symbols). The effective cluster size  $K(t)$  converges to a dynamical equilibrium determined by the localization size  $K_{\text{loc}}$  independently of the initial cluster size  $K_i$ . The presented model correctly predicts the effective cluster size evolution towards the dynamical equilibrium (dashed lines).

with imperfect echo experiments. We consider the MQC protocol based on imperfect time reversion echoes as described in Fig. 1(b), following the technique introduced in Refs. [15, 59]. In this experimental protocol, a controlled perturbation is introduced to the forward Hamiltonian  $\mathcal{H}_F = (1-p)\mathcal{H}_0 + p\Sigma$ , where the non-reverted term  $\Sigma$  is weighted by the dimensionless parameter  $p$  using average Hamiltonian techniques that can control the perturbation strength  $p$ . This allows performing quantum simulations to evaluate the effect of non-reverted interactions that are inherent to any echo experiment.

We perform the experimental quantum simulations on a Bruker Avance III HD 9.4 T WB NMR spectrometer with a  $^1\text{H}$  resonance frequency of  $\omega_z = 400.15$  MHz. We consider the nuclear spins  $^1\text{H}$  of a powdered adamantane sample as the system, which constitutes a dipolar interacting many-body system of equivalent  $1/2$  spins. The imperfect echo protocol of Fig. 1(b) is implemented using the perturbation Hamiltonian  $\Sigma = \mathcal{H}_{dd}$  as the raw dipolar interaction of the system. The perturbation Hamiltonian is introduced using the NMR sequence described in Refs. [15, 59]. This protocol provides a magnetization echo at the end of the sequence that is proportional to the fidelity function  $f(\phi, t, p)$  of Eq. (16), from which we experimentally monitor the dynamics of the effective cluster size  $K(t, p)$  based on Eq. (19) (see Fig. 5). We also calculate the instantaneous decay rate  $\chi'(t) = \frac{d\chi(t)}{dt}$  with  $\chi = \log(f)$  of the echo fidelity  $f(\phi, t, p)$ , which is shown in Fig. 6(a). It is seen from the experiments that the decoherence rate scales with a power law func-




Figure 6. Decoherence scaling as a function of the perturbation strength. (a) Experimentally determined decoherence rates (scatter) and fitting curves (solid lines)  $\chi'(K) = \chi'(1)K^\alpha$  for different perturbation strengths  $p$ . This power law behavior determines the decay rates  $\Gamma(L)$  of the presented model. The scaling exponent  $\alpha$  and the scaling factor  $\chi'(1)$  are shown in panels (b) and (c). Black squares are the fitting parameters of curves  $\chi'(K)$ , and the red circles are the parameters determined by assuming the single-parameter Ansatz of Eq. (40).

tion  $\chi'(K) = \chi'(1)K^\alpha$ . This scaling behavior indicates the sensibility of the controlled quantum dynamics to the perturbation as a function of the instantaneous effective cluster size  $K$  [18].

For an unperturbed echo experiment ( $p = 0$ ), the cluster size  $K_0(t)$  grows indefinitely following a power law  $K_0(t) \propto t^{4.3}$  up to the experimentally accessible timescales (black squares in Fig. 5). If the perturbation to the control Hamiltonian  $p \neq 0$ , the effective cluster size  $K(t)$  growth is reduced. The decay rate  $\chi'(K)$  and its power law exponent  $\alpha$  increase with the perturbation strength  $p$  (Fig. 6). As seen in Fig. 5(a),  $K(t)$  reaches a localization value  $K_{loc}$  that remains constant in time for large perturbations  $p > 0.02$  [9, 15]. This localization size reduces by increasing the perturbation strength  $p$  consistently with the predictions of Sec. IV.

The experimental results evidence that  $K_{loc}$  is determined by a dynamical equilibrium of  $K(t)$  that converges to the same stationary value, independently of the initial cluster size [15, 59, 60]. Figure 5(b) shows a series of experiments following the protocol implemented in Refs. [15, 59, 60]. Here, an initial cluster size of correlated spins  $K_i$  is prepared by an unperturbed evolution with the propagator  $U_0 = e^{-it_{ini}\mathcal{H}_0}$ , where  $t_{ini}$  is the initial-ization time required for preparing the initial cluster size. Then, the cluster size  $K_i$  is used as the initial information state for the perturbed evolution  $U_p = e^{-it[(1-p)\mathcal{H}_0 + p\Sigma]}$ . As shown in Refs. [15, 59, 60], the cluster size converges to a localization size  $K_{loc}(p)$  independently of the initial value of  $K_i$ .

## B. Quantitative evaluation of the decoherent model for the information scrambling dynamics

We perform here a quantitative comparison between the predictions of our model and the experimental results of the information scrambling dynamics in imperfect echo experiments. We determine the parameters  $d$  and  $\delta$  of Eq. (27) to reproduce the cluster size dynamics at  $p = 0$  shown in Fig. 5(a). We observe from the experimental data that the cluster size is  $K_0(t) \approx (7.41 \frac{1}{\text{ms}} t)^{4.2}$ . We found that the power law exponent  $a = 4.2$  is obtained if  $\delta = 0.78$ . This value is slightly larger than the one expected  $1 - 1/D = 0.66$  for a 3-dimensional system as in our case, according to the original Levy-Gleason model. This result is consistent with a cluster size that keeps growing for a long time, at a rate that is faster than in normal diffusion [60]. This might be related with a “super-diffusion” mechanism due to the complex long-range nature of the dipolar interaction in our system [87, 88]. Setting  $d = 13\text{kHz}$  defined by the width of the resonance line of adamantane, we obtain an excellent agreement with the experimental evolution of  $K_0(t)$  as shown by black dashed lines in Fig. 5(a).

To predict the evolution of the effective cluster size  $K(t)$  for every perturbation strength  $p$ , we need to define the decay rates  $\Gamma(L)$  in Eq. (31) for the average  $L$ -spin operators  $P_L$ . The experimentally observed decay rates  $\chi'(K) = \chi'(1)K^\alpha$  [Fig. 6 (a)] of the fidelity  $f$  have a power law dependence on the instantaneous cluster size  $K(t)$  with a power law exponent  $\alpha$  and proportionally constant  $\chi'(1)$  that depend on the perturbation strength [Black squares in Fig. 6 (b) and (c) respectively]. We assume that this decoherence rates determine the decay rates of the model as  $\Gamma(L) = \chi'(L)$  for each perturbation strength. The exponent  $\alpha$  presents a transition as a function of  $p$ , between a low-scaling regimen with  $\alpha \sim 0.5$  for weak perturbations and a high-scaling regimen with  $\alpha \sim 1$  for large perturbations [18].

We calculate then  $K(t)$ , using a system size  $N = 2 \times 10^4$  large enough to avoid finite size effects on the experimentally accessible temporal scales of Fig. 5(a). We compare the experimental results for  $K(t)$  in Fig. 5(a) with their predictions for several perturbation strengths. The calculated  $K(t)$  correctly predicts its time evolution and the achieved localization size  $K_{loc}$  for large  $p$ , although  $K_{loc}$  is slightly overestimated in all cases. The model predicts with high accuracy the cluster size growth for the long-time behavior of the weakest perturbation strengths.

Our model also quantitatively predicts well the dynamical equilibrium localization size  $K_{loc}$  predicted in subsection VA as show in Fig. 5(b), for the perturbation strength  $p = 0.108$ . The dynamical equilibrium value for the localization size  $K_{loc}$  is determined from Eq. (39) as the eigenvector matrix  $c_{Lk}$  is independent of the initial condition.

Figure 7 shows with black squares the prediction for the localization size  $K_{loc}$  as a function of the pertur-




Figure 7. Localization cluster size  $K_{loc}$  as a function of the perturbation strength for the perturbation Hamiltonian  $\Sigma = \mathcal{H}_{dd}$ . Yellow diamonds shows the experimentally determined localization sizes only for the largest perturbation strengths where localization effects are observed. We observe a power-law dependence  $K_{loc} \propto p^{-2}$  shown with a dotted line as a guide to the eye. The black squares are the localization size predicted by our model for every studied perturbation using the measured parameters  $\chi'(1)$  and  $\alpha$  from the black symbols in Fig. 6 (b) and (c). The red circles are the predicted localization size determined from our model assuming the single-parameter Ansatz of Eq. (40) using the experimentally determined parameters  $s, \nu, \alpha_0$  and  $\alpha_\infty$ . The predicted localization size  $K_{loc}$  exhibits two different scaling regimes as a function of the perturbation strength, for weak and strong perturbations. The determined critical perturbation  $p_c$ , where this transition occurs, is shown with a vertical dashed line.

bation strength  $p$  compared with the ones determined from the experimental data (blue diamonds scatters). We determined  $K_{loc}$  from Eq. (39) which only depends on the eigenvector coefficients  $c_{L1}$  associated to the smallest eigenvalue  $\lambda_1$ . The calculation of a single eigenvector can be performed in shorter times compared with solving the full system dynamics, allowing us to calculate  $K_{loc}$  for systems with  $N = 2 \cdot 10^5$ . The computed values of  $K_{loc}$  evidence two different regimes. For large perturbations ( $p > 0.025$ ) our model predicts a scaling for the localization size  $K_{loc}(p) \propto p^{-2}$ . For weak perturbations ( $p < 0.025$ ), we observe a weaker dependence of the localization size as a function of the perturbation that can be described approximately by  $K_{loc}(p) \propto p^{-0.5}$ . This transition is consistent with the transition on the decoherence scaling exponent  $\alpha$  shown in Fig. 6(b). Experimental evidence of localization effects are only observed for  $p > 0.026$ , therefore only this experimental data is shown in Fig. 7. Within this perturbation regime the experimental values exhibits a functional dependence  $K_{loc}(p) \propto p^{-2}$  consistently with our predictions from the model. We do not observe experimental evidence of localization effects for weaker perturbations within the accessible time which is limited by the decoherence decay of the NMR signal. The largest cluster sizes observed ex-

perimentally for the weak perturbations are always lower than the predicted localization sizes.

In a previous work [18], we showed experimental evidence that the scaling behavior of decoherence as a function of the cluster size is consistent with the single-parameter Ansatz for the asymptotic functional dependence at long times

$$\chi'(p, K) \sim \begin{cases} (p - p_c)^s K^{\alpha_0} & p < p_c \\ (p - p_c)^{-2\nu} K^{\alpha_\infty} & p > p_c, \end{cases} \quad (40)$$

where we obtained the critical exponents  $s = (-0.911 \pm 0.004)$  and  $\nu = (-0.57 \pm 0.03)$ . The scaling exponents are  $\alpha_\infty = 0.96 \pm 0.02$ , near to a linear scaling, and  $\alpha_0 = (0.48 \pm 0.03)$ . The determined critical perturbation is  $p_c = (0.026 \pm 0.006)$ . Due to the finite evolution time of the experimental data, the transition from one regime to another is smooth (see Fig. 6). The experimental data give a smooth functional behavior also for the decay rate  $\Gamma(K) = \chi'(K)$  that we introduced in our model [black squares in Fig. 6(b,c)]. To further evaluate the consistency of the scaling behavior determined by Eq. (40) and extrapolate its predicted behavior for the weak perturbations where we do not have experimental data, we assume this functional behavior for  $\Gamma(K) = \chi'(K)$  using the extracted parameters  $s, \nu, \alpha_0$  and  $\alpha_\infty$ . Figure 7 shows with red circles the prediction for the localization size  $K_{loc}$  as a function of the perturbation strength  $p$ . The localization-size  $K_{loc}$  determined from Eq. (39) now fit better the experimental data for the strong perturbations ( $p > 0.025$ ). Again for weak perturbations ( $p < 0.025$ ), we observe a weaker dependence of the localization size as a function of the perturbation strength. However, our model now predicts a finite localization size when  $p \rightarrow 0$ .

Both assumptions for  $\Gamma(K) = \chi'(K)$  manifest different scaling laws for  $K_{loc}$  comparing the weak and strong perturbation regimes. For weak perturbations, it seems from the predicted curves that there might exist a limiting value for  $K_{loc}$  for  $p \rightarrow 0$ . Further experimental designs need to be implemented to verify if the experimental behavior match these predictions, but we expect that this limiting value is due to intrinsic perturbations on the experimental implementation that are not accounted in our microscopic model described in Sec. II and in Refs. [9, 18].

## VI. CONCLUSION AND DISCUSSIONS

We developed a model for studying the quantum information scrambling dynamics measured with out-of-time-order correlations by Loschmidt echoes combined with Multiple Quantum Coherence experiments. Experimental implementations ubiquitously contain imperfections in the quantum operations that leads to the presence of non-reversed interactions in the LE procedure for

measuring the OTOC. Based on an imperfect experimental protocol, we derived expressions for OTOC functions connected with the effective number of correlated spins that are active on quantum superpositions generated by coherence transfers of a local information, that survived the perturbation effects. Decoherence effects induced by the time reversal imperfection arise naturally by the derived OTOCs as a leakage of the ideal unitary dynamics. The derived OTOCs quantify the observable degree of scrambling of information based on an inner-product between the ideal and the perturbed scrambling dynamics. The main prediction of our model is the existence of localization effects on the measurable information scrambling that bound the effective cluster size  $K(t)$  where the ideal information was propagated. We also found that whether the initial information on  $K(0)$  is local or not, the dynamics of the effective cluster size  $K(t)$  tends to a dynamical equilibrium value  $K_{loc}$ . Our prediction were contrasted with quantum simulations performed with NMR experiments on a solid state adamantane sample, showing excellent quantitative agreement with the experimental observations.

Levy and Gleason originally proposed a model to describe the dynamics of the spin-cluster of correlated spins in MQC experiments in solid-state samples. The model was based on simplifying the spin quantum dynamics to a phenomenological equation depending on two physical parameters: the mean value of the spin-spin interactions and the number of dimensions of the system. It correctly described how the number of correlated spins grows in NMR solid-state experiments with only these parameters. Our work revisit this model and adapt it to treat information scrambling. We also introduced a decoherence process induced by imperfection on the experimental implementations into the model. The decoherence process is described by a leakage rate that depends on the number of active spins involved on the states that describe the dynamics of the system. Determining this rates from experimental data, we obtained accurate predictions of the cluster size growth as a function of time and the localization values that were experimentally observed. Our results indicates that quantum information scrambling dynamics and its localization effects due to perturbations are phenomena that can be predicted in terms of few physical parameters by the presented model.

Treating many-body dynamics with exact numerical solutions is not possible with present technology. Therefore, the results shown in this work provide a framework for describing the quantum information scrambling dynamics of many-body systems determined from experiments affected by non-unitary decays, either by imperfections on the control or from interaction with external degrees of freedom. In this article, we have focused on modeling the dynamic of quantum information based on OTOC functions that give the cluster size of correlated spins generated by the scrambling dynamics. However, the model building block is based on finding the density matrix evolution of the system by solving the Liouville-

von Neumann equation reduced to average operator of the active spins in the dynamics. Therefore, one can envisage that the approach can be adapted to study the dynamic behavior of relevant physical quantities such other types of correlations in the presence of decoherence effects [89–93]. The framework presented here can be a useful tool to predict the quantum information dynamics in large quantum systems, and address the effect of imperfections on the control Hamiltonian that drives the quantum evolutions.

## ACKNOWLEDGMENTS

We thank M.C. Rodriguez for helpful discussions. This work was supported by CNEA, ANPCyT-FONCyT PICT-2017-3447, PICT-2017-3699, PICT-2018-04333, PIP-CONICET (11220170100486CO), UNCUIYO SIIP Tipo I 2019-C028, and Instituto Balseiro. F.D.D. and G.A.A. acknowledge support from CONICET.

### Appendix A: Quantum information scrambling as the cluster size of active spins

We here demonstrate that the norm of the commutator  $[I_z, \sigma^0(t)]$  quantifies the average number of active spins in the operator  $\sigma^0(t) = U_0(t)\sigma U_0^\dagger(t)$  with  $\sigma$  an arbitrary operator of the system. Then, when the operator  $\sigma = I_z$ , the following demonstration relates the cluster size of correlated spins of Eq. (13) and the OTO commutator of Eq. (11). We first calculate the commutator  $[I_z, \sigma^0(t)]$  of the OTOC expression of Eq. (11). We find that  $[I_z, \sigma^0(t)] = \sum_j^N \sum_{\vec{u}} C_{\vec{u}}^0(t) [I_z^j, P_{\vec{u}}]$ , where we used that  $\sigma^0(t) = \sum_{\vec{u}} C_{\vec{u}}^0(t) P_{\vec{u}}$  and  $I_z = \sum_j^N I_z^j$ . The operator  $I_z$  is a mixture of local spin operators  $I_z^j$ ,

$$I_z^j = \underbrace{\mathbb{I} \otimes \dots \otimes}_{j-1} \underbrace{I_z^{(1)}}_j \otimes \underbrace{\dots \otimes \mathbb{I}}_{N-j}. \quad (\text{A1})$$

The operators  $P_{\vec{u}}$  are the elements of the orthonormal product basis  $\left\{ (\sqrt{2})^N \otimes_{k=1}^N I_{u_k}^{(1)} \right\}$ , where the coefficients of the vector  $\vec{u}$  are  $u_i \in \{x, y, z, 0\}$  (0 for identity operator  $I_0 = \mathbb{I}$ ). By using the following property of the Kronecker product

$$[A \otimes B, C \otimes D] = [A, C] \otimes BD + CA \otimes [B, D] \quad (\text{A2})$$

and the expression for  $I_z^j$  of Eq. (A1), the commutator  $[I_z^j, P_{\vec{u}}]$  results

$$[I_z^j, P_{\vec{u}}] = \left( \sqrt{2} \right)^N \underbrace{I_{u_1} \otimes \dots \otimes [I_z^{(1)}, I_{u_j}] \otimes \dots \otimes I_{u_N}}_{j-1 \quad N-j} \\ = \begin{cases} iP_{\vec{u}'(j)} & \text{if } u_j = x \\ -iP_{\vec{u}'(j)} & \text{if } u_j = y \\ 0 & \text{if } u_j = 0, z \end{cases}. \quad (\text{A3})$$

The operator  $P_{\vec{u}'(j)}$  belongs to the product basis  $\{P_{\vec{u}}\}$ , where the vector  $\vec{u}'(j)$  is a vector identical to  $\vec{u}$ , except for the  $j$ -th element (e.g. if  $u_j = x$  then  $u'(j) = y$  and vice versa). We have used that  $[I_z^{(1)}, I_x^{(1)}] = iI_y^{(1)}$ ,  $[I_z^{(1)}, I_y^{(1)}] = -iI_x^{(1)}$  and  $[I_z^{(1)}, I_z^{(1)}] = [I_z^{(1)}, \mathbb{I}] = 0$ . This means that if  $[I_z^j, P_{\vec{u}}] \neq 0$ , then  $[I_z^j, P_{\vec{u}}]$  is proportional to other element  $P_{\vec{u}'(j)}$  of the product basis  $\{P_{\vec{u}}\}$ . The sign of  $[I_z^j, P_{\vec{u}}]$  depends on whether  $u_j = x$  (+) or  $u_j = y$  (-).

We then find that the commutator

$$[I_z, P_{\vec{u}}] = \sum_{j=1}^N [I_z^j, P_{\vec{u}}] = \sum_{j \in \mathcal{A}(\vec{u})} \pm iP_{\vec{u}'(j)},$$

where  $\mathcal{A}(\vec{u})$  is the set of indexes  $\{j\}$  corresponding to active spins in  $\vec{u}$ , this is, those which satisfy  $u_j = x, y$ . The set  $\mathcal{A}(\vec{u})$  has  $L(\vec{u})$  elements, where  $L(\vec{u})$  is the number of elements  $u_j$  equal to  $x$  and  $y$ , which implies that  $[I_z, P_{\vec{u}}]$  has  $L(\vec{u})$  non-zero terms  $\pm iP_{\vec{u}'(j)}$ . The resulting expression for the OTOC and therefore for cluster size  $K_0(t)$  is then

$$K_0(t) = \text{Tr} \left\{ [I_z, \sigma^0(t)] [I_z, \sigma^0(t)]^\dagger \right\} \\ = \sum_{\vec{u}, \vec{v}} C_{\vec{u}}^0 C_{\vec{v}}^{0*} \times \\ \times \sum_{i \in \mathcal{A}(\vec{u})} \sum_{j \in \mathcal{A}(\vec{v})} \text{Tr} \left[ (\pm iP_{\vec{u}'(i)}) \cdot (\pm iP_{\vec{v}'(j)})^\dagger \right]. \quad (\text{A4})$$

Finally, we prove that the factor of the previous equation is equal to the function  $\mathcal{L}(\vec{u}, \vec{v})$  introduced in Eq. (14)

$$\mathcal{L}(\vec{u}, \vec{v}) = \sum_{i \in \mathcal{A}(\vec{u})} \sum_{j \in \mathcal{A}(\vec{v})} \text{Tr} \left[ (\pm iP_{\vec{u}'(i)}) \cdot (\pm iP_{\vec{v}'(j)})^\dagger \right]. \quad (\text{A5})$$

To prove this, we divide the demonstration in five different propositions as we discuss below. We introduce the notation  $h_{0z}(\vec{u}, \vec{v})$  for the Hamming distance between  $\vec{u}$  and  $\vec{v}$  considering only the elements  $0, z$ , and  $h_{xy}(\vec{u}, \vec{v})$  for the Hamming distance between  $\vec{u}$  and  $\vec{v}$  considering only the elements  $x, y$ . In Propositions 1 and 2, we deduce the necessary conditions that  $\vec{u}$  and  $\vec{v}$  must satisfy for obtaining  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ . Then, Propositions 3, 4 and 5 provides the values of  $\mathcal{L}(\vec{u}, \vec{v})$  when the conditions deduced in Propositions 1 and 2 are satisfied.

*Proposition 1* If  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ , then  $h_{0z}(\vec{u}, \vec{v}) = 0$ .

*Proof:* Since  $\{P_{\vec{u}}\}$  is an orthonormal base  $\text{Tr}(P_{\vec{u}'(i)} \cdot P_{\vec{v}'(j)}) = \delta_{\vec{u}'(i), \vec{v}'(j)}$ , to obtain  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ , there must be states  $\vec{u}'(i) = \vec{v}'(j)$  for some  $i \in \mathcal{A}(\vec{u})$ ,  $j \in \mathcal{A}(\vec{v})$ . Since the vectors  $\vec{u}'(i)$  and  $\vec{v}'(j)$  are identical to  $\vec{u}$  and  $\vec{v}$  respectively, in those elements that are equal to  $z, 0$ , it is necessary that  $u_l = v_l$  if  $l = 0, z$  for  $\vec{u}'(i) = \vec{v}'(j)$  to exist. Therefore, the Hamming distance  $h_{0z}(\vec{u}, \vec{v}) = 0$ . ■

*Proposition 2* If  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ , then  $h_{xy}(\vec{u}, \vec{v}) = 0$  or  $h_{xy}(\vec{u}, \vec{v}) = 2$ .

*Proof:* Again, to obtain  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ , there must be states  $\vec{u}'(i) = \vec{v}'(j)$  for some  $i \in \mathcal{A}(\vec{u})$ ,  $j \in \mathcal{A}(\vec{v})$ . We prove Proposition 2 in two steps.

(i) We can see first that if  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ ,  $h_{xy}(\vec{u}, \vec{v}) \leq 2$ . From the definition of the vectors  $\vec{u}'(i)$ , we know that  $\vec{u}'(i)$  only differs from  $\vec{u}$  in the  $i$ -th element, and that  $u_i$  must be  $x$  or  $y$ . We thus deduce that  $h_{xy}(\vec{u}, \vec{u}'(i)) = h_{xy}(\vec{v}, \vec{v}'(j)) = 1$ , for every  $i \in \mathcal{A}(\vec{u})$ ,  $j \in \mathcal{A}(\vec{v})$ . If  $\vec{u}'(i) = \vec{v}'(j)$  for some  $i, j$ , we then deduce from the triangle inequality that

$$h_{xy}(\vec{u}, \vec{v}) \leq h_{xy}(\vec{u}, \vec{u}'(i)) + h_{xy}(\vec{v}, \vec{v}'(i)) \\ = h_{xy}(\vec{u}, \vec{u}'(i)) + h_{xy}(\vec{v}, \vec{v}'(j)) \\ = 2. \quad (\text{A6})$$

(ii) We can also see that  $h_{xy}(\vec{u}, \vec{v}) \neq 1$ . Let us suppose that  $h_{xy}(\vec{u}, \vec{v}) = 1$ , then there is a unique index  $l$  such that  $u_l \neq v_l$ . Therefore  $\vec{v} = \vec{u}'(l)$ , and then it is impossible to find  $\vec{v}'(j)$  such that  $\vec{u}'(l) = \vec{v}'(j)$ , because  $\vec{v}'(j) \neq \vec{v} = \vec{u}'(l)$ . If we consider  $\vec{u}'(i)$  (with  $i \neq l$ ) then  $h_{xy}(\vec{u}'(i), \vec{v}) = 2$  and again it is impossible to find  $\vec{v}'(j)$  such that  $\vec{u}'(i) = \vec{v}'(j)$ . We have proven that if  $h_{xy}(\vec{u}, \vec{v}) = 1$  then  $\mathcal{L}(\vec{u}, \vec{v}) = 0$ . Therefore, if  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ , then  $h_{xy}(\vec{u}, \vec{v}) \neq 1$ . Therefore, if  $\mathcal{L}(\vec{u}, \vec{v}) \neq 0$ ,  $h_{xy}(\vec{u}, \vec{v}) = 0$  or  $h_{xy}(\vec{u}, \vec{v}) = 2$ . ■

*Proposition 3*  $\mathcal{L}(\vec{u}, \vec{u}) = L(\vec{u})$ , where  $L(\vec{u})$  is the number of elements in  $\vec{u}$  that are equal to  $x, y$ . Notice that in this case when  $\vec{u} = \vec{v}$ , the Hamming distances are  $h_{xy}(\vec{u}, \vec{v}) = 0$  and  $h_{0z}(\vec{u}, \vec{v}) = 0$ .

*Proof:* Since  $\{P_{\vec{u}}\}$  is an orthonormal basis,  $\text{Tr}(P_{\vec{u}'(i)} \cdot P_{\vec{v}'(j)}) = \delta_{\vec{u}'(i), \vec{v}'(j)}$ . As the number of elements in  $\mathcal{A}(\vec{u})$  is  $L(\vec{u})$ , we get

$$\mathcal{L}(\vec{u}, \vec{u}) = \sum_{i \in \mathcal{A}(\vec{u})} \sum_{j \in \mathcal{A}(\vec{u})} \text{Tr} \left[ (\pm iP_{\vec{u}'(i)}) \cdot (\pm iP_{\vec{u}'(j)})^\dagger \right] \\ = -i^2 \sum_{j \in \mathcal{A}(\vec{u})} 1 \\ = L(\vec{u}). \quad (\text{A7})$$

*Proposition 4* If  $h_{xy}(\vec{u}, \vec{v}) = 2$ ,  $h_{0z}(\vec{u}, \vec{v}) = 0$  and  $\vec{u}$  is not a permutation of  $\vec{v}$ , i.e.  $\vec{u} \neq \Pi(\vec{v})$ , then  $\mathcal{L}(\vec{u}, \vec{v}) = -2$ .

*Proof:* Since  $h_{xy}(\vec{u}, \vec{v}) = 2$  and  $h_{0z}(\vec{u}, \vec{v}) = 0$ , there are only two indexes  $l_1$  and  $l_2$  for which  $u_{l_1} \neq v_{l_1}$  and  $u_{l_2} \neq v_{l_2}$ . Since  $l_1, l_2 \in \{x, y\}$  and  $\vec{u} \neq \Pi(\vec{v})$ , there must be the following conditions  $u_{l_1} = x$ ,  $u_{l_2} = x$  and  $v_{l_1} = y$ ,  $v_{l_2} = y$ , or vice versa that is equivalent to exchange  $\vec{u}$  and  $\vec{v}$ . Then, we have that  $\vec{u}'(l_1) = \vec{v}'(l_2)$  and  $\vec{u}'(l_2) = \vec{v}'(l_1)$ . These are the only two terms different from zero in Eq. (A5). If  $u_{l_1} = u_{l_2} = x$ , then the operators  $P_{\vec{u}'(l_1)}$  and  $P_{\vec{u}'(l_2)}$  have a multiplicative factor  $i$ . Analogously, since  $v_{l_1} = v_{l_2} = y$ , then the operators  $P_{\vec{v}'(l_1)}$  and  $P_{\vec{v}'(l_2)}$  have a multiplicative factor  $-i$ . Therefore, the operator products  $P_{\vec{u}'(l_1)} \cdot P_{\vec{v}'(l_2)}$  and  $P_{\vec{u}'(l_2)} \cdot P_{\vec{v}'(l_1)}$  are proportional to  $i^2 = -1$ , and the function  $\mathcal{L}(\vec{u}, \vec{v})$  is

$$\begin{aligned} \sum_{i \in \mathcal{A}(\vec{u})} \sum_{j \in \mathcal{A}(\vec{v})} \text{Tr} \left[ (\pm i P_{\vec{u}'(i)}) \cdot (\pm i P_{\vec{v}'(j)})^\dagger \right] &= \\ &= i \cdot (-i)^* \text{Tr} (P_{\vec{u}'(l_1)} \cdot P_{\vec{v}'(l_2)}) + \\ &+ (-i)^* \cdot i \text{Tr} (P_{\vec{v}'(l_1)} \cdot P_{\vec{u}'(l_2)}) \\ &= -2. \quad (\text{A8}) \end{aligned}$$

*Proposition 5* If  $h_{xy}(\vec{u}, \vec{v}) = 2$ ,  $h_{0z}(\vec{u}, \vec{v}) = 0$  and  $\vec{u}$  is a permutation of  $\vec{v}$ , i.e.  $\vec{u} = \Pi(\vec{v})$ , then  $\mathcal{L}(\vec{u}, \vec{v}) = 2$ .

*Proof:* Since  $h_{xy}(\vec{u}, \vec{v}) = 2$  and  $h_{0z}(\vec{u}, \vec{v}) = 0$ , there are only two indexes  $l_1$  and  $l_2$  for which  $u_{l_1} \neq v_{l_1}$  and  $u_{l_2} \neq v_{l_2}$ . Since  $l_1, l_2 \in \{x, y\}$  and  $\vec{u} = \Pi(\vec{v})$ , it must hold that  $u_{l_1} = x$ ,  $u_{l_2} = y$  and  $v_{l_1} = y$ ,  $v_{l_2} = x$ . Then, we have that  $\vec{u}'(l_1) = \vec{v}'(l_2)$  and  $\vec{u}'(l_2) = \vec{v}'(l_1)$ . These are the only two terms different from zero in Eq. (A5). Since  $u_{l_1} = v_{l_2} = x$ , then the operators  $P_{\vec{u}'(l_1)}$  and  $P_{\vec{v}'(l_2)}$  have a multiplicative factor  $i$ . Analogously, since

$v_{l_1} = u_{l_2} = y$ , then the operators  $P_{\vec{v}'(l_1)}$  and  $P_{\vec{u}'(l_2)}$  have a multiplicative factor  $-i$ . Therefore, the operator products  $P_{\vec{u}'(l_1)} \cdot P_{\vec{v}'(l_2)}$  and  $P_{\vec{u}'(l_2)} \cdot P_{\vec{v}'(l_1)}$  are proportional to  $-i^2 = 1$ , and the function  $\mathcal{L}(\vec{u}, \vec{v})$  is

$$\begin{aligned} \sum_{i \in \mathcal{A}(\vec{u})} \sum_{j \in \mathcal{A}(\vec{v})} \text{Tr} \left[ (\pm i P_{\vec{u}'(i)}) \cdot (\pm i P_{\vec{v}'(j)})^\dagger \right] &= \\ &= i \cdot (i)^* \text{Tr} (P_{\vec{u}'(l_1)} \cdot P_{\vec{v}'(l_2)}) + (i)^* \cdot i \text{Tr} (P_{\vec{v}'(l_1)} \cdot P_{\vec{u}'(l_2)}) \\ &= 2. \end{aligned} \quad (\text{A9})$$

These five propositions thus demonstrate the expression of Eq. (14) in the main text for the function  $\mathcal{L}(\vec{u}, \vec{v})$ .

For the perturbed case the effective cluster size  $K(t)$  of Eq. (21) is derived directly from the previous demonstration. But, we now write  $\sigma^0(t) = \sum_{\vec{u}} C_{\vec{u}}^0(t) P_{\vec{u}}$  and  $\sigma(t) = \sum_{\vec{u}} C_{\vec{u}}(t) P_{\vec{u}}$ , and then it results

$$\begin{aligned} K(t) &= \text{Tr} [[I_z, \sigma^0(t)][I_z, \sigma(t)]^\dagger] \\ &= \sum_{\vec{u}, \vec{v}} C_{\vec{u}}^0(t) C_{\vec{v}}^*(t) \left\{ \sum_{i \in \mathcal{A}(\vec{u})} \sum_{j \in \mathcal{A}(\vec{v})} \text{Tr} \left[ (\pm i P_{\vec{u}'(i)}) \cdot (\pm i P_{\vec{v}'(j)})^\dagger \right] \right\}, \end{aligned} \quad (\text{A10})$$

where the function  $\mathcal{L}(\vec{u}, \vec{v}) = \sum_{i \in \mathcal{A}(\vec{u})} \sum_{j \in \mathcal{A}(\vec{v})} \text{Tr} \left[ (\pm i P_{\vec{u}'(i)}) \cdot (\pm i P_{\vec{v}'(j)})^\dagger \right]$  is the same of Eq. (14) for the non-perturbed case. The expression for  $K(t)$  of Eq. (21) is obtained when  $\sigma^0(t) = I_z^0(t)$  and  $\sigma(t) = I_z(t)$ .

- 
- [1] Y. Sekino and L. Susskind, *J. High Energy Phys.* **2008**, 065.
- [2] N. Lashkari, D. Stanford, M. Hastings, T. Osborne, and P. Hayden, *J. High Energy Phys.* **2013**, 22.
- [3] E. A. Martinez, C. A. Muschik, P. Schindler, D. Nigg, A. Erhard, M. Heyl, P. Hauke, M. Dalmonte, T. Monz, P. Zoller, and R. Blatt, *Nature* **534**, 516 (2016).
- [4] N. Friis, O. Marty, C. Maier, C. Hempel, M. Holzäpfel, P. Jurcevic, M. B. Plenio, M. Huber, C. Roos, R. Blatt, and B. Lanyon, *Phys. Rev. X* **8**, 21012 (2018).
- [5] B. Swingle, *Nat. Phys.* **14**, 988 (2018).
- [6] R. J. Lewis-Swan, A. Safavi-Naini, A. M. Kaufman, and A. M. Rey, *Nat. Rev. Phys.* **1**, 627 (2019).
- [7] J. Eisert, M. Friesdorf, and C. Gogolin, *Nat. Phys.* **11**, 124 (2015).
- [8] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, *Rev. Mod. Phys.* **91**, 21001 (2019).
- [9] G. A. Álvarez, D. Suter, and R. Kaiser, *Science* **349**, 846 (2015).
- [10] T. Schweigler, V. Kasper, S. Erne, I. Mazets, B. Rauer, F. Cataldini, T. Langen, T. Gasenzer, J. Berges, and J. Schmiedmayer, *Nature* **545**, 323 (2017).
- [11] A. Lukin, M. Rispoli, R. Schittko, M. E. Tai, A. M. Kaufman, S. Choi, V. Khemani, J. Léonard, and M. Greiner, *Science* **364**, 256 (2019).
- [12] K. A. Landsman, C. Figgatt, T. Schuster, N. M. Linke, B. Yoshida, N. Y. Yao, and C. Monroe, *Nature* **567**, 61 (2019).
- [13] T. Brydges, A. Elben, P. Jurcevic, B. Vermersch, C. Maier, B. P. Lanyon, P. Zoller, R. Blatt, and C. F. Roos, *Science* **364**, 260 (2019).
- [14] H. G. Krojanski and D. Suter, *Phys. Rev. Lett.* **93**, 090501 (2004).
- [15] G. A. Álvarez and D. Suter, *Phys. Rev. Lett.* **104**, 230403 (2010).
- [16] C. M. Sánchez, R. H. Acosta, P. R. Levstein, H. M. Pastawski, and A. K. Chattah, *Phys. Rev. A* **90**, 042122 (2014).
- [17] M. Niknam, L. F. Santos, and D. G. Cory, *Phys. Rev. Research* **2**, 13200 (2020).

- [18] F. D. Domínguez, M. C. Rodríguez, R. Kaiser, D. Suter, and G. A. Álvarez, *Phys. Rev. A* **104**, 012402 (2021).
- [19] P. Hosur, X.-L. Qi, D. A. Roberts, and B. Yoshida, *J. High Energy Phys.* **2016**, 4.
- [20] M. Garttner, J. G. Bohnet, A. Safavi-Naini, M. L. Wall, J. J. Bollinger, and A. M. Rey, *Nat. Phys.* **13**, 781 (2017).
- [21] J. Li, R. Fan, H. Wang, B. Ye, B. Zeng, H. Zhai, X. Peng, and J. Du, *Phys. Rev. X* **7**, 031011 (2017).
- [22] D. A. Roberts, D. Stanford, and L. Susskind, *J. High Energy Phys.* **2015**, 1.
- [23] A. Larkin and Y. N. Ovchinnikov, *Sov. Phys. JETP* **28**, 1200 (1969).
- [24] S. H. Shenker and D. Stanford, *J. High Energy Phys.* **2014**, 67.
- [25] J. Maldacena, S. H. Shenker, and D. Stanford, *J. High Energy Phys.* **2016**, 106.
- [26] I. García-Mata, M. Saraceno, R. A. Jalabert, A. J. Roncaglia, and D. A. Wisniacki, *Phys. Rev. Lett.* **121**, 210601 (2018).
- [27] K. X. Wei, C. Ramanathan, and P. Cappellaro, *Phys. Rev. Lett.* **120**, 070501 (2018).
- [28] K. X. Wei, P. Peng, O. Shtanko, I. Marvian, S. Lloyd, C. Ramanathan, and P. Cappellaro, *Phys. Rev. Lett.* **123**, 090605 (2019).
- [29] C. M. Sánchez, A. K. Chattah, K. X. Wei, L. Buljubasich, P. Cappellaro, and H. M. Pastawski, *Phys. Rev. Lett.* **124**, 030601 (2020).
- [30] M. K. Joshi, A. Elben, B. Vermersch, T. Brydges, C. Maier, P. Zoller, R. Blatt, and C. F. Roos, *Phys. Rev. Lett.* **124**, 240505 (2020).
- [31] X. Nie, B. B. Wei, X. Chen, Z. Zhang, X. Zhao, C. Qiu, Y. Tian, Y. Ji, T. Xin, D. Lu, and J. Li, *Phys. Rev. Lett.* **124**, 250601 (2020).
- [32] X. Mi, P. Roushan, C. Quintana, S. Mandra, J. Marshall, C. Neill, F. Arute, K. Arya, J. Atalaya, R. Babbush, J. C. Bardin, R. Barends, A. Bengtsson, S. Boixo, A. Bourassa, M. Broughton, B. B. Buckley, D. A. Buell, B. Burkett, N. Bushnell, Z. Chen, B. Chiaro, R. Collins, W. Courtney, S. Demura, A. R. Derk, A. Dunsworth, D. Eppens, C. Erickson, E. Farhi, A. G. Fowler, B. Foxen, C. Gidney, M. Giustina, J. A. Gross, M. P. Harrigan, S. D. Harrington, J. Hilton, A. Ho, S. Hong, T. Huang, W. J. Huggins, L. B. Ioffe, S. V. Isakov, E. Jeffrey, Z. Jiang, C. Jones, D. Kafri, J. Kelly, S. Kim, A. Kitaev, P. V. Klimov, A. N. Korotkov, F. Kostritsa, D. Landhuis, P. Laptev, E. Lucero, O. Martin, J. R. McClean, T. McCourt, M. McEwen, A. Megrant, K. C. Miao, M. Mohseni, W. Mruczkiewicz, J. Mutus, O. Naaman, M. Neeley, M. Newman, M. Y. Niu, T. E. O'Brien, A. Opremcak, E. Ostby, B. Pato, A. Petukhov, N. Redd, N. C. Rubin, D. Sank, K. J. Satzinger, V. Shvarts, D. Strain, M. Szalay, M. D. Trevithick, B. Villalonga, T. White, Z. J. Yao, P. Yeh, A. Zalcman, H. Neven, I. Aleiner, K. Kechedzhi, V. Smelyanskiy, and Y. Chen, Information scrambling in computationally complex quantum circuits (2021), [arXiv:2101.08870](https://arxiv.org/abs/2101.08870).
- [33] A. Peres, *Phys. Rev. A* **30**, 1610 (1984).
- [34] R. A. Jalabert and H. M. Pastawski, *Phys. Rev. Lett.* **86**, 2490 (2001).
- [35] P. Jacquod and C. Petitjean, *Adv. Phys.* **58**, 67 (2009).
- [36] T. Gorin, T. Prosen, T. H. Seligman, and M. Žnidarič, *Phys. Rep.* **435**, 33 (2006).
- [37] A. Goussev, R. A. Jalabert, H. M. Pastawski, and D. A. Wisniacki, *Scholarpedia* **7**, 11687 (2012).
- [38] D. Suter and G. A. Álvarez, *Rev. Mod. Phys.* **88**, 041001 (2016).
- [39] B. Swingle and N. Yunger Halpern, *Phys. Rev. A* **97**, 062113 (2018).
- [40] S. V. Syzranov, A. V. Gorshkov, and V. Galitski, *Phys. Rev. B* **97**, 161114 (2018).
- [41] J. R. González Alonso, N. Yunger Halpern, and J. Dreschel, *Phys. Rev. Lett.* **122**, 040404 (2019).
- [42] J. Tuziemski, *Phys. Rev. A* **100**, 062106 (2019).
- [43] P. Zanardi and N. Anand, *Phys. Rev. A* **103**, 062214 (2021).
- [44] J. Baum, M. Munowitz, A. N. Garroway, and A. Pines, *J. Chem. Phys.* **83**, 2015 (1985).
- [45] M. Munowitz, A. Pines, and M. Mehring, *J. Chem. Phys.* **86**, 3172 (1987).
- [46] S. Lacelle, On the Growth of Multiple Spin Coherences in NMR of Solids, in *Adv. Magn. Opt. Reson.*, Vol. 16 (Elsevier, 1991) pp. 173–263.
- [47] S. Lacelle, S. J. Hwang, and B. C. Gerstein, *J. Chem. Phys.* **99**, 8407 (1993).
- [48] C. E. Hughes, *Prog. Nucl. Magn. Reson. Spectrosc.* **45**, 301 (2004).
- [49] J. B. Murdoch, W. S. Warren, D. P. Weitekamp, and A. Pines, *J. Magn. Reson.* **60**, 205 (1984).
- [50] D. Levy and K. Gleason, *J. Phys. Chem.* **96**, 8125 (1992).
- [51] V. E. Zobov and A. A. Lundin, *J. Exp. Theor. Phys.* **103**, 904 (2006).
- [52] V. E. Zobov and A. A. Lundin, *Russ. J. Phys. Chem. B* **2**, 676 (2008).
- [53] Y. Mogami, Y. Noda, H. Ishikawa, and K. Takegoshi, *Phys. Chem. Chem. Phys.* **15**, 7403 (2013).
- [54] V. E. Zobov and A. A. Lundin, *J. Exp. Theor. Phys.* **112**, 451 (2011).
- [55] A. A. Lundin and V. E. Zobov, *Appl. Magn. Reson.* **47**, 701 (2016).
- [56] G. Cho and J. P. Yesinowski, *Chem. Phys. Lett.* **205**, 1 (1993).
- [57] G. Cho and J. P. Yesinowski, *J. Phys. Chem.* **100**, 15716 (1996).
- [58] H. Cho, T. D. Ladd, J. Baugh, D. G. Cory, and C. Ramanathan, *Phys. Rev. B* **72**, 054427 (2005).
- [59] G. A. Álvarez and D. Suter, *Phys. Rev. A* **84**, 012320 (2011).
- [60] G. A. Álvarez, R. Kaiser, and D. Suter, *Ann. Phys.* **525**, 833 (2013).
- [61] C. P. Slichter, *Principles of magnetic resonance* (Springer-Verlag Berlin Heidelberg, 1990).
- [62] U. Haeberlen and J. S. Waugh, *Phys. Rev.* **175**, 453 (1968).
- [63] M. Gärttner, P. Hauke, and A. M. Rey, *Phys. Rev. Lett.* **120**, 040402 (2018).
- [64] B. Yan, L. Cincio, and W. H. Zurek, *Phys. Rev. Lett.* **124**, 160603 (2020).
- [65] A. K. Khitrin, *Chem. Phys. Lett.* **274**, 217 (1997).
- [66] M. Munowitz and A. Pines, Principles and applications of multiple-quantum NMR, in *Adv. Chem. Phys.*, Vol. 66 (John Wiley & Sons, Ltd, 1986) pp. 1–152.
- [67] G. A. Álvarez, E. P. Danieli, P. R. Levstein, and H. M. Pastawski, *Phys. Rev. Lett.* **101**, 120503 (2008).
- [68] O. W. Sørensen, M. H. Levitt, and R. R. Ernst, *J. Magn. Reson.* **55**, 104 (1983).
- [69] C. Griesinger, O. W. Sørensen, and R. R. Ernst, *J. Chem. Phys.* **85**, 6837 (1986).
- [70] B. E. Scraggs and K. Gleason, *Chem. Phys.* **166**, 367



- (1992).
- [71] S. I. Doronin, E. B. Fel'dman, I. Y. Guinzbourg, and I. I. Maximov, *Chem. Phys. Lett.* **341**, 144 (2001).
- [72] M. Munowitz, *Mol. Phys.* **71**, 37 (2006).
- [73] S. I. Doronin, A. V. Fedorova, and A. I. Zenchuk, *J. Chem. Phys.* **131**, 104109 (2009).
- [74] S. I. Doronin, E. B. Fel'dman, and A. I. Zenchuk, *J. Chem. Phys.* **134**, 034102 (2011).
- [75] H. De Raedt and K. Michielsen, Computational methods for simulating quantum computers (2004), [arXiv:quant-ph/0406210](https://arxiv.org/abs/quant-ph/0406210).
- [76] W. Zhang, N. Konstantinidis, K. A. Al-Hassanieh, and V. V. Dobrovitski, *J. Condens. Matter Phys.* **19**, 083202 (2007).
- [77] G. M. Palma, K.-A. Suominen, and A. K. Ekert, *Proc. Math. Phys. Eng. Sci.* **452**, 20 (1997).
- [78] L. M. Duan and G. C. Guo, *Phys. Rev. A* **57**, 737 (1998).
- [79] J. H. Reina, L. Quiroga, and N. F. Johnson, *Phys. Rev. A* **65**, 032326 (2002).
- [80] J. Jing and X. Hu, *Sci. Rep.* **5**, 17013 (2015).
- [81] G. A. Álvarez, E. P. Danieli, P. R. Levstein, and H. M. Pastawski, *J. Chem. Phys.* **124**, 194507 (2006).
- [82] E. Danieli, G. Álvarez, P. Levstein, and H. Pastawski, *Solid State Commun.* **141**, 422 (2007).
- [83] I. Rotter, *J. Phys. A Math. Theor.* **42**, 153001 (2009).
- [84] I. Rotter, *J. Opt.* **12**, 065701 (2010).
- [85] I. Rotter and J. P. Bird, *Rep. Prog. Phys.* **78**, 114001 (2015).
- [86] V. M. Martínez Álvarez, J. E. Barrios Vargas, and L. E. F. Foa Torres, *Phys. Rev. B* **97**, 121401 (2018).
- [87] R. Metzler and J. Klafter, *Phys. Rep.* **339**, 1 (2000).
- [88] N. Mercadier, W. Guerin, M. Chevrollier, and R. Kaiser, *Nat. Phys.* **5**, 602 (2009).
- [89] J. Maziero, L. C. Céleri, R. M. Serra, and V. Vedral, *Phys. Rev. A* **80**, 044102 (2009).
- [90] J.-S. Xu, X.-Y. Xu, C.-F. Li, C.-J. Zhang, X.-B. Zou, and G.-C. Guo, *Nat. Commun.* **1**, 7 (2010).
- [91] A. Touil and S. Deffner, *PRX Quantum* **2**, 010306 (2021).
- [92] Z. Xu, A. Chenu, T. c. v. Prosen, and A. del Campo, *Phys. Rev. B* **103**, 064309 (2021).
- [93] G. Styliaris, N. Anand, and P. Zanardi, *Phys. Rev. Lett.* **126**, 030601 (2021).