



Abstracts from the

## **IV JOINT MEETING OF THE BIOLOGY SOCIETIES OF ARGENTINA**

**(Cuarta Reunión Conjunta de  
Sociedades de Biología de la  
República Argentina)**

**XXXVII Annual Scientific Meeting of the Tucumán Biology Association  
XXIII Annual Scientific Meeting of the Córdoba Biology Society  
XXXVIII Annual Scientific Meeting of the Cuyo Biology Society  
Argentine Biology Society  
Rosario Biology Society  
Chilean Society of Reproduction and Development**

September 9–15, 2020  
Online edition

*The abstracts have been revised and evaluated by a Scientific Committee prior to publication*

## SCIENTIFIC COMMITTEE

### ASOCIACIÓN DE BIOLOGÍA DE TUCUMÁN

Dra. María Teresa Ajmat  
Dra. Patricia L. Albornoz  
Dr. Mario Fortuna  
Dra. Lucrecia Iruzubieta Villagra  
Mag. Analía Salvatore  
Dr. Federico Bonilla  
Dra. Liliana I. Zelarayán  
Dra. María Eugenia Pérez  
Dra. Elisa Ofelia Vintiñi

### SOCIEDAD DE BIOLOGÍA DE CÓRDOBA

Dra. Graciela Borioli  
Dra. Paola Boeris  
Dra. Cecilia Conde  
Dra. Marta Dardanelli  
Dra. Elena Fernández  
Dr. Leonardo Fruttero  
Dra. Susana Genti-Raimondi  
Dr. Alejandro Guidobaldi  
Dr. Edgardo Jofré  
Dra. Melina Musri  
Dra. Graciela Panzetta-Dutari  
Dr. Germán Robert  
Dra. Luciana Torre  
Dra. Cristina Torres

### SOCIEDAD DE BIOLOGÍA DE CUYO

#### *MENDOZA*

Dra. Nora Arenas  
Dra. Silvia Belmonte  
Dra. Alejandra Camargo  
Dr. Diego Cargnelutti  
Dra. María Teresa Damiani  
Dra. María Inés Echeverría

Dr. Carlos Gamarra-Luques  
Vet. Paula Ginevro  
Dr. Diego Grilli  
Dr. Eduardo Koch  
Dra. Myriam Laconi  
Dr. Luis López  
Dra. Alejandra Mampel  
Dr. Walter Manucha  
Dr. Ricardo Masuelli  
Dra. Marcela Michaut  
Dra. Adriana Telechea  
Dr. Roberto Yunes

#### *SAN LUIS*

Dra. Silvina Álvarez  
Dra. Cristina Barcia  
Dra. María Eugenia Ciminari  
Dr. Juan Gabriel Chediack  
Dr. Fabricio Cid  
Dra. Gladys Ciuffo  
Lic. Óscar Córdoba Mascali  
Dra. María Esther Escudero  
Dra. Susana Ferrari  
Dra. Lucia Fuentes  
Esp. Mónica Laurentina Gatica  
Dra. Nidia Noemí Gomez  
Dra. Marta Moglia  
Esp. Facundo Morales  
Dra. Edith Pérez  
Dra. María Verónica Pérez Chaca  
Dra. Hilda Elizabeth Pedranzani  
Dra. Graciela Wendel  
Dra. Alba Edith Vega  
Dra. Liliana Villegas

#### *SAN JUAN*

Dra. Gabriela Feresín

## **SOCIEDAD DE BIOLOGÍA DE ROSARIO**

Dra. Ariana Diaz  
Méd. Vet. Melina Gay  
Dra. Graciela Klekailo  
Dra. Milagros López Hiriart  
Dra. Stella Mattaloni  
Dra. Nidia Montechiarini  
Dra. Alejandra Peruzzo  
Dr. Claudio Luis Pidone  
Dra. Marta Posadas  
Dra. Mariana Raviola  
Dra. María Elena Tosello  
Dra. Silvina Villar

## **SOCIEDAD ARGENTINA DE BIOLOGÍA**

Dra. Fernanda Parborell  
Dra. Débora Cohen  
Dra. Griselda Irusta  
Dra. Isabel María Lacau  
Dra. Silvina Pérez Martínez  
Dra. Evelin Elia  
Dra. Clara I. Marín Briggiler  
Dr. Leandro Miranda  
Dr. Pablo Cetica

were separated into three experimental groups: young *ad libitum* (3-month-old, Y-AL group, N = 10), older *ad libitum* (22-month-old, O-AL group, N = 10), and older subjected to a 40% CR treatment during the last 3 months prior to the 22 months of age (O-CR group, N = 5). Cognitive performance was assessed using the Barnes Maze (BM) test for spatial learning and memory and the New Object Recognition (NOR) test for contextual learning. In the BM test, we observed that the O-CR rats presented a shorter distance traveled on the platform, similar to Y-AL group. We did not find significant differences between O-CR and O-AL animals in the rest of the parameters analyzed with BM test (exploratory frequency of the target region, total exploratory activity, numbers of errors in reaching around the target hole, escape box latencies, percentage of exploration of the meta holes). In the NOR test, again we did not find significant differences between old animals and those subjected to CR. To date, the studies carried out on the effects of CR on cognitive functions are inconclusive and depend on the used protocol. The effects of CR depend on its intensity, the period of life in which the treatment begins and its duration. Our studies are preliminary, with a first group of animals in CR (N = 5), therefore increasing the number of studied animals could provide more conclusive data. CR could be a non-pharmacological alternative for maintaining mental and cognitive health during aging.

## A251

### CONTENT OF ZINC, MACRONUTRIENTS AND FIBER IN MENUS OFFERED IN ELDERLY HOMES.

#### “MACA ANDINA” AS A SUPPLEMENTATION PROPOSAL IN THE FACE OF ZINC DEFICIENCY

Vallejos AL<sup>1</sup>, Biasi AM<sup>1,2</sup>, Ramirez MM<sup>1</sup>, Luna FG<sup>1</sup>, Perez Chaca MV<sup>3</sup>, Gomez NN<sup>2,3</sup>

<sup>1</sup>Facultad de Ciencias de la Salud, UNSL. <sup>2</sup>IMIBIO-SL CONICET, San Luis. <sup>3</sup>Laboratorio de Morfofisiología, Facultad de Química, Bioquímica y Farmacia, UNSL. <sup>4</sup>Universidad Nacional de Villa Mercedes. E-mail: anto.mbiasi@gmail.com

The biochemical and physiological changes that accompany the aging process, associated with the components of institutionalized life, have clear implications for the nutritional status of the elderly, making them more susceptible to nutritional deficiencies. The purpose of this research was to know the average nutritional values of zinc, macronutrients and total fiber of lunches and dinners offered to the elderly between 75 and 90 years old, who reside in long-term accommodation centers in the province of San Luis, Argentina, and furthermore, in case of zinc deficiency, suggest supplementation with “maca andina” (*Lepidium meyenii*). The study was carried out in two stages, one of them included an observational design with a cross-sectional correlational descriptive scope, and the other one involved an experimental design, being the sampling of probabilistic and multistage type. The sample was made up of 44 menus, which were classified into menus without meat (N = 22) and menus with meat (N = 22). The nutritional composition of all the menus (with and without meat) was analyzed using the SARA software. In addition, an experimental analysis of the meat-free menus was carried out (for which 3 types of menus were taken) and “maca andina” was also analyzed in triplicate, using the corresponding analytical techniques. The composition per serving of the meat-free menus was: 497.69 kcal, 58.26 g of carbohydrates (CHO), 15.81 g of proteins, 21.59 g of total lipids, 6.31 g of total fiber, and 2.29 mg of zinc. In the menus with meat, the average nutritional composition per serving was: 542.97 kcal, 49 g CHO, 27.76 g of proteins, 22.23 g of total lipids, 5.29 g of total fiber, and 4.31 mg of zinc. In relation to the recommendations established for dining rooms for the elderly, the meat-free menus covered 86.88% of kcal, 70.12% of CHO, 95.91% of proteins, 142.1% of lipids, 77.01% of total fiber, and 80.09% of zinc. Menus with meat contributed 86.17% of kcal, 53.72% of CHO, 160.7% of proteins, 113% of lipids, 63.69% of total fiber, and 151.3% of zinc. It was observed that both the composition and the percentage of protein and zinc adequacy was significantly higher in the meat menus ( $P < 0.05$ ). Zinc deficiency was observed on meat-free menus. The experimental composition of the analyzed meat-free menu was as follows: menu n° 1 showed a deficit of all its components except for lipids, menu n° 2 showed adequate fiber and zinc coverage and menu n° 3 showed adequate caloric and zinc intake. Finally, when analyzing the zinc content in “maca andina”, it was found that it provides 18.58 mg zinc/100 g; therefore, zinc deficient menus would cover the recommendations for this trace element with one tablespoon (15 g) of it. Due to its easy access and high nutritional value, “maca andina” could be beneficial to supplement diets deficient in this trace element.

## A252

### EFFECT OF A PPAR $\gamma$ SYNTHETIC AGONIST ASSOCIATED WITH RETINOIC ACID ON 24-HOUR RHYTHMS IN THE HIPPOCAMPUS OF AN EXPERIMENTAL MODEL OF ALZHEIMER'S DISEASE

Castro A, Mazaferro P, Golini R, Navigatore Fonzo L, Anzulovich AC

Laboratorio de Cronobiología, IMIBIO-SL, CONICET-UNSL, San Luis. E-mail: lnavigatore@unsl.edu.ar

Alzheimer's disease (AD) is the most frequent cause of dementia in the older adults. The main pathogenic mechanism in sporadic AD is the decrease in amyloid beta peptide (A $\beta$ ) clearance. It is known that Apolipoprotein E (Apo E) modulates A $\beta$  deposition and clearance. ApoE expression is transcriptionally induced by PPAR $\gamma$  in coordination with RXRs. Previously, we found that an intracerebroventricular injection of A $\beta$ (1-42) modified the daily rhythms of Apo E, Bmal 1, and A $\beta$  in the rat hippocampus. Taking into account those observations, the objective of this work was to investigate the effects of synthetic PPAR $\gamma$  agonist, pioglitazone, and retinoic acid (Pio-RA) on the 24-h rhythms of Apo E, BMAL1 and A $\beta$  protein levels, as well as on the daily rhythms of brain-derived neurotrophic factor (Bdnf) and its receptor (TrkB) expression in the rat hippocampus. In this study, male Holtzman rats from control, A $\beta$ -injected (A $\beta$ ) and A $\beta$ -injected treated with Pio-RA groups were euthanized throughout a 24-h period and hippocampus samples were isolated every 6 h. Apo E, BMAL1 and A $\beta$  proteins levels were analyzed by immunoblotting and Bdnf and TrkB mRNA levels were determined by RT-PCR. Regulatory regions of Apo E and clock genes were scanned for E-box, RORE, RXRE and PPRE sites. We observed that the treatment of Pio-RA reestablished the daily rhythms of Apo E, A $\beta$ , BMAL1 protein, and Bdnf mRNA levels. This treatment also increased Bdnf and TrkB levels. We found E-box, RXRE, and PPRE sites on regulatory regions of Apo E and Bmal1 genes. The results of the present study could suggest that the treatment of Pio-RA would not only restore the altered rhythmic patterns of the clock genes and their target genes observed in animals injected with A $\beta$  aggregates, but also, interestingly, would increase the levels of cognition-related genes, which are decreased in Alzheimer's patients.