


Limiting Values of the ^{15}N Chemical Shift of the Imidazole Ring of Histidine at High pH

Jorge A. Vila*

Baker Laboratory of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853-1301, United States
Universidad Nacional de San Luis, IMASL-CONICET, Ejército de Los Andes, 950-5700 San Luis, Argentina

ABSTRACT: Tautomeric identification by direct observation of ^{15}N chemical shifts of the imidazole ring of histidine (His) has become a common practice in NMR spectroscopy. However, such applications require knowledge of the “canonical” limiting values of the ^{15}N chemical shift of the imidazole ring of His in which each form of His, namely, the protonated (H^+) and the tautomeric $\text{N}^{\epsilon 2}\text{—H}$ and $\text{N}^{\delta 1}\text{—H}$ forms, respectively, is present to the extent of 100%. So far, the adopted canonical limiting values of the ^{15}N chemical shift have been those available from model compounds. Whether these canonical values reflect those of the individual pure forms of His is investigated here by carrying out an analysis of the second-order shielding differences, $\Delta\Delta = |\Delta^\epsilon - \Delta^\delta|$, with Δ^ξ ($\xi = \epsilon$ or δ) being the density functional theory (DFT)-computed average shielding differences between the two nitrogens of the imidazole ring of His in each pure tautomeric form. In the high-pH limit, the results indicate that (i) the $\Delta\Delta$ values from the DFT-computed shielding, but not from the commonly used canonical limiting values, are in closer agreement with those obtained with experimental chemical shift data from model compounds in solution and solid-state NMR; and (ii) the commonly used canonical limiting values of the ^{15}N chemical shifts lead to an *average* tautomeric equilibrium constant that differs by a factor of ~ 2.6 from the one computed by using DFT-based ^{15}N limiting values, raising concern about the practice of using canonical limiting ^{15}N values. This can be avoided by reporting tautomeric equilibrium constants computed by using *only* limiting ^{15}N values for the $\text{N}^{\epsilon 2}\text{—H}$ tautomer.



INTRODUCTION

Since chemical shifts were first observed by Arnold et al. in 1951,¹ using nuclear magnetic resonance (NMR) spectroscopy, Mandel,² in a pioneering NMR experiment, in 1965, first detected the imidazole (C2) protons of histidine (His) residues in Ribonuclease A, and, soon after that in 1966, Bradbury & Scheraga³ in a seminal NMR spectroscopic work were able to distinguish between the His residues of Ribonuclease A; namely, they resolved the peaks of three out of four His residues of this enzyme. Afterward, use of NMR spectroscopy, X-ray crystallography, and theoretical studies, based on quantum-chemistry calculations, continuously evolved in the ability to determine the properties of the His residues in solution and in the solid state.^{4–27} The reason for this persistent interest in His is due to the fact that this residue is unique among all 20 naturally occurring amino acids, and, among other reasons, because $\sim 50\%$ of all enzymes use His in their active sites.¹⁹ This is mainly because of the versatility of its imidazole His ring, which includes two neutral, chemically distinct forms ($\text{N}^{\delta 1}\text{—H}$ and $\text{N}^{\epsilon 2}\text{—H}$ tautomers) and a charged H^+ form, with one form favored over the other by the protein environment and pH. In addition, His with a $\text{p}K^{\circ}$ of 6.6^{28} is the *only* ionizable residue that titrates around neutral pH, allowing the nonprotonated nitrogen of its imidazole ring to serve as an effective ligand for metal binding,²³ or to play a crucial role in the proton-transfer process.²⁶

As noted by Markley,⁶ there are mainly three factors that contribute to the observed chemical shifts of the nuclei imidazole ring of His, namely the following: (a) The “intrinsic” chemical shifts, which can be determined theoretically, at the density functional theory (DFT) level of theory, from model peptides, e.g., by using a model tripeptide with the sequence Ac-GHG-NMe.²⁹ The conformational-average intrinsic values of the ^{15}N chemical shifts, for a given His form, are averages over *all* conformations, i.e., $\sim 35\,000$, for which the shieldings were computed here at the DFT level of theory, namely, for each nitrogen in the imidazole ring of His. (b) The “local environment” of each His residue in the protein, which can be very different for different His residues in the sequence, with the tautomeric coexistence being a function of the “local environment” of each His residue.²⁷ (c) The “proton binding/release equilibrium” of the imidazole His ring which, at a given fixed pH, is influenced by the presence of nearby positively or negatively charged side chains that will lower or raise the $\text{p}K'$ of a given His; overall, this factor determines the fraction of the protonated His form and, hence, those of the neutral form.²⁷ Here, focus is directed to determining the conformational-

Special Issue: Harold A. Scheraga Festschrift

Received: November 21, 2011

Revised: February 2, 2012

average intrinsic values of the ^{15}N chemical shifts for each of the two tautomeric forms of the imidazole ring of His as representative of the canonical values at high-pH; in other words, a theoretically determined set with which to represent the limiting values of the ^{15}N chemical shift. Hereafter, the computed conformational-average values of the intrinsic ^{15}N chemical shift will be referred to us as DFT-computed limiting ^{15}N values.

The fact that NMR spectroscopy is a powerful tool with which to investigate the role of His residues located in active sites of proteins is an observation that goes back to pioneering NMR experiments on His by Mandel,² who concluded "...This provides us with a window to observe the active site under various conditions...". Since this prediction was made, several methods are currently used to identify the protonation states of the imidazole ring of His as well as to distinguish between different tautomers.^{13,22} Most of these methods rely on the observation of ^{15}N chemical shifts of the imidazole ring of His, although an alternative to the use of ^{15}N chemical shifts has recently been presented,²⁷ namely, a new method based on the observed $^{13}\text{C}^{\gamma}$ and $^{13}\text{C}^{\delta 2}$ chemical shifts for the imidazole ring of His. Nevertheless, for a quantitative analysis of the tautomeric structure of the imidazole ring of His, use of ^{15}N chemical-shift data is very desirable for a number of reasons, among others because ^{15}N chemical shifts of the imidazole His ring are spread over a wide range and, even more important, are very sensitive to the presence or absence of a directly bonded proton.¹³ However, such applications require an accurate determination of the canonical limiting value of the chemical shift of each nitrogen in the imidazole ring of His, in each of the pure tautomers.¹⁰ Such determination based on model compounds is not an easy task since the difference between the chemical shifts of the nitrogens of the imidazole ring of His may vary significantly among different solvents.¹² In fact, for the neutral imidazole ring of $\text{N}^{\delta 1}$ -methylimidazole at 25 °C, values of 83, 96, and 104 ppm⁸ have been observed in water, chloroform, and benzene, respectively. In addition, it is not yet clear which solvent represents the His environment better in a protein. Together, such evidence raises the question about the accuracy of the canonical limiting values of the ^{15}N chemical shift obtained from model compounds, e.g., those listed by Pelton et al.¹³ as commonly used in the literature.²⁴

Consequently, it is our interest here to explore whether the gas-phase DFT-computed ^{15}N values, from $\sim 35\,000$ conformations of each tautomeric form of His²⁷ at high-pH, could provide a standard with which to compare the canonical limiting values of the ^{15}N chemical shift obtained from model compounds (from here on referred to us as canonical ^{15}N values). The existence of possible differences between DFT-computed and canonical ^{15}N values together with existing experimental evidence will enable us to determine the magnitude of changes of some properties of His residues, such as the tautomeric distribution at high pH, as a function of the adopted ^{15}N limiting values of the chemical shift.

METHOD

Computation of the ^{15}N Shielding of the Imidazole Ring of His at High pH. Each form of His is represented as a terminally blocked model tripeptide with the sequence Ac-GH $^{\xi}$ G-NMe,²⁷ with H $^{\xi}$ in the $\text{N}^{\delta 1}$ -H or the $\text{N}^{\epsilon 2}$ -H tautomer form, respectively. The analysis of the charged form, H $^+$, was omitted because this form does not exist at high pH. For each tautomer, a set of more than $\sim 35\,000$ conformations, each

representing a uniform sampling of the whole Ramachandran map as a function of the ϕ , ψ , ω , $\chi 1$, and $\chi 2$ torsional angles, was generated. For each of the $\sim 35\,000$ conformations, the gas-phase, isotropic shielding value of the His residue was calculated with the Gaussian 03 package³⁰ following the identical procedure that was already used for the computation of the $^{13}\text{C}^{\alpha}$ shielding database.²⁷ Finally, the distribution of the computed shielding for each ^{15}N of the imidazole ring of His was analyzed. For each ^{15}N nucleus, namely, $^{15}\text{N}^{\delta 1}$ and $^{15}\text{N}^{\epsilon 2}$, the histogram of the shielding distribution, among all $\sim 35\,000$ conformations, is shown in Figure 1a,b, together with the average values and standard deviations.




Figure 1. (a) The red-filled and blue-filled profiles for the shielding of the $^{15}\text{N}^{\epsilon 2}$ and $^{15}\text{N}^{\delta 1}$ nuclei, respectively, of the imidazole ring of His, in the $^{15}\text{N}^{\epsilon 2}$ -H tautomer represent the histogram of the $\sim 35\,000$ DFT-computed shielding distribution from the model tripeptide Ac-GHG-NMe; the corresponding average value and standard deviation of the red-filled and blue-filled profiles are 121 ± 3 ppm and -27 ± 10 ppm, respectively (b) Same as panel a, with an average value and standard deviation of -22 ± 5 ppm and 105 ± 9 ppm, respectively, for the $^{15}\text{N}^{\delta 1}$ -H tautomer.

Analysis of the Computed ^{15}N Shieldings for the Tautomeric Forms. For each tautomeric form, the average-shielding value, and its standard deviation, for each nitrogen of the imidazole ring of His were computed from the generated $\sim 35\,000$ conformations, and the results are shown in Figure 1a,b. These average values were used to compute the shielding differences between nitrogens of the imidazole ring of His (Δ^{ξ} values shown in Table 1), which is a common way to analyze

Table 1. Values of the Chemical Shift Difference, Δ^ξ

Δ^ξ ^a	canonical ^b (ppm)	DFT-computed ^c (ppm)	observed (ppm)
Δ^δ	82	83	79 ^d ; 77 ^e ; 76 ^f
Δ^ϵ	82	94	92 ^d ; 83 ^e ; 83 ^f

^a $\Delta^\xi = \|\zeta_{15N\delta^1} - \zeta_{15N\epsilon^2}\|$, with ζ denoting the average-shielding (σ) DFT-computed value, shown in the caption of Figure 1, or alternatively, the observed chemical shift (δ) for each of the imidazole ring nitrogens,¹³ and $\xi = \delta$ or ϵ denoting each of the tautomeric forms of the imidazole ring of His, namely, the N ^{δ^1} -H and N ^{ϵ^2} -H form, respectively. ^bUsing the ¹⁵N chemical shift limiting values listed by Pelton et al.¹³ ^cFrom average DFT-computed ¹⁵N shielding values (see Figure 1a,b). ^dFrom solid-state NMR, at pH 8.5, from Hu et al.²⁶ ^eFrom Farr-Jones et al.¹² for His at -55 °C at pH 11.2 in ethanol. ^fFrom Farr-Jones et al.¹² for N ^{ϵ} -methylhistidine at 25 °C in H₂O.

¹⁵N chemical shifts.¹¹ Chemical shift differences are equivalent to shielding differences because of cancelation of the reference value, thereby avoiding any source of error in the reference value.²¹ Consequently, the nitrogen shielding differences are $\Delta^\xi = \|\sigma_{15N\epsilon^2} - \sigma_{15N\delta^1}\|$, with σ denoting the average-shielding and $\xi = \epsilon$ or δ denoting each of the tautomeric forms of the imidazole ring of His, namely, the N ^{ϵ^2} -H and N ^{δ^1} -H forms, respectively.

RESULTS

Analysis of the Differences between Nitrogens in the Imidazole Ring of His: Δ^ξ . From Table 1, we conclude that the following inequality holds: $\Delta^\epsilon > \Delta^\delta$, if DFT-computed data, or data from model compounds or solid-state NMR, are used. On the other hand, the use of standard¹² canonical ¹⁵N values (see Table 1) gives the following equality: $\Delta^\epsilon = \Delta^\delta$.

We have also computed the second-order difference as $\Delta\Delta = |\Delta^\epsilon - \Delta^\delta|$. In general, use of Δ^ξ values, with $\xi = \epsilon$ or δ , calculated with DFT-computed shieldings, agrees qualitatively with data from model compounds and solid-state NMR (see Table 1), i.e., all of them predicting $\Delta\Delta = |\Delta^\epsilon - \Delta^\delta| > 0$. In particular, DFT-computed and solid-state NMR-data lead to close quantitative agreement, namely, $\Delta\Delta = 11$ and 13 ppm, respectively. Since observed isotropic ¹⁵N chemical shifts for His and imidazole in the solid state do not differ appreciably from those measured in solution,¹⁰ the latter result is significant. By contrast, a similar analysis carried out using canonical ¹⁵N values (see Table 1), gives $\Delta\Delta = 0$.

Overall, the experimental evidence and DFT-based calculations, but not canonical ¹⁵N values, predict $\Delta^\epsilon \neq \Delta^\delta$. The consequences of this finding will be discussed in the next sections.

Determining DFT-Computed Limiting Values of ¹⁵N Chemical Shifts in the High-pH Limit. A question of central importance, that goes beyond the existence of the differences in terms of Δ^ξ or $\Delta\Delta$ mentioned above, is related to the relevance of such discrepancies, for example, in the prediction of the tautomeric fractions at high-pH. In order to answer this question, we first need to determine a set of DFT-computed limiting values of ¹⁵N chemical shift differences.

The chemical shifts (δ) can be calculated by employing the equation $\delta = \sigma_{\text{ref}} - |\sigma_{\text{subst}}|$, where σ denotes the shielding of the reference substance (σ_{ref}) or the substance of interest (σ_{subst}), respectively. Use of this definition enables us to find an effective-reference value (σ_{ref}) with which *all* DFT-computed average-shielding values can be converted into chemical shifts

and, hence, a set of DFT-computed ¹⁵N limiting values of chemical shifts can be determined.

Therefore, in order to compute an effective reference, the following assumption was adopted: the DFT-computed chemical shift for the ¹⁵N ^{ϵ^2} nucleus ($\delta_{15N\epsilon^2}$), in the N ^{ϵ^2} -H tautomeric form of the imidazole ring of His, is assumed to be the canonical limiting value reported for this nucleus,¹³ namely, $\delta_{15N\epsilon^2} = 167.5$ ppm (see Table 2). We choose this chemical

Table 2. Limiting Values of ¹⁵N Chemical Shift at High pH^a

tautomer	canonical (ppm)		DFT-computed (ppm)	
	¹⁵ N ^{ϵ^2}	¹⁵ N ^{δ^1}	¹⁵ N ^{ϵ^2}	¹⁵ N ^{δ^1}
N ^{δ^1} -H	249.5	167.5	266.5	183.5
N ^{ϵ^2} -H	167.5	249.5	167.5	261.5

^aThe canonical limiting values are those listed by Pelton et al.¹³ See the text for details about the derivation of the DFT-computed limiting values.

shift as a starting point because the ¹⁵N ^{ϵ^2} nucleus shows the lowest standard deviation (3 ppm) among all four DFT-computed ¹⁵N average shielding values (see caption of Figure 1). Consequently, the effective reference is $\sigma_{\text{ref}} = 288.5$ ppm, i.e., computed as $\sigma_{\text{ref}} = (\delta_{15N\epsilon^2} + \sigma_{15N\epsilon^2})$, with $\sigma_{15N\epsilon^2} = 121$ ppm (see caption of Figure 1a) representing the DFT-computed average-shielding value for the ¹⁵N ^{ϵ^2} nucleus. Hence, the chemical shift of the ¹⁵N ^{δ^1} nucleus in the N ^{δ^1} -H tautomeric form is $\delta_{15N\delta^1} = 183.5$ ppm; this value was obtained by using the computed average-shielding value ($\sigma_{15N\delta^1} = 105$ ppm; see caption of Figure 1b) together with the above-determined effective reference. Up to this point, we have been able to determine the chemical shifts for the *protonated* ¹⁵N nucleus of the imidazole ring of His in the N ^{ϵ^2} -H and the N ^{δ^1} -H tautomeric form, respectively.

The chemical shift for the *nonprotonated* ¹⁵N nucleus in each tautomeric form can be computed straightforwardly by using the definition of Δ^ξ together with the constraints given in Table 1, i.e., by using the following relations: $\Delta^\delta = |\delta_{15N\delta^1} - 183.5| = 83$ ppm and $\Delta^\epsilon = |\delta_{15N\delta^1} - 167.5| = 94$ ppm. As a result, the following chemical-shift values were obtained: $\delta_{15N\epsilon^2} = 266.5$ ppm and $\delta_{15N\delta^1} = 261.5$ ppm, i.e., for the *nonprotonated* nucleus in the N ^{δ^1} -H and N ^{ϵ^2} -H tautomeric forms, respectively. Table 2 lists all DFT-computed and the canonical ¹⁵N limiting values for comparison.

Calculation of the Tautomeric Fraction, at High pH, from ¹⁵N Chemical Shifts. The significance of the use of the DFT-computed rather than the canonical ¹⁵N value is illustrated by calculating the tautomeric fractions for His61 in plastocyanin²⁴ at high pH as the average ratio between the tautomeric populations,²⁴ namely as $(P_\delta/P_\epsilon) = (\delta\epsilon - \delta_0)/(\delta_0 - \delta_\delta)$, where P_δ and P_ϵ are the populations of the N ^{ϵ^2} -H and N ^{δ^1} -H tautomeric forms, respectively; $\delta\epsilon$ and δ_δ are the corresponding limiting values of the ¹⁵N chemical shift, and δ_0 is the observed ¹⁵N chemical shift at high pH. It is worth noting that, at high pH, computation of the ratio (P_δ/P_ϵ) is equivalent to determining the tautomeric equilibrium constant K_T ,²² i.e., with $K_T = (\text{N}^{\delta^1}\text{-H tautomer})/(\text{N}^{\epsilon^2}\text{-H tautomer})$, which is 0.25 in aqueous solution.¹¹ Use of the canonical ¹⁵N value led Hass et al.²⁴ to obtain $K_{T,\text{canonical}} \sim 0.102$, as an *average* of the 0.096 and 0.108 values obtained for the (P_δ/P_ϵ) ratio computed from the fits of the ¹⁵N ^{ϵ^2} and ¹⁵N ^{δ^1} titration curves, respectively. Thus, the computed average $K_{T,\text{canonical}} \sim 0.102$ would correspond to a fraction of ~ 0.91 and ~ 0.09 for the N ^{ϵ^2} -H

and $N^{\delta 1}$ -H tautomers of His61, respectively. This result was obtained by using chemical shift data, at high pH, from the titration curve of the imidazole ring of His61,²⁴ namely, by using δ_0 observed values of ~ 241 and ~ 175 ppm for the $^{15}N^{\delta 1}$ and $^{15}N^{\epsilon 2}$ chemical shift values, respectively.

This set of δ_0 observed values (~ 241 and ~ 175 ppm) obtained at high pH can be used to recompute the average tautomeric equilibrium constant for His61 in plastocyanin by using DFT-computed rather than canonical limiting ^{15}N values (see Table 2). In this case, we obtain $K_{T,DFT} \sim 0.265$, as an average of the 0.086 and 0.443 values obtained for the $(P_{\delta}/P_{\epsilon})$ ratio computed by using ^{15}N limiting values for the $N^{\epsilon 2}$ -H and $N^{\delta 1}$ -H tautomeric forms, respectively. Hence, the computed average $K_{T,DFT} \sim 0.265$ corresponds to a fraction of ~ 0.79 and ~ 0.21 for the $N^{\epsilon 2}$ -H and $N^{\delta 1}$ -H tautomers, respectively.

Therefore, the commonly used canonical limiting values of the ^{15}N chemical shifts lead to an average tautomeric equilibrium constant (~ 0.102) that differs, by a factor of ~ 2.6 , from the average one computed by using DFT-computed ^{15}N limiting values (~ 0.265). The origin of this difference lies in the fact that DFT-computed ^{15}N limiting values, but not the canonical ones, lead to a very different ratio between the tautomeric populations $(P_{\delta}/P_{\epsilon})$ depending on the chosen tautomer. In other words, the DFT-computed, but not the canonical, limiting ^{15}N values are asymmetrical between tautomers (see Table 2). As a result, the difference in the equilibrium constants computed by using canonical and DFT-computed limiting ^{15}N values shows a minimum when only data from the $N^{\epsilon 2}$ -H tautomer are used to estimate the $(P_{\delta}/P_{\epsilon})$ ratio. At this minimum, as shown above, $K_{T,canonical}$ is ~ 0.096 and $K_{T,DFT}$ is ~ 0.086 ; hence, these fractions lead to a close estimation of the population of the $N^{\epsilon 2}$ -H and $N^{\delta 1}$ -H tautomers in solution, namely, ~ 0.92 and 0.08 , for the $N^{\epsilon 2}$ -H and $N^{\delta 1}$ -H tautomers, respectively.

Overall, considering that (i) the $^{15}N^{\epsilon 2}$ nucleus shows the lowest standard deviation among all DFT-computed ^{15}N average shielding values (see caption of Figure 1), (ii) the second-order difference values, $\Delta\Delta$, from the DFT-computed shielding, but not from the commonly used canonical limiting values, are in closer agreement with those obtained with the experimental chemical shift data from model compounds in solution and solid-state NMR; and (iii) the tautomeric predictions computed using canonical and DFT-computed ^{15}N values for the $^{15}N^{\epsilon 2}$, but not for the $^{15}N^{\delta 1}$, nucleus are in close agreement, it is recommended that the equilibrium constant, K_T , computed by using the ^{15}N limiting values for only the $N^{\epsilon 2}$ -H tautomer be reported. In other words, this option would help to reduce inaccuracies associated with the existing uncertainty in the set of limiting ^{15}N values.²² Whether this conclusion depends on the $N^{\delta 1}$ -H tautomeric fraction remains to be proved.

CONCLUSIONS

Overall, our results suggest that a considerable difference for the average tautomeric equilibrium constant, K_T , can be obtained if DFT-computed ^{15}N limiting values rather than canonical values are used. Furthermore, since the DFT-computed, but not the canonical, second-order shielding differences, $\Delta\Delta$, are in agreement with those obtained with experimental chemical shift data from model compounds in solution and solid-state NMR, our results also raise concerns about the magnitude of the uncertainty associated with the

usual predictions of the tautomeric fractions and, hence, their impact in studies of the mechanism of action of enzymes in which His dynamics may play a central role. To minimize the magnitude of such uncertainty, it is recommended here that the equilibrium constant, and then the tautomeric fractions, computed by using only limiting ^{15}N canonical or DFT-computed values for the $N^{\epsilon 2}$ -H tautomer be reported.

AUTHOR INFORMATION

Corresponding Author

*E-mail: jv84@cornell.edu or vila@unsl.edu.ar.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

I thank all spectroscopists and crystallographers who have deposited their coordinates at the PDB, and chemical shifts at the BMRB; without their effort, this work would not be possible. This research was supported by grants from the National Institutes of Health (GM-14312 and GM-24893), and the National Science Foundation (MCB05-41633). Support was also received from CONICET and from the UNSL (P-328402), Argentina. The research was conducted by using the resources of Pople and Blacklight, facilities of the National Science Foundation Terascale Computing System at the Pittsburgh Supercomputer Center.

REFERENCES

- (1) Arnold, J. T.; Dharmatti, S. S.; Packard, M. E. *J. Chem. Phys.* **1951**, *19*, 507.
- (2) Mandel, M. *J. Biol. Chem.* **1965**, *240*, 1586–1592.
- (3) Bradbury, J. H.; Scheraga, H. A. *J. Am. Chem. Soc.* **1966**, *88*, 4240–4246.
- (4) Meadows, D. H.; Jardetzky, O.; Epand, R. M.; Ruterjans, H. H.; Scheraga, H. A. *Proc. Natl. Acad. Sci. U.S.A.* **1968**, *60*, 766–772.
- (5) Reynolds, W. F.; Peat, I. R.; Freedman, M. H.; Lyerla, J. R. Jr. *J. Am. Chem. Soc.* **1973**, *95*, 328–331.
- (6) Markley, J. L. *Acc. Chem. Res.* **1974**, *8*, 70–80.
- (7) Wüthrich, K. *NMR in Biological Research: Peptides and Proteins*; North-Holland: Amsterdam, 1976.
- (8) Schuster, I. I.; Roberts, J. D. *J. Org. Chem.* **1979**, *44*, 3864–3867.
- (9) Harbison, G.; Herzfeld, J.; Griffin, R. G. *J. Am. Chem. Soc.* **1981**, *103*, 4752–4754.
- (10) Munowitz, M.; Bachovchin, W. W.; Herzfeld, J.; Dobson, C. M.; Griffin, R. G. *J. Am. Chem. Soc.* **1982**, *104*, 1192–1196.
- (11) Bachovchin, W. W. *Biochemistry* **1986**, *25*, 7751–7759.
- (12) Farr-Jones, S.; Wong, W. Y. L.; Gutheil, W. G.; Bachovchin, W. W. *J. Am. Chem. Soc.* **1993**, *115*, 6813–6819.
- (13) Pelton, J. G.; Torchia, D. A.; Meadow, N. D.; Roseman, S. *Protein Sci.* **1993**, *2*, 543–558.
- (14) Steiner, T. *Acta Crystallogr.* **1996**, *C52*, 2554–2556.
- (15) Steiner, T.; Koellner, G. *Chem. Commun.* **1997**, *13*, 1207–1208.
- (16) Shimba, N.; Takahashi, H.; Sakakura, M.; Fuji, I.; Shimada, I. *J. Am. Chem. Soc.* **1998**, *120*, 10988–10989.
- (17) Sudmeier, J. L.; Bradshaw, E. M.; Haddad, K. E. C.; Day, R. M.; Thalhauser, C. J.; Bullock, P. A.; Bachovchin, W. W. *J. Am. Chem. Soc.* **2003**, *125*, 8430–8431.
- (18) Shimba, N.; Serber, Z.; Ledwidge, R.; Miller, S. M.; Craik, C. S.; Dotsch, V. *Biochemistry* **2003**, *42*, 9227–9234.
- (19) Strohmeier, M.; Stueber, D.; Grant, D. M. *J. Phys. Chem. A* **2003**, *107*, 7629–7642.
- (20) Chen, X.; Zhan, C. G. *THEOCHEM* **2004**, *682*, 73–82.
- (21) Cheng, F.; Sun, H.; Zhang, Y.; Mukkamala, D.; Oldfield, E. J. *Am. Chem. Soc.* **2005**, *127*, 12544–12554.

- (22) Shimahara, H.; Yoshida, T.; Shibata, Y.; Shimizu, M.; Kyogoku, Y.; Sakiyama, F.; Kakazawa, T.; Tate, S-i.; Kato, T.; Moriyama, H.; et al. *J. Biol. Chem.* **2007**, *282*, 9646–9656.
- (23) Jensen, M. R.; Has, M. A. S.; Hansen, D. F.; Led, J. J. *Cell. Mol. Life Sci.* **2007**, *64*, 1085–1104.
- (24) Hass, M. A. S.; Hansen, D. F.; Christensen, H. E. M.; Led, J. J.; Kay, L. E. *J. Am. Chem. Soc.* **2008**, *130*, 8460–8470.
- (25) Hass, M. A. S.; Yilmaz, A.; Christensen, H. E. M.; Led, J. J. *J. Biomol. NMR* **2009**, *44*, 225–233.
- (26) Hu, F.; Wenbin, L.; Hong, M. *Science* **2010**, *330*, 505–508.
- (27) Vila, J. A.; Arnautova, Y. A.; Vorobjev, Y.; Scheraga, H. A. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 5602–5607.
- (28) Demchuk, E.; Wade, R. C. *J. Phys. Chem.* **1996**, *100*, 17373–17387.
- (29) Vila, J. A.; Scheraga, H. A. *Acc. Chem. Res.* **2009**, *42*, 1545–1553.
- (30) Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Zakrzewski, V. G., Montgomery, J. A., Stratmann, R. E., Burant, J. C., et al. *Gaussian 03*, revision E.01; Gaussian, Inc.: Wallingford, CT; 2004.