

Lithospheric-scale 3D model of the southern Central Andes (<https://doi.org/10.5880/GFZ.4.5.2020.001>)

Constanza Rodriguez Piceda^{1,2}, Magdalena Scheck-Wenderoth^{1,3}; Maria Laura Gomez Dacal¹, Judith Bott¹, Claudia Prezzi⁴, Manfred Strecker²

1. *GFZ German Research Centre for Geosciences, Potsdam, Germany*
2. *University of Potsdam. Potsdam, Germany*
3. *RWTH Aachen. Aachen, Germany*
4. *IGEBA-(UBA-CONICET); University of Buenos Aires, Argentina*

1. Licence

Creative Commons Attribution 4.0 International License (CC BY 4.0)



2. Citation

When using the data please cite:

Rodriguez Piceda, C.; Scheck-Wenderoth, M.; Gomez Dacal, M. L.; Bott, J.; Prezzi, C.; Strecker, M. (2020): Lithospheric-scale 3D model of the Southern Central Andes. GFZ Data Services. <https://doi.org/10.5880/GFZ.4.5.2020.001>

The data are supplementary material to:

Rodriguez Piceda, C., Scheck Wenderoth, M., Gomez Dacal, M. L., Bott, J., Prezzi, C. B., & Strecker, M. R. (2020). Lithospheric density structure of the southern Central Andes constrained by 3D data-integrative gravity modelling. In *International Journal of Earth Sciences* (Vol. 110, Issue 7, pp. 2333–2359). Springer Science and Business Media LLC. <https://doi.org/10.1007/s00531-020-01962-1>

3. Model area

Model coordinates are given in UTM map projection (UTM Zone 19S)

Datum: World Geodetic System 1984.

Model bounds in UTM Zone 19S:

Easting: from 200000 m to 900000 m

Northing: from 5700000 m to 6800000 m

Model bounds in longitude/latitude (WGS 84):

Longitude: from -72.4543 to -64.3973

Latitude: from -38.8488 to -28.86550

4. Abstract

The Central Andean orogeny is caused by the subduction of the Nazca oceanic plate beneath the South-American continental plate. In Particular, the southern Central Andes (SCA, 27°-40°S) are characterized by a strong N-S and E-W variation in the crustal deformation style and intensity. Despite being the surface geology relatively well known, the information on the deep structure of the upper plate in terms of its thickness and density configurations is still scarcely constrained. Previous seismic studies have focused on the crustal structure of the northern part of the SCA (~27°-33°S) based upon 2D cross-sections, while 3D crustal models centered on the South-American or the Nazca Plate have been published with lower resolution.

To gain insight into the present-day state of the lithosphere in the area, we derived a 3D model that is consistent with both the available geological and seismic data and with the observed gravity field (Rodríguez-Piqueda et al., 2020). The model consists on a continental plate with sediments, a two-layer crust and the lithospheric mantle being subducted by an oceanic plate. The model extension covers an area of 700 km x 1100 km, including the orogen (i.e. magmatic arc, main orogenic wedge), the forearc and the foreland.

5. Relevance

The presented 3D structural model resolves the thickness and density distribution of the main layers that make up the SCA lithosphere. Through the integration of a large amount of diverse data, we are able to link the crustal configuration with different tectonic domains which were previously defined based on the (near)-surface geology. The present-day crustal configuration most likely reflects differences in the tectonic evolution of each tectonic domain.

This model can be used as a base to investigate the regional 3D temperature distribution and the rheological field in the SCA. Therefore, it contributes to discussions about the role of the upper plate on the short- (i.e. earthquake-related) and long-term (i.e. mountain-building related) deformation pattern, as well as on the subduction geometry (i.e. slab dip).

6. Methods

Different data sets were integrated to derive the lithospheric features:

- We used the global relief model of ETOPO1 (Amante and Eakins 2009) for the topography and bathymetry.
- The sub-surface structures were defined by integrating seismically constrained models, including the South-American crustal thickness of Assumpção et al. (2013; model A; 0.5-degree resolution), the sediment thickness of CRUST1 (Laske et al. 2013) and the slab geometry of SLAB2 (Hayes et al. 2018).
- Additionally, we included seismic reflection and refraction profiles performed on the Chile margin (Araneda et al. 2003; Contreras-Reyes et al. 2008, 2014, 2015; Flueh et al. 1998; Krawzyk et al. 2006; Moscoso et al. 2011; Sick et al. 2006; Von Huene et al. 1997).
- Besides, we used sediment thickness maps from the intracontinental basin database ICONS (6 arc minute resolution, Heine 2007) and two oceanic sediment compilations: one along the southern trench axis (Völker et al. 2013) and another of global-scale (GlobSed; Straume et al. 2019).

To build the interfaces between the main lithospheric features, we compiled and interpolated these datasets on a regular grid with a surface resolution of 25 km. For that purpose, the convergent algorithm of the software Petrel was used. We assigned constant densities within each layer, except for the lithospheric mantle. In this case, we implemented a heterogeneous distribution by converting s-wave velocities from the SL2013sv seismic tomography (Schaeffer and Lebedev 2013) to densities. The python tool VelocityConversion was used for the conversion (Meeßen 2017).

To further constrain the crustal structure of the upper plate, a gravity forward modelling was carried out using IGMAS+ (Schmidt et al. 2010). The gravity anomaly from the model (calculated gravity) was compared to the free-air anomaly from the global gravity model EIGEN-6C4 (observed gravity; Förste et al 2014; Ince et al. 2019). Subsequently, the crystalline crust of the upper plate was split vertically into two layers of different densities. We inverted the residual between calculated and observed gravity to compute the depth to the interface between the two crustal layers. For the inverse modelling of the gravity residual, the Python package Fatiando a Terra was used (Uieda et al. 2013)

7. Technical Information

For each layer, the depth to the top surface, thickness and density can be found as separate files. All files contain identical columns:

- Northing as "X Coord (UTM zone 19S)";
- Easting as "Y Coord (UTM zone 19S)";
- depth to the top surface as "Top (m.a.s.l)" and
- thickness of each layer as "Thickness (m)".

The header 'Density' indicates the bulk density of each unit in kg/m³. For the oceanic and continental mantle units, a separate file is provided with a regular grid of the density distribution with a lateral resolution of 8 km x 9 km and a vertical resolution of 5 km. The containing columns are: Northing as "X Coord (UTM zone 19S)"; Easting as "Y Coord (UTM zone 19S)"; depth as "Depth (m.a.s.l)" and density as "Density (kg/m³)"

8. Acknowledgements

This research was funded by the Deutsche Forschungsgemeinschaft (DFG) and the Federal State of Brandenburg under the auspices of the International Research Training Group IGK2018 "SuRfAce processes, TEctonics and Georesources: The Andean foreland basin of Argentina" (STRATEGy DFG 373/34-1). We are grateful to Christian Meeßen for providing scripts for the conversion of velocities to densities, and for the gravity inversion. We also thank Hans-Jürgen Götze and Sabine Schmidt for permission to use the 3D gravity modelling software (IGMAS+) as well as for providing us help with this software.

9. Related Work

Data presented here were derived from an extensive list of related works (Amante and Eakins, 2009; von Huene et al. 1997; Flueh et al. 1998; Araneda et al. 2003; Krawczyk et al. 2006; Sick et al. 2006; Heine 2007; Contreras-Reyes et al. 2008, 2014, 2015; Moscoso et al. 2011; Laske et al. 2013; Schaeffer and Lebedev 2013; Völker et al. 2013; Assumpção et al. 2013; Förste et al. 2014; Hayes et al. 2018; Ince et al. 2019; Straume et al. 2019)

10. References

- Amante, C. and Eakins, B. W. (2009). ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis [Data set]. National Geophysical Data Center, NOAA. <https://doi.org/10.7289/V5C8276M>
- Araneda, M., Asch, G., Bataille, K., Bohm, M., Bruhn, C., Giese, P., ... Wigger, P. (2003). A crustal model along 39°S from a seismic refraction profile- ISSA 2000. *Revista Geológica de Chile*, 30(1). <https://doi.org/10.4067/s0716-02082003000100006>
- Assumpção, M., Feng, M., Tassara, A., & Julià, J. (2013). Models of crustal thickness for South America from seismic refraction, receiver functions and surface wave tomography. *Tectonophysics*, 609, 82–96. <https://doi.org/10.1016/j.tecto.2012.11.014>
- Contreras-Reyes, E., Becerra, J., Kopp, H., Reichert, C., & Díaz-Naveas, J. (2014). Seismic structure of the north-central Chilean convergent margin: Subduction erosion of a paleomagmatic arc. *Geophysical Research Letters*, 41(5), 1523–1529. <https://doi.org/10.1002/2013gl058729>
- Contreras-Reyes, E., Grevenmeyer, I., Flueh, E. R., & Reichert, C. (2008). Upper lithospheric structure of the subduction zone offshore of southern Arauco peninsula, Chile, at ~38°S. *Journal of Geophysical Research*, 113(B7). <https://doi.org/10.1029/2007jb005569>
- Contreras-Reyes, E., Ruiz, J. A., Becerra, J., Kopp, H., Reichert, C., Maksymowicz, A., & Arriagada, C. (2015). Structure and tectonics of the central Chilean margin (31°–33°S): implications for subduction erosion and shallow crustal seismicity. *Geophysical Journal International*, 203(2), 776–791. <https://doi.org/10.1093/gji/ggv309>
- Flueh, E. R., Vidal, N., Ranero, C. R., Hojka, A., von Huene, R., Bialas, J., ... Zelt, C. (1998). Seismic investigation of the continental margin off- and onshore Valparaiso, Chile. *Tectonophysics*, 288(1-4), 251–263. [https://doi.org/10.1016/s0040-1951\(97\)00299-0](https://doi.org/10.1016/s0040-1951(97)00299-0)
- Förste, Christoph; Bruinsma, Sean.L.; Abrikosov, Oleg; Lemoine, Jean-Michel; Marty, Jean Charles; Flechtner, Frank; Balmino, G.; Barthelmes, F.; Biancale, R. (2014): EIGEN-6C4 the latest combined global gravity field model including GOCE data up to degree and order 2190 of GFZ Potsdam and GRGS Toulouse. GFZ Data Services. <https://doi.org/10.5880/icgem.2015.1>
- Hayes, G. P., Moore, G. L., Portner, D. E., Hearne, M., Flamme, H., Furtney, M., & Smoczyk, G. M. (2018). Slab2, a comprehensive subduction zone geometry model. *Science*, 362(6410), 58–61. <https://doi.org/10.1126/science.aat4723>
- Heine, C. (2007) Formation and Evolution of intracontinental basins. Dissertation. School of Geosciences. University of Sydney. <https://mds.sydney.edu.au/redbox/published/detail/947d1178df46a48887ca09e255513814>
- Ince, E. S., Barthelmes, F., Reißland, S., Elger, K., Förste, C., Flechtner, F., & Schuh, H. (2019). ICGEM – 15 years of successful collection and distribution of global gravitational models, associated services, and future plans. *Earth System Science Data*, 11(2), 647–674. <https://doi.org/10.5194/essd-11-647-2019>

- Krawczyk, C. M., Mechie, J., Lüth, S., Tašárová, Z., Wigger, P., Stiller, M., ... Bataille, K. (2006). Geophysical Signatures and Active Tectonics at the South-Central Chilean Margin. *Frontiers in Earth Sciences*, 171–192. https://doi.org/10.1007/978-3-540-48684-8_8
- Laske, G.; Masters, G.; Ma, Z.; Pasyanos, M.E. (l. 2013) CRUST1.0: An Updated Global Model of Earth's Crust. *Geophysical Research Abstracts* Vol. 14, EGU2012-3743-1, EGU General Assembly 2012, <https://meetingorganizer.copernicus.org/EGU2012/EGU2012-3743-1.pdf>
- Meeßen, C. (2017). *VelocityConversion* (Version v1.0.1). GFZ Data Services. <https://doi.org/10.5880/gfz.6.1.2017.001>
- Moscoso, E., Grevemeyer, I., Contreras-Reyes, E., Flueh, E. R., Dzierma, Y., Rabbel, W., & Thorwart, M. (2011). Revealing the deep structure and rupture plane of the 2010 Maule, Chile earthquake (Mw=8.8) using wide angle seismic data. *Earth and Planetary Science Letters*, 307(1-2), 147–155. <https://doi.org/10.1016/j.epsl.2011.04.025>
- Rodriguez Piceda, C., Scheck Wenderoth, M., Gomez Dacal, M. L., Bott, J., Prezzi, C. B., & Strecker, M. R. (2020). Lithospheric density structure of the southern Central Andes constrained by 3D data-integrative gravity modelling. In *International Journal of Earth Sciences* (Vol. 110, Issue 7, pp. 2333–2359). Springer Science and Business Media LLC. <https://doi.org/10.1007/s00531-020-01962-1>
- Schaeffer, A. J., & Lebedev, S. (2013). Global shear speed structure of the upper mantle and transition zone. *Geophysical Journal International*, 194(1), 417–449. <https://doi.org/10.1093/gji/ggt095>
- Sick, C., Yoon, M.-K., Rauch, K., Buske, S., Lüth, S., Araneda, M., ... Wigger, P. (n.d.). Seismic Images of Accretive and Erosive Subduction Zones from the Chilean Margin. *Frontiers in Earth Sciences*, 147–169. https://doi.org/10.1007/978-3-540-48684-8_7
- Straume, E. O., Gaina, C., Medvedev, S., Hochmuth, K., Gohl, K., Whittaker, J. M., ... Hopper, J. R. (2019). GlobSed: Updated Total Sediment Thickness in the World's Oceans. *Geochemistry, Geophysics, Geosystems*, 20(4), 1756–1772. <https://doi.org/10.1029/2018gc008115>
- Uieda, L., Oliveira, V., & Barbosa, V. (2013). Modeling the Earth with Fatiando a Terra. *Proceedings of the 12th Python in Science Conference*. <https://doi.org/10.25080/majora-8b375195-010>
- Völker, D., Geersen, J., Contreras-Reyes, E., & Reichert, C. (2013). Sedimentary fill of the Chile Trench (32–46°S): volumetric distribution and causal factors. *Journal of the Geological Society*, 170(5), 723–736. <https://doi.org/10.1144/jgs2012-119>
- von Huene, R., Corvalán, J., Flueh, E. R., Hinz, K., Korstgard, J., Ranero, C. R., & Weinrebe, W. (1997). Tectonic control of the subducting Juan Fernández Ridge on the Andean margin near Valparaiso, Chile. *Tectonics*, 16(3), 474–488. <https://doi.org/10.1029/96tc03703>