

# Settlement and recruitment of the crab *Halicarcinus planatus* (Crustacea: Decapoda: Hymenosomatidae) in Golfo San Jorge, Argentina

JULIO H. VINUESA<sup>1,2</sup>, MARTÍN VARISCO<sup>1,2,3</sup> AND FLORENCIA ESCRICHE<sup>1</sup>

<sup>1</sup>Instituto de Desarrollo Costero, Universidad Nacional de la Patagonia San Juan Bosco (UNPSJB), Campus Universitario, Ruta No. 1, Km 4 (9000) Comodoro Rivadavia, Chubut, Argentina, <sup>2</sup>Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), <sup>3</sup>Facultad de Ciencias Naturales UNPSJB

*The settlement and recruitment of the small brachyuran crab Halicarcinus planatus from the central area of Golfo San Jorge, south-western Atlantic Ocean, was studied. Three different artificial collectors were evaluated: smooth-surface panels, panels covered with synthetic lawn and ballasted plastic boxes filled with remains of fishing nets. In addition, plankton samples were taken monthly and water salinity and temperature were recorded. Between July 2006 and June 2007, the collectors were deployed monthly in the subtidal zone of two nearby localities and collected two months later. The recovered crabs were classified into four stages: settlers, recruits, advanced juveniles and mature crabs. There were differences in the presence of stages among collector types and seasons but not between localities. Settlers and recruits were abundant on the panels with synthetic lawn, while late juvenile and mature females were more frequent in the boxes. Larval stages occur between July and February, and only zoeas I and II were found. The males, which were only represented by early juveniles, were exclusively found on the panels with synthetic lawn, from where they would subsequently migrate to cryptic habitats such as the holdfast of *Macrocystis pyrifera*. Settlement occurs between September and April. Settlers showed a peak in abundance during November–December, while the recruits showed two peaks, one in December and the other in February–March. For late juveniles, the maximum abundance was recorded in April–May. Ovigerous females were found between May and September, when the experiment ended. Settlement and recruitment of *H. planatus* in Golfo San Jorge occur over a prolonged period. Individuals may undergo shifts in microhabitat use during this period, probably related to the search for food and shelter, and avoidance of predation and cannibalism.*

**Keywords:** larvae, settlement, juveniles, sub-Antarctic crab, Patagonia

Submitted 22 September 2009; accepted 16 September 2010; first published online 15 December 2010

## INTRODUCTION

*Halicarcinus planatus* (Fabricius, 1775) is a small brachyuran crab of the family Hymenosomatidae, which is only present in the extreme south of America. Its geographical distribution is closely associated with the Antarctic Circumpolar Current; it is found on Chilean coasts from Bahía Taltal (35°S) to the Cabo de Hornos, New Zealand and sub-Antarctic Islands in the Pacific Ocean; along Fuegian coasts in Patagonia until 38°S in the Atlantic Ocean; and in the Prince Edward, Crozet and Kerguelen Islands in the Indian Ocean (Boschi *et al.*, 1969; Melrose, 1975).

*Halicarcinus planatus* is a coastal species common in rocky, lower intertidal and sublittoral bottoms, and it occurs down to depths of about 10–15 m in Golfo San Jorge (Vinuesa, 2005). Populations are more abundant in sheltered areas such as bays and inlets and can also be found on exposed coasts, but in places protected from the high wave energies and where

water is retained. In Golfo San Jorge and other areas, crabs often shelter under rocks in the lower intertidal and in sublittoral beds of Mytilidae or in the holdfasts of *Macrocystis pyrifera* (L.) Agardh, 1820 (Zaixso & Pastor, 1977; López Gappa *et al.*, 1982).

Information on different aspects of the biology of *H. planatus* has been provided by populations from the Kerguelen Islands (Richer de Forges, 1977) and Puerto Deseado (Vinuesa & Ferrari, 2008a, b). However, data on settlement and recruitment of this species are scarce. Richer de Forges (1977) reported the periods of recruitment and aggregation of juveniles. Vinuesa & Ferrari (2008a), who described temporal changes in the structure of the *H. planatus* population in Puerto Deseado, suggested that recruitment occurs in the subtidal environment.

Benthic species with planktonic larvae experience high mortality during settlement and recruitment, thus these stages appear as critical in their life-cycle. In this context, the regulatory mechanisms play a major role in conditioning population dynamics (Beck, 1995; Caley *et al.*, 1996). Settlement and recruitment are complex processes influenced by factors intrinsic to the population and by numerous environmental

### Corresponding author:

J.H. Vinuesa

Email: vinuesa@unpata.edu.ar

variables (Rodríguez *et al.*, 1993). In particular, the type of substrate and accessibility to refuge are of great importance for coastal crabs.

The objective of the present work was to study some aspects of the settlement and recruitment of *H. planatus*. The temporal patterns of these processes, the type of microhabitat and post-settlement movement of juveniles were investigated. In addition, the seasonal variations in the presence of the planktonic stages were evaluated.

## MATERIALS AND METHODS

### Study area

The study was performed in the central area of Golfo San Jorge, Argentina, in two nearby rocky shores, one located in Caleta del Fondo, a province of Santa Cruz (46°03'03"S and 67°36'13"W) and the other in front of Comodoro Rivadavia City (45°51'13"S and 67°28'35"W). These localities show similar characteristics in terms of orientation and exposure to waves, and depth.

The area is influenced by cold-temperate waters of the Patagonian Current which flow along the Patagonian coasts from the Estrecho de Magellanes northwards (Brandhorst & Castello, 1971). A branch of this current entering in north-north-west direction predominates in the studied area.

### Sampling

Four plankton samples, two from each of the above-mentioned localities, were taken monthly between September 2006 and August 2007. Samples were collected with a conical net of 70-cm mouth diameter and 185- $\mu$ m mesh opening. Oblique tows were conducted from near the bottom (7 m) to the surface. The volume of water filtered was estimated considering the distance of the haul and the mouth diameter of the plankton net. The samples were preserved in 4% formalin-seawater until further analysis in the laboratory. The temperature and salinity of surface water were recorded during each sampling period.

Three types of collectors were offered as settlement substrates:

- (1) fibre concrete panels of 10 × 20 cm and 1 cm of thickness, provided with synthetic lawn of 10 × 10 cm and 1 cm of height on the surface. Panels had two holes in the ends to allow placement;
- (2) fibre concrete panels of 10 × 20 cm and 1 cm of thickness, with a smooth surface of 10 × 10 cm without synthetic lawn. Panels had two holes in the ends to allow placement; and
- (3) 5 l plastic boxes (20 × 25 × 10 cm) with an upper aperture of about 10 × 10 cm closed with a plastic net of 10 mm in diameter, pierced with many 10-mm holes, filled with approximately 1 kg of fishing nets, and ballasted with concrete.

The collectors were placed in the sea for 20 days to eliminate the initial effects of chemical compounds of the materials that became collectors. Later they were mounted on 1 × 2 m iron frames attached to the rocky bottom. Six smooth-surface panels, six panels with synthetic lawn and four collecting

boxes were installed every month between July 2006 and June 2007. All were removed two months after immersion. The installation and recovery of the collectors were performed by a diver, who placed the collectors in sealed bags *in situ*. The panels were frozen at -14°C until analysis, and the boxes inspected within the following 20 hours.

### Laboratory analysis

*Halicarcinus planatus* larvae were sorted from the rest of the zooplankton and the zoeal stages were identified according to Boschi *et al.* (1969). The larvae were determined and counted under stereo zoom microscope. When needed, to avoid misidentifications, the observations were also made at higher magnifications with a compound microscope.

The collectors were washed and sieved with a mesh size of 0.5 mm to remove sediment. Taxonomic identification of the accompanying fauna and algae was made to the lowest possible taxonomic level. The crabs were sorted from the rest of the organisms, sexed and their carapace width (CW) was measured to the nearest 0.01 mm. Males are recognized by the narrowing of the last three abdominal segments. Females were first classified based on the morphology of the abdomen into the following five categories: immature (IMM2, IMM3 and IMM4); adolescent (ADO); and adult or mature (MAT), according to a previous work (Vinuesa & Ferrari, 2008a). Then, they were grouped into four stages as follows: (a) settlers were IMM1 individuals, who included both small males and females since sex cannot be determined at this stage; (b) recruits were small males and IMM2 and IMM3 females; (c) late juveniles included IMM4 and ADO females; and (d) adults were ovigerous and post-ovigerous females.

### Statistical analyses

The comparison of the total number of *H. planatus* individuals among collector types was performed by factorial analysis of variance. Data were checked for homocedasticity and normality using the Cochran C test and the Lilliefors test, respectively (Sokal & Rohlf, 1995). Significant differences in the distribution of postlarval stages were tested with permutational analysis of variance (PERMANOVA), using the Bray-Curtis dissimilarity index and presence-absence data transformation (Anderson, 2001, 2005). PERMANOVA is a univariate or multivariate analysis of variance using permutations procedures to obtain *P* values. It is suitable for any multifactorial ANOVA design allowing for all pairwise multiple comparisons by permutation. Similarity percentage analysis (SIMPER) was performed to quantify the contribution of each category to the different collector types. Bray-Curtis dissimilarity index and presence-absence data transformation were used for this test (Clarke & Warwick, 2001).

## RESULTS

The temperatures and salinities were similar in both sites. The minimum temperature, recorded in August, was 7.6°C and the maximum temperature, recorded in January, was 17.3°C. Salinity ranged between 33.1 and 34.2‰.

The *H. planatus* zoeae were collected between September and February 2006 and in July and August 2007, and were not found between March and June (Figure 1). No larvae were collected in November, probably due to the rough sea conditions during the sampling. The temporal distribution of larvae was similar between localities. Zoeae I were present throughout the months mentioned above, while zoeae II were only found in December and February.

A total of 412 crabs and several exuviae were removed from all the collectors recovered ( $N = 192$ ), with an overall sex-ratio of 1:25 (male:female). For recruits, the male:female ratio was 1:7. Adult males were not registered. The settlers (IMM<sub>1</sub>) represented 24.8% of total crabs ( $N = 105$ ), with sizes ranging between 0.8 and 2.1 mm CW. The recruits included males between 1.8 and 3.0 mm CW and females between 1.8 and 3.5 mm CW ( $N = 139$ ). Late juveniles ( $N = 103$ ) and adults ( $N = 60$ ) were only represented by females (Table 1). Exuviae of IMM I were recorded also, in synthetic lawn panels.

In the smooth-surface panels, the number of individuals ( $N = 5$ ) was much lower than in the rest of the collectors, and then were excluded from further analysis. There were significant differences in the distribution of the developmental stages among collector types and seasons but not between localities. No interactions were found among the analysed factors (PERMANOVA; Table 2).

Settlers had a higher contribution to dissimilarities among collectors compared to the other developmental stages (SIMPER analysis; Table 3). This stage was almost exclusively found on the panels covered with synthetic lawn, and recruits were also more abundant on this collector type. Late juveniles and adults predominated in the boxes (Figure 2). Ten exuviae of settlers were recovered from the collector panels. The accompanying fauna and flora also differed between panels and boxes. The panels were colonized by diverse organisms such as more than ten algal species, colonial benthic diatoms, foraminifers, nematodes, polychaetes, harpacticoid copepods, amphipods, isopods, cumaceans and other juvenile decapods. The boxes contained mainly polychaete worms, small fissurellids and octopuses, juvenile sea urchins, ophiuroids and starfish, isopods, caridean shrimps and fish.

The settlers were found throughout spring and summer, showing an abundance peak in December; the recruits were

more abundant during December 2006 and February 2007; and the abundance peak of late juveniles was observed in April 2006. Mature females occurred from May throughout the winter months (Figure 3). The decrease in the number of crabs observed in January may have been due to severe storms some days before the collectors were recovered.

## DISCUSSION

Settlement and recruitment are important events to understand the dynamics and structure of a population (Beck, 1995; Caley *et al.*, 1996). These are complex processes determined by the interaction between physical (e.g. light, water flow and substrate surface), chemical (e.g. secondary metabolites) and biological factors (e.g. larval behaviour, predation or juvenile dispersion) (Rodríguez *et al.*, 1993).

The larval supply, its distribution and behaviour are important pre-settlement factors (Rodríguez *et al.*, 1993; Moksnes & Wennhage, 2001). Larvae of *H. planatus* occur during eight months of the year. No larvae were captured in November. This is due to weather conditions during sampling. The absence of zoeae in the latter month is unexpected since it is when the local adult population undergoes larval hatching (Vinuesa, unpublished results). Zoeae I were the most abundant stage throughout the study period, indicating a continuous supply of larvae to the water column. The rare presence of zoeae II and the absence of zoea III may result from underestimation by an increase in positive geotaxis during larval ontogeny. Similar results were obtained with late zoeae in *H. planatus* from the Kerguelen Islands (Richer de Forges, 1977). The absence of later stages due to larval dispersal seems unlikely; according to Lucas (1980) a limited dispersal is a common strategy among the Hymenosomatidae.

Post-settlement factors have an important role in survival and distribution of juvenile crabs. These factors include: shelter, food supply, predation and cannibalism. The habitat availability is important in settlement and recruitment. Many works on the role of shelter and substrate in these processes conclude that species select complex habitats to minimize mortality (e.g. Botero & Atema 1982; Marx & Herrnkind, 1985; Herrnkind & Butler, 1986; Beck, 1995; Moksnes, 2002). Mytilid and oyster beds, seagrass and shell-hash beds, kelp forests, and other structurally complex microhabitat provide refuge for early crab stages from predation (Palacios *et al.*, 1985; Fernández *et al.*, 1993; Forward *et al.*, 1996; Stevens & Kittaka, 1998).

Richer de Forges (1977) reported the aggregation of *H. planatus* juveniles < 4 mm CW in holdfasts of *M. pyrifera*, highlighting their nursery role. In Puerto Deseado ( $\sim 48^\circ$  SL), the smallest juveniles (between 1.1 and 1.3 mm CL) were also exclusively found in the holdfasts of this kelp (Vinuesa & Ferrari, 2008a). In this work, settlers and recruits were more




Fig. 1. Abundance of larvae of *Halicarcinus planatus* between September 2006 and August 2007. Values from both localities are pooled.

Table 1. Mean and standard deviation (SD) of the carapace width (CW) at each developmental stage of *Halicarcinus planatus*.

	CW mean (mm)	SD
Settlers	1.37	0.28
Recruits	3.38	1.2
Late juveniles	6.55	1.03
Matures	8.33	1.33

**Table 2.** Permutational analysis of variance based on the Bray–Curtis dissimilarity index. df, degrees of freedom; SS, sum of squares; MS, mean square; F, F ratio; *P* (f), probability values associated with F values; *P* (MC), probability values obtained using Monte Carlo permutations.


	df	SS	MS	F	<i>P</i> (f)	<i>P</i> (MC)
Locality (L)	1	829.86	829.86	0.5513	0.619	0.628
Season (S)	3	28590.04	9530.01	6.3314	0.001	0.001
Collector (C)	1	12185.87	12185.87	8.0958	0.002	0.002
L × S	3	4159.02	1386.34	0.9210	0.511	0.511
L × C	1	4296.52	4296.52	2.8544	0.054	0.054
S × C	3	2547.45	849.15	0.5641	0.757	0.757
L × C × S	3	5923.84	1974.61	1.3119	0.301	0.301
Residual	32	48166.66	1505.20			

abundant on collectors with synthetic lawn. The smooth-surface collectors with unprotected, exposed areas were unfavourable for crab settlement. The early postlarval stages of *H. planatus* showed preference for complex structural collectors.

According to Rodríguez *et al.* (1993) the selection of microhabitats by benthic marine organisms at settlement may be related to food availability. In the spiny lobster *Panulirus argus* (Latreille, 1804), the abundance of food resources is an important factor for habitat selection in juveniles (Herrnkind & Butler, 1986). *Halicarcinus planatus* is a generalist crab; polychaete annelids, copepods, algae and diatoms were found in its diet (Arnaud, 1974; Richer de Forges, 1977). The high number of algae and animals found on the panels suggests that settlers and recruits used them not only as shelters but also as food source.

Crustaceans have discontinuous growth and therefore the shelter is particularly important during moulting and post-moult stages. Moreover, decapods are able to delay moulting in the absence of an appropriate shelter (Hopkins, 1992). The occurrence of two successive developmental stages and the presence of exuviae on the panels provide evidence that the settlers remain for some time in the settlement microhabitat, where they undergo moulting.

Predation is responsible for high mortality during the post-settlement of numerous decapod species (Herrnkind & Butler, 1986; Eggleston & Armstrong, 1995; Moksnes, 2002). Moreover, predation contributes to the dispersal of early juvenile stages of some species, e.g. *P. argus* (Herrnkind & Butler, 1986), and the blue king crab *Paralithodes platypus* Brandt, 1850 (Stevens & Swiney, 2005). The low number of settlers and recruits in the boxes could be related to the presence of predators. In fact, some predators like the starfish *Anasterias minuta* Perrier, 1875 (Gil & Zaixso, 2007), notothenid fish (Hureau, 1970), and potential predators like the shrimp *Betaeus truncatus* Dana, 1852 and the octopus



**Fig. 2.** Abundance (as percentage) of each stage of *Halicarcinus planatus* by type of collector. Grey and black bars represent the collecting panels and boxes, respectively. Individuals from both localities are grouped.

*Enteractopus megalocyathus* (Gould, 1852) were found in the boxes.

In some species, the presence of adult conspecific specimens or parental odour is likely to act as a stimulus for settlement (Jensen 1991; Rodríguez *et al.*, 1993; Gebauer *et al.*, 2003), but would induce intercohort cannibalism. Intraspecific predation is important in regulating recruitment and has been thoroughly studied in the blue crab *Callinectes sapidus* Rathbun, 1896 (Moksnes *et al.*, 1997), the ‘Dungeness crab’ *Cancer magister* Dana 1852 (Fernández *et al.*, 1993), the estuarine crabs *Chasmagnatus* (= *Neohelice*) *granulata* Dana 1851 and *Cyrtograpsus angulatus* Dana 1851 (Luppi *et al.*, 2001) and *Paralithodes camtschaticus* (Tilesius, 1815) (Stevens & Swiney, 2005), among others. The presence of larger conspecific individuals in the boxes may also account for the small number of early stages found in them. *Halicarcinus planatus* was observed to feed on recently moulted or smaller conspecific individuals under laboratory conditions (Vinuesa, unpublished results). The presence of larger conspecific individuals in the boxes may also account for the small number of early stages found in them.

Post-settlement movements may result from ontogenetic shifts in diet, behaviour and shelter requirements (Herrnkind & Butler, 1986). This behaviour has been described for several decapods, e.g. *P. argus* (Herrnkind & Butler, 1986), *C. sapidus* (Moksnes *et al.*, 1998), *Carcinus maenas* L. (Moksnes, 2002), *C. granulata* and *C. angulatus* (Luppi *et al.*, 2002) and *P. platypus* (Tapella *et al.*, 2009). The spatial separation among different developmental stages of *H. planatus* may imply a microhabitat shift during recruitment, which seems to be particularly pronounced in males. The panels may become less safe as individuals increase in size, compelling juveniles to occupy more protected habitats. The *H. planatus* population from the Golfo San Jorge showed a strongly female-biased sex-ratio, which could be

**Table 3.** Similarity percentage analysis based on the Bray–Curtis dissimilarity index and transformed absence–presence data. The percentages of presence in the collectors are shown. All values are expressed as percentage of presence.

	Presence in panels (%)	Presence in boxes (%)	Dissimilarity contribution	Accumulated dissimilarity
Settlers	65	4	27.54	27.54
Recruits	74	63	24.68	52.21
Late juveniles	43	63	24.65	76.87
Mature crabs	17	50	23.13	100.00




Fig. 3. Abundance (number of crabs per month) of the developmental stages of *Halicarcinus planatus*. Crabs from both localities are grouped.

explained by the movement of males to more cryptic habitats at early stages. In the Deseado estuary, all crabs collected between low intertidal and upper subtidal rocky shore (Vinuesa & Ferrari, 2008a) and holdfasts of *Macrocystis pyrifera* (López Gappa *et al.*, 1982) were exclusively females.

The recruitment may be mainly limited by settlement when there is a good correlation between larval settlers, recruits and the subsequent stages (Connell, 1985; Doherty & Williams, 1988; Palma *et al.*, 1998) or regulated, if the population density is affected by post-settlement processes (Eggleston & Armstrong, 1995; Moksnes *et al.*, 1998; Luppi *et al.*, 2002; Sainte-Marie & Lafrance, 2002). Temporal distribution of larvae, settlers and recruits of *H. planatus* showed a clear correlation. Prolonged periods of occurrence with peaks of abundance were observed for these stages. Post-settlement factors (shelter, food, predation and cannibalism), may have important roles in microhabitat selection and distribution of juveniles of *H. planatus*. On this basis, the species in Golfo San Jorge would follow a mixed pattern of recruitment, during which individuals undertake movements leading to differences in the spatial distribution of the developmental stages and sexes.

## ACKNOWLEDGEMENTS

Thanks are due to the diver Mr Héctor Durbas for the installation and recovery of the collectors. We appreciate the helpful comments of two anonymous referees on the manuscript. This work was supported by the Project PNUD ARG 02/018: 'Coastal Contamination Prevention and Marine Biodiversity Management' Proyecto Marino Patagónico, Donación GEF/BIRF (Subproject AB-67).

## REFERENCES

Anderson M.J. (2001) A new method of non-parametric multivariate analysis of variance. *Austral Ecology* 26, 32–46.

Anderson M.J. (2005) *PERMANOVA: a FORTRAN computer program for permutational multivariate analysis of variance*. Department of Statistics, University of Auckland, New Zealand.

Arnaud P.M. (1974) Contribution à la bionomie marine benthique des régions antarctiques et sub-antarctiques. *Téthys* 6, 465–656.

Beck M.W. (1995) Size-specific shelter limitation in stone crabs: a test of the demographic bottleneck hypothesis. *Ecology* 76, 968–980.

Boschi E.E., Scelzo M.A. and Goldstein B. (1969) Desarrollo larval del cangrejo, *Halicarcinus planatus* (Fabricius) (Crustacea, Decapoda, Hymenosomidae), en el laboratorio, con observaciones sobre la distribución de la especie. *Bulletin of Marine Science* 19, 225–242.

Botero L. and Atema J. (1982) Behavior and substrate selection during larval settling in the lobster *Homarus americanus*. *Journal of Crustacean Biology* 2, 59–69.

Brandhorst R. and Castello J. (1971) *Evaluación de los recursos de anchoíta (Engraulis anchoíta) frente a la Argentina y Uruguay*. Informe Técnico Proyecto Desarrollo Pesquero, FAO 29, 63 pp.

Caley M.J., Carr M.H., Hixon M.A., Hughes T.P., Jones G.P. and Menge B.A. (1996) Recruitment and the local dynamics of open marine populations. *Annual Review of Ecology and Systematics* 27, 477–500.

Clarke K.R. and Warwick R.M. (2001) *Change in marine communities: an approach to statistical analysis and interpretation*. 2nd edition. Bournemouth: Bourne Press Ltd.

Connell J.H. (1985) The consequences of variation in initial settlement vs. post-settlement mortality in rocky intertidal communities. *Journal of Experimental Marine Biology and Ecology* 93, 11–45.

Doherty P.J. and Williams D.McB. (1988) The replenishment of populations of coral reef fishes, recruitment surveys, and the problems of variability manifest on multiple scales. *Bulletin of Marine Science* 41, 411–422.

Eggleston D. and Armstrong D. (1995) Pre- and post-settlement determinants of estuarine Dungeness crab recruitment. *Ecological Monographs* 65, 193–216.

Fernández M., Iribarne O. and Armstrong D. (1993) Habitat selection by young-of-the-year Dungeness crab *Cancer magister* and predation risk in intertidal habitats. *Marine Ecology Progress Series* 92, 171–177.

- Forward Jr R.B., De Vries M.C., Ritschoff D., Frankel D.A., Bischof J.P., Fisher C.M. and Welch J.M.** (1996) Effects of environmental cues on metamorphosis of the blue crab *Callinectes sapidus*. *Marine Ecology Progress Series* 131, 165–177.
- Gebauer P., Paschke K. and Anger K.** (2003) Delayed metamorphosis in decapod crustaceans: evidence and consequences. *Revista Chilena de Historia Natural* 76, 169–175.
- Gil D.G. and Zaixso H.** (2007) The relation between feeding and reproduction in *Anasterias minuta* (Asteroidea: Forcipulata). *Marine Biology Research* 3, 256–264.
- Herrnkind W.F. and Butler M.J.** (1986) Factors regulating postlarval settlement and juvenile microhabitat use by spiny lobsters *Panulirus argus*. *Marine Ecology Progress Series* 34, 23–30.
- Hopkins P.M.** (1992) Hormonal control of the molt cycle in the fiddler crab *Uca pugnator*. *American Zoologist* 32, 450–458.
- Hureau J.** (1970) Biologie comparée de quelques Poissons antarctiques (Nototheniidae). *Bulletin de l'Institut Océanographique, Monaco* 68, 1–250.
- Jensen G.C.** (1991) Competency, settling behavior, and postsettlement aggregation by porcelain crab megalopae (Anomura: Porcellanidae). *Journal of Experimental Marine Biology and Ecology* 153, 49–61.
- López Gappa J.J., Romanello E.E. and Hernández D.A.** (1982) Observaciones sobre la macrofauna y flora asociadas a los grampones de *Macrocystis pyrifera* (L.) C.Agh. en la ría Deseado (Santa Cruz, Argentina). *Ecosur* 9, 67–106.
- Lucas J.S.** (1980) Spider crabs of the family Hymenosomatidae (Crustacea; Brachyura) with particular reference to Australian species: systematics and biology. *Records of the Australian Museum* 33, 148–247.
- Luppi T., Spivak E. and Anger K.** (2001) Experimental studies of predation and cannibalism in recruits of *Chasmagnathus granulata* and *Cyrtograpsus angulatus* (Brachyura: Grapsidae). *Journal of Experimental Marine Biology and Ecology* 265, 29–48.
- Luppi T., Spivak E., Anger K. and Valero J.** (2002) Patterns and processes of *Chasmagnathus granulata* and *Cyrtograpsus angulatus* (Brachyura: Grapsidae) recruitment in Mar Chiquita coastal lagoon, Argentina. *Estuarine, Coastal and Shelf Science* 55, 287–297.
- Marx J. and Herrnkind W.** (1985) Factors regulating microhabitat use by young juvenile spiny lobsters, *Panulirus argus*: food and shelter. *Journal of Crustacean Biology* 5, 650–656.
- Melrose M.J.** (1975) The marine fauna of New Zealand: family Hymenosomatidae (Crustacea, Decapoda, Brachyura). *Memoirs of the New Zealand Oceanographic Institute* 34, 1–123.
- Moksnes P.O.** (2002) The relative importance of habitat-specific settlement, predation and juvenile dispersal for distribution and abundance of young juvenile shore crabs *Carcinus maenas*. *Journal of Experimental Marine Biology and Ecology* 215, 157–187.
- Moksnes P., Lipcius R., Pihl L. and van Montfrans J.** (1997) Cannibal-prey dynamics in young juveniles and postlarvae of the blue crab. *Journal of Experimental Marine Biology and Ecology* 215, 157–187.
- Moksnes P.O., Pihl L. and van Montfrans J.** (1998) Predation on postlarvae and juveniles of the shore crab *Carcinus maenas*: importance of shelter, size, and cannibalism. *Marine Ecology Progress Series* 166, 211–225.
- Moksnes P.O. and Wennhage H.** (2001) Methods for estimating decapod larval supply and settlement: importance of larval behavior and development stage. *Marine Ecology Progress Series* 209, 257–273.
- Palacios R., Armstrong D.A., Armstrong J.L. and Williams G.** (1985) Community analysis applied to characterization of blue king crab habitat around the Pribilof Islands. In Melteff B. (ed.) *Proceedings of the International King Crab Symposium, Anchorage, Alaska, January 22–24, 1985*, pp. 193–209.
- Palma A.T., Wahle R.A. and Steneck R.S.** (1998) Different early postsettlement strategies between American lobsters *Homarus americanus* and rock crabs *Cancer irroratus* in the Gulf of Maine. *Marine Ecology Progress Series* 162, 215–225.
- Richer de Forges B.** (1977) Étude du crabe des Iles Kerguelen: *Halicarcinus planatus* (Fabricius). *Comité National Française des Recherches Antarctiques, Paris, France* No. 42, 71–133.
- Rodríguez S.R., Ojeda F.P. and Inestrosa C.** (1993) Settlement of marine benthic invertebrates. *Marine Ecology Progress Series* 97, 193–207.
- Sainte-Marie B. and Lafrance M.** (2002) Growth and survival of recently settled snow crab *Chionoecetes opilio* in relation to intra- and intercohort competition and cannibalism: a laboratory study. *Marine Ecology Progress Series* 244, 191–203.
- Stevens B.G. and Kittaka J.** (1998) Postlarval setting behaviour, substrate preference, and time to metamorphosis for red king crab *Paralithodes camtschaticus*. *Marine Ecology Progress Series* 167, 197–206.
- Stevens B.G. and Swiney K.M.** (2005) Post-settlement effects of habitat type and predator size on cannibalism of glaucothoe and juveniles of red king crab *Paralithodes camtschaticus*. *Journal of Experimental Marine Biology and Ecology* 321, 1–11.
- Sokal R.R. and Rohlf F.J.** (1995) *Biometry. The principles and practice of statistics in biological research*. 3rd edition. New York: W.H. Freeman and Co.
- Tapella F., Romero M.C., Stevens B. and Buck C.** (2009) Substrate preferences and redistribution of blue king crab *Paralithodes platypus* glaucothoe and first crab on natural substrates in the laboratory. *Journal of Experimental Marine Biology and Ecology* 372, 31–35.
- Vinuesa J.H.** (2005) Distribución de crustáceos decápodos y estomatópodos del Golfo San Jorge, Argentina. *Revista de Biología Marina y Oceanografía* 40, 7–21.
- Vinuesa J.H. and Ferrari L.** (2008a) Postlarval development of *Halicarcinus planatus* females (Crustacea, Decapoda, Hymenosomatidae) in the estuary of Deseado River, South Western Atlantic Ocean. *Scientia Marina* 72, 127–132.
- Vinuesa J.H. and Ferrari L.** (2008b) Reproductive biology of the crab *Halicarcinus planatus* (Decapoda, Hymenosomatidae) in the estuary of the Deseado River. *Marine Biology* 154, 345–354.
- and
- Zaixso H.E. and Pastor C.T.** (1977) Observaciones sobre la ecología de los mitílicos de la ría Deseado. I. Distribución y análisis biocenótico. *Ecosur* 4, 1–46.

#### Correspondence should be addressed to:

J.H. Vinuesa  
 Instituto de Desarrollo Costero  
 Universidad Nacional de la Patagonia  
 San Juan Bosco (UNPSJB) Campus Universitario  
 Ruta No. 1, Km 4 (9000) Comodoro Rivadavia  
 Chubut, Argentina  
 email: vinuesa@unpata.edu.ar