



# XI CONGRESO ARGENTINO DE MICROBIOLOGÍA GENERAL

5 al 7 de Agosto de 2015  
Córdoba, Argentina

**SAMGE**

Asociación Civil de Microbiología General

health, as well as for the agricultural and food industry; although inhibition of phytopathogenic fungi was not observed. Maximum inhibitory activity was obtained in CSM medium at 20°C, showing that bacterial antagonism was neither related to siderophore- nor biosurfactant-like compounds, although both activities were detected for living-cell and cell-free supernatants. The antimicrobial showed similar properties, such as positive net charge, resistance to proteases and the UV-Vis spectrum, to those of pseudobactin, a well-known siderophore produced by species of *P. fluorescens*. However, the non-siderophore nature of the produced antimicrobial makes difficult to relate it with pseudobactin. Further studies including HPLC-MS/MS and NMR are expected to help at elucidating the *P. yamanorum* 8H1<sup>T</sup> antimicrobial structure and its potential novelty.

Código de Resumen: BF-022

Sección: Biotecnología y Fermentaciones

Modalidad: Poster

### **IMPACT OF MANGANESE ON THE PRODUCTION OF A BIOMASS ASSOCIATED β-D-GLUCOSIDASE ACTIVITY USING A THERMOPHILIC *Bacillus licheniformis* STRAIN**

M.B. Abdulhamid<sup>1</sup>, M.D. Baigori<sup>1,2</sup>, L. Pera<sup>1</sup>.

<sup>1</sup>PROIMI-CONICET. <sup>2</sup>Universidad Nacional de Tucumán.

beluabdul@hotmail.com

The β-Glucosidase (EC 3.2.1.21) constitutes a group of well-studied hydrolases. This enzyme catalyzes the hydrolysis of arylglucosides, alkylglucosides, cellobiose, and cellooligosaccharides. The interest in this biocatalyst centers on its roles in the enzymatic hydrolysis of cellulose. The rate of cellulose hydrolysis can be improved by supplementing commercial cellulases with immobilized β-D-glucosidase, which usually has high stability and can be recovered and reused. In addition, for industrial saccharification of cellulosic materials, β-glucosidases from thermophilic bacteria are also of particular interest due to their increased stability. In this work, we study the influence of manganese on the production of a biomass associate β-D-glucosidase activity using a thermophilic *Bacillus licheniformis* strain. Assays were performed at 45 °C in 500 ml Erlenmeyer containing 200 ml of LB medium supplemented with 0 - 1.0 mM MnCl<sub>2</sub>. The β-D-glucosidase activity was also determined at 45 °C using 3.6 mM p-nitrophenyl-β -D-glucopyranoside (Sigma) as substrate. Cells were harvested by centrifugation and washed twice with 100 mM Tris-HCl buffer (pH 7). The pellet was resuspended in the same buffer, and it was directly used as the β-D-glucosidase source. The mixture was shaken at 1000 rpm. Then, the absorbance of the supernatant was measured at 405 nm and the enzyme activity calculated and related to the biomass dry weight. One unit of the enzyme was defined as the amount needed to release 1 μmol p-nitrophenol per min. Thus, dose-response experiments showed that in the presence of 0.3 mM MnCl<sub>2</sub> the enzyme production was increased by about 20%. Under this culture condition, a specific activity value of 19.99 U per mg of dry weight was obtaining after 4 h of cultivation. Finally, these results could be of relevance to the bioethanol industry where lignocellulosic material is used as feedstock for fermentation and, which should be treated enzymatically. The use of naturally bound enzymes is an important immobilization technique. This type of biocatalyst system is potentially cost-effective because the biomass can be directly utilized in the treatment.

Código de Resumen: BF-023

Sección: Biotecnología y Fermentaciones

Modalidad: Poster

### ***Aspergillus terreus* STRAIN IMPROVEMENT FOR ENHANCED LOVASTATIN PRODUCTION**

J.D. Babot<sup>1</sup>, F.C. Caro<sup>1</sup>, O.D. Delgado<sup>2,3</sup>, J.I. Fariña<sup>1,3</sup>.

<sup>1</sup>PROIMI CONICET. <sup>2</sup>CITCA CONICET. <sup>3</sup>FACEN UNCA.

florcecaro@gmail.com

Cholesterol plays a vital role in body metabolism and membrane transport, and acts as precursor for the synthesis of several key biomolecules. Nevertheless, changes in cholesterol level lead to cardiovascular disorders, like atherosclerosis and hypercholesterolemia, which are currently the main causes of death. This is why controlling cholesterol by inhibition of its biosynthesis is a promising approach. Cholesterol is synthesized from acetyl-CoA through a complex pathway, where the rate-limiting step is the conversion of HMG-CoA to mevalonate, catalyzed by HMG-CoA reductase. This key enzyme is selectively and competitively inhibited by lovastatin, a fungal secondary metabolite used as a hypocholesterolemic which can therefore reduce the risk of cardiovascular diseases. Lovastatin production is normally carried out using selected *Aspergillus terreus* strains, however industrial process yields may be improved by strain manipulation. Accordingly, the aim of this work was to develop a lovastatin-hyperproducing *A. terreus* strain. To this end, 10<sup>7</sup>-spores/mL suspensions of lovastatin-producer *A. terreus* MEC were exposed to UV radiation for different times ranging from 5 to 15 min. Spores were kept in the dark for 30 min, plated onto PDA plates and incubated at 25°C for 48 h. Isolated colonies were transferred to an optimized lovastatin production medium (SQop) containing cheese-whey as substrate and incubated at 25°C for 14 days. Lovastatin was extracted from fungal colonies by using ethyl acetate and converted to its β-hydroxyacid form by alkaline hydrolysis. Organic extracts were preliminary