



First isolation of *Mycobacterium avium* subsp. *paratuberculosis* in a dairy goat in Argentina: Pathology and molecular characterization

M.A. Fiorentino^{a,*}, A. Gioffré^b, K. Cirone^a, C. Morsella^a, B. Alonso^c, F. Delgado^b, F. Paolicchi^a

^a Laboratorio de Bacteriología, Departamento de Producción Animal, Estación Experimental Agropecuaria Balcarce, INTA, Argentina

^b CICVyA, Castelar, INTA, Argentina

^c GELAB, SENASA, Argentina

ARTICLE INFO

Article history:

Received 16 March 2012

Received in revised form 6 June 2012

Accepted 19 June 2012

Available online 6 July 2012

Keywords:

Paratuberculosis

Dairy goats

Mycobacterium avium subsp. *paratuberculosis*

ABSTRACT

No data concerning the presence of caprine paratuberculosis (PTBC) in Argentina are currently available. In this work, a Saanen dairy goat herd located in Buenos Aires province was sampled and tested for PTBC. Feces and blood samples were collected from 210 goats. Sera were analyzed by ELISA and fecal samples were pooled for culture. Necropsy was performed in two affected adult goats. Colonies were characterized and genotyped by IS900 PCR and IS1311 PCR-REA, respectively. Gross pathology and histopathology of necropsied goats were consistent with PTBC. Bacteriology, serology, and PCR confirmed the diagnosis. The *Mycobacterium avium* subsp. *paratuberculosis* isolates showed a restriction pattern characteristic of type C strains. This is the first report on the isolation and molecular characterization of *M. avium* subsp. *paratuberculosis* in dairy goats in Argentina.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

Paratuberculosis (PTBC) or Johne's disease is a chronic, contagious and emaciating gastrointestinal disease of ruminants caused by infection with *Mycobacterium avium* subsp. *paratuberculosis* (Map). The disease is not only troublesome for the cattle and sheep industries, but also for breeders of other ruminants, such as elk, deer, and goats, as well as for managers of captive wildlife (Ayele et al., 2001). In addition, the etiologic agent has also been implicated as a possible cause of Crohn's disease in humans (Thompson, 1994). In cattle, transmission of Map has been shown to occur by multiple routes, including the fecal–oral route, intrauterine infection, and via milk or

colostrum (Sweeney et al., 1992). Infected animals may excrete Map in their feces and milk and spread the infection, which can cause considerable financial losses due to premature culling and death (Ayele et al., 2001). It is assumed that this organism infects animals in the first months of life and elicits a slowly progressive inflammatory response in the gastrointestinal tract that is not clinically evident until months to years later. Clinical signs of this infection are the same as for many other diseases: chronic weight loss and unremitting diarrhea. Diarrhea is usually absent in infected sheep and infrequent in infected goats and deer (Stehman, 1996). In advanced cases, the organism can disseminate beyond the gastrointestinal tract to other organ systems (Whitlock et al., 1997). The disease in small ruminants is widespread in the world (Harris and Barletta, 2001). In Argentina, although caprine PTBC was early reported by Ubach (1941), the etiologic agent has not been isolated yet. The present work constitutes the first report of the isolation and molecular characterization of Map in dairy goats in Argentina.

* Corresponding author. Tel.: +54 02266439121; fax: +54 02266439101.

E-mail address: mafiorentino@balcarce.inta.gov.ar (M.A. Fiorentino).

2. Materials and methods

2.1. Population under study

Goats from a Saanen dairy goat herd located in Buenos Aires province, Argentina, were sampled and tested for PTBC as part of a research project. The herd consisted of 210 adult animals, all of which were sampled for this study. Goats grazed on pasture during the day. At night, goats were closed into the shed. Goats were milked mechanically twice daily.

2.2. Samples

Feces and blood samples were simultaneously collected once from each goat. Feces were collected *per rectum*, transferred to a sterile plastic bag and kept at -20°C until processing. Blood samples (10 mL) were obtained from the jugular vein and, after clotting and centrifugation, the serum was harvested and frozen at -20°C until tested. Fecal samples were pooled for culture. Pools were formed from samples of five animals of the same age group. Nineteen animals with apparent symptoms of PTBC were cultured individually.

2.3. Fecal smears and culture

Fecal smears were heat-fixed and stained by the Ziehl Neelsen's (ZN) method (OIE Manual, 2011). Culture from fecal samples was performed after applying a decontamination process. Briefly, samples were homogenized and decontaminated with hexadecylpyridinium chloride (Sigma, USA) (final concentration 0.75%) and inoculated onto three slants of Herold's egg yolk medium (pH 7.1–7.4), with and without mycobactin J (Allied Monitor Inc., MO, Fayette, USA). The cultures were incubated at 37°C for up to 6 months and examined biweekly for bacterial growth. Typical Map colonies were primarily characterized on the basis of mycobactin J dependency, acid fastness and slow growth. Colonies were then characterized and genotyped by IS900 PCR and IS1311 PCR-REA (restriction endonuclease analysis), respectively (Collins et al., 1993; Marsh et al., 1999).

2.4. IS900 PCR from cultures

PCR was used for the detection of IS900, as described by Collins et al. (1993). A loopful of the culture was suspended in distilled water and subjected to freeze-thaw cycles to lyse bacterial cells. The supernatant was subjected to PCR and the following conditions of amplification were used, generating a 218-bp amplification product. The mixture for the amplification consisted of: 10 mM Tris-HCl buffer, pH 9.0, 50 mM KCl and 0.1% Triton X-100, 1.5 mM MgCl_2 , 0.2 mM of each deoxynucleotide triphosphate (dNTP), 250 ng of each primer, 5 μL of template, and 1.25 U of Taq polymerase (Promega, Madison, WI, USA) and water to a final volume of 50 μL . PCR products were analyzed by submerged electrophoresis on a 2% agarose gel.

2.5. IS1311 PCR-REA

In order to identify the subspecies of *M. avium* isolated and strain type, PCR and PCR-REA of IS1311 gene were performed. The protocol according to Marsh et al. (1999) was used. Briefly, the IS1311 was amplified using the primers M56 and M119 under the following conditions: one cycle of denaturation at 94°C for 3 min, 35 cycles of denaturation at 94°C for 30 s, annealing at 62°C for 15 s and extension at 72°C for 1 min. The 608-bp product was digested using 2 U of *Hinf* (NEB) and *MseI* (NEB) using the buffer provided by the manufacturer and adding 1.6 μL of 100 $\mu\text{g}/\text{mL}$ bovine serum albumin per reaction. The restriction products were resolved in 4% agarose gels stained with ethidium bromide. The presence of 67-bp, 218-bp, 285-bp and 323-bp products is indicative of type C strains, while the presence of restriction fragments of 285 bp and 323 bp is indicative of type S strains.

2.6. Pathological examination

After euthanasia, necropsy was performed in two affected adult goats (#1 and #2). Samples of the ileocecal valve, ileum, jejunum, mesenteric lymph node (MLN), and cecum were collected for histopathological examination and culture. Samples for histopathology were fixed in 10% buffered formalin and processed following routine procedures. Sections (2 μm in

thickness) were cut from each sample and stained with hematoxylin and eosin (HE) and by the ZN method for acid-fast bacteria (AFB). Sections of the ileum and MLN were used for immunohistochemistry. The technique was performed following procedures previously described (Delgado et al., 2009).

The smears were heat-fixed and stained with the ZN method. The presence of clumps of three or more strongly AFB was recorded as positive (OIE Manual, 2011).

Tissue samples for bacteriological examination were homogenized with 5 mL sterile physiological saline and processed as described for fecal culture.

2.7. Serology

The sera of all the animals were analyzed by indirect enzyme immunoassay (ELISA) using Map protoplasmic antigen (PPA-3; Allied Monitor, Fayette, MO, USA). The indirect ELISA technique used was a slight modification of that described by Verna et al. (2007). Briefly, the antigen was diluted to a concentration of 0.07 mg/mL in carbonate buffer and a volume of 100 μL per well was allowed to fix overnight at 4°C in flat-bottomed polystyrene plates (Immulon I, Dynatech, USA). To eliminate unspecific antibodies, sera were previously absorbed with a 1:1 suspension of *Mycobacterium phlei* for 2 h at room temperature under stirring and then centrifuged. After centrifugation, the supernatant fraction of each sample was recovered and diluted 1:100 in phosphate-buffered-Tris-glycerine solution (PBS-TG) and then 100 μL was added to each well. Each sample was tested in duplicate. The plates were then incubated for 2 h at room temperature. After washing with PBS-TG, a 1:1500 dilution of a horseradish peroxidase-labeled Protein G (Bio-Rad Laboratories, Hercules, California, USA) was added to each well. After 2 h of incubation at room temperature, the plates were washed thoroughly, and 100 μL of substrate solution (0.04 M ABTS in citrate buffer (0.05 M pH 5) and 1 mM H_2O_2 100 vol) was added to each well. After 10 min of incubation at room temperature, the A450 was recorded with a microplate reader (Titertek, Multiskan, ICN, Finland). The cut-off value for a positive test was based on the absorbance values at 450 nm of 326 serum samples from goats testing negative to the Johnin test, fecal culture and from a region of Argentina where no animals with symptoms of PTBC have been observed. Samples having at least twice the mean absorbance of the negative serum were recorded as positive (Rajukumar et al., 2001). Accordingly, the optical density (OD) value of 0.375 was fixed as the cut-off value for deciding the positive sera in the ELISA. The OD results were transformed to an index value by division of the mean OD obtained with the serum samples from the animals by the mean OD of the negative controls.

3. Results and discussion

The clinical signs reported in the goats studied in this work were mainly those of a chronic wasting disease which leads to emaciation, severe weakness, prostration and death and the disease was characterized mainly by a progressive loss of weight but without diarrhea.

Culture is considered the only definitive and critical technique to diagnose Johne's disease (Whittington et al., 1999). Pooling sheep fecal samples has been reported by Australian investigators as an efficient method to detect PTBC in flocks and has been used as an excellent tool to monitor sheep flocks over time. This same approach could also be used for deer and goat flocks (Whitlock, 2005). In agreement with these findings, in our work, 15 out of 34 (44.1%) pools of fecal samples tested in the infected herd were culture-positive. The use of pooled fecal samples was a useful tool, since Map excretion was demonstrated in at least 15 out of 170 (8.8%) animals without clinical signs. The highest number of positive pools was that of the animals aged 4 years (Table 1). Nevertheless, clinical signs and fecal culture-positive samples were also observed in 1-year-old goats. Typically, sheep and goats are 2–4 years old when

Table 1

Isolation of *Mycobacterium avium* subsp. *paratuberculosis* by culture of pools of fecal samples vis-à-vis age of goats.

Age (years)	Pools	Positive culture
1	3	1 (33%)
2	5	2 (40%)
3	7	3 (42.8)
4	7	6 (85.7%)
>4	12	3 (25%)

Table 2

Mycobacterium avium subsp. *paratuberculosis* fecal culture and ELISA in goats with apparent symptoms of paratuberculosis.

Age (years)	Goats sampled	Fecal culture-positive	ELISA-positive
1	2	0/2	2/2
2	1	1	1
3	4	3/4	4/4
4	6	4/6	5/6
>4	6	0/6	5/6

culled for PTBC. However, it has been observed that goats are more susceptible to PTBC than other species, and that, in environments with heavy exposure to Map, they may develop clinical signs as early as 6 months of age (Lybeck et al., 2010). Eight out of the 19 (42%) goats with apparent symptoms of PTBC were Map fecal culture-positive and 18 out of 19 (94.7%) were ELISA-positive (Table 2). The herd sero-incidence of PTBC was 48%.

The two necropsied goats showed classical clinical signs of caprine PTBC: emaciation despite willingness to eat, dry and rough hair coat, and no evidence of diarrhea. The MLNs were strongly enlarged and edematous, and focal granulomatous lymphadenitis with central necrosis and calcification was detected in one of the goats. The lymphatic vessels draining the intestine were distended. Severe atrophy of perirenal, mesenteric and pericardial fat was observed in the two goats. Map was also isolated from the intestine, lymph nodes and mammary glands of the two goats. Typical Map colonies developed after 20 and 30 days of culture of samples from organs and feces, respectively.

Two forms of PTBC have been described: multibacillary and paucibacillary (Clarke, 1997). The cases presented in this study showed both types of PTBC. The gross pathology, histopathology, bacteriology, and serology of the necropsied goat #1 were consistent with the multibacillary form, which is characterized by granulomatous enteritis by




Fig. 1. Goat #1. Ileum. Diffuse granulomatous enteritis. Numerous acid fast bacilli were detected within the cytoplasm of macrophages in the lamina propria (arrowheads). Ziehl Nielsen staining, Bar: 100 μm.

macrophages filled with abundant AFB (Fig. 1) and which is associated with a strong humoral immune response (Catton, 2002). However, in goat #2, paucibacillary PTBC was observed and the serology by ELISA was negative, no AFB were observed in tissue sections stained with the ZN method, and Map was isolated from the ileum and MLN. The paucibacillary form is seen less often and may occur more frequently in sheep than in other animal species (Corpa et al., 2000). Other authors have reported paucibacillary PTBC in goats but its prevalence is unknown (Kruze et al., 2006). This is the first report of caprine PTBC in Argentina that confirms this less common paucibacillary form of PTBC in small ruminants.

All cultures were confirmed as Map by IS900-PCR. Map isolates have been classified into two main groups based on culture characteristics and molecular characterization: sheep (also called S or Type I/Type III) and cattle (also called C or Type II) (Stevenson et al., 2002). The PCR-REA based on polymorphisms in IS1311 allows differentiating Map from *M. avium* subsp. *avium* (Marsh et al., 1999). In addition, a polymorphism at base 223 in the IS1311 sequence is useful to type S and C strains. In our work, all the strains tested resulted C type (Fig. 2). Since C-type strains are commonly isolated in many species including cattle, goats and sheep




Fig. 2. PCR-restriction analysis of mycobacterial isolates. Samples from 5 representative isolates are shown. (A) Lanes 2–6, IS1311-positive isolates (608-bp); lane 7, negative control; lane 1, molecular marker. (B) Lanes 2–6, PCR product digested with *Hinf-Mse I*. Lane 1, molecular marker. Arrowheads show the fragments of 67-bp, 218-bp, 285-bp and 323-bp, characteristic of type C strains. Molecular marker: 50-bp DNA step ladder (Promega).

(De Juan et al., 2005), we cannot conclude which was the source of infection of this flock.

Goats are most commonly infected through the fecal–oral route: by ingestion of contaminated milk or food products or by accidental ingestion of the organism from contaminated surfaces (Harris and Barletta, 2001). Practical implementation of management practices, such as removal of kids from their dams, segregation of young offspring from adults, and exclusive use of milk replacer to feed kids, is difficult and even sometimes financially impossible for many dairy goat herd owners (Kruze et al., 2006).

4. Conclusion

In Argentina, caprine production has been historically based on raising kids to slaughter and, in some cases, meat and milk production. In the last years, milk production has grown and, at present, there are 213 dairy goat herds, 27 of which are in the province of Buenos Aires (Ginés de Gea et al., 2005). In the present work, the clinical suspicion of PTBC disease in a dairy goat herd in Buenos Aires province was confirmed by serological, pathological, bacteriological and molecular tools. These findings confirm that the disease occurs in goats from Argentina, as it had been early suggested (Ubach, 1941). Due to the lack of awareness about PTBC among goat owners, lack of regulations on animal trade and standard goat husbandry practices, a continuous spread of PTBC among and within goat herds should be expected. The safety of raw milk derived from Map infected dairy goats becomes an important consideration for public health, because Map could be associated with Crohn's disease in humans. However, more exhaustive work is needed to estimate the herd-level prevalence of Map infection in dairy goat herds in Argentina.

Acknowledgements

Thanks to Laura Mendez and Claudia Moreno for their collaboration with the laboratory tests. Financial support was provided by Tuberculosis and Paratuberculosis Project AESA 52-202831 from Instituto Nacional de Tecnología Agropecuaria, Argentina.

References

- Ayele, W.Y., Machackova, M., Pavlik, I., 2001. The transmission and impact of paratuberculosis infection in domestic and wild ruminants. *Vet. Med.* 46, 205–224.
- Catton, B.M., 2002. Paucibacillary paratuberculosis in a goat. *Can. Vet. J.* 43, 787–788.
- Clarke, C.J., 1997. The pathology and pathogenesis of paratuberculosis in ruminants and other species. *J. Comp. Pathol.* 116, 217–261.
- Collins, D.M., Stephens, D.M., de Lisle, G.W., 1993. Comparison of polymerase chain reaction tests and faecal culture for detecting *Mycobacterium paratuberculosis* in bovine faeces. *Vet. Microbiol.* 36, 289–299.
- Corpa, J.M., Garrido, J., García-Marín, J.F., Pérez, V., 2000. Classification of lesions observed in natural cases of paratuberculosis in goats. *J. Comp. Pathol.* 122, 255–265.
- De Juan, L., Mateos, A., Dominguez, L., Sharp, J.M., Stevenson, K., 2005. Genetic diversity of *Mycobacterium avium* subspecies *paratuberculosis* isolates from goats detected by pulsed-field gel electrophoresis. *Vet. Microbiol.* 106, 249–257.
- Delgado, F., Etchechoury, D., Gioffre, A., Paolicchi, F., Blanco Viera, F., Mundo, S., Romano, M.I., 2009. Comparison between two in situ methods for *Mycobacterium avium* subsp. *paratuberculosis* detection in tissue samples from infected cattle. *Vet. Microbiol.* 134, 383–387.
- Ginés de Gea, S., Petryna, A.M., Mellano, A., Bonvillani, A., Turiello, P., 2005. La situación caprina en las provincias. In: *El ganado caprino en Argentina*, 1ª Edición. UNRC, Río Cuarto, Argentina, pp. 19–27.
- Harris, N.B., Barletta, R.G., 2001. *Mycobacterium avium* subsp. *paratuberculosis* in veterinary medicine. *Clin. Microbiol. Rev.* 14, 489–512.
- Kruze, J., Salgado, M., Paredes, E., Mella, A., Collins, M.T., 2006. Goat paratuberculosis in Chile: first isolation and confirmation of *Mycobacterium avium* subspecies *paratuberculosis* infection in a dairy goat. *J. Vet. Diagn. Invest.* 18, 476–479.
- Lybeck, K.R., Storset, A.K., Djønnø, B., Valheim, M., Olsen, I., 2010. Faecal shedding detected earlier than immune responses in goats naturally infected with *Mycobacterium avium* subsp. *paratuberculosis*. *Res. Vet. Sci.* 91, 32–39.
- Marsh, I., Whittington, R., Cousins, D., 1999. PCR–restriction endonuclease analysis for identification and strain typing of *Mycobacterium avium* subsp. *paratuberculosis* and *Mycobacterium avium* subsp. *avium* based on polymorphisms in IS1311. *Mol. Cell. Probes* 13, 115–126.
- Office International des Epizooties (OIE), 2011. Paratuberculosis (Johne's disease). In: *Manual of Diagnostic Test and Vaccines for Terrestrial Animals 2011*. Available at <http://www.oie.int/en/international-standard-setting/terrestrial-manual/access-online>.
- Rajukumar, K., Tripathi, B.N., Kurade, N.P., Parihar, N.S., 2001. An enzyme linked immunosorbent assay using immunoaffinity purified antigen in the diagnosis of caprine paratuberculosis and its comparison with conventional ELISAs. *Vet. Res. Commun.* 25, 539–553.
- Stehman, S.M., 1996. Paratuberculosis in small ruminants, deer, and South American camelids. *Vet. Clin. North Am. Food Anim. Pract.* 12, 441–455.
- Stevenson, K., Hughes, V., de Juan, L., Inglis, N., Wright, F., Sharp, J., 2002. Molecular characterization of pigmented and nonpigmented isolates of *Mycobacterium avium* subsp. *paratuberculosis*. *J. Clin. Microbiol.* 40, 1804–1978.
- Sweeney, R.W., Whitlock, R.H., Rosenberger, A.E., 1992. *Mycobacterium paratuberculosis* cultured from milk and supramammary lymph nodes of infected asymptomatic cows. *J. Clin. Microbiol.* 30, 166–171.
- Thompson, D.E., 1994. The role of mycobacteria in Crohn's disease. *J. Med. Microbiol.* 41, 74–94.
- Ubach, F.A., 1941. Observaciones sobre la enfermedad de Johne y la enteritis coccidiósica de los bovinos, ovinos y caprinos identificadas en la República Argentina. *Rvta. Med. Vet. Bs. As.* 23, 3–45.
- Verna, A.E., García-Pariente, C., Muñoz, M., Moreno, O., García-Marín, J.F., Romano, M.I., Paolicchi, F., Pérez, V., 2007. Variation in the immuno-pathological responses of lambs after experimental infection with different strains of *Mycobacterium avium* subsp. *paratuberculosis*. *Zoonosis Public Health* 54, 243–252.
- Whitlock, R.H., Rosenberger, A.E., Sweeney, R.W., Spencer, P.A., 1997. Distribution of *Mycobacterium paratuberculosis* in tissues from cattle herds infected with Johne's disease. In: Chiodini, R.J., Hines II, M.E., Collins, M.T. (Eds.), *Proceedings of the 5th International Colloquium on Paratuberculosis*. International Association for Paratuberculosis, Rehoboth, Massachusetts, pp. 200–210.
- Whitlock, R.H., 2005. Johne's disease in small ruminants and other species. In: *Proceeding of the NAVC North American Veterinary Conference*, January 8–12, 2005, Orlando, Florida.
- Whittington, R.J., Marsh, I., McAllister, S., Turner, M.J., Marshall, D.J., Fraser, C.A., 1999. Evaluation of modified BACTEC 12B radiometric medium and solid media for culture of *Mycobacterium avium* subsp. *paratuberculosis* from sheep. *J. Clin. Microbiol.* 37, 1077–1083.