

Trends in shark bycatch research: current status and research needs

Juan M. Molina & Steven J. Cooke

Reviews in Fish Biology and Fisheries

ISSN 0960-3166


Rev Fish Biol Fisheries

DOI 10.1007/s11160-012-9269-3

Reviews in Fish Biology and Fisheries

VOLUME 18 NUMBER 4 2008

Editor **Jennifer Nielsen**



 Springer



 Springer

Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media B.V.. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.

Trends in shark bycatch research: current status and research needs

Juan M. Molina · Steven J. Cooke

Received: 23 December 2011 / Accepted: 21 May 2012
© Springer Science+Business Media B.V. 2012

Abstract Over the last few decades, much effort has been devoted towards quantifying and reducing bycatch in marine fisheries. Of late, there has been a particular focus on sharks given that bycatch is a frequently listed threat for sharks on the International Union for the Conservation of Nature Red List. However, currently there are no quantitative reviews or syntheses that explore the issue of shark bycatch globally which is problematic given that such a synthesis could inform conservation actions and identify pressing research gaps. We performed a qualitative and quantitative survey of the peer-reviewed literature to characterize trends in shark bycatch research with a particular goal of identifying research needs and opportunities. Using a structured literature review we identified 103 papers that met our

search criteria, with the first one published in 1993. Early research efforts focused on documenting the scope of bycatch (i.e., determining that sharks were indeed captured as bycatch), but more recently there have been increased efforts devoted to developing and evaluating bycatch reduction strategies for sharks. Research activity was most common in the North Atlantic (~40 % of the total articles analysed) with comparatively less research in other areas such as the Indo-Pacific region where shark bycatch is regarded as particularly common and problematic. Most studies were observational with comparatively fewer experimental and modeling studies, and even fewer that combined research approaches. Gear modifications (e.g., hook size and type for long lines, net size and mesh design for nets) were the most commonly evaluated strategy for reducing shark bycatch; however, development and use of techniques like repellents, or seasonal area closures, or a combination of strategies, offer interesting possibilities that require further study. In addition, although many sharks are discarded, little is known about post-release survival or sub-lethal consequences of fisheries interactions, or evaluations of different fish handling strategies, making it difficult to quantify the true cost of bycatch or to recommend handling strategies to fishers. Although there are some inherent challenges with developing and testing shark bycatch reduction strategies, there is an urgent need to do so and this would be best achieved through interdisciplinary research that spans field, laboratory, and modeling realms.

J. M. Molina (✉)
Consejo Nacional de Investigaciones Científicas y
Técnicas (CONICET), Buenos Aires, Argentina
e-mail: ichtyostega@gmail.com

J. M. Molina
Departamento de Biología Bioquímica y Farmacia,
Universidad Nacional del Sur, Bahía Blanca,
Buenos Aires, Argentina

S. J. Cooke
Fish Ecology and Conservation Physiology Laboratory,
Department of Biology, Institute of Environmental
Science, Carleton University, Ottawa, ON, Canada

Keywords Sharks · Bycatch · Trends · Conservation · Review

Introduction

Sharks and other chondrichthyans are a relatively conserved group of fishes that have successfully functioned in diverse marine ecosystems for over 400 million years (Camhi et al. 2007). Sharks in particular play an important role in structuring marine communities by influencing mortality rates and behaviour of mesoconsumers and other organisms (Heithaus et al. 2008). The life history of these fishes is characterized by late maturity, long life spans, long gestation periods, and few well developed offspring (Dulvy et al. 2008). Additionally, some sharks gather in schools by age, sex, and reproduction states, while others have restricted distributions (Barker and Schluessel 2005). These life history traits confer a low intrinsic rate of population increase and growth, rendering sharks extremely vulnerable to fishing mortality (Dulvy et al. 2008). Indeed, globally shark populations are in decline (Camhi et al. 2007) as a result of a variety of stressors and threats including fishing mortality arising from targeted harvest and incidental capture (i.e., bycatch). From the 176 species of shark monitored by the International Union for the Conservation of Nature (IUCN), 27.27 % were reported to be Near Threatened, 22.72 % Vulnerable, 6.25 % Endangered, and 5.68 % Critically Endangered (based on an analysis conducted by J. Molina on June 1, 2011). This means that approximately 62 % of the total shark species are facing major conservation threats. In addition, there are also 63 species of sharks that are categorized as data deficient, for which there is simply insufficient validated information available about their status and threats but for which surely more of them are imperiled.¹

Bycatch is the accidental capture of non-targeted species or undesirable size ranges of the target species by non-selective fishing gear (Hall 1996; Crowder and

Murawski 1998). Species with low resilience to fishing practices can be severely affected by bycatch, which poses a serious conservation concern in a variety of ecosystems and for a variety of taxa (e.g., Barker and Schluessel 2005; Raby et al. 2011). Moreover, much of the discarded catch is not documented, which makes a proper evaluation of bycatch more difficult (Romanov 1999; Barker and Schluessel 2005). Early work on bycatch identified that between 17.9 and 39.5 million tons of fish were discarded annually in marine commercial fisheries (Alverson et al. 1994), but more recent estimates suggest that the number may be closer to 6.8 million tons (Kelleher 2005). Although it is unclear why this apparent reduction has occurred, there is some evidence that awareness of bycatch as a conservation concern and the associated development and implementation of bycatch reduction strategies may be contributing factors (Kelleher 2005). There have been a variety of syntheses on bycatch (e.g., Davis 2002; Broadhurst et al. 2002; Hall and Mainprize 2005; Soykan et al. 2008) and even some that briefly discuss sharks (e.g., Lewison et al. 2004; Zydalis et al. 2009) or focus entirely on sharks (e.g., Ward and Hindmarsh 2007; Mandelman et al. 2008; Watson et al. 2008; Blaber et al. 2009). Of the reviews available that focus shark bycatch, they tend to be limited to a particular species (e.g., Watson et al. 2008), certain location (Mandelman et al. 2008; Blaber et al. 2009), or specific type of fishing practice (Ward and Hindmarsh 2007; Gilman et al. 2007), with no global synthesis of trends in shark bycatch research. Bycatch is the most frequent threat for sharks, accounting for 66.9 % of shark species reported by the IUCN that are facing conservation threats (see footnote 1). However, currently there are no quantitative reviews or syntheses that explore the issue of shark bycatch globally. An authoritative and contemporary overview of the threats from non-retention shark fisheries could inform conservation actions and identify pressing research gaps.

The aim of this study was characterize trends in shark bycatch research and identify key research needs and opportunities. Given that shark bycatch is already recognized as a problem, we focus much of our analysis on the evaluation of strategies to reduce bycatch or to improve the fate of sharks that are captured and discarded. To do so, we performed a qualitative and quantitative survey of the peer-reviewed literature dealing with shark bycatch.

¹ All shark species listed in the IUCN red list of endangered species until 2011 were analyzed. Species conservation category, population status and threat information was extracted and used to create a database, from which percentage of each category, and occurrence of each mayor population threat was calculated, in order to describe the conservation status, and find at which extent bycatch represents the major threat for sharks.

Materials and methods

Structured literature searches were conducted to generate a database of shark bycatch literature for the quantitative review. We recognize all elasmobranchs are susceptible to bycatch but for the purpose of this review we focus on sharks (i.e., Selachii) and exclude the rays and skates (i.e., Batoidea). In addition, we focused our synthesis on bycatch associated with commercial or artisanal fisheries and excluded recreational fisheries where fish are discarded but where they may be intentionally targeted as part of a practice known as catch-and-release angling (Arlinghaus et al. 2007). There is certainly a growing interest and body of research on recreational catch-and-release angling impacts on sharks (e.g., Gurshin and Szedlmayer 2004; Lynch et al. 2010), but we consider those issues to be sufficiently different such that we excluded recreational fisheries from the current review. Some papers, especially those that were experimental, used generalized techniques to induce stress and did not specify whether their study was focused on recreational or commercial fisheries (e.g., Brill et al. 2008) so those studies were not excluded. Searches were performed using Web of Science, Scopus, and Google Scholar, with the Boolean search terms: bycatch, by-catch, shark*, incidental catch, and discard*. We recognize that this approach has the potential to miss articles that could otherwise be found by using cited reference searches, trolling through websites for individual researchers, or looking through literature cited lists but this structured approach means that our search could be replicated which is an essential element of quantitative literature reviews (Pullin and Stewart 2006). As noted below, we do refer to and cite additional relevant papers that we found via means other than our structured search, but they were relatively sparse such that we are confident that the trends characterized in the structured review is indeed representative of the broader literature and knowledge base. Only peer-reviewed articles written in English were included in the structured literature review. We recognize the potential bias associated with this approach but the majority of international journals are published in English. We excluded non-peer reviewed literature (e.g., thesis papers, technical reports from governments) because their likelihood of being available in English was much reduced and thus we were concerned with further biasing the sample

towards areas such as North America, Australia and Europe. The literature search was conducted between January and April 2011 and updated in April 2012. The quantitative literature review focused on papers prior to 2012, however, papers from 2012 (including those emanating from special “shark focused” issues of *Journal of Fish Biology and Comparative Physiology and Biochemistry A*) were cited where appropriate. After the structured database was generated, each paper was examined individually and those that were false positives (i.e., did not have to deal with shark bycatch; ~30 %) were excluded yielding a total of 103 papers on shark bycatch. These papers were then read and information extracted and entered into a database (MS Access 2007) by the same individual to ensure consistency. For each article, multiple data elements were extracted including the type (e.g., laboratory experiment, field experiment, modeling study, review, observational study) and general objective of the article (e.g., reduce capture, reduce mortality, supply information, analyze species’ bycatch, analyze fleet bycatch, provide management strategies), type of fishery (e.g., subsistence, commercial) and the scale (artisanal small-scale or industrial), location of the fishery (e.g., north and south Atlantic, north and south Pacific, Indian and Mediterranean oceans), fishing gear (e.g., longline, trawling nets, gill nets, purse seine, etc.), target species, number of shark bycatch species reported/studied, scale of the consequences (e.g., community/assemblage, population, or individual mortality, as well as sub lethal effects, like injury, behaviour, and physiological effects), and proposed changes to reduce bycatch (e.g., changes in gear, fishing practices, handling, use of repellents, etc.). Where summary statistics are presented as means the error term represents one standard deviation.

Findings

Characteristics of shark bycatch literature

In total, we found 103 papers on shark bycatch (Table 1). The first paper (Russell 1993) on shark bycatch was published in 1993 with a reasonably steady increase in publications through to current day. The maximum number of studies per year was in 2011 when 15 articles on shark bycatch were published.

Table 1 Literature (organized alphabetically by author last name) located by the structured literature search and synthesized as part of qualitative and quantitative review of trends in shark bycatch research

Author (year)	Article type	Article objective	Scale of the fishing fleet	Location of the fishery	Fishing gear used	Target species	Scale of the consequences	Proposed changes
Afonso et al. (2011)	Field experiment	Reduce mortality	Large scale	South Atlantic	Long lines	Large osteicht	Mortality	Type/Size of hooks
Alverson et al. (1994)	Review	Supply management strategies	Large scale	Multiple	Multiple types	Large osteicht	Assemblages	Multiple
Arocha et al. (2002)	Observational	Supply information	Large scale	North Atlantic	Long lines	Large osteicht	Population	No change suggested
Baeta et al. (2010)	Observational	Fleet bycatch analysis	Artisanal	North Atlantic	Trammel nets	Polyspecific	Assemblages	No change suggested
Barker and Schluessel (2005)	Review	Supply management strategies	Large scale	Multiple	Multiple types	Small elasmobranch	Assemblages	Type/Size of nets
Beerkircher et al. (2000)	Observational	Supply information	Large scale	North Atlantic	Long lines	Large osteicht	Mortality	Fishing time
Benjamins et al. (2010)	Review	Supply information	Large scale	North Atlantic	Gill nets	Polyspecific	Population	No change suggested
Blaber et al. (2009)	Review	Supply management strategies	Artisanal	Indian	Multiple types	Small elasmobranch	Assemblages	Fishing practice
Belcher and Jennings (2011)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Trawling nets	Shrimp	Assemblages	No change suggested
Bonfil (1995)	Review	Supply management strategies	Large scale	Multiple	Multiple types	Polyspecific	Assemblages	No change suggested
Brewer et al. (1998)	Field experiment	Reduce capture	Large scale	Indian	Trawling nets	Shrimp	Assemblages	Type/Size of Nets
Brill et al. (2008)	Laboratory experiment	Reduce capture	Large scale	North Atlantic	Long lines	Large osteicht	Sub-lethal: behaviour	Repelets
Broadhurst et al. (2002)	Field experiment	Reduce capture	Large scale	Indian	Trawling nets	Shrimp	Population	Type/Size of Nets
Burgess et al. (2005)	Review	Species bycatch analysis	Large scale	North Atlantic	Multiple types	Polyspecific	Population	No change suggested
Camhi et al. (2007)	Observational	Supply information	Large scale	Multiple	Multiple types	Polyspecific	Assemblages	No change suggested
Campana et al. (2009a, b)	Review	Fleet bycatch analysis	Large scale	North Atlantic	Long lines	Large osteicht	Sub-lethal: injury	No change suggested
Carbonell (2003)	Observational	Species bycatch analysis	Large scale	Mediterranean	Trawling nets	Shrimp	Population	No change suggested
Carruthers et al. (2009)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Long lines	Large osteicht	Mortality	Type/Size of Hooks
Carruthers et al. (2011)	Field experiment	Reduce capture	Large scale	North Atlantic	Long lines	Large osteicht	Population	Fishing practice
Cartamil et al. (2011)	Observational	Fleet bycatch analysis	Artisanal	North Pacific	Multiple types	Small osteicht	Assemblages	No change suggested
Clarke (2002)	Observational	Species bycatch analysis	Large scale	North Atlantic	Multiple types	Polyspecific	Population	No change suggested
Coelho and Erzini (2008)	Observational	Supply information	Large scale	Mediterranean	Multiple types	Polyspecific	Population	No change suggested
Coelho et al. (2003)	Field experiment	Fleet bycatch analysis	Artisanal	North Atlantic	Long lines	Small osteicht	Assemblages	Fishing practice
Connolly and Kelly (1996)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Multiple types	Polyspecific	Assemblages	No change suggested

Table 1 continued

Author (year)	Article type	Article objective	Scale of the fishing fleet	Location of the fishery	Fishing gear used	Target species	Scale of the consequences	Proposed changes
Cortés et al. (2009)	Modeling	Supply management strategies	Large scale	North Atlantic	Long lines	Large osteicht	Mortality	No change suggested
Cramer et al. (1998)	Observational	Supply information	Large scale	North Atlantic	Long lines	Large osteicht	Population	No change suggested
Curran and Bigelow (2011)	Field experiment	Reduce capture	Large scale	North Pacific	Long lines	Large osteicht	Assemblages	Type/Size of hooks
de Silva et al. (2001)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Purse-seine	Small osteicht	Assemblages	No change suggested
Diaz and Serafy (2005)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Long lines	Large osteicht	Mortality	No change suggested
Dulvy et al. (2008)	Review	Supply management strategies	Large scale	Multiple	Multiple types	Polyspecific	Population	No change suggested
Fennessy and Isaksen (2007)	Field experiment	Reduce capture	Large scale	Indian	Trawling nets	Shrimp	Population	Type/Size of nets
Fogarty and Murawski (1998)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Multiple types	Polyspecific	Assemblages	No change suggested
Fortuna et al. (2010)	Observational	Fleet bycatch analysis	Large scale	Mediterranean	Trawling nets	Small osteicht	Population	No change suggested
Francis and Duffy (2002)	Observational	Species bycatch analysis	Artisanal	North Atlantic	Trawling nets	Polyspecific	Population	No change suggested
Francis et al. (2001)	Observational	Fleet bycatch analysis	Large scale	South Pacific	Long lines	Large osteicht	Population	No change suggested
Frick et al. (2010)	Laboratory experiment	Supply information	Large scale	South Pacific	Trawling nets	Polyspecific	Sub-lethal: physiological	No change suggested Fishing practice
Gilman (2011)	Review	Supply information	Large scale	Multiple	Purse-seine	Large osteicht	Mortality	Multiple
Gilman et al. (2007)	Observational	Supply management strategies	Large scale	Multiple	Long lines	Polyspecific	Assemblages	Multiple
Godin and Worm (2010)	Review	Supply management strategies	Large scale	Multiple	Multiple types	Polyspecific	Population	Multiple
Grantham et al. (2008)	Field experiment	Supply management strategies	Large scale	South Atlantic	Long lines	Large osteicht	Assemblages	Fishing closure
Gurshin and Szedlmayer (2004)	Field experiment	Supply information	Artisanal	North Atlantic	Long lines	Large osteicht	Sub-lethal: behaviour	No change suggested
Hall et al. (2000)	Review	Supply information	Multiple	Multiple	Multiple types	Polyspecific	Assemblages	No change suggested
Heithaus et al. (2008)	Modeling	Supply information	Large scale	Multiple	Multiple types	Polyspecific	Assemblages	Multiple
Hill and Wassenberg (2000)	Observational	Fleet bycatch analysis	Large scale	Indian	Trawling nets	Shrimp	Assemblages	Type/Size of nets
Huang (2011)	Review	Supply management strategies	Large scale	North Pacific	Long lines	Large osteicht	Assemblages	Multiple
Huang and Liu (2010)	Observational	Fleet bycatch analysis	Large scale	Indian	Long lines	Large osteicht	Population	Type/Size of hooks
Jones et al. (2011)	Observational	Fleet bycatch analysis	Large scale	Indian	Multiple types	Polyspecific	Population	No change suggested
Jordan et al. (2011)	Laboratory experiment	Reduce capture	Large scale	North Pacific	Long lines	Large osteicht	Sub-lethal: behaviour	Repelents

Table 1 continued

Author (year)	Article type	Article objective	Scale of the fishing fleet	Location of the fishery	Fishing gear used	Target species	Scale of the consequences	Proposed changes
Kaplan et al. (2007)	Modeling	Reduce capture	Large scale	North Pacific	Long lines	Large osteicht	Assemblages	Type/Size of hooks
Kersterter and Graves (2006)	Field experiment	Reduce mortality	Large scale	North Atlantic	Long lines	Large osteicht	Mortality	Type/Size of hooks
Krogh and Reid (1996)	Observational	Fleet bycatch analysis	Artisanal	South Pacific	Gill nets	Large elasmobranch	Population	No change suggested
Levesque (2008)	Review	Supply information	Large scale	North Atlantic	Long lines	Large osteicht	Population	Multiple
Lewison et al. (2004)	Review	Supply information	Large scale	Multiple	Multiple types	Polyspecific	Assemblages	No change suggested
Mandelman and Farrington (2007)	Laboratory experiment	Species bycatch analysis		Multiple	Trawling nets		Sub-lethal: injury	Fishing practice
Mandelman et al. (2008)	Review	Fleet bycatch analysis	Large scale	North Atlantic	Long lines	Large osteicht	Assemblages	Multiple
Maunder and Punt (2004)	Review	Supply information	Large scale		Multiple types	Polyspecific	Assemblages	No change suggested
McKinnell (1998)	Observational	Fleet bycatch analysis	Large scale	North Pacific	Driftnets	Mollusc	Population	No change suggested
Megalofonou and Deflorio (2005)	Observational	Fleet bycatch analysis		Mediterranean	Long lines		Mortality	No change suggested
Minami et al. (2007)	Modeling	Supply information	Large scale	North Pacific	Purse-seine	Large osteicht		No change suggested
Morgan et al. (2009)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Long lines	Large elasmobranch	Assemblages	No change suggested
Musyl et al. (2009)	Observational	Supply information	Large scale	North Atlantic	Long lines	Large osteicht	Mortality	No change suggested
O'Connell et al. (2011)	Field experiment	Reduce capture	Large scale	North Atlantic	Multiple types		Sub-lethal: behaviour	Repelets
Ordines et al. (2006)	Field experiment	Reduce capture	Artisanal	Mediterranean	Trawling nets	Polyspecific	Population	Type/Size of nets
Pajuelo et al. (2010)	Observational	Fleet bycatch analysis	Artisanal	Mediterranean	Long lines	Small osteicht	Population	No change suggested
Perez and Wahrlich (2005)	Observational	Fleet bycatch analysis	Large scale	South Atlantic	Gill nets	Small osteicht	Assemblages	No change suggested
Petersen et al. (2009)	Observational	Fleet bycatch analysis	Large scale	South Atlantic	Long lines	Large osteicht	Population	No change suggested
Philippart (1998)	Modeling	Fleet bycatch analysis	Large scale	North Atlantic	Trawling nets		Population	No change suggested
Poisson et al. (2009)	Observational	Fleet bycatch analysis	Large scale	North Pacific	Long lines	Large osteicht	Sub-lethal: injury	Fish handling
Polovina et al. (2002)	Observational	Supply information	Large scale	North Pacific	Long lines	Large osteicht	Assemblages	No change suggested
Punt et al. (2006)	Modeling	Supply information	Large scale	South Pacific	Long lines	Polyspecific	Population	No change suggested
Revill et al. (2005)	Field experiment	Fleet bycatch analysis	Large scale	North Atlantic	Trawling nets	Polyspecific	Mortality	No change suggested
Richards (1994)	Review	Fleet bycatch analysis	Large scale	South Pacific	Driftnets	Polyspecific	Mortality	No change suggested
Robbins et al. (2011)	Field experiment	Reduce capture	Artisanal	South Pacific	Hook and line	Polyspecific	Sub-lethal: behaviour	Repelets
Rodríguez-Cabello et al. (2005)	Observational	Species bycatch analysis	Large scale	North Atlantic	Trawling nets	Polyspecific	Mortality	No change suggested
Rodríguez-Cabello et al. (2005)	Observational	Supply information	Large scale	North Atlantic	Long lines	Large osteicht	Mortality	No change suggested
Rogan and Mackey (2007)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Driftnets	Polyspecific	Mortality	No change suggested
Romanov (1999)	Observational	Fleet bycatch analysis	Large scale	Indian	Purse-seine	Large osteicht	Assemblages	No change suggested

Table 1 continued

Author (year)	Article type	Article objective	Scale of the fishing fleet	Location of the fishery	Fishing gear used	Target species	Scale of the consequences	Proposed changes
Romine et al. (2009)	Modeling	Species bycatch analysis	Large scale	North Atlantic	Long lines	Polyspecific	Mortality	Fishing closure
Russell (1993)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Long lines	Large oseticht	Population	No change suggested
Samaranyanka et al. (1997)	Field experiment	Fleet bycatch analysis	Artisanal	Indian	Driftnets	Large oseticht	Population	No change suggested
Schindler et al. (2002)	Modeling	Supply information	Large scale	North Pacific	Long lines	Large oseticht	Population	No change suggested
Shepherd and Myers (2005)	Field experiment	Supply information	Large scale	North Atlantic	Multiple types	Polyspecific	Assemblages	No change suggested
Skomal (2007)	Review	Supply information	Large scale	North Atlantic	Long lines	Large oseticht	Sub-lethal: physiological	No change suggested
Stobutzki et al. (2001)	Observational	Fleet bycatch analysis	Large scale	Indian	Trawling nets	Shrimp	Assemblages	Multiple
Stobutzki et al. (2002)	Observational	Fleet bycatch analysis	Large scale	Indian	Trawling nets	Shrimp	Assemblages	No change suggested
Stoner and Kaimmer (2008)	Laboratory experiment	Reduce capture	Large scale	North Atlantic	Long lines	Large oseticht	Sub-lethal: physiological	Repelets
Storai et al. (2011)	Observational	Fleet bycatch analysis	Artisanal	Mediterranean	Multiple types	Large oseticht	Assemblages	No change suggested
Swimmer et al. (2011)	Field experiment	Reduce capture	Large scale	North Pacific	Long lines	Large oseticht	Assemblages	Type/Size of hooks
Thorpe and Frierson (2009)	Field experiment	Reduce capture	Artisanal	North Atlantic	Gill nets	Small oseticht	Assemblages	Fishing practice
Trent et al. (1997)	Observational	Fleet bycatch analysis	Large scale	North Atlantic	Driftnets	Small elasmobranch	Assemblages	No change suggested
Valenzuela et al. (2008)	Observational	Supply information	Artisanal	South Pacific	Long lines	Polyspecific	Population	No change suggested
Vega and Licandeo (2009)	Field experiment	Fleet bycatch analysis	Large scale	South Pacific	Long lines	Large oseticht	Assemblages	Fishing practice
Walker et al. (2005)	Field experiment	Fleet bycatch analysis	Large scale	South Pacific	Multiple types	Small elasmobranch	Assemblages	No change suggested
Ward and Hindmarsh (2007)	Review	Fleet bycatch analysis	Large scale	North Pacific	Long lines	Large oseticht	Assemblages	No change suggested
Ward et al. (2008)	Field experiment	Reduce capture	Large scale	South Pacific	Long lines	Large oseticht	Population	Type/Size of hooks
Ward et al. (2009)	Field experiment	Reduce capture	Large scale	South Pacific	Long lines	Large oseticht	Population	Type/Size of hooks
Watson et al. (2008)	Modeling	Supply management strategies	Large scale	North Pacific	Purse-seine	Large oseticht	Population	Fishing closure
Whoriskey et al. (2011)	Observational	Fleet bycatch analysis	Large scale	North Pacific	Long lines	Small oseticht	Assemblages	No change suggested
Yamandú et al. (1998)	Observational	Fleet bycatch analysis	Large scale	South Atlantic	Long lines	Large oseticht	Population	No change suggested
Yokota et al. (2006)	Field experiment	Reduce capture	Large scale	North Pacific	Long lines	Large oseticht	Population	Type/Size of hooks
Zeeberg et al. (2006)	Field experiment	Reduce mortality	Large scale	North Atlantic	Trawling nets	Small oseticht	Mortality	Type/Size of nets
Zhou et al. (2011)	Modeling	Supply information	Large scale	Multiple	Multiple types	Polyspecific	Assemblages	No change suggested
Zollett (2009)	Observational	Supply information	Large scale	North Atlantic	Multiple types	Large oseticht	Assemblages	No change suggested

We found several temporal trends in the literature reviewed. In general, the early studies tended to focus on trawl fisheries, while more recent studies have focused on long line fisheries. Moreover, early studies tended to not explore reduction strategies, while nearly all current studies explore potential solutions to bycatch rather than just emphasizing the problems. Early studies mainly analyzed levels and composition of bycatch (i.e., observational studies aboard fishing vessels) while recent studies have been more diverse focus, with modeling, laboratory, and field experiments. Also in recent years, a variety of biological endpoints have been studied. Early literature focused exclusively on population-level parameters or mortality consequences whereas recent papers include sublethal endpoints such as physiological stress (e.g.,

Mandelman and Farrington 2007), behaviour (e.g., Brill et al. 2008), and injury (e.g., Poisson et al. 2009; Campana et al. 2009a) (Fig. 1a).

Published research on shark bycatch that has been conducted in the North Atlantic Ocean represents almost 40 % of the total articles analysed, while the North Pacific Ocean accounted for 14.4 %. Studies that involved multiple locations comprised 14.4 % of papers. FAO (food and agricultural organization of the United Nations) statistics as presented by Barker and Schluessel (2005) show that most elasmobranch catches occur in developed countries of the Indian (10.6 %) and Pacific Oceans (South Pacific Ocean comprised 10.6 % of the studies). Interestingly, comparatively few peer reviewed studies dealing with shark bycatch the Indo-Pacific region were found.




Fig. 1 Number of articles for **a** each year, **b** each article type, **c** each fishing gear by article objective (□ Supply management strategies, □ Supply information, □ Species bycatch analysis, □ Reduce mortality, ■ Reduce capture, ■ Fleet bycatch analysis) and **d** for each fishing gear by suggested bycatch reduction strategies (□ Multiple types, □ Type/Size of nets, □ Type/Size of

hooks, ■ Repelents, ■ Fishing practice, ■ Fishing closure, ■ No change suggested). *Note:* For better interpretation, Trammel nets and hook and line gear, and Fishing time and fish handling bycatch reduction strategies were excluded since they only comprised one article each

Literature collected for South Atlantic and Mediterranean Oceans was scarce; 4.8 and 6.7 % respectively.

Studies of shark bycatch tended to focus on either one species (e.g., Campana et al. 2009a, b) or on many species (e.g., Petersen et al. 2009; Morgan et al. 2009; Godin and Worm 2010; described as many as 25, 26 and 28 shark bycatch species, respectively). Because of the great disparity in the number of species studied, measures of central tendency (i.e., the average was 6.13 ± 6.61 species) fail to characterize the pattern appropriately. The modal number of species was 1, accounting for 25 % of all studies.

Most studies found dealt with the analysis and description of shark bycatch in a particular fishery (38.4 % of the total studies). Observational studies where “observers” monitored and recorded shark bycatch information that was subsequently synthesized in the context of a real fishery (on occasion as part of government research fishery activities) comprised a substantial amount of the published shark bycatch research (28.4 % of the total studies; Fig. 1b). In general, observational studies predominated (43.3 %) although a fair amount of field experiments (23.1 %) and reviews (19.2 %) were also conducted.

There was a particular focus on large-scale long line fisheries for tunas and swordfish (44.8 % of the total studies) seconded by multispecies fisheries (31.2 %). Other types of fisheries represented less than a 10 % of the total studies. The focus on the tuna fishery can be explained by the importance of tuna landings, being the third most landed fish worldwide, with values over six million tons per year (FAO 2008).

Long line fisheries reviews and studies accounted for 44.6 % of the studies analysed, of which 34.7 % were aimed to reduce bycatch. Trawling fisheries were the focus of 16.5 % of the studies examined of which 29.4 % were aimed at reducing bycatch (Fig. 1c). Trawling, specially bottom trawling, represents one of the most destructive fishing methods, as it not only represents the bulk of the near 30 million tons of bycatch discarded annually, it also destroys bottom structures and communities, which have key roles in the recruiting of young fish (Pauly 2002). Cook (2003) found that trawl fisheries, in particular shrimp trawl fisheries (3 million tons per year; FAO 2008), have high bycatch rates compared to long line fisheries. Hill and Wassenberg (2000) found that 98 % of the total fish discarded from trawling vessels were dead. This is of interest because research by Poisson et al. (2009)

shows that the mortality of sharks (at haul-back—no information on delayed mortality) in long line fisheries for tunas and swordfish is relatively low for the species they analyzed. Mandelman et al. (2008) and Beerkircher et al. (2000) also conducted mortality assessments on longline fisheries in the USA, and found only three species of sharks (namely *Carcharhinus signatus*, *C. falciformis* and *Sphyrna lewini*) of the ten species assessed to have mortality rates over 50 %. Our results suggest that studies dealing with shark bycatch are often focused on tuna and swordfish longline fisheries. Although relevant given the apparently variable at-vessel mortality among shark species for longline fisheries and the fact that pelagic longline fisheries are the dominant mode of capture for sharks globally, trawl fisheries deserve future attention given the high bycatch and mortality rates, reduction in recruit biomass and habitat loss associated with that gear. Mortality rates on trawling vessels for some species of sharks seems to be lower when tow duration remains lower than 30 min, and fish are returned to the sea in less than 30 min (Rodríguez-Cabello et al. 2005).

Literature reviewed tended to focus on large-scale industrial fisheries (86.1 % of the total studies), while artisanal fisheries were less frequently studied (13.9 % of the total studies). At this point it is unclear if that pattern signifies there are fewer incidences of shark bycatch for small-scale artisanal fisheries or if it simply reflects the fact that industrial fisheries tend to operate out of developed countries where governance structures require bycatch reduction research and research capacity exists to conduct such studies. Considering the study by Pauly (2002) the case seems to be the former.

Trends in research on strategies for reducing/mitigating bycatch

Bycatch reduction strategies were suggested in 41.6 % of the reviewed articles. Net size (e.g., Sobrino et al. 2000) and mesh design (e.g., Broadhurst et al. 2002; Ordines et al. 2006) were the most frequent change suggested for trawl fisheries (66.7 % for studies on this fishing gear). For longline fisheries, hook size and type (e.g., circle hooks; Kaplan et al. 2007; Ward et al. 2009) were the most frequently suggested change (45.4 % for studies on this fishing gear; Fig. 1d), but some studies also noted an important role of bait type (e.g., Watson et al. 2005). Historically, gear selectivity

has been an issue for fisheries managers who attempt to minimize the capture of undersized target species, providing escapement opportunities in the gear design (Armstrong et al. 1990; Dawson 1991; Millar 1992). This issue has been addressed by many authors (MacLennan 1995; Sobrino et al. 2000; Smith and Jeremy 2005), and different solutions exist. However, reducing bycatch is a more complex problem, since size is not the only variable to be taken into account. For instance, sharks taken as bycatch are often very different in shape and often several times larger than target species. This has been observed in the New Zealand commercial trawl fishery (Francis and Duffy 2002). Design changes, such as the ones proposed by Brewer et al. (1998), Broadhurst et al. (2002), Zeeberg et al. (2006) and Chosid et al. (2012) for trawl nets which recognize the differences between target species and bycatch such as sharks could be effective for reducing shark bycatch. In general, there is little work on strategies to reduce the high mortality rates associated with trawl nets caused by fish packing in the net and fish handling once on board (Frick et al. 2010).

Regulations on fishing zones (i.e., area closures/restrictions) and the timing (i.e., temporal closures/restrictions) of commercial fishing activities were only reported in 9.9 % of the studies that we reviewed (e.g., Beerkircher et al. 2000; Watson et al. 2008; Romine et al. 2009). Regulations involving closure of fishing zones in combination with regulated fishing times are common practices used to reduce bycatch levels (Alverson et al. 1994). In this type of scenario, fishing fleets are allowed to operate only in a certain area at a certain time, in order to avoid high concentrations of bycatch species, or to encounter high concentration of target species. However, information about the seasonal and diel variation in fish distribution (both of target and bycatch species) is needed for this strategy to be implemented properly, which is not the case for many fisheries. Hoffmayer et al. (2012) studied the seasonal variation in stress response of *Rhizoprionodon terraenovae* and found summer to be the most vulnerable season for this species. This type of information combined with regulations on fishing zones and timing might prove an effective bycatch reduction strategy. However, basic information on movement and spatial ecology is lacking for all but a few species making use of area and time closures challenging and ineffective.

Repellent agents (including electricity, magnets and surfactants) have been used worldwide to protect bathing beaches, divers, and aquaculture gear from shark attack (e.g., Kalmijn 1982; Sisneros and Donald 2001). There is also potential for such deterrent devices to be used on fishing gear to reduce shark bycatch (discussed in Gilman et al. 2007), however, there remain challenges with the implementation of such tools. Many repellents that involve electricity or magnets are large, expensive, and not practical for wide-scale use in fisheries (Frick et al. 2010). As part of the quantitative literature review we only found 5 papers (4.9 % of total studies analysed) focused on exploring the potential use of rare-earth (lanthanide) metals and alloys and permanent magnets that could be incorporated into hooks to repel sharks (Brill et al. 2008; Kaimmer and Stoner 2008; Jordan et al. 2011; O'Connell et al. 2011; Robbins et al. 2011). However, we found additional 2 papers outside of the structured literature review (e.g., Stoner and Kaimmer 2008; Tallack and Mandelman 2009) emphasizing that some of those studies do not use keywords or title elements related to bycatch so were missed by our structured search. The results from most of the studies on rare-earth (lanthanide) metals and alloys are promising, although the use of this bycatch reducing strategy is still rather undeveloped, and further studies on the subject needed to assess its applicability in large-scale commercial fisheries. Use of chemical-based repellents also has promise but it is necessary to understand how such repellents influence target species.

Research needs and opportunities

Our synthesis revealed a number of limitations with current knowledge on shark bycatch which leads to a number of research needs (i.e., in the context of management and conservation) and opportunities. Below we provide a list of what we regard as critical research needs as well as a rationale and discussion of how these research needs could be addressed.

Need for studies that combine approaches (e.g., field, lab and modeling)

In our survey we found few articles combining laboratory and field experiments with modeling studies on sharks. Combined approaches have led to

important findings in the field of bycatch assessment. For example, Goldsworthy and Page (2007) identified important mitigation opportunities to reduce pinniped bycatch produced by Australian fisheries by performing a risk assessment using data from different realms. Also working with a marine mammal, the harbour porpoise (*Phocoena phocoena*), Orphanides (2010) revealed that bycatch estimates were more stable when using a combined approach, than using only the ratio estimate approach. Of particular benefit would be the ability to scale experimental laboratory and field research to the ecosystem level using modeling techniques. There is also much benefit from combining controlled laboratory experiments with field observations whereby mechanisms and detailed aspects can be evaluated in the laboratory and then the innovations (e.g., say in a gear modification) can be tested in the field. Not surprisingly, given the size of sharks there have been relatively few laboratory studies on fisheries interactions but there are a number of large research aquaria around the world where studies could occur. It is also possible to use field mesocosms such as large pen enclosures as has been used to study lemon shark energetics at Bimini, The Bahamas (Shepard et al. 2008) or adult whitetip reef sharks (*Triaenodon obesus*) in a lagoon in Kaneohe Bay at the Hawaii Institute of Marine Biology (Whitney et al. 2007). In such facilities captive sharks can be studied to examine the long-term consequences of fisheries interactions including healing of injuries, delayed mortality, and to generate stress profiles. Moreover, it would be possible to design experiments to evaluate post-release behaviour (and physiology using devices like heart rate loggers) relative to different types of fisheries interactions.

Need for experimental studies on shark repellents or other aspects of sensory physiology relevant to shark bycatch reduction

The Ampullae of Lorenzini in sharks is a unique organ that is used for electroreception. This organ is not found on bony fishes; therefore, a repellent that affects this sensory organ of sharks would be selective. Permanent magnets have been shown to be deterrent agents for sharks, and do not require power input, making them potentially ideal for use in fisheries and as bycatch reduction devices (Rigg et al. 2009). Other rare-earth (lanthanide) metals and alloys are also being

investigated. However, few studies outside of strictly laboratory experiments exist that assess the applicability of this deterrent agent to reduce bycatch. Chemical agents have also been studied, although their effectiveness is limited (Sisneros and Donald 2001). An effective repellent for sharks with applicability on large-scale fishing fleets would be a key tool for the conservation of sharks worldwide. Research that combines laboratory and field experiments, along with physiological studies on shark behaviour is needed to develop such a tool. As recently done for sea turtles (reviewed in Southwood et al. 2008), linking sensory physiology with bycatch research for sharks could produce interesting results and help to advance the field.

Need for studies that explore the handling component of bycatch

Studies dealing with the handling component of bycatch were poorly represented in the shark bycatch literature. In the recreational fisheries literature (e.g., Cooke and Suski 2005; Arlinghaus et al. 2007) and bycatch work on other non-shark fisheries (see Davis 2002), there is a strong recognition that how animals are handled by fishers has an influence on their eventual fate. Indeed, appropriate handling of specimens onboard is vital for some species' survivability, although other species, like the lesser spotted dogfish (*Scyliorhinus canicula*), shows great resilience to onboard handling (Revill et al. 2005). Although there is benefit in preventing bycatch in the first place, the reality is that it does occur and it is unrealistic to assume that it can be totally eliminated so fishers will continue to play an important role in the handling and release of sharks. Ideally, research efforts would inform fishers on proper handling techniques (e.g., hook removal, when to cut the line, air exposure thresholds) for minimizing post release mortality. An example of tow duration, air exposure and crowding effects on mortality can be found in Frick et al. (2010). There is undoubtedly much variation among gear types and species so there is much opportunity for research on this topic. In the recreational catch-and-release literature, Cooke and Suski (2005) called for species-specific research on how fish respond to fisheries interactions given the inter-specific variation in size, dentition, feeding mode, environmental tolerances, physiological capacity, etc. that exists.

Also relevant here is the growing interest in fish welfare (Diggle et al. 2011) and the recognition that the actions (or inactions) of fishers influence the outcome of the event for discarded sharks. Hyatta et al. (2012) is a good and recent example, providing handling suggestions to reduce the level of stress in three species of sharks. The provision of science-based best handling practices would go a long way towards improving welfare status of discarded sharks as has occurred in the recreational sector for a variety of gamefish that are angled and released (Arlinghaus et al. 2007).

Need to examine the condition and fate of sharks that are discarded

To date, very few studies have addressed post-release mortality of sharks that are discarded from fisheries. Indeed, most mortality estimates are based solely on quantifying initial mortality (i.e., mortality at time of haul-back/landing) and assume that discards survive. Clearly not all sharks survive upon release. Not incorporating the post release mortality component in bycatch assessments risks an underestimation of real mortality produced by discards (Coggins et al. 2007) and thus undermines conservation and management actions. Evaluating post-release mortality can be done by holding sharks in cages, pens, or tanks or alternatively by releasing sharks with electronic tags. Donaldson et al. (2008) reviewed the various ways in which electronic devices such as pop-up satellite tags (PSAT) or acoustic transmitters can be used to assess post release mortality in a recreational catch-and-release context and those techniques could readily be applied to shark bycatch studies. Although published after we conducted our quantitative review and thus excluded from formal analysis, a recent paper by Musyl et al. (2011) provided an example of where PSAT tags were used to document post release mortality of five species of pelagic sharks. A challenge with shark work will be obtaining control fish but that can be overcome by contrasting different gear types and methods (e.g., compare mortality using hook timers to identify sharks on long lines for brief vs long duration periods; Brooks et al. (2012) used hook timers to evaluate various levels of physiological disturbance for Caribbean reef sharks (*Carcharhinus perezi*) captured on experimental long-lines). Documenting mortality post-release could identify the

mechanisms of mortality including physiological exhaustion, metabolic scope and capacity, environmental correlates (e.g., water temperature, water depth), severe injury, and when one or several of those mechanisms facilitate post-release predation by other sharks and animals (Skomal and Bernal 2010).

Beyond mortality, there are also a range of sublethal consequences associated with injury and stress of capture. For example, sharks could experience behavioural impairments post release which could influence their ability to find prey or avoid predators. Physiological disturbances in the short term could restrict exercise capacity and thus ability to escape and in the long term could contribute to problems with immune function and disease (reviewed in Skomal and Mandelman 2012). Presence of hooks in fish that are released could also lead to a variety of pathogenic problems as has been documented in the recreational sectors (Borucinska et al. 2002). There are a number of studies that we documented that compared the physiological consequences of different types of capture stressors or on multiple species (e.g., Mandelman and Skomal 2009; note—not a reference that was part of the quantitative review) but none of these studies evaluated the longer term consequences of those physiological disturbances (but see Frick et al. 2012). A suite of tools (such as field physiology techniques, molecular markers, oxidative stress measures) now exist that could be applied to studies of sharks to advance the study of sublethal effects and a better understanding of the factors that contribute to mortality (Gallagher et al. 2010; Hyatta et al. 2012; Marshall et al. 2012; Renshaw et al. 2012; Skomal and Mandelman 2012). In fact, a recent special issue of the journal *Comparative Biochemistry and Physiology A* contains numerous papers on sublethal aspects of shark-fisheries interactions (e.g., Brooks et al. 2012; Frick et al. 2012; Renshaw et al. 2012; Skomal and Mandelman 2012) emphasizing the opportunity in that realm. Injury produced by the fishing gear is a factor that has been better documented in the reviewed literature; however, only 11 (10.9 %) studies dealt with sublethal effects on shark bycatch so there is a pressing need for more research on this topic. Relatedly, there is also a need to develop and validate tools to predict mortality of released sharks (Renshaw et al. 2012) which may involve using novel techniques such as reflex indicators (Davis 2010). Moyes et al. (2006) provided the first attempt to do so for sharks.

Knowledge of the factors that lead to mortality can be used to identify thresholds for stress and injury and to estimate post-release mortality in the absence of actual post-release mortality studies. The conservation physiology toolbox (Wikelski and Cooke 2006) has the potential to inform shark bycatch research and would best do so if able to establish relationships between stress and relevant population-level endpoints (Cooke and O'Connor 2010).

Need for research that covers more regions, fisheries (gear types), and species

As revealed by our review, there are many research gaps with respect to regions, gear types and species. For example, from a regional perspective, the Indian, South Atlantic and both north and south Pacific oceans have seen comparatively little research activity on shark bycatch relative to the north Atlantic. This is particularly problematic as there is intense and growing fishing pressure in these regions including industrial shrimp trawling, squid driftnet, and coastal varied bottom trawling fisheries, all of which could result in shark bycatch. With respect to species, shark bycatch has been thoroughly studied for several charismatic species such as white, basking and blue sharks, species which distributional ranges include developed countries (mainly North Atlantic Ocean), or species for which the accidental catch affects the commercial profit of certain fisheries (i.e., spiny dogfish, *Squalus acanthias*). Many commercially unimportant shark species (most of the species of the order Hexanchiformes and Orectolobiformes) remain unstudied despite being captured as bycatch. While the importance of charismatic umbrella species is undeniable for incentivizing conservation efforts that help protect less charismatic species, specific research on these species is needed to understand how to reduce mortality arising from bycatch.

Large scale fisheries were the focus of most of the literature reviewed (>80 % of the articles). Although it is assumed that these are the ones that produce the largest amount of bycatch, small-scale fisheries can also yield a significant amount of bycatch (Thorpe and Frierson 2009; Alfaro-Shigueto et al. 2010; Baeta et al. 2010). Among large scale fisheries, several authors have described and highlighted the importance of trawl fisheries bycatch on shark populations, pinpointing it as an issue for shark conservation worldwide

(e.g., Frick et al. 2010; Hill and Wassenberg 2000). Providing adequate management strategies, incentivizing research on shark bycatch, and publishing of the results in peer reviewed journals would help to identify in which region shark bycatch predominates, which species are more affected, and why, and what type of fishery contributes mainly to the discards.

Need for human dimensions studies related to shark bycatch

Bycatch of sharks and bycatch in general, has important social, economic, and cultural components. We have addressed so far the more technical problems and up to date information on the ecological, biological and physiological impacts for sharks from bycatch. We have also explored the range of bycatch reduction technologies for sharks. However, the existence of a good technological means to reduce bycatch does not mean that it can be implemented on fisheries worldwide, even if it technically possible. The human dimension such as fisher adoption of gears is critical to their success (Gilman et al. 2007; Campbell and Cornwell 2008). Moreover, fishers themselves can be the source of innovations in bycatch reduction. Engaging them in the process of bycatch reduction can actually yield practical solutions that may not have been considered by managers or researchers. In addition, the development of management plans for conservation of bycatch species such as sharks face particular issues in the social, cultural, and economic realms. Fishers might not be prone to adopt a given BRD (Bycatch Reduction Device) if they are not convinced that the bycatch species are really threatened. They also may resent resources spent in the conservation of what they may regard as uncharismatic species (i.e., in their minds, sharks; See Simpfendorfer et al. 2011). Implementing BRD might also put the fishers at an economic disadvantage, if the adoption of that BRD is not shared internationally. Defence of traditional fishing practices, fear of losing independence, and resentment of the marginalization of their activity are all cultural factors influencing the acceptance or rejection of a particular BRD. When addressing these variables, fisheries management agencies often make assumptions to fill in the void of information and data regarding these important components of bycatch. Recent efforts to incorporate human dimensions research into bycatch reduction

have been encouraged (Campbell and Cornwell 2008) but there remain few examples in the literature. There is much opportunity in this realm for research on the human dimensions of shark bycatch.

Conclusion

Bycatch in general is a serious threat for biodiversity worldwide, and sharks seem to be particularly vulnerable to bycatch. Indeed, shark bycatch threatens to wipe out these top predators that structure many marine habitats, with possible catastrophic consequences for ecosystems and the services that they provide. The fact that shark bycatch is regarded as the primary threat facing so many of the imperiled shark species provides direction to those interested in or responsible for their conservation and management. We have described the characteristics and trends in the published research on shark bycatch, and conclude that there are a number of knowledge gaps that make it difficult to reduce shark bycatch or to understand the consequences of that activity. New technology offers potential solutions to reduce bycatch, but studies concerning them are scarce. Also, most research is focused on the impact of large industrial-scale fisheries, which undoubtedly contribute in great measure to global bycatch of shark, but failure to consider the impact of artisanal fisheries would be unwise. Our study has identified what we believe to be the most pressing research needs that must be addressed including the need for multidisciplinary studies that combine different approaches, development and testing of bycatch reduction devices, understanding the fate of discarded sharks, expanding research coverage to other regions, fisheries, and gear types (including more work on the recreational sector which we did not cover here), and finally incorporating human dimensions science into the study of shark bycatch reduction. Although there are some inherent challenges with developing and testing shark bycatch reduction strategies, there is an urgent need to do so and this would be best achieved through interdisciplinary research that spans field, laboratory, and modeling realms. It is our desire that this review stimulates research activities that will improve our understanding of shark bycatch, but more importantly yields practical advances related to bycatch reduction and improving the survival and welfare of those sharks that are captured and

discarded, that can be readily adopted by the fishing community and/or incorporated into regulations and policies.

Acknowledgments Molina was supported by the Emerging Leaders in the Americas Program during a visit to Carleton University in 2010. Cooke was supported by the Canada Research Chairs Program and the Ontario Ministry of Research and Innovation. We thank members of the Cooke Lab, Bs. M.E. Croce, Dr. A.C. Lopez Cazorla, and two anonymous referees for critiques on the paper.

References

- Afonso AS, Hazin FHV, Carvalho F, Pacheco JC, Hazin H, Kerstetter DW, Murie D, Burgess GH (2011) Fishing gear modifications to reduce elasmobranch mortality in pelagic and bottom longline fisheries off Northeast Brazil. *Fish Res* 108:336–343
- Alfaro-Shigueto J, Mangel JC, Pajuelo M, Dutton PH, Seminoff JA, Godley BJ (2010) Where small can have a large impact: structure and characterization of small-scale fisheries in Peru. *Fish Res* 106:8–17
- Alverson D, Freeberg M, Murawski S, Pope J (1994) A global assessment of fisheries bycatch and discards. United Nations Food and Agriculture Organization Fisheries Technical Paper 339
- Arlinghaus R, Cooke SJ, Schwab A, Cowx IG (2007) Fish welfare: a challenge to the feelings-based approach, with implications for recreational fishing. *Fish Fish* 8:57–71
- Armstrong DW, Ferro RST, MacLennan DN, Reeves SA (1990) Gear selectivity and conservation of fish. *J Fish Biol* 37: 261–262
- Arocha F, Arocha O, Marcano LA (2002) Observed shark bycatch from the Venezuelan tuna and swordfish fishery from 1994 through 2000. *Collect Vol Sci Papers ICCAT* 54:1123–1131
- Baeta F, Batista M, Maia A, Costa MJ, Cabral H (2010) Elasmobranch bycatch in a trammel net fishery in the Portuguese west coast. *Fish Res* 102:123–129
- Barker MJ, Schluessel V (2005) Managing global shark fisheries: suggestions for prioritizing management strategies. *Aquat Conserv* 15:325–347
- Beerkircher LR, Cortés E, Shivji M (2000) Characteristics of shark bycatch observed on pelagic longlines off the southeastern United States, 1992–2000. *Mar Fish Rev* 64:40–49
- Belcher CN, Jennings CA (2011) Identification and evaluation of shark bycatch in Georgia's commercial shrimp trawl fishery with implications for management. *Fish Manage Ecol* 18:104–112
- Benjamins S, Kulka D, Lawson J (2010) Recent incidental catch of sharks in gillnet fisheries of Newfoundland and Labrador, Canada. *Endanger Species Res* 11:133–146
- Blaber SJM, Dichmont CM, White W, Buckworth R, Sadiyah L, Iskandar B, Nurhakim S, Pillans R, Andamari R (2009) Elasmobranchs in southern Indonesian fisheries: the fisheries, the status of the stocks and management options. *Rev Fish Biol Fisheries* 19:367–391

- Bonfil R (1995) World bycatches of sharks in high-seas fisheries: appraising the waste of a resource. Fisheries Centre Research Reports 2
- Borucinska J, Kohler N, Natanson L, Skomal G (2002) Pathology associated with retained fishing hooks in blue sharks, *Prionace glauca* (L.), with implications for their conservation. J Fish Dis 25:515–521
- Brewer D, Rawlinson N, Eayrs S, Burrige C (1998) An assessment of bycatch reduction devices in a tropical Australian prawn trawl fishery. Fish Res 36:195–215
- Brill R, Bushnell P, Smith L, Speaks C, Sundaram R, Stroud E, Wang J (2008) The repulsive and feeding-deterrent effects of electropositive metals on juvenile sandbar sharks (*Carcharhinus plumbeus*). Fish Bull 107:298–307
- Broadhurst M, Kangas M, Damiano C, Bickford S, Kennelly S (2002) Using composite square-mesh panels and the Nordmøre-grid to reduce bycatch in the Shark Bay prawn-trawl fishery, Western Australia. Fish Res 58:349–365
- Brooks EJ, Mandelman JW, Sloman KA, Liss S, Danylchuk AJ, Cooke SJ, Skomal GB, Philipp DP, Sims DW, Suski CD (2012) The physiological response of the Caribbean reef shark (*Carcharhinus perezi*) to longline capture. Comp Biochem Physiol A 162(2):94–100
- Burgess GH, Beerkircher LR, Cailliet GM, Carlson JK, Cortés E, Goldman KJ, Grubbs RD, Musick JA, Musyl MK, Simpfendorfer CA (2005) Is the collapse of shark populations in the Northwest Atlantic Ocean and Gulf of Mexico real? Fisheries 30:19–26
- Camhi MD, Valenti SV, Fordham SV, Fowler SL, Gibson C (eds) (2007) The conservation status of pelagic sharks and rays. IUCN Species Survival Commission's Shark Specialist Group
- Campana SE, Joyce W, Manning MJ (2009a) Bycatch and discard mortality in commercially caught blue sharks *Prionace glauca* assessed using archival satellite pop-up tags. Mar Ecol Prog Ser 387:241–253
- Campana SE, Warren J, Malcom PF, Manning MJ (2009b) Comparability of blue shark mortality estimates for the Atlantic and Pacific longline fisheries. Mar Ecol Prog Ser 396:161–164
- Campbell L, Cornwell M (2008) Human dimensions of bycatch reduction technology: current assumptions and directions for future research. Endanger Species Res 5:325–334
- Carbonell A (2003) The by-catch of sharks in the western Mediterranean (Balearic Islands) trawl fishery. Fish Res 61:7–18
- Carruthers EH, Schneider DC, Neilson JD (2009) Estimating the odds of survival and identifying mitigation opportunities for common bycatch in pelagic longline fisheries. Biol Conserv 142:2620–2630
- Carruthers EH, Neilson JD, Smith SC (2011) Overlooked by-catch mitigation opportunities in pelagic longline fisheries: soak time and temperature effects on swordfish (*Xiphias gladius*) and blue shark (*Prionace glauca*) catch. Fish Res 108:112–120
- Cartamil D, Santana-Morales O, Escobedo-Olvera M, Kacev D, Castillo-Geniz L, Graham JB, Rubin RD, Sosa-Nishizaki O (2011) The artisanal elasmobranch fishery of the Pacific coast of Baja California, Mexico. Fish Res 108:393–403
- Chosid DM, Pol M, Szymanski M, Mirarchi F, Mirarchi A (2012) Development and observations of a spiny dogfish *Squalus acanthias* reduction device in a raised footrope silver hake *Merluccius bilinearis* trawl. Fish Res 114:66–75
- Clarke M (2002) Catch, discarding, age estimation, growth and maturity of the squalid shark *Deania calceus* west and north of Ireland. Fish Res 56:139–153
- Coelho R, Erzini K (2008) Effects of fishing methods on deep water shark species caught as by-catch off southern Portugal. Hydrobiologia 606:187–193
- Coelho R, Bentes L, Goncalves JM, Lino PG, Ribeiro J, Erzini K (2003) Reduction of elasmobranch by-catch in the hake semipelagic near-bottom longline fishery in the Algarve (Southern Portugal). Fish Sci 69:293–299
- Coggins LG, Catalano MJ, Allen MS, Pine WE, Walters CJ (2007) Effects of cryptic mortality and the hidden costs of using length limits in fishery management. Fish Fish 8:196–210
- Connolly PL, Kelly CJ (1996) Catch and discards from experimental trawl and longline fishing in the deep water of the Rockall Trough. J Fish Biol 49:132–144
- Cook R (2003) The magnitude and impact of by-catch mortality by fishing gear. In: Valdimarsson G, Sinclair M (eds) Responsible fisheries in the marine ecosystem. FAO, Rome
- Cooke SJ, O'Connor CM (2010) Making conservation physiology relevant to policy makers and conservation practitioners. Conserv Lett 3:159–166
- Cooke SJ, Suski CD (2005) Do we need species-specific guidelines for catch-and-release recreational angling to effectively conserve diverse fishery resources? Biodivers Conserv 14:1195–1209
- Cortés E, Arocha F, Beerkircher L, Carvalho F, Domingo A, Heupel M, Holtzhausen H, Santos MN, Ribera M, Simpfendorfer C (2009) Ecological risk assessment of pelagic sharks caught in Atlantic pelagic longline fisheries. Aquat Living Resour 23:25–34
- Cramer J, Bertolino A, Scott GP (1998) Estimates of recent shark bycatch by U.S. vessels fishing for Atlantic tuna and tuna-like species. Collect Vol Sci Papers ICCAT 48:117–128
- Crowder LB, Murawski SA (1998) Fisheries bycatch: implications for management. Fisheries 23:8–17
- Curran D, Bigelow K (2011) Effects of circle hooks on pelagic catches in the Hawaii-based tuna longline fishery. Fish Res 109:265–275
- Davis MW (2002) Key principles for understanding fish bycatch discard mortality. Can J Fish Aquat Sci 59:1834–1843
- Davis MW (2010) Fish stress and mortality can be predicted using reflex impairment. Fish Fish 11:1–11
- Dawson SM (1991) Modifying gillnets to reduce entanglement of cetaceans. Mar Mamm Sci 7:274–282
- de Silva JA, Condrey RE, Thompson BA (2001) Profile of shark bycatch in the U.S. Gulf of Mexico menhaden fishery. North Am J Fish Manag 21:111–124
- Diaz G, Serafy JE (2005) Longline-caught blue shark (*Prionace glauca*): factors affecting the numbers available for live release. Fish Bull 103:720–724
- Diggles BK, Cooke SJ, Rose JD, Sawynok W (2011) Ecology and welfare of aquatic animals in wild capture fisheries. Rev Fish Biol Fisheries 21:739–765
- Donaldson MR, Arlinghaus R, Hanson KC, Cooke SJ (2008) Enhancing catch-and-release science with biotelemetry. Fish Fish 9:79–105

- Dulvy KN, Baum JK, Clarke S, Compagno LJV, Cortes E, Domingo A, Fordham S, Flower S, Francis M, Gibson C, Martinez J, Musick JA, Soldo A, Stevens JD, Valenti S (2008) You can swim but you can't hide: the global status and conservation of oceanic pelagic sharks and rays. *Aquat Conserv* 482:459–482
- Fennessy S, Isaksen B (2007) Can bycatch reduction devices be implemented successfully on prawn trawlers in the Western Indian Ocean? *Afr J Mar Sci* 29:453–463
- Fogarty MJ, Murawski SA (1998) Large-scale disturbance and the structure of marine systems: fishery impacts on georges. *Ecol Appl* 8:6–22
- Fortuna C, Vallini C, Filidei E, Ruffino M, Consalvo I, Muccio SD, Gion C, Scacco U, Tarulli E, Giovanardi O, Mazzola A (2010) By-catch of cetaceans and other species of conservation concern during pair trawl fishing operations in the Adriatic Sea (Italy). *Chem Ecol* 26:65–76
- Francis M, Duffy C (2002) Distribution, seasonal abundance and bycatch of basking sharks (*Cetorhinus maximus*) in New Zealand, with observations on their winter habitat. *Mar Biol* 140:831–842
- Francis MP, Griggs LH, Baird SJ (2001) Pelagic shark bycatch in the New Zealand tuna longline fishery. *Mar Freshw Res* 52:165–178
- Frick LH, Walker TI, Reina RD (2010) Trawl capture of Port Jackson sharks, *Heterodontus portusjacksoni*, and gummy sharks, *Mustelus antarcticus*, in a controlled setting: effects of tow duration, air exposure and crowding. *Fish Res* 106:344–350
- Frick LH, Walker TI, Reina RD (2012) Immediate and delayed effects of gill-net capture on acid–base balance and intramuscular lactate concentration of gummy sharks, *Mustelus antarcticus*. *Comp Biochem Physiol A* 162(2):88–93
- Gallagher AJ, Frick LH, Bushnell PG, Brill RW, Mandelman JW (2010) Blood gas, oxygen saturation, pH, and lactate values in elasmobranch blood measured with a commercially available portable clinical analyzer and standard laboratory instruments. *J Aquat Anim Health* 22:229–234
- Gilman EL (2011) Bycatch governance and best practice mitigation technology in global tuna fisheries. *Mar Policy* 35:590–609
- Gilman C, Clarke S, Brothers N, Alfaro-Shigueto J, Mandelman J, Mangel J, Petersen S, Piovano S, Thomson N, Dalzell P, Donoso M, Goren M, Werner T (2007) Shark depredation and unwanted bycatch in Pelagic longline fisheries: industry practices and attitudes, and shark avoidance strategies. Western Pacific Regional Fishery Management Council, Honolulu
- Godin AC, Worm B (2010) Keeping the lead: how to strengthen shark conservation and management policies in Canada. *Mar Policy* 34:995–1001
- Goldsworthy SD, Page B (2007) A risk-assessment approach to evaluating the significance of seal bycatch in two Australian fisheries. *Biol Conserv* 139:269–285
- Grantham H, Petersen S, Possingham H (2008) Reducing bycatch in the South African pelagic longline fishery: the utility of different approaches to fisheries closures. *Endanger Species Res* 5:291–299
- Gurshin CWD, Szedlmayer ST (2004) Short-term survival and movements of Atlantic sharpnose sharks captured by hook-and-line in the north-east Gulf of Mexico. *J Fish Biol* 65:973–986
- Hall M (1996) On bycatches. *Rev Fish Biol Fisheries* 6:319–352
- Hall SJ, Mainprize BM (2005) Managing by-catch and discards: how much progress are we making and how can we do better? *Fish Fish* 6:134–155
- Hall M, Alverson DL, Metzuzals K (2000) By-catch: problems and solutions. *Mar Pollut Bull* 41:204–219
- Heithaus MR, Frid A, Wirsing AJ, Worm B (2008) Predicting ecological consequences of marine top predator declines. *Trends Ecol Evol* 23:202–210
- Hill B, Wassenberg TJ (2000) The probable fate of discards from prawn trawlers fishing near coral reefs A study in the northern Great Barrier Reef, Australia. *Fish Res* 48:277–286
- Hoffmayer ER, Hendon JM, Parsons GR (2012) Seasonal modulation in the secondary stress response of a carcharhinid shark, *Rhizoprionodon terraenovae*. *Comp Biochem Physiol A* 162(2):81–87
- Huang HW (2011) Bycatch of high sea longline fisheries and measures taken by Taiwan: actions and challenges. *Mar Policy* 35:712–720
- Huang HW, Liu KM (2010) Bycatch and discards by Taiwanese large-scale tuna longline fleets in the Indian Ocean. *Fish Res* 106:261–270
- Hyatta MW, Anderson PA, O'Donnell PM, Berzins IK (2012) Assessment of acid–base derangements among bonnethead (*Sphyrna tiburo*), bull (*Carcharhinus leucas*), and lemon (*Negaprion brevirostris*) sharks from gillnet and longline capture and handling methods. *Comp Biochem Physiol A* 162(2):113–120
- Jones AA, Hall NG, Potter IC (2011) Species compositions of elasmobranchs caught by three different commercial fishing methods off southwestern Australia, and biological data for four abundant bycatch species. *Fish Bull* 108:365–382
- Jordan LK, Mandelman JW, Kajiura SM (2011) Behavioral responses to weak electric fields and a lanthanide metal in two shark species. *J Exp Mar Biol Ecol* 409:345–350
- Kaimmer S, Stoner AW (2008) Field investigation of rare-earth metal as a deterrent to spiny dogfish in the Pacific halibut fishery. *Fish Res* 94:43–47
- Kalmijn A (1982) Electric and magnetic field detection in elasmobranch fishes. *Science* 218:916–918
- Kaplan IC, Cox SP, Kitchell JF (2007) Circle hooks for pacific longliners: not a panacea for marlin and shark bycatch, but part of the solution. *Trans Am Fish Soc* 136:392–401
- Kelleher K (2005) Discards in the world's marine fisheries an update. United Nations Food and Agriculture Organization Fisheries Technical Paper 470
- Kerstetter DW, Graves JE (2006) Effects of circle versus J-style hooks on target and non-target species in a pelagic longline fishery. *Fish Res* 80:239–250
- Krogh M, Reid D (1996) Bycatch in the protective shark meshing programme off south-eastern New South Wales, Australia. *Biol Conserv* 77:219–226
- Levesque J (2008) International fisheries agreement: review of the International Commission for the Conservation of Atlantic Tunas: case study-shark management. *Mar Policy* 32:528–533

- Lewison R, Crowder L, Read A, Freeman S (2004) Understanding impacts of fisheries bycatch on marine megafauna. *Trends Ecol Evol* 19:598–604
- Lynch A-MJ, Sutton SG, Simpfendorfer CA (2010) Implications of recreational fishing for elasmobranch conservation in the Great Barrier Reef Marine Park. *Aquat Conserv* 20:312–318
- MacLennan DN (1995) Gear selectivity and the variation of yield. *ICES J Mar Sci* 52:827–836
- Mandelman J, Farrington M (2007) The estimated short-term discard mortality of a trawled elasmobranch, the spiny dogfish (*Squalus acanthias*). *Fish Res* 83:238–245
- Mandelman JW, Skomal G (2009) Differential sensitivity to capture stress assessed by blood acid-base status in five carcharhinid sharks. *J Comp Physiol B* 179:267–277
- Mandelman JW, Cooper PW, Werner TB, Lagueux KM (2008) Shark bycatch and depredation in the U.S. Atlantic pelagic longline fishery. *Rev Fish Biol Fisheries* 18:427–442
- Marshall H, Field L, Afadada A, Sepulveda C, Skomal G, Bernal D (2012) Hematological indicators of stress in longline-captured sharks. *Comp Biochem Physiol A* 162(2):121–129
- Maunder M, Punt A (2004) Standardizing catch and effort data: a review of recent approaches. *Fish Res* 70:141–159
- McKinnell S (1998) Shark bycatch in the Japanese high seas squid driftnet fishery in the North Pacific Ocean. *Fish Res* 39:127–138
- Megalofonou P, Deflorio M (2005) Incidental catch and estimated discards of pelagic sharks from the swordfish and tuna fisheries in the Mediterranean Sea. *Fish Bull* 103:620–634
- Millar RB (1992) Estimating the size selectivity of fishing gear by conditioning of the total catch. *J Am Stat Assoc* 87:962–968
- Minami M, Lennertcody C, Gao W, Romanverdesoto M (2007) Modeling shark bycatch: the zero-inflated negative binomial regression model with smoothing. *Fish Res* 84:210–221
- Morgan A, Carlson J, Ford T, Siceloff L, Hale L, Allen MS, Burgess G (2009) Temporal and spatial distribution of finfish bycatch in the U.S. Atlantic bottom longline shark fishery. *Mar Fish Rev* 72:34–38
- Moyes CD, Fragoso N, Musyl MK, Brill RW (2006) Predicting postrelease survival in large pelagic fish. *Trans Am Fish Soc* 135:1389–1397
- Musyl MK, Moyes CD, Brill RW, Fragoso NM (2009) Factors influencing mortality estimates in post release survival studies. *Mar Ecol Prog Ser* 396:157–159
- Musyl MK, Brill RW, Curran DS, Fragoso NM, McNaughton LM, Nielsen A, Kikkawa BS, Moyes CD (2011) Postrelease survival, vertical and horizontal movements, and thermal habitats of five species of pelagic sharks in the central Pacific Ocean. *Fish Bull* 109:341–368
- O'Connell C, Abel D, Stroud E, Rice P (2011) Analysis of permanent magnets as elasmobranch bycatch reduction devices in hook-and-line and longline trials. *Fish Bull* 109:394–401
- Ordines F, Massutí E, Guijarro B, Mas R (2006) Diamond vs. square mesh codend in a multi-species trawl fishery of the western Mediterranean: effects on catch composition, yield, size selectivity and discards. *Aquat Living Resour* 19:329–338
- Orphanides CD (2010) Protected species bycatch estimating approaches: estimating harbor porpoise bycatch in U.S. Northwestern Atlantic gillnet fisheries. *J Northwest Atl Fish Sci* 42:55–76
- Pajuelo JG, González JA, Santana JI (2010) Bycatch and incidental catch of the black scabbardfish (*Aphanopus* spp.) fishery off the Canary Islands. *Fish Res* 106:448–453
- Pauly D (2002) Fisheries management: sustainability vs. reality. p 5–11. In: Pauly D, Palomares ML (eds) Fisheries centre research reports 10(8):28p
- Perez J, Wahrlich R (2005) A bycatch assessment of the gillnet monkfish fishery off southern Brazil. *Fish Res* 72:81–95
- Petersen S, Honig M, Ryan P, Underhill L, Compagno L (2009) Pelagic shark bycatch in the tuna- and swordfish-directed longline fishery off southern Africa. *Afr J Mar Sci* 31: 215–225
- Philippart C (1998) Long-term impact of bottom fisheries on several by-catch species of demersal fish and benthic invertebrates in the south-eastern North Sea. *ICES J Mar Sci* 55:342–352
- Poisson F, Gaertner J-C, Taquet M, Durbec J-P, Bigelow K (2009) Effects of lunar cycle and fishing operations on longline-caught pelagic fish: fishing performance, capture time, and survival of fish. *Fish Bull* 108:268–281
- Polovina JJ, Abecassis M, Howel EA, Woodworth P (2002) Increases in the relative abundance of mid-trophic level fishes concurrent with declines in apex predators in the subtropical North Pacific, 1996–2006. *Fish Bull* 107:523–531
- Pullin AS, Stewart GB (2006) Guidelines for systematic review in conservation and environmental management. *Conserv Biol* 20:1647–1656
- Punt A, Smith D, Tuck G, Methot R (2006) Including discard data in fisheries stock assessments: two case studies from south-eastern Australia. *Fish Res* 79:239–250
- Raby GD, Colotelo AH, Blouin-demers G, Cooke SJ (2011) Freshwater commercial bycatch: an understated conservation problem. *Bioscience* 61:271–280
- Renshaw GMC, Kutek AK, Grant GD, Anoopkumar-Dukie S (2012) Forecasting elasmobranch survival following exposure to severe stressors. *Comp Biochem Physiol A* 162(2):101–112
- Revill A, Dulvy N, Holst R (2005) The survival of discarded lesser-spotted dogfish (*Scyliorhinus canicula*) in the Western English Channel beam trawl fishery. *Fish Res* 71:121–124
- Richards A (1994) Problems of drift-net fisheries in the South Pacific. *Mar Pollut Bull* 29:106–111
- Rigg DP, Peverell SC, Hearndon M, Seymour JE (2009) Do elasmobranch reactions to magnetic fields in water show promise for bycatch mitigation? *Mar Freshw Res* 60:942–948
- Robbins WD, Peddemors VM, Kennelly SJ (2011) Assessment of permanent magnets and electropositive metals to reduce the line-based capture of Galapagos sharks, *Carcharhinus galapagensis*. *Fish Res* 109:100–106
- Rodríguez-Cabello C, Sánchez F, Velasco F (2005) Growth of lesser spotted dogfish (*Scyliorhinus canicula* L., 1758) in the Cantabrian Sea, based on tag-recapture data. *J Northwest Atl Fish Sci* 37:131–140
- Rogan E, Mackey M (2007) Megafauna bycatch in drift nets for albacore tuna (*Thunnus alalunga*) in the NE Atlantic. *Fish Res* 86:6–14

- Romanov EV (1999) Bycatch in the tuna purse-seine fisheries of the western Indian Ocean. *Fish Bull* 100:90–105
- Romine JG, Musick JA, Burgess GH (2009) Demographic analyses of the dusky shark, *Carcharhinus obscurus*, in the Northwest Atlantic incorporating hooking mortality estimates and revised reproductive parameters. *Environ Biol Fishes* 84:277–289
- Russell S (1993) Shark bycatch in the northern Gulf of Mexico tuna longline fishery, 1988–91, with observations on the nearshore directed shark fishery. In: Branstetter S (ed) Conservation biology of elasmobranchs. NOAA Technical Report NMFS, pp 19–330
- Samaranyanka A, Engas A, Jorgensen T (1997) Effects of hanging ratio and fishing depth on the catches rates of drifting tuna gillnets in Sri Lankan waters. *Fish Res* 29:1–12
- Schindler DE, Essington TE, Kitchell JF, Boggs C, Hilborn R (2002) Sharks and tunas: fisheries impacts on predators with contrasting life histories. *Ecol Appl* 12:735–748
- Shepard E, Wilson R, Halsey LG, Quintana F, Gomez Laich A, Gleiss AC, Liebsch N, Myers AE, Norman B (2008) Derivation of body motion via appropriate smoothing of acceleration data. *Aquat Biol* 4:235–241
- Shepherd TD, Myers RA (2005) Direct and indirect fishery effects on small coastal elasmobranchs in the northern Gulf of Mexico. *Ecol Lett* 8:1095–1104
- Simpfendorfer CA, Heupel MR, White WT, Dulvy NK (2011) The importance of research and public opinion to conservation management of sharks and rays: a synthesis. *Mar Freshw Res* 62:518–527
- Sisneros J, Donald NR (2001) Surfactants as chemical shark repellents: past, present, and future. *Environ Biol Fishes* 60:117–130
- Skomal GB (2007) Evaluating the physiological and physical consequences of capture on post-release survivorship in large pelagic fishes. *Fish Manage Ecol* 14:81–89
- Skomal GB, Bernal D (2010) Physiological responses to stress in sharks. In: Carrier JC, Musick JA, Heithaus MR (eds) Sharks and their relatives II—Biodiversity, adaptive physiology and conservation. CRC Press Taylor Francis Group, Boca Raton, pp 457–488
- Skomal GB, Mandelman JW (2012) The physiological response to anthropogenic stressors in marine elasmobranch fishes: a review with a focus on the secondary response. *Comp Biochem Physiol A* 162(2):146–155
- Smith T, Jeremy HS (2005) Assessing effects of diel period, gear selectivity and predation on patterns of microhabitat use by fish in a mangrove dominated system in SE Australia. *Mar Ecol Prog Ser* 294:257–270
- Sobrino I, García T, Baro J (2000) Trawl gear selectivity and the effect of mesh size on the deep-water rose shrimp (*Parapenaeus longirostris*, Lucas, 1846) fishery off the gulf of Cádiz (SW Spain). *Fish Res* 44:235–245
- Southwood A, Fritsches K, Brill R, Swimmer Y (2008) Sound, chemical, and light detection in sea turtles and pelagic fishes: sensory-based approaches to bycatch reduction in longline fisheries. *Endanger Species Res* 5:225–238
- Soykan C, Moore J, Zydalis R, Crowder L, Safina C, Lewison R (2008) Why study bycatch? An introduction to the theme section on fisheries bycatch. *Endanger Species Res* 5:91–102
- Statistics and Information Service of the Fisheries and Aquaculture Department/Service des statistiques et de l'information du Département des pêches et de l'aquaculture/Servicio de Estadísticas y Información del Departamento de Pesca y Acuicultura. *FAO yearbook. Fishery and Aquaculture Statistics. 2008/FAO annuaire. Statistiques des pêches et de l'aquaculture. 2008/FAO anuario. Estadísticas de pesca y acuicultura. 2008. Rome/Roma, FAO. 2010. 72p*
- Stobutzki IC, Miller MJ, Jones P, Salini JP (2001) Bycatch diversity and variation in a tropical Australian penaeid (Shrimp) fishery; the implications for monitoring. *Fish Res* 53:293–301
- Stobutzki IC, Miller MJ, Heales D, Brewer D (2002) Sustainability of elasmobranchs caught as bycatch in a tropical prawn (shrimp) trawl fishery. *Fishery* 100:800–821
- Stoner A, Kaimmer S (2008) Reducing elasmobranch bycatch: laboratory investigation of rare earth metal and magnetic deterrents with spiny dogfish and Pacific halibut. *Fish Res* 92:162–168
- Storai T, Zinzula L, Repetto S, Zuffa M, Morgan A, Mandelman J (2011) Bycatch of large elasmobranchs in the traditional tuna traps (*tonnare*) of Sardinia from 1990 to 2009. *Fish Res* 109:74–79
- Swimmer Y, Suter J, Arauz R, Bigelow K, López A, Zanela I, Bolaños A, Ballesterio J, Suárez R, Wang J, Boggs C (2011) Sustainable fishing gear: the case of modified circle hooks in a Costa Rican longline fishery. *Mar Biol* 158:757–767
- Tallack SML, Mandelman JW (2009) Do rare-earth metals deter spiny dogfish? A feasibility study on the use of electro-positive “mischmetal” to reduce the bycatch of *Squalus acanthias* by hook gear in the Gulf of Maine. *ICES J Mar Sci* 66:315–322
- Thorpe T, Frierson D (2009) Bycatch mitigation assessment for sharks caught in coastal anchored gillnets. *Fish Res* 98:102–112
- Trent L, Parshley DE, Carlson JK (1997) Catch and bycatch in the shark drift gillnet fishery off Georgia and East Florida. *Mar Fish Rev* 59:19–28
- Valenzuela A, Bustamante C, Lamilla J (2008) Morphological characteristics of five bycatch sharks caught by southern Chilean demersal longline fisheries. *Sci Mar* 72:231–237
- Vega R, Licandeo R (2009) The effect of American and Spanish longline systems on target and non-target species in the eastern South Pacific swordfish fishery. *Fish Res* 98:22–32
- Walker TI, Hudson RJ, Gason AS (2005) Catch evaluation of target, by-product and by-catch species taken by gillnets and longlines in the shark fishery of South-eastern Australia. *J Northwest Atl Fish Sci* 37:505–530
- Ward P, Hindmarsh S (2007) An overview of historical changes in the fishing gear and practices of pelagic longliners, with particular reference to Japan's Pacific fleet. *Rev Fish Biol Fisheries* 17:501–516
- Ward P, Lawrence E, Darbyshire R, Hindmarsh S (2008) Large-scale experiment shows that nylon leaders reduce shark bycatch and benefit pelagic longline fishers. *Fish Res* 90:100–108
- Ward P, Epe S, Kreutz D, Lawrence E, Robins C, Sands A (2009) The effects of circle hooks on bycatch and target catches in Australia's pelagic longline fishery. *Fish Res* 97:253–262

- Watson JW, Epperly SO, Shah AK, Foster DG (2005) Fishing methods to reduce sea turtle mortality associated with pelagic longlines. *Can J Fish Aquat Sci* 62:965–981
- Watson JT, Essington TE, Lennert-Cody CE, Hall MA (2008) Trade-offs in the design of fishery closures: management of silky shark bycatch in the eastern Pacific Ocean tuna fishery. *Conserv Biol* 23:626–635
- Whitney NM, Papastamatiou YP, Holland KN, Lowe CG (2007) Use of an acceleration data logger to measure diel activity patterns in captive whitetip reef sharks, *Triaenodon obesus*. *Aquat Living Resour* 20:299–305
- Whoriskey S, Arauz R, Baum JK (2011) Potential impacts of emerging mahi-mahi fisheries on sea turtle and elasmobranch bycatch species. *Biol Conserv* 144:1841–1849
- Wikelski M, Cooke SJ (2006) Conservation physiology. *Trends Ecol Evol* 21:38–46
- Yamandú HM, Brum F, Barea LC, Chocca JF (1998) Incidental catch associated with swordfish longline fisheries in the south-west Atlantic Ocean. *Mar Freshw Res* 49:633–639
- Yokota K, Kiyota M, Minami H (2006) Shark catch in a pelagic longline fishery: comparison of circle and tuna hooks. *Fish Res* 81:337–341
- Zeeberg J, Corten A, Degraaf E (2006) Bycatch and release of pelagic megafauna in industrial trawler fisheries off Northwest Africa. *Fish Res* 78:186–195
- Zhou S, Smith ADM, Fuller M (2011) Quantitative ecological risk assessment for fishing effects on diverse data-poor non-target species in a multi-sector and multi-gear fishery. *Fish Res* 112:168–178
- Zollett E (2009) Bycatch of protected species and other species of concern in US east coast commercial fisheries. *Endanger Species Res* 9:49–59
- Zydelis R, Wallace BP, Gilman EL, Werner TB (2009) Conservation of marine megafauna through minimization of fisheries bycatch. *Conserv Biol* 23:608–616