

Ecology and conservation of grassland birds in southeastern South America: a review

Adrián B. Azpiroz,^{1,8} Juan Pablo Isacch,² Rafael A. Dias,^{3,4} Adrián S. Di Giacomo,⁵ Carla Suertegaray Fontana,⁶ and Cristina Morales Palarea⁷

¹Laboratorio de Genética de la Conservación, Instituto de Investigaciones Biológicas Clemente Estable, Av. Italia 3318 (C.P. 11600), Montevideo, Uruguay

²Instituto de Investigaciones Marinas y Costeras (IIMyC), Facultad de Ciencias Exactas y Naturales, Consejo Nacional de Investigaciones Científicas y Técnica, Universidad Nacional de Mar del Plata, Funes 3250, (7600) Mar del Plata, Argentina

³LABECO, Universidade Católica de Pelotas, R. Félix da Cunha 412, CEP 96010–000, Pelotas, RS, Brazil

⁴Programa de Pós-Graduação em Ecologia, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, CEP 91501–970, Porto Alegre, RS, Brazil

⁵Instituto de Ecología, Genética y Evolución de Buenos Aires (IEGEB-CONICET), Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Int. Güiraldes y Av. Cantilo s/n, Pabellón II, Ciudad Universitaria, C1428EHA, Buenos Aires, Argentina

⁶Laboratório de Ornitologia, Museu de Ciências e Tecnologia, Curso de Pós-graduação em Zoologia, Pontifícia Universidade Católica do Rio Grande do Sul, PUCRS. Av. Ipiranga, 6681. CEP 90619-900. Porto Alegre, RS, Brazil

⁷Asociación Guyra Paraguay, Gaeano Martino 215 c/Te. Ross, C, C:1132, Asunción, Paraguay

Received 4 January 2012; accepted 1 May 2012

ABSTRACT. The grasslands of southeastern South America (SESA), comprising one of the most extensive grassland ecosystems in the Neotropics, have been negatively impacted by the development of the livestock industry, arable agriculture, and forestry. SESA grasslands have a rich avifauna that includes 22 globally threatened and near-threatened species, and many other species have suffered local population extinctions and range reductions. In addition to habitat loss and fragmentation, grassland birds in SESA are threatened by improper use of agrochemicals, unfavorable fire management regimes, pollution, and illegal capture and hunting. Studies to date have provided information about the distribution of grassland birds, the threats populations face, and the habitat requirements of some threatened species, but more information is needed concerning dispersal and migration patterns, genetics, and factors that influence habitat use and species survival in both natural and agricultural landscapes. There are few public protected areas in the region (1% of original grasslands), and many populations of threatened grassland birds are found on private lands. Therefore, efforts to preserve grassland habitat must reconcile the interests of land owners and conservationists. Current conservation efforts include establishment of public and private reserves, promotion of agricultural activities that reconcile production with biodiversity conservation, development of multilateral conservation projects across countries, and elaboration of action plans. Measures that result in significant losses to private land owners should include economic compensation, and use of economic incentives to promote agriculture and forestry in native grassland areas should be discouraged, especially in priority areas for grassland birds. Although more studies are needed, some actions, particularly habitat protection and improved management of public and private lands, should be taken immediately to improve the conservation status of grassland birds in SESA.

RESUMEN. **Ecología y conservación de aves de pastizal en el SE de Sudamérica: una revisión**

Los pastizales del sureste de Sudamérica (SESA), que conforman uno de los mayores ecosistemas de pastizales en el Neotrópico, han sufrido transformaciones importantes debido al desarrollo de la industria ganadera, la agricultura y la forestación. Los pastizales del SESA tienen una rica avifauna que incluye 22 especies amenazadas o casi amenazadas y muchas otras han sufrido extinciones poblacionales locales y reducciones muy sustanciales de sus rangos de distribución. Además de la pérdida de hábitat y la fragmentación, las aves de pastizal del SESA están amenazadas por el uso inapropiado de agroquímicos, regímenes desfavorables de manejo del fuego, contaminación, captura y caza ilegales. Los estudios al día de hoy han generado información sobre la distribución de aves de pastizal, las amenazas que sus poblaciones enfrentan y de los requerimientos de hábitat de algunas especies amenazadas, pero aun es necesaria más información sobre patrones de dispersión y migración, genética y los factores que afectan los patrones de uso de hábitat y la sobrevivencia de especies en paisajes naturales y agrícolas. Hay muy pocas áreas

⁸Corresponding author. Email: azpiroz@iibce.edu.uy

protegidas públicas en la región (1% de los pastizales originales) y muchas poblaciones de aves de pastizal amenazadas se encuentran en tierras privadas. Por ende, los esfuerzos que buscan preservar características de hábitat similares a las de los pastizales naturales deben reconciliar los intereses de productores y conservacionistas. Los actuales esfuerzos de conservación, incluyen, el establecimiento de reservas privadas y públicas, la promoción de prácticas agrícolas que reconcilien producción y conservación, el desarrollo de proyectos multilaterales de conservación entre varios países, y la elaboración de planes de acción. Las medidas que resultan en pérdidas significativas para los productores privados deberían incluir compensaciones económicas, y el empleo de incentivos económicos para fomentar la agricultura y forestación en pastizales nativos deberían ser desalentado, especialmente en el caso de áreas de prioridad para aves amenazadas. A pesar de que todavía se necesitan mas estudios, algunas medidas, en particular la protección y manejo de hábitat tanto en tierras públicas como privadas deberían aplicarse inmediatamente para mejorar el estado de conservación de las aves de pastizal en el SESA.

Key words: Argentina, Brazil, Brazilian Upland grasslands, Campos, Humid Chaco grasslands, Pampas, Paraguay, Río de la Plata grasslands, Uruguay

Grassland ecosystems are found on every continent except Antarctica, covering between 31 and 43% of the total land area of the world (World Resources 2000–2001). Grasslands have been identified as the terrestrial biome where biodiversity and ecosystem services are most at risk on a global scale due to the large disparity between loss of habitat and the low degree of protection (Hoekstra et al. 2005, Henwood 2010). Grasslands in the southeastern South America (SESA) region are one of the most extensive grassland ecosystems in the Neotropics and have been transformed by the development of a vast livestock industry, arable agriculture, and afforestation (Soriano et al. 1991, Overbeck et al. 2007). About 40 to 45% of the original cover remains, but even most grassland remnants have been modified by livestock grazing (Demaría et al. 2003, Bilenca and Miñarro 2004, Paruelo et al. 2004, Herrera et al. 2009). Only about 1% of the original extent of the Pampas and the Campos region is currently protected (Henwood 2010).

Loss and degradation of grasslands have negatively affected bird populations worldwide. In North America, 75% of grassland birds have experienced population declines over the past 30 years due to changes in land use (Askins et al. 2007). In Europe, 70% of grassland and steppe species have suffered drastic reductions of their populations due to changes in agricultural practices (Donald et al. 2001, 2006). The negative effects of native habitat replacement on grassland birds have also been reported in Asia, Oceania, and Africa (Goriup 1988). In the Neotropics, populations of several resident grassland birds have declined markedly (Fraga et al. 1998, Tubaro and Gabelli 1999, Fraga 2003, Di Giacomo and Di Giacomo 2004), and those of long-distance Nearctic migrants

have also decreased since the 1800s (Lanctot et al. 2002, Vickery et al. 2010). Because nearly 10% of globally threatened bird species in South America inhabit grasslands, compared to 6.3% for the world as a whole (Collar et al. 1994), conservation of grassland birds in the Neotropics is a critical issue.

Here we review the status of grassland bird ecology and conservation in SESA. First, we briefly describe SESA grasslands and their associated avifauna, as well as land-use patterns and evidence of declining bird populations. We analyze factors affecting grassland birds in light of data collected during the last 15 years and highlight knowledge gaps. Finally, we describe current conservation strategies (and their limitations) and provide recommendations for further action.

We used Vickery et al. (1999) as a primary source for defining grassland habitat and grassland birds. According to these authors, grasslands are extensive areas with >50% grass (Poaceae) or sedge (Cyperaceae) cover and few scattered shrubs (<4 m high) and trees. We included habitats where the soil is saturated with water, namely wet-mesic grasslands in upland swales, but excluded those that remain flooded for long periods, such as freshwater, brackish, and saltwater wetlands. Following Vickery et al. (1999), we define a grassland bird as “any species that has become adapted to and reliant on some variety of grassland habitats for part or all of its life cycle.” We used the list of grassland birds provided by Vickery et al. (1999) as a baseline for identifying species in our study area. We modified our list based on new information or based on our personal experience.

Our review is divided into six sections. The first three topics (SESA grassland birds, Declining populations of SESA grassland birds, and

Factors affecting SESA grassland birds) were developed primarily by reviewing all available literature we could access. Information was accessed using Google Scholar and the electronic databases ISI Web of Science, SCOPUS, and SCIELO. We also contacted experts to include unpublished information or clarify existing data, and used our own unpublished data when necessary.

For the “South American Grassland Birds: Knowledge and Gaps” section, we reviewed the literature to locate papers focusing on grassland birds in our study area. Papers were identified by searching all databases in ISI Web of Knowledge and SCOPUS spanning all available years up to 2011. We followed this method to limit our sampling to high-impact journals expected to contain the most relevant information. We conducted searches using the key words “bird pampas” and “bird grassland South America.” We read all abstracts and available key words in the resulting papers to limit the information to our geographical and ecological areas of interest. Papers focusing on fossil and wetland birds were excluded. We did, however, include papers on waterbirds using grassland habitat. We also examined possible geographic and topic biases in the literature. To do so, we classified papers into 11 categories, including assemblages and community ecology, biology (including feeding, breeding, and social behavior), climate and weather impacts, conservation, distribution, genetics, habitat and land-use associations, methods, morphology, population monitoring (including migration), and toxicology, and compared the number of papers in each category for each country. In the last two sections (“Current bird conservation initiatives” and “Additional proposed conservation measures”), we summarize available information concerning grassland conservation issues in SESA as well as our own experiences developed through direct involvement with grassland bird conservation in the region.

DESCRIPTION OF SESA GRASSLANDS

The geographic extent of our review covers grasslands in Southeastern South America from 28° to 39° S. This region forms an arc around the Río de la Plata covering eastern and northeastern Argentina, southern Paraguay, southern Brazil, and Uruguay. The region is treated here as the

SESA Grasslands and includes: (1) the “Río de la Plata grasslands” (*sensu* Soriano et al. 1991), (2) the upland grasslands of southern Brazil (*sensu* Overbeck et al. 2007; hereafter “Brazilian Upland grasslands”), and (3) the savanna grasslands in the Humid Chaco along the Paraguay river in southeastern Paraguay and northeastern Argentina (*sensu* Olson et al. 2001 and references therein; hereafter “Humid Chaco grasslands”; Fig. 1). All these grassland subregions have similar habitats and vegetation types, and support an assemblage of bird species that faces similar threats and conservation needs.

The Río de la Plata grasslands (Fig. 1) occupy a region of more than 700,000 km² in central-eastern Argentina (eastern Corrientes, Entre Ríos, southern Córdoba, southern Santa Fe, Buenos Aires, eastern La Pampa, and eastern San Luis), Uruguay, and southern Brazil (southern and western Rio Grande do Sul; Soriano et al. 1991). These grasslands are characterized by flat to undulating relief with elevations < 900 m. The region has mesothermic features mostly regulated by the tempering effect of the Atlantic Ocean, with annual mean temperatures ranging from 14° in the south to 18° in the north (Soriano et al. 1991). There is also a rainfall gradient (1600 mm in the northeast to 500 mm in the southwest) with high interannual variability that determines incipient water excess and water deficits, respectively (Burgos and Vidal 1951, Overbeck et al. 2007). Paleontological and palynological evidence suggest that grasslands have dominated the region at least since the Neogene (Prieto 2000, Behling et al. 2005). The southern half of this region is a seral graminoid steppe of medium height (~0.4–1 m), where dominant genera are *Stipa*, *Poa*, *Piptochaetium*, and *Sorghastrum*; in the northern half, cespitose grasslands with grasses of the genera *Andropogon*, *Aristida*, *Briza*, *Erianthus*, *Piptochaetium*, *Poa*, and *Stipa* predominate (Soriano et al. 1991, Overbeck et al. 2007). Broadleaf genera such as *Eryngium* and *Baccharis* are widespread in the Southern and Northern Campos (Soriano et al. 1991, Overbeck et al. 2007; Fig. 1). Grassland regions adjacent to areas originally covered by Atlantic Forest, especially in the northern and eastern sectors of the Northern Campos, are characterized by forest-grassland mosaics, parkland, and shrubland (Pillar and Quadros 1997, Cordeiro and Hasenack 2009, Boldrini et al. 2009). Wetlands occur in coastal




Fig. 1. Southeastern South American Grasslands (modified from Soriano et al. 1991, Olson et al. 2001 and references therein, Guyra Paraguay 2005, and Overbeck et al. 2007), including the Río de la Plata Grasslands (RP1: Southern Pampa, RP2: Flooding Pampa, RP3a West Inland Pampa, RP3b: Flat Inland Pampa, RP4: Rolling Pampa, RP5: Mesopotamic Pampa, RP6: Northern Campos in Argentina, RP7: Southern Paraguayan grasslands, RP8: Northern Campos in Brazil, RP9: Northern Campos in Uruguay, and RP10: Southern Campos), Brazilian Upland grasslands (BR), and Humid Chaco grasslands (CH). Bold black lines in main picture represent approximate country limits and fine black lines show relevant province and state limits for Argentina and Brazil, respectively.

areas and floodplains along rivers and lagoons (Soriano et al. 1991, Cordeiro and Hasenack 2009). Grasslands in southern Paraguay have recently been considered as an extension of the Río de la Plata grasslands (*sensu* Soriano et al. 1991) on the basis of close vegetation and avifaunal affinities with the Northern Campos of Argentina (Guyra Paraguay 2005, Clay et al. 2008). These grasslands occupy an area of ~2,035,400 ha in central-southern Paraguay (Misiones, western Itapúa and Caazapá, and south of Paraguari).

The Brazilian Upland grasslands are upland formations (700–1800 m asl) that covered ~60,000 km² in northeastern Rio Grande do Sul, Santa Catarina, and Paraná states in Brazil (Overbeck et al. 2007, H. Hasenack,

pers. comm.). These grasslands are distributed as large discrete patches on gently rolling terrain amid Araucaria forest in the Southern Brazilian Plateau (Klein 1978, Leite 2002, Cordeiro and Hasenack 2009). Annual precipitation is high (1500–2000 mm) and evenly distributed throughout the year; mean annual temperatures range from 10–22°C, with frequent frosts and occasional snow during winter (Overbeck et al. 2007). These grasslands are characterized by a high diversity of Asteraceae and Poaceae and high levels of endemism (Boldrini et al. 2009). Cespitose grasses predominate in hilltops, especially *Andropogon lateralis*, whereas drainages are covered with Cyperaceae and the spiny *Eryngium pandanifolium*, a ubiquitous element in boggy swales (Boldrini et al. 2009).

Humid Chaco grasslands are part of a complex mosaic of palm and tree savannas, forests, and marshes. In Argentina, they occupy the eastern portion of the provinces of Formosa, Chaco, Santa Fe, and western and central Corrientes. In Paraguay, these grasslands are found mainly in the departments of Cordillera, Central, Paraguari, and Ñeembucú, and eastern portions of Pte. Hayes. Relief is flat and the climate is humid temperate with high interannual variability (Gorleri 2005). Mean annual temperature drops from north to south (23 to 18°C) and annual rainfall varies from 1300 mm in the east to 750 mm in the west. Vegetation in Chaco grasslands is dominated by tall grasses such as *Paspalum*, *Sorghastrum*, *Andropogon*, *Imperata*, and other medium-size grasses such as *Elionurus* and *Schizachyrium*. In marshy areas, the dominant vegetation is represented by *Panicum* and *Cyperus* species (Maturro et al. 2005).

LAND-USE PATTERNS AND HABITAT ALTERATION IN SESA GRASSLANDS

Conversion of SESA native grasslands to cultivated fields and grazing rangelands has varied with land-use and land-cover patterns in different areas (Baldi and Paruelo 2008, Cordeiro and Hasenack 2009). Although climate shifts and the arrival of humans at the beginning of the Holocene brought major alterations to grasslands (Behling et al. 2005), current impacts are related to European colonization that started in the late 15th century. Horses and cattle were introduced in the Río de la Plata grasslands during the 16th century (Soriano et al. 1991). Livestock flourished in these grasslands, and fire was commonly used as a management tool to increase forage quality of grasslands. By the end of the 19th century, substantial areas of native grasslands had been converted to pastures sown with non-native species (e.g., *Festuca* sp., *Lolium* sp., and *Medicago* sp.) to further improve forage quality, and crops (Soriano et al. 1991). When Darwin arrived at the Río de la Plata grasslands (1831–1832), the landscape had changed substantially and large native herbivores such as guanacos (*Lama guanicoe*), Pampas deer (*Ozotoceros bezoarticus*), and Greater Rheas (*Rhea americana*) had disappeared from several areas (Soriano et al. 1991).

During the 20th century, many original natural grasslands in the Pampas' subregions and a smaller proportion in the Campos subregions were replaced by cropland (wheat, maize, sorghum, and soybean) and sown pastures for livestock. During the last few decades, in particular, agricultural intensification has been supported by technological developments (tillage, transgenic seeds, and high agrochemical input) and market factors (Paruelo et al. 2005, Baldi et al. 2006). This has favored the further expansion of crops over lands traditionally devoted to cattle ranching. In fact, 6% of the native grassland area in the Río de la Plata (*sensu* Soriano et al. 1991) was lost between 1985 and 2005, mainly due to expansion of annual crops (Baldi and Paruelo 2008). In Argentina, soybean-cultivated areas have multiplied and currently represent 50% of the total crop area in that country (mainly in the Pampas and Chaco ecoregions; Aizen et al. 2009). In the Inland Pampa in western Argentina, habitat alteration has resulted in a dramatic loss of grasslands. During a 16-year period (1985–2001), total grassland cover in this subregion was reduced from 84% to 38% at a regional scale (Demaría et al. 2008). This loss of native vegetation is probably irreversible (Demaría et al. 2008).

In Buenos Aires, nearly 33% of the total area of the province (~10,000,000 ha) consisted of cultivated annual crops in 2003–2004 (FAUBA 2011). In the Northern Campos of southern Brazil, agricultural practices have followed a pattern similar to that in Argentina and soybeans, maize, and rice are the dominant crops (Crawshaw et al. 2007, Overbeck et al. 2007). In Brazilian Upland grasslands, in addition to soybeans and maize, potatoes, wheat, and apples are dominant crops. In Rio Grande do Sul, grassland conversion has been substantial, with 15.6% of native grasslands lost in a 27-year period (1976–2002) at an annual conversion rate of ~1000 km² (Cordeiro and Hasenack 2009). By 2002, only 25% of the original grasslands remained in the state (Cordeiro and Hasenack 2009). In western Uruguay, croplands have also increased, especially soybeans, wheat, and corn, while the vast ricelands in the east have remained stable (Paruelo et al. 2006). In Paraguay, ~50% of the grassland area has been transformed from traditional cattle ranching to monocultures of rice, soybeans, and eucalyptus (*Eucalyptus* spp.; Morales et al. 2010).

In extensive areas of grasslands of north-eastern Argentina, Uruguay, southern Paraguay, and southern Brazil, afforestation with non-native species has expanded over traditional cattle production areas (Overbeck et al. 2007, Baldi et al. 2008, Morales et al. 2010). In Argentina, afforestation has particularly occurred in Corrientes province, with ~500,000 ha of pines planted in natural grasslands, and this is expected to reach 1,000,000 ha (12% of the provincial area) in the next several years. In southern Brazil, exotic pine (*Pinus* spp.) plantations used for timber and resin extraction cover large expanses of upland grasslands over the three southern states and the coastal grasslands in Rio Grande do Sul. Afforestation with pines is probably the most serious threat to Brazilian Upland grasslands (Bristot 2001, Fontana et al. 2009) and grasslands in Corrientes province, Argentina (Di Giacomo and Krapovickas 2001). In both regions, there are several species of threatened grassland birds sensitive to forestry activity (Di Giacomo and Krapovickas 2001, Bencke et al. 2003, Di Giacomo 2005b, Fontana et al. 2009, Di Giacomo et al. 2010, BirdLife International 2012). In Corrientes, afforestation has been subsidized by the government, but legislation has not considered its impacts on grassland biodiversity. In Brazil, eucalyptus grown for the pulp-wood industry is planted largely in grasslands in the southern and midwestern portions of Rio Grande do Sul. About 4% of the Southern and Northern Campos of Uruguay and southern and western Rio Grande do Sul (1,362,000 ha), most of which originally were covered with grasslands, are currently afforested (Martino and Methol 2008, H. Hasenack, pers. comm.). In both regions, afforestation is encouraged by the government, which markets large areas of grassland remnants as afforestation-prone (Hasse 2006, Paruelo et al. 2006). Afforestation has altered additional grassland areas because plantations can be established even in areas where conventional farming is impractical due to soil limitations.

Because habitat loss and modification usually result in fragmentation (Viglizzo et al. 2001, Demaría et al. 2008), natural grasslands in several areas of SESA are substantially fragmented (Viglizzo et al. 2001). Agriculture-related disturbances have resulted in significant changes in area and patch configuration (Baldi et al. 2006). Again, patterns reflect agriculture suit-

ability that have resulted in significant spatial heterogeneity. Within the Río de la Plata Grasslands, the Flat Inland and Rolling Pampas in Argentina (Fig. 1) show the greatest grassland conversion and fragmentation index, with 34.1–55.7% of grassland cover scattered within an agricultural matrix (Baldi et al. 2006, Baldi and Paruelo 2008, Demaría et al. 2008). In contrast, the Flooding Pampa is the least fragmented subregion of the Argentinean Pampas, with larger (>200 ha) grassland patches making up the landscape matrix (Bilenca and Miñarro 2004). The Southern and Northern Campos of Uruguay and Brazil as well as the Southern and Mesopotamic Pampas of Argentina have intermediate levels of fragmentation (Baldi et al. 2006, Baldi and Paruelo 2008). Large grassland remnants in Uruguay are found primarily in the northern half of the country, and are restricted to the western part of the state in Rio Grande do Sul (Cordeiro and Hasenack 2009, H. Hasenack, pers. comm.). In the Humid Chaco grasslands of Paraguay, agriculture expansion, deforestation, and the invasion of alien species (De Egea et al. 2012) represent serious threats to SESA grassland birds.

SESA GRASSLAND BIRDS

In the SESA grasslands, 109 species of birds have been recorded, with 82 using grasslands on a regular basis (Appendix 1; nomenclature and sequence follow Remsen et al. 2012). These include 22 globally threatened and near-threatened species (BirdLife International 2012; Appendix 1). In addition, many long-distance migrants use the SESA grasslands as wintering or breeding grounds. Non-breeding migrants include 12 Patagonian species present mainly during austral fall and winter, and eight Nearctic visitors that stay during the austral spring and summer (Appendix 1). Breeding migrants are represented by 13 species, with all moving north in the Neotropics during austral winter (Narosky and Di Giacomo 1993, Belton 1994, Stotz et al. 1996). Pampas Pipits (*Anthus chacoensis*) are also migrants, but their breeding and wintering grounds are restricted to SESA (Casañas et al. 2007), so they are treated as a resident in terms of their presence in the region. Populations of some SESA grassland species make local or regional movements (e.g., Bearded Tachuris, *Polystictus pectoralis*, and Spectacled Tyrants, *Hymenops*

perspicillata; Narosky and Di Giacomo 1993), but, because some of these movements are still not clearly characterized, we did not identify any partial migrants.

Based on Narosky and Di Giacomo (1993), Belton (1994), Vickery et al. (1999), and our own observations, we placed grassland birds in SESA in three categories based on use of grassland habitats (Appendix 1): (1) species restricted to grassland habitats, (2) species that make extensive use of grassland habitats, but also use wetlands (including beaches), shrublands, savannas, and row-crop habitats in the region, and (3) species that make extensive use of grassland habitats in some subregions, but not others. The 12 species in the first group are thought to be particularly sensitive to grassland modification. Some birds in the second group can also be locally sensitive to grassland loss, especially those unable to use cropland as a substitute for grasslands or species that depend on non-substitutable resources available both in grasslands and other natural habitats, such as wetlands. Examples of the latter are Black-and-white Monjitas (*Xolmis dominicanus*) and Saffron-cowled Blackbirds (*Xanthopsar flavus*) that forage in grasslands, but depend on adjacent wetlands for breeding and roosting, and many *Sporophila* seedeaters that regularly use herbaceous wetlands amid grasslands for feeding and nesting (Bencke et al. 2003). However, we do not expect all species among or within any of these groups to respond uniformly to transformation of native grasslands because sensitivity to habitat loss and fragmentation is species-specific (Manning et al. 2004, Fischer and Lindenmayer 2007, Johnson et al. 2011). Species in the third group include birds whose dependence on grasslands varies regionally. Black-and-rufous Warbling-Finches (*Poospiza nigrorufa*), for example, inhabit mostly tall grasslands in the Flooding Pampa of Argentina, but make extensive use of other habitats (marsh edges and brushy vegetation) outside this subregion (Narosky and Di Giacomo 1993, Belton 1994, JPI, pers. observ.).

Vegetation height is one of the main drivers of bird diversity and composition in grasslands (Fisher and Davis 2010). Thus, we further discriminated species restricted to homogeneous short grass (<20 cm) or tall grass (>40 cm) and those that use both short and tall grasslands or require patches of short grass within a matrix of

tall grass (Appendix 1). For this classification, we followed Belton (1994), Comparatore et al. (1996), Isacch et al. (2005), Azpiroz and Blake (2009), Isacch and Cardoni (2011), and our own personal observations (Appendix 1). In SESA, 23 species of birds are restricted to short grass grasslands, 25 to tall grass, and 61 make broader use of grassland types. Tall-grass species tended to be of greater conservation concern than short-grass species (Appendix 1).

DECLINING POPULATIONS OF SESA GRASSLAND BIRDS

Human-induced habitat homogenization in SESA agroecosystems has negatively affected bird diversity, leading to local extinctions (Comparatore et al. 1996, Bencke et al. 2003, Di Giacomo and Di Giacomo 2004, Codesido et al. 2011). Populations of several species have declined over the last 100 years in association with land-use changes or hunting (Collar et al. 1992). As a result, 22 species are considered globally threatened or near-threatened (BirdLife International 2012; Appendix 1) and many others are considered in peril at national or regional levels (Bencke et al. 2003, Straube et al. 2004, López-Lanús et al. 2008, Fontana et al. 2009, Santos and Scherer-Neto 2009, IGNIS 2010).

In the absence of long-term monitoring efforts, evidence of declining populations of SESA birds comes from chronological records of the disappearance of local or regional populations. Saffron-cowled Blackbirds (*Xanthopsar flavus*), Black-and-white Monjitas (*Xolmis dominicanus*), Strange-tailed Tyrants (*Alectrurus risora*), and Pampas Meadowlarks (*Sturnella defilippii*) have been extirpated from large areas of their historical ranges and these range contractions have been related to changes in land-use patterns.

In Argentina, Saffron-cowled Blackbirds and Black-and-white Monjitas have suffered range reductions of 50% and 31%, respectively (Fraga et al. 1998, Fraga 2003). Although there are no specific estimates for other countries, both species may be declining throughout their entire ranges, despite no obvious range contractions (Collar et al. 1992, Azpiroz 2000, Dias and Mauricio 2002, Bencke et al. 2003, ICMBio, in press). Population strongholds for these two grassland birds currently include a few small,

disjunct areas in eastern Argentina and neighboring western Uruguay, southeastern Paraguay, and especially in eastern Uruguay, southeastern Rio Grande do Sul, and the Brazilian Upland grasslands of northeastern Rio Grande do Sul and southeastern Santa Catarina (Bencke et al. 2003, Fontana et al. 2009, Santos and Scherer-Neto 2009, IGNIS 2010, BirdLife International 2012, ICMBio, in press).

Strange-tailed Tyrants currently inhabit only 10% of their historical range in Argentina, and current populations are restricted to provinces in the northeast and neighboring southern Paraguay (Di Giacomo and Di Giacomo 2004, Codesido and Fraga 2009, BirdLife International 2012). In Argentina, populations of summer breeders have been extirpated in Buenos Aires, Cordoba, Santa Fe, and Entre Ríos provinces (Di Giacomo and Di Giacomo 2004). In Paraguay, populations have declined locally (BirdLife International 2012) and there are few recent records (Arballo and Gambarotta 1987) in Uruguay, where these tyrants were relatively common summer breeders in the 19th century (Gibson 1885). In southern Brazil, Strange-tailed Tyrants were last recorded in 1914 (Bencke et al. 2010) and may now be extirpated.

Pampas Meadowlarks have suffered a 90% contraction of their historical range (Tubaro and Gabelli 1999). In Argentina, the largest population is in southern Buenos Aires province, with a few scattered records in San Luis, La Pampa, and Corrientes provinces (Chebez 2008). In Uruguay, these meadowlarks were widespread in the past, but only a single population (~300 pairs) remains in southeastern Salto department (Azpiroz 2005, unpubl. data). In Brazil, there are two records from south Rio Grande do Sul that probably date from the 19th century (Ihering 1899, Belton 1994, Bencke 2001, Bencke et al. 2010), and two doubtful observations from Paraná and Santa Catarina (BirdLife International 2012).

Cock-tailed Tyrants (*Alectrurus tricolor*), Sharp-tailed Tyrants (*Culicivora caudacuta*), and Black-masked Finches (*Coryphaspiza melanotis*) were previously considered near-threatened (Collar and Andrew 1988, Collar et al. 1992), but have been up-listed to vulnerable because of local extinctions in Argentina, Paraguay, and Brazil (BirdLife International 2012). In Argentina, Cock-tailed Tyrants were recorded in the provinces of Corrientes and Misiones,

but are now thought to be extirpated, with no confirmed records in the last several decades. In southern Brazil, Cock-tailed Tyrants were reported from Rio Grande do Sul in the 19th century (Wied 1831, Bencke et al. 2010) and currently only a small population subsists in the Brazilian Upland grasslands of Paraná state (Straube et al. 2004, ICMBio, in press). The species is also rare in Paraguay (BirdLife International 2012), with only a few scattered populations (Del Castillo and Clay 2004). Sharp-tailed Tyrants were formerly present in several provinces of Argentina, but are now found at scattered sites in Misiones, Corrientes, Santa Fe, and Formosa (Di Giacomo 2005b). In southern Brazil, the species occurs in Brazilian Upland grasslands of Paraná, Santa Catarina, and northeastern Rio Grande do Sul states, where they appear to be numerous, and in a few localities in the western sector of the latter state, where they are rare (Straube et al. 2004, Fontana et al. 2008, Bencke et al. 2010, Repenning et al. 2010, ICMBio, in press). Black-masked Finches were formerly present in the provinces of Chaco, Santa Fe, Corrientes, and Misiones in Argentina, but recent records are few and limited to northern Corrientes (Chebez 2008). In Paraguay, there are only a few records from Ñeembucu and San Pedro departments (C. Morales, unpubl. data).

Buff-breasted Sandpipers (*Tryngites subruficollis*) are near-threatened migratory grassland shorebirds that have shown a gradual, but marked, decline in numbers (Lancot et al. 2002, Isacch and Martínez 2003a). Historical evidence indicates that this species was abundant, with flocks of 200 to 500 individuals regularly observed during migration in the southern SESA grasslands (Hudson 1920). Regular counts at the most important wintering site in Argentina (Medaland Ranch) reveal a marked decrease in numbers during the last 30 years. Myers and Myers (1979) estimated a maximum of 2000 individuals at that ranch in 1975, but only ~360 individuals were observed in 1991 (Blanco et al. 1993) and, more recently (1996–2000), numbers rarely exceeded 100 individuals at the site (Isacch and Martínez 2003a). Data for other grassland shorebirds obtained in 1992–1993 and 2006–2007 in Argentina suggest range contractions and declining populations of American Golden Plovers (*Pluvialis dominica*), and Upland (*Bartramia longicauda*) and

Pectoral (*Calidris melanotos*) sandpipers (Isacch and Martínez 2003a, A. S. Di Giacomo, unpubl. data).

Populations of other long-distance migrant grassland birds also show evidence of recent declines in wintering areas in SESA. Di Giacomo et al. (2005) compared historical and current records of Bobolinks (*Dolichonyx oryzivorus*) and found a 25% range reduction in wintering areas in Argentina. Similarly, Swainson's Hawks (*Buteo swainsoni*) suffered massive mortality due to insecticide poisoning during the 1990s in Argentina (Woodbridge et al. 1995, Goldstein et al. 1996, 1999a,b). For some *Sporophila* seedeaters, such as Black-bellied Seedeaters (*Sporophila melanogaster*), "Yellow-billed" Plumbeous Seedeaters (*Sporophila* aff. *plumbea*), Marsh Seedeaters (*Sporophila palustris*), and Chestnut Seedeaters (*Sporophila cinnamomea*), SESA grasslands contain key breeding and feeding areas (BirdLife International 2012). Morphological and behavioral evidence suggests that the 'Yellow-billed' Plumbeous Seedeater, currently restricted to grasslands between 700 and 950 m in altitude on mountain slopes near some rivers in the Brazilian Highland Grasslands is a valid species (Repenning and Fontana, unpubl. data). Although published information is lacking, preliminary data indicates that the 'Yellow-billed' Plumbeous Seedeater has the smallest population of the *Sporophila* seedeaters in southern Brazil (CSF, pers. observ.). As with other seedeaters, the population decline is due to habitat destruction and alteration, as well as capture for the illegal bird trade (Repenning 2012). The entire breeding population of Black-bellied Seedeaters is restricted to the Brazilian Upland Grasslands and faces similar threats (Fontana et al. 2009, Repenning et al. 2010, BirdLife International 2012, Rovedder 2011, ICMBio, in press). Although information about population trends is largely lacking for seedeaters, local declines are also suspected for the remaining species (Bencke et al. 2003, ICMBio, in press).

Among Patagonian migrants, three sheldgeese species (Ruddy-headed Geese *Chloephaga rubidiceps*, Upland Geese *Chloephaga picta*, and Ashy-headed Geese *Chloephaga poliocephala*) that winter in southern Buenos Aires have suffered marked population declines (Chebez 2008) due to hunting (Rumboll 1975, 1979). The mainland population of Ruddy-headed

Geese is thought to consist of <1000 individuals (Blanco et al. 2003, Madsen et al. 2003). Consequently, all three species are considered threatened in Argentina (López-Lanús et al. 2008).

Other grassland birds of SESA are probably also declining due to habitat loss and modification, but lack of data precludes thorough conservation assessments. For example, based on historic and current records for Buenos Aires province, Narosky and Di Giacomo (1993) identified several species thought to have either declined in numbers or disappeared from certain areas, including Greater Rheas, Red-winged Tinamou (*Rynchotus rufescens*), Elegant Crested-Tinamou (*Eudromia elegans*), Spotted Nothuras (*Nothura maculosa*), Plumbeous Ibises (*Theristicus caerulescens*), Rufous-chested Dotterels (*Charadrius modestus*), Least Seedsnipes (*Thinocorus rumicivorus*), and Hudson's Canasteros (*Asthenes hudsoni*). Comparison of the results presented by Narosky and Di Giacomo (1993; prior to the massive planting of transgenic soybeans, Aizen et al. 2009) and information gathered more recently in the Pampas of central Argentina (Codesido et al. 2011) also reveals substantial range contractions of some common grassland species, such as Spectacled Tyrants (*Hymenops perspicillatus*), Great Pampa-Finches (*Embernagra platensis*), and Brown-and-yellow Marshbirds (*Pseudoleistes virescens*).

FACTORS AFFECTING SESA GRASSLAND BIRDS

Factors driving declines in bird populations throughout SESA are diverse and act at various scales, and studies have revealed that grassland birds show different responses to land transformation in the region (Comparatore et al. 1996, Lanctot et al. 2002, Isacch et al. 2005, Azpiroz and Blake 2009, Di Giacomo et al. 2010, Isacch and Cardoni 2011, Codesido et al. 2012). Particularly during the last century, suitable soil and climatic conditions resulted in the transformation of SESA native grasslands into croplands and this has resulted in substantial grassland alteration and fragmentation (Viglizzo et al. 2001, Baldi et al. 2006, Crawshaw et al. 2007).

Crop cultivation and afforestation. Throughout the SESA grasslands, the area devoted to annual crops, sown pastures, and

forestry is increasing (Bilenca and Miñarro 2004, Viglizzo et al. 2005, Martino and Methol 2008). Over the past 15 years, conversion of grasslands to cropland has resulted in loss of habitat for many grassland birds (Vickery et al. 2003, Gabelli et al. 2004, Sarasola et al. 2007, Thompson and Carroll 2009, Codesido et al. 2011). These activities usually result in the total replacement of native vegetation that can trigger changes in bird species richness, composition, and abundance (Fillooy and Bellocq 2007, Pedrana et al. 2008). Conversion to cropland can also affect the persistence of species through increased population isolation (Giordano et al. 2009). Croplands and sown pastures have lower species richness than natural grasslands (Leveau and Leveau 2004, Isacch et al. 2005, Di Giacomo et al. 2010). In other cases, greater numbers of bird species in mixed grazing-cropping systems have been explained by the presence of different vegetation characteristics associated with alternative land uses such as crops and sown pastures (Azpiroz and Blake 2009).

The composition of bird assemblages in croplands and sown pastures differs from that in natural grasslands. In the Northern Campos of Uruguay, facultative grassland species tended to be associated with mixed grazing-cropping systems, but this was not the case for most grassland obligate species (Azpiroz and Blake 2009). In the West Inland Pampa, however, where bird richness is relatively low, replacement of natural grasslands with sown pastures may not have produced substantial changes in the composition of grassland bird assemblages (Isacch et al. 2005).

Grassland birds differ in their sensitivity to conversion of grasslands to cropland and pasture. Several species respond negatively to conversion, including Greater Rheas (Giordano et al. 2009), Long-winged Harriers (*Circus buffoni*; Pedrana et al. 2008), Pampas Pipits (Isacch et al. 2005), Hellmayr's Pipits (*Anthus hellmayri*), Ochre-breasted Pipits (*Anthus nattereri*; Azpiroz and Blake 2009), and Pampas Meadowlarks (Fernandez et al. 2003, Gabelli et al. 2004). Alteration of vegetation structure and composition of native grasslands in croplands and exotic pasturelands (Isacch et al. 2005) can affect bird survival and reproduction (Johnson et al. 2006, Rovedder and Fontana, unpubl. data). Associated activities such as crop harvesting

and mowing can also disturb and kill birds or destroy nests (Fernandez et al. 2003, Giordano et al. 2009). In contrast, other species of birds seem more tolerant of intensive farming systems. For example, Darwin's Nothuras (*Nothura darwinii*), Correndera Pipits (*Anthus correndera*), and Grassland Yellow-Finches (*Sicalis luteola*) use sown pastures in the Inland or Southern Pampas (Leveau and Leveau 2004, Isacch et al. 2005).

Effects of habitat modification due to agriculture on species such as Spotted Nothuras and Swainson's Hawks remain unclear. Some authors have suggested that these species are well adapted to or benefit from such changes (Bucher and Nores 1988, Canavelli et al. 2003), whereas others believe otherwise (Narosky and Di Giacomo 1993, Pinheiro and Lopez 1999, Sarasola et al. 2007, Thompson and Carroll 2009). In any case, habitat use patterns do not necessarily reflect habitat preferences or habitat quality (Krausman 1999). In fact, studies of grassland birds in North America have shown that bird densities may not correlate positively with reproductive success (Vickery et al. 1992).

Use of land for crop production seems particularly detrimental to birds in SESA (Johnson et al. 2011). Whereas 70% of the species that responded to the agricultural gradient were negatively affected by the intensity of crop production, only 22% increased in abundance with a growing percentage of land used for crop production, and none of the latter were typical grassland species (Fillooy and Bellocq 2007). The expansion of agriculture in the last 15 years and simultaneous modification of native grasslands traditionally used for cattle ranching (Bilenca and Miñarro 2004, Martino and Methol 2008) will probably result in further changes in bird population dynamics, including declines in the populations of many species (Isacch et al. 2005, Fillooy and Bellocq 2007, Azpiroz and Blake 2009, Codesido et al. 2011).

Another negative result of agriculture expansion is related to an indirect effect on neighboring cattle-raising areas. In southern Brazil, despite the continuing conversion of grasslands for soybean, wheat, and maize cultivation since 1970, the size of the cattle herd has increased (Crawshaw et al. 2007). Because the area of sown pastures and conversion of other vegetation types into pastures was negligible, Crawshaw

et al. (2007) concluded that the density of cattle increased 20% in the remaining natural grasslands. High cattle densities alter vegetation structure with important consequences for birds (Isacch and Cardoni 2011).

Afforestation on natural grasslands can have adverse effects on water and soil characteristics as well as biodiversity (Jobbágy and Jackson 2004, Brockerhoff et al. 2008, Berthrong et al. 2009, Silveira and Alonso 2009). Plantations of non-native trees can have detrimental effects on birds, especially in grassland-dominated landscapes such as the southern Río de la Plata grasslands (Fillooy et al. 2010). Alternatively, species such as Swainson's Hawks can benefit from availability of novel resources such as stands of introduced exotic trees for use as roost sites (Sarasola and Negro 2006). Introduction of exotic trees may have resulted in the recent expansion of this raptor in the Argentine Pampas (Sarasola and Negro 2006). Other endemic and threatened grassland species, however, can be negatively affected by even small stands of planted trees (Allan et al. 1997). Not surprisingly, the composition of bird assemblages in grasslands and tree plantations differs, with beneficial effects for edge species and negative effects for grassland taxa (Lantschner et al. 2008). Throughout the region, afforestation for pulp-wood affects well-drained grasslands along hillcrests and slopes, threatening a series of common grassland birds and eliminating foraging habitat for Saffron-cowled Blackbirds, Black-and-white Monjitas, and Black-bellied Seedeaters, among others (Fontana et al. 2009, RAD, pers. observ.). Although swales used by these and other species of conservation concern are legally protected and remain free from conversion, little is known about how indirect effects of neighboring plantations affect the physical and biotic characteristics of these wet grasslands, especially alteration of water regimes, facilitation of shrub encroachment, and direct disturbance of open-area species. In South Africa, areas spared from afforestation in plantations are often too small and suffer alteration of vegetation enhanced by the surrounding tree stands and forestry management practices that result in unsuitable habitat for grassland specialists (Lipseý and Hockey 2010). Furthermore, rates of nest predation in open vegetation areas adjacent to tree plantations increase in response to changes in the composition and configuration of the land-

scape, and perhaps also by an increase in abundance of generalist, forest-associated predators (Reino et al. 2010). Licensing and certification of tree plantations in priority areas for conservation, such as Important Bird Areas (Devenish et al. 2009), should be discouraged.

Inappropriate use of agrochemicals.

The quality of agricultural land as bird habitat is reduced by the effects of agrochemicals (Thompson 2004, Zaccagnini 2004, as cited in Bernardos and Zaccagnini 2008). In the Argentinean Pampas, most cropland is subject to misuse of agrochemicals (Zaccagnini 2004). A survey conducted in 1998 indicated that 32% of agricultural lands treated with pesticides represented some risk of mortality for birds (Rivera Milán et al. 2004). Herbicides such as glyphosate are widely used in soy crops to eliminate "weeds" used by birds for perching or foraging and thus can further reduce their presence by simplifying vegetation structure (Leveau and Leveau 2004). In addition, there have been many cases of bird mortality where organophosphate insecticides such as monocrotophos were likely involved (Hooper et al. 2003). Direct mortality has been documented for several grassland birds and, for some long-distance migrants such as Ruddy-headed Geese, Swainson's Hawks, Upland Sandpipers, and Bobolinks, agrochemicals may be one of the main threats in non-breeding areas (Woodbridge et al. 1995, Blanco et al. 2003, Vickery et al. 2003, 2010, Bernardos and Zaccagnini 2008). Swainson's Hawks have probably been affected most because they feed on grasshoppers targeted for chemical control (Woodbridge et al. 1995). During 1995–1996, ~20,000 individuals (5% of the world's population) were killed by inappropriate use of organophosphate insecticides (Goldstein et al. 1999a, b). Although monocrotophos was banned in 1999, other highly toxic organophosphate products are still used in the Río de la Plata grasslands (Hooper et al. 1999, 2003). Bobolinks are considered agricultural pests and are actively killed with pesticides in rice fields in Argentina (López-Lanús and Marino 2010). Migratory shorebirds that also use rice fields are exposed to agrochemicals (Blanco et al. 2006). Although evidence of acute exposure has not been recorded, sub-lethal effects were observed in Buff-breasted Sandpipers exposed to organophosphate and carbamate in Argentinean rice fields (Strum et al. 2010).

Livestock ranching and fire management. Because cattle grazing changes vegetation structure and composition (Sala et al. 1986, Pillar and Quadros 1997), grazing management can affect bird diversity (Isacch and Martínez 2001, 2003b, Zalba and Cozzani 2004, Isacch et al. 2005, Azpiroz and Blake 2009, Di Giacomo et al. 2010, Isacch and Cardoni 2011). In the West Inland Pampa, moderate grazing of native grasslands can increase structural heterogeneity and plant species richness relative to undisturbed grasslands, resulting in greater bird species richness and abundance (Isacch et al. 2003). Apart from the effects of grazing on habitat quality, livestock can also affect bird populations through direct disturbance of individuals (Thompson and Carroll 2009) and nest trampling (Azpiroz 2000, Zalba and Cozzani 2004, A. Di Giacomo, unpubl. data).

Effects of cattle grazing on SESA grassland birds vary. In the Flooding Pampa, ungrazed or lightly grazed sites are dominated by tall-grass species of birds such as Bay-capped Wren-Spinetails (*Spartonoica maluroides*) and Grass Wrens (*Cistothorus platensis*), whereas moderately grazed areas favor generalist tall grass species such as Great Pampa-Finches, Grassland Yellow-Finches, and Brown-and-yellow Marshbirds (Isacch and Cardoni 2011). In heavily grazed grasslands, a short-grass assemblage of birds is present, including Bar-winged Cinclodes (*Cinclodes fuscus*), Rufous-backed Negritos (*Lessonia rufa*), and Southern Lapwings (*Vanellus chilensis*; Comparatore et al. 1996, Isacch and Cardoni 2011). The presence of Buff-breasted Sandpipers and other Patagonian shorebirds in coastal halophytic short grasslands is directly linked to the presence of livestock grazing (Lancot et al. 2002, 2004, Isacch and Martínez 2003b, Blanco et al. 2004, Alfaro et al. 2008, Isacch and Cardoni 2011).

Grassland specialists such as Hudson's Canasteros, Black-and-white Monjitas, and Pampas Meadowlarks seem to favor a mosaic of tall and short vegetation that is usually maintained by moderate grazing pressure. These species feed in patches of short grass and nest in tall herbaceous vegetation (Fontana 1994, Isacch et al. 2001, Fraga 2003, Isacch and Cardoni 2011, Azpiroz, unpubl. data).

Several globally threatened grassland birds in our study region depend on tall herbaceous vegetation for feeding, breeding, and roosting.

Strange-tailed Tyrants are scarce in grazed grasslands of northeast Argentina, and the absence of livestock in some protected areas such as Rio Pilcomayo National Park and Iberá Natural Reserve has resulted in recovery of their populations (Di Giacomo 2005a, Di Giacomo et al. 2010). Areas with light or no grazing are necessary to ensure viable populations of specialized tall-grass species (Cardoni 2011, Isacch and Cardoni 2011) and these areas could act as "source" areas able to repopulate neighboring sites (Zalba and Cozzani 2004). Ungrazed or lightly grazed grasslands are also important for other tall-grass birds such as Bearded Tachuris (*Polystictus pectoralis*), Sharp-tailed Grass Tyrants, and seedeaters (*Sporophila* spp.) throughout SESA (Codesido and Fraga 2009).

In some SESA grasslands, especially in northern Argentina and the Brazilian Upland grasslands, fire is used as a management tool, mainly to improve forage quality for cattle (Isacch et al. 2004, Boldrini et al. 2009, Di Giacomo et al. 2011a). Fire frequency and intensity also affect vegetation and have indirect effects on bird assemblages, especially on species composition (Comparatore et al. 1996, Isacch and Martínez 2001). The timing of burns is also important because burning during the breeding season can have negative effects on reproductive success (Petry and Kruger 2010, Di Giacomo et al. 2011a). Burning can also affect availability of roost sites of gregarious species (López-Lanús and Marino 2010). In addition, excessive burning may reduce habitat availability for threatened species such as Bay-capped Wren-Spinetails (Isacch et al. 2004). Other species (Southern Lapwings and Correndera Pipits), however, use burned fields shortly after fires (Isacch et al. 2004). Fire can also affect habitat use during the breeding season. Strange-tailed Tyrants and Black-bellied and "Yellow-billed" Plumbeous Seedeaters avoid burned plots for nesting (Di Giacomo et al. 2011a, Rovedder 2011, Repening 2012). For Strange-tailed Tyrants, relocation to nesting areas in nearby unburned sites does not affect reproductive success, suggesting the species is adapted to regular fires (Di Giacomo et al. 2011a). During the third breeding season after a fire, individuals did not discriminate between burned and unburned sites. Thus, for Strange-tailed Tyrants, burning regimes in a given grassland patch must guarantee availability of unburned areas, and burning intervals longer

than two years should be considered. Unfortunately, most current fire management practices in northeast Argentina involve annual burning.

Change in interspecific relationships.

Grassland birds do not always use all available suitable habitat (Fernández et al. 2003, Gabelli et al. 2004, Giordano et al. 2009). The presence of competitors, predators, or brood parasites may limit use of sites otherwise suitable (Fernández et al. 2003, Gabelli et al. 2004). For example, the recent expansion of the ranges of Pampas Meadowlark congeners (White-browed Blackbirds, *Sturnella superciliaris*, and Long-tailed Meadowlarks, *S. loyca*) may have had adverse effects on the distribution and abundance of this threatened species (Gabelli et al. 2004). In addition, intense agriculture and livestock grazing have contributed to an expansion of the range and increase in numbers of Shiny Cowbirds (*Molothrus bonariensis*), and this brood parasite may exert further pressure on declining populations of grassland specialists such as Pampas Meadowlarks, Black-and-white Monjitas, and Saffron-cowled Blackbirds (Fraga et al. 1998, Azpiroz 2000, Gabelli et al. 2004, Zalba and Cozzani 2004). Because cowbirds associate with cattle and can move long distances between feeding and breeding areas, the negative consequences of brood parasitism may reduce the value of small grassland patches (Zalba and Cozzani 2004). Grazing and crop production can also facilitate tree propagation in grasslands and this may also contribute to increased numbers of predators and brood parasites (Isacch et al. 2005).

Urban development and sewage pollution. Although the area covered by urban settlements and transportation networks is relatively small (Baldi et al. 2006), habitat loss due to urbanization appears to have resulted in the local extinction of populations of some grassland birds. This is probably the case for Black-and-white Monjitas, with extensive urban development along the Río de la Plata coastline destroying habitats and eliminating local populations. The current isolation of populations in Buenos Aires Province may have been the result of urban development (Fraga 2003).

An experimental study looking at the behavior of grassland birds in a suburban setting (*Cortaderia selloana* grassland in the Pampas) showed that grassland specialists tend to avoid feeding near unfamiliar objects

(Echeverría et al. 2006). In addition, grassland generalists exhibit greater neophobia than generalist urban species (Echeverría et al. 2006). These factors may explain in part the greater vulnerability of grassland birds to urban development.

Bird species richness and abundance may increase near nutrient-discharge sources in cordgrass-dominated communities. In the Southern Pampa, nutrient enrichment from sewage input resulted in alterations to vegetation structure that positively affected bird diversity, especially common and generalist grassland species (Cardoni et al. 2011). Nutrient management, however, is key to avoiding habitat degradation that would have negative effects on birds, such as favoring conditions for parasites (Cardoni et al. 2011). Ecological trap conditions may also be an issue. Although these nutrient-rich sites can be attractive for grassland birds because of increased food availability, they may provide poor breeding conditions for species nesting near or on the ground and vulnerable to flooding (Cardoni et al. 2011).

Trapping and hunting. Some SESA grassland birds are illegally captured and hunted (Chebez 2008). In Rio Grande do Sul, populations of 18% of threatened bird species are affected by hunting and trapping (Bencke et al. 2003). These activities are conducted even in protected areas. Although the impact of illegal hunting and trapping has not been assessed, these are common and widespread activities throughout the region (Chebez 2008, BirdLife International 2012, RAD and CSF, pers. observ.).

Greater Rheas are hunted for their meat, feathers, and skin and are also hunted as agricultural pests in some areas because of their negative impacts on crops and the assumption that they compete with cattle for forage (Pereira et al. 2003, Bellis et al. 2004a, b, Herrera et al. 2004). Diet studies have shown, however, that the latter assumption is not justified (Martella et al. 1996, Comparatore and Yagueddú 2007). In Rio Grande do Sul and Uruguay, rhea eggs are gathered for consumption (CSF and ABA, pers. observ.).

Three species of sheldgeese (Ruddy-headed Geese, Upland Geese, and Ashy-headed Geese) have also been persecuted as agricultural pests in southern Buenos Aires province (Blanco et al. 2003). Birds have been hunted and even

scared away with aircraft (Blanco et al. 2003). Although sheldgeese are currently protected by law, hunting is still a threat (Chebez 2008).

Other grassland birds are illegally captured for the cage-bird trade and sold at wildlife markets. Saffron-cowled Blackbirds are targeted in Argentina and Uruguay, and trappers have been responsible for the disappearance of entire colonies (Chebez 2008). Several species of *Sporophila* seedeaters are globally threatened because their small or geographically restricted populations are subject to high rates of capture for the illegal wildlife market (BirdLife International 2012). An important issue is that mature male seedeaters are targeted and this may have direct demographic implications (e.g., *S. aff. plumbea* in southern Brazil; Fontana et al. 2009).

Other threats. Grassland birds face additional threats, the effects of which have not been adequately quantified. Transportation networks may represent a potential hazard to tall-grass specialists in areas where suitable habitat is largely available as vegetation strips along roads. Grassland birds that forage and nest near roads are prone to accidental collisions with vehicles. Additionally, collisions with towers and transmission lines can affect negatively resident and, especially, migratory grassland birds in Rio Grande do Sul (Fontana et al., unpubl. data). Greater Rheas have also suffered losses due to entanglement in wire fences (ABA and RAD, pers. observ.).

Severe weather sometimes causes breeding failure and additional mortality. Rainstorms can affect ground-nesting grassland birds such as Greater Rheas, Spotted Nothuras, Ochre-breasted Pipits, and Pampas Meadowlarks (ABA, pers. observ.). Severe hailstorms are frequent in Pampas subregions of Argentina and have been responsible for mass mortality of Swainson's Hawks (Sarasola et al. 2005). The impact of these causes of mortality may be significant when combined with the effects of other negative factors (Sarasola et al. 2005).

In northern Argentina and southern Paraguay, construction of the Yacyretá Dam has affected local populations of threatened grassland birds, such as Saffron-cowled Blackbirds, Cock-tailed Tyrants, and Black-masked Finches, that have almost disappeared from the Argentine side of the Paraná River and nearby islands after extensive areas of grassland were inundated (Di Giacomo 2005b, Chebez 2008, Codesido and

Fraga 2009). In southern Brazil, construction of a series of dams is also expected in the next few years (Hüffner and Engel 2011). Construction of the Paiquerê hydroelectric dam on the Pelotas River would be especially problematic because it would seriously imperil populations of "Yellow-billed" Plumbeous Seedeaters and other threatened grassland birds (Fontana et al. 2009).

Construction of small impoundments is also widespread in Rio Grande do Sul and southeastern Uruguay to ensure water availability for agriculture and cattle ranching (Azpiroz and Rilla 2007, RAD, pers. observ.). These impoundments are frequently built in grassy swales used by Black-and-white Monjitas, Saffron-cowled Blackbirds, and *Sporophila* seedeaters (Dias and Maurício 2002, Bencke et al. 2003).

Expansion of wind energy in southern Brazil may also have negative impacts. Installation of several wind farms is planned throughout the region, especially in the upland grasslands of Santa Catarina and Rio Grande do Sul and in coastal grasslands in the latter state (Camargo 2002, RAD, pers. observ.). Birds are known to collide with wind turbines and power lines and can be negatively affected by the disturbance caused by turbines (Langston and Pullan 2003, Barrios and Rodríguez 2004). Because most negative impacts of wind farms on birds can be prevented by carefully planning their locations (Langston and Pullan 2003), adequate zoning of wind-development facilities is needed in southern Brazil.

SOUTH AMERICAN GRASSLAND BIRDS: KNOWLEDGE AND GAPS

We identified 60 papers about SESA grassland birds published in high-impact journals from 1999 to 2011, with most (52, or 87%) published since 2004. However, geographic and topic biases were evident. Most studies (48, or 80%) were conducted in Argentina. Studies dealing with habitat and land-use associations predominated (23, or 38%). Other topics that were the focus of multiple papers included biology (including feeding, breeding and social behaviors; 13, or 21.7%), conservation (6, or 10%), geographic distribution (4, or 6.7%), assemblages and community ecology (3, or 5%), genetics (3, or 5%), population monitoring and migration (3, or 5%), and toxicology (2, or 3.3%). Climate and weather impacts, methods,

and morphology were each the focus of one paper.

Studies in SESA have provided information about the distribution of grassland birds, factors affecting their populations, and habitat requirements of some threatened species. Studies of the effects of habitat fragmentation and area sensitivity patterns are, however, lacking. These topics are especially relevant to reserve design and land-management planning. Because SESA grasslands have been subject to large-scale modification, the possible role of small grassland patches in sustaining bird assemblages must be evaluated (Zalba and Cozzani 2004). These analyses should consider land-use matrix characteristics, which can have a significant influence on processes within patches (Prevedello and Vieira 2010). More data concerning how habitat transformation and edge effects impact reproductive success are also needed (Fernández et al. 2003, Zalba and Cozzani 2004, Johnson et al. 2011), as well as assessments of the genetic variability of isolated populations of threatened birds.

Given the current state of grasslands in some subregions of SESA, programs to enhance recovery of almost extinct vegetal communities and their associated fauna should be implemented. Data are available concerning how grassland vegetation responds to different land uses and environmental conditions, and this knowledge could be used to assess the possibility of restoring native conditions (Aguilera et al. 1998, Laterra et al. 1998, León and Burkart 1998). Unfortunately, information concerning how grassland birds might respond to restoration efforts and how factors such as distance to source areas and size of restored patches might influence outcomes is lacking (Fuhlendorf and Engle 2001).

Most studies of SESA grassland birds have been short-term, resulting in a lack of long-term data about species and their populations. Some studies, however, have revealed significant inter-annual variation in grassland bird populations (Isacch and Martínez 2003a), highlighting the need for volunteer-based, long-term monitoring programs such as North America's Breeding Bird Surveys and Christmas Bird Counts. Such initiatives could provide data needed to distinguish between natural fluctuations and human-induced declines (Butcher et al. 1990), and could also be useful for monitoring populations of species of conservation concern

(Gabelli et al. 2004, Sarasola et al. 2007). A good example of this is the Nearctic Grassland Shorebird Survey that targets plovers and sandpipers that winter in the SESA region and is sponsored by the Alliance for the Grasslands (<http://www.pastizalesdelconosur.org>). Another example is the annual monitoring of birds in agroecosystems of east-central Argentina initiated in 2002 by the National Agency for Agricultural Technology – INTA (Canavelli et al. 2004, Schrag et al. 2009). More monitoring programs are also needed in protected areas to determine population trends and estimate the viability of threatened species in reserves through population modeling (Beissinger and Westphal 1998, Brook et al. 2000, Prugh et al. 2008).

Population and ecological studies of threatened species that expand our knowledge of dispersal and migration patterns, mating systems, and genetic variability, among others, will be key for conservation management. Factors that influence habitat use patterns and species survival in both natural and agricultural landscapes in SESA are poorly known (Thompson 2004, Filloy and Bellocq 2007, Medan et al. 2011). In addition, most available information is based solely on presence/absence data or species' density estimates. Such data, however, may not be informative about habitat preferences or quality for grassland birds (Vickery et al. 1992, Thompson 2004). With the exception of a few species (e.g., Strange-tailed Tyrants, Di Giacomo et al. 2011b), basic information about the population status and reproduction of SESA grassland birds is lacking (Vickery et al. 1999, Medan et al. 2011), precluding an adequate assessment of population viability in given habitats or sites (Morris and Doak 2002, Giordano et al. 2009).

Ecology-oriented molecular studies of Neotropical grassland birds are scarce. These could shed light on several issues directly related to management and conservation. For example, for small, isolated populations, increased variance in reproductive success may reduce effective population size (Nunney 1993, 1996, Hedrick 2005). More taxonomic studies are also needed because the status of several grassland species, particularly in the genus *Sporophila*, is still unclear (Areta 2008, Machado and Silveira 2011).

By considering the threatened status of SESA grasslands, future research programs should also prioritize studies that promote establishment of

creative and practical measures for conserving grassland birds in economically productive contexts. Large-scale experimental studies focusing on the effects of land-use practices (e.g., fire regimes, cattle grazing rotation rates, and urbanization) should be developed (Thompson 2004, Sarasola *et al.* 2007) by cooperating researchers, non-governmental organizations, and rural and territorial planners (Isacch and Cardoni 2011). There is also a need to determine the real magnitude of crop and pasture losses caused by birds considered pests (Blanco *et al.* 2003) and to quantify impacts of agrochemicals on bird populations (Vickery *et al.* 2010).

Human impacts affecting grassland birds in SESA are often the same irrespective of borders. Thus, there is an urgent need to promote and implement coordinated studies designed to assess these impacts and offer management tools in a broader context throughout the entire SESA region.

CURRENT BIRD CONSERVATION INITIATIVES

Current conservation efforts include elaboration of action plans, establishment of public and private reserves, promotion of agricultural activities that reconcile production with biodiversity conservation, and development of multilateral conservation projects across countries. The Brazilian government has recently completed an action plan for the conservation of grassland birds must be implemented in five years (ICMBio, *in press*). Because species share common threats, the plan assumes that well-planned actions can optimize conservation and benefit entire assemblages (ICMBio, *in press*). Actions include the implementation of reserves, empowerment of law reinforcement agencies, and development of land-use ordination schemes (ICMBio, *in press*).

The restricted network of public protected areas in the region has made a limited contribution to grassland conservation within SESA (Bilenca and Miñarro 2004, Azpiroz and Rilla 2007, Overbeck *et al.* 2007). Together, these areas protect ~1% of the land originally covered by grasslands. In Uruguay, no current or proposed sites of the National Protected Area Systems (DINAMA/MVOTMA 1999, Ghione and Martino 2008) include large continuous tracts (>5000 ha) of native grasslands. Only

1.48% and 2% of the areas originally encompassed by grasslands are protected in Rio Grande do Sul and Paraná states, Brazil, respectively (Brandão *et al.* 2007, Santos and Scherer-Neto 2009). Protected grasslands in southern Brazil are not managed adequately and suffer from shrub encroachment, reducing their value for grassland species (Pillar and Vélez 2010). There have been several initiatives to identify sites of conservation priority in grassland ecosystems (Bilenca and Miñarro 2004, Di Giacomo 2005b, MMA 2007, Fontana *et al.* 2008, Di Giacomo and Parera 2008, Cartes and Clay 2009, Develey and Goerck 2009), but there has been little progress toward increasing the amount of protected habitat. In 2010, Brazil established a goal of conserving 10% of all its biomes (Vélez *et al.* 2009). To achieve this goal, creation of 21 new protected areas encompassing 979,966 ha was recommended in grassland regions of southern and western Rio Grande do Sul state (E. Vélez, *pers. comm.*), but the process has been placed on hold by the Brazilian Environmental Ministry.

In southern Brazil, licensing renewal for the Barra Grande Dam required creation of an ecological corridor along the Pelotas River (Refugio de Vida Silvestre Corredor do Rio Pelotas). The goal is to connect the trough region of the river and its main tributaries with protected areas such as São Joaquim and Aparados da Serra National Parks through creation of a 270,000-ha conservation unit. The area would protect about 46% of the natural Brazilian Upland grasslands in Rio Grande do Sul and Santa Catarina states (APREMAVI 2012), and would include three of four grassland-bird priority areas identified by Fontana *et al.* (2008) in southeastern Brazil. This initiative is now under consideration by the Brazilian Government.

Apart from habitat protection in public reserves, grassland bird conservation is also occurring on private lands. There are 73,229 ha of private reserves in southern Brazil, but most protect forests and other vegetation types (Vélez *et al.* 2009). Furthermore, management plans are seldom elaborated and implemented (Vélez *et al.* 2009), and private reserves tend to be concentrated in certain regions. A few private reserves, however, play important roles in the conservation of threatened grassland birds. For example, El Bagual Reserve in Formosa Province, Argentina, has breeding populations

of several globally threatened species, and these birds have been studied and monitored since the 1990s through an agreement between landowners and the NGO Aves Argentinas (Di Giacomo 2005a). In Corrientes, Argentina, The Conservation Land Trust acquired >150,000 ha of critical habitat for globally threatened grassland birds to be appended to the Reserva Provincial del Iberá that covers ~700,000 ha of mainly flooded areas (Di Giacomo 2005a).

Nearly all large grassland remnants in SESA are privately owned, including most priority areas for grassland bird conservation. Most are rangelands used for cattle, sheep, and horse ranching, where native grasslands are threatened by inappropriate management or risk of conversion to cropland or forests (Bilenca and Miñarro 2004).

There are legal measures that protect natural vegetation on private lands, but most are not followed or enforced (Vélez et al. 2009). In Brazil, the “Código Florestal” (Forest Code) is perhaps the most important legal instrument to regulate land use on private lands (Vélez et al. 2009). Economic use of legal reserves (one of the main conservation instruments of the code) is allowed as long as natural vegetation is not altered; initiatives to promote sustainable cattle production in these areas are being developed (IBAMA 2009). Unfortunately, the code is currently being revised to address issues raised by agricultural interests and its future is uncertain (Metzger et al. 2010).

There is currently interest in promoting wildlife-friendly practices in agricultural lands by governmental and non-governmental institutions throughout SESA (Marino 2008). This is especially important in SESA grasslands, where establishment of state-owned reserves is precluded by financial and social issues related to land expropriation. In the case of livestock ranching, sustainable practices can increase profitability of the beef industry and ensure maintenance of rangelands and grassland biodiversity (Netto 2009).

An example of this type of initiative at a regional level is the Alliance for the Grasslands (www.pastizalesdelconosur.org) created by several NGOs in 2006. Today, the Alliance involves more than 40 institutions representing different social sectors, with an emphasis on ranchers and conservationists from all countries in the SESA grasslands. One objective of the Alliance is to

promote certified beef production in natural grasslands.

Other initiatives with similar objectives are also being encouraged. Researchers and technicians from Argentina, Brazil, and Uruguay recently formed a technical working-group to develop guidelines for sustainable livestock ranching on natural grasslands (Vélez et al. 2009). In Uruguay, the Responsible Rural Production Program (PPR Program; www.cebra.com.uy/presponsable/) is sponsored by the Uruguayan government and promotes application of integrated and efficient management systems that use natural resources, including biological diversity. For grassland birds, the latter has provided financial support to private ranchers for actions that favor habitat features important to threatened species (ABA, pers. observ.). In Argentina, a recent (2010) GEF-funded project with similar aims has already produced management guides and educational material for farmers. In Brazil, there are also at least seven programs focusing on promoting sustainable livestock raising on natural grasslands (Vélez et al. 2009).

Another multilateral initiative is a grassland bird conservation agreement recently developed under the auspices of the Convention on Migratory Species (CMS) and signed by governments of all four countries of the SESA region (and Bolivia). The aim of this agreement is to address the needs of threatened migratory grassland birds that cross international borders in southern South America. Through development of an action plan, specific research and conservation priorities have already been identified. These activities will require substantial funding, and it is unclear how financial support will be obtained.

Despite the many conservation projects and initiatives discussed above, prospects for native grasslands remain grim and few concrete actions have been taken to prevent loss of grassland birds. Although there are many gaps in our knowledge of grassland bird biology and ecology that could improve the situation, we agree with Vélez et al. (2009) that the lack of programs to promote conservation and public and political support are probably the main reasons why little has been accomplished. More must be done to reverse the existing negative trends of grassland bird populations and guarantee the persistence of threatened birds over the long term.

ADDITIONAL PROPOSED CONSERVATION MEASURES

Several grassland birds depend on grasslands unaltered by agriculture or cattle grazing (Zalba and Cozzani 2004, Giordano et al. 2009, Isacch and Cardoni 2011). The current public reserve system, however, offers limited opportunities for preservation of such grasslands (Fraga 2003). As discussed above, opportunities to create new reserves are generally limited. In addition, the characteristics of species can limit the effectiveness of protected areas. For example, Saffron-cowled Blackbirds move over extensive areas in search of foraging sites during the non-breeding season. These erratic movements represent a challenge to protection efforts based on establishing small reserves restricted to breeding areas (Fraga et al. 1998). The size of new reserves is also critical because small patches of grassland may not guarantee maintenance of bird diversity (Zalba and Cozzani 2004).

Because most populations of threatened grassland birds in SESA are on private lands, recent interest in promoting improved agricultural practices is encouraging. Land management programs in agricultural areas need to focus on reconciling the interests of producers and conservationists to preserve habitat features similar to those of native grasslands (Giordano et al. 2009). Several species depend on protection of nesting sites and on adequate field and grazing management policies, including careful use of fire (Fernandez et al. 2003, Vickery et al. 2010, Di Giacomo et al. 2011a,b). There is also a need for regional strategies and planning because, at large scales, grassland bird diversity can be enhanced by maintaining a mosaic of patches with different cattle-grazing intensities (Isacch and Cardoni 2011, Johnson et al. 2011).

Government agencies should also promote best cultivation practices, with special attention on reducing use of agrochemicals (Vickery et al. 2010). Use of the Ecotoxicologic Risk Calculator (Zaccagnini et al. 2005, Bernardos et al. 2007), developed by the National Institute of Agricultural Technology (INTA) of Argentina, should be promoted. This software tool calculates the risk of poisoning for birds exposed to agrochemicals by considering the kind of agrochemical, its dosage and concentration, and crop type (Bernardos and Zaccagnini 2008).

Management measures that result in significant losses to landowners, such as land set-asides, should include mechanisms for economic compensation (Isacch 2008). Management alternatives that benefit targeted grassland species should also be promoted. The Uruguayan PPR project provides a good example of public-private partnerships working towards native grassland conservation. Conversely, economic incentives for agriculture and forestry in grassland priority areas should be revised.

Governments should also evaluate the possibility of establishing grassland easements like those in North America (Vickery et al. 2010, Johnson et al. 2011). These agreements could be particularly useful in largely modified landscapes and areas with highly threatened bird populations. One way to compensate for land set-asides could be through ecotourism development. Some private initiatives are being developed on Argentinean ranches (JPI, pers. observ.), and others have been recommended for application in Brazilian Upland Grasslands (Fontana et al. 2009). In Rio Grande do Sul, for example, eco-touristic regions and routes are being identified by the state's tourism agency with support from the private sector (Vélez et al. 2009; RAD, pers. observ.). There is a huge potential for expansion of bird-watching and bird photography in the SESA grasslands, but much remains to be done to encourage these activities. Training and capacity-building are also needed to provide local people with the necessary tools to apply such bird-friendly alternative practices (Johnson et al. 2011).

Land-use planning is still incipient in the region. Many activities that alter grasslands could be constrained if land-use regulations are implemented. In Rio Grande do Sul, an environmental zoning scheme for the wood industry was recently completed. Despite some important limitations, this scheme has incorporated objective technical criteria to ensure the environmental viability of this activity in the state (Vélez et al. 2009). Studies are also needed to better solve energy and water demands in a way that avoids irreversible damage to bird populations such as that caused by the Barra Grande Dam in southern Brazil (Fontana et al. 2009).

Given the conservation status of many long-distance migrants, development and implementation of effective conservation strategies will

also require international collaboration (Woodbridge et al. 1995). For example, several SESA grassland passerines winter in the Cerrado region of central South America where they suffer similar threats (Vickery et al. 1999, Lopes et al. 2010). It would be particularly helpful if governments of the SESA countries could provide much-needed funding that would allow application of specific measures detailed in the new CMS action plan that focus on threatened migratory birds. Most of these actions would also benefit many other grassland species.

Many aspects of bird conservation in SESA grasslands would also benefit from public awareness programs. These could not only reduce threats posed by hunting, trapping, and agrochemical abuse, but would help build political support for conservation initiatives. Communication networks (e.g., web-based) between researchers, conservationists, managers, and farmers are needed to guide decision-making about agricultural practices, promote law enforcement, and manage wildlife in agro-ecosystems (Rivera Milán et al. 2004, Medan et al. 2011, Johnson et al. 2011). Public campaigns, targeted at farmers and other relevant stakeholders, would help raise awareness about the importance of conserving grasslands and grassland birds (Vickery et al. 2010).

CONCLUSIONS

Although grassland birds do not respond uniformly to habitat change, current land-use trends toward conversion to high-intensity croplands in some regions will likely result in further declines in bird populations (Fillooy and Bellocq 2007). In the case of livestock grazing, SESA grassland birds likely evolved in ecosystems subjected to moderate grazing regimes and frequent fires, and these conditions must be recreated to provide adequate habitat for certain species (Isacch et al. 2004, Zalba and Cozzani 2004, Isacch and Cardoni 2011). At large scales, grassland bird diversity can be enhanced in some regions by maintaining a mosaic of patches with different cattle-grazing intensities (e.g., Bahía Samborombón grasslands, Isacch and Cardoni 2011; Iberá grasslands, A. S. Di Giacomo, unpubl. data). Several grassland species have suffered substantial range contractions and subsist as small isolated populations. Conservation of these species will require effective action

on both public and private lands. The burden of protecting grassland birds, however, should not be put solely on farmers; national governments must assume responsibility for long-term conservation of grassland-associated biodiversity (Isacch 2008, Johnson et al. 2011).

The response of species to diverse habitat-altering factors and threats can vary geographically so management actions should consider these regional differences. In addition, many species do not use all available suitable habitats. This has important implications for reserve design and management because unoccupied habitats may still be valuable. In addition, presence/absence data or even abundance patterns may be misleading regarding long-term persistence of bird populations, especially for small patches of native vegetation or highly modified grasslands (e.g., crops). In fact, it is not clear if small areas of grassland are areas with high breeding success (sources) or whether they are ecological traps (Zalba and Cozzani 2004). Agro-ecosystems could also be acting as ecological traps for grassland-specialist birds (Giordano et al. 2009).

Although our knowledge of grassland birds in SESA is increasing, additional studies are needed to improve our understanding of the ecology of grassland birds in the region. However, we are also aware that recent land-use changes threaten bird populations as never before. Successful recovery of threatened populations will depend in part on our ability to generate key ecological data. More importantly, rapid and effective management strategies are needed to improve the status of threatened species and grassland bird assemblages. We hope this review will stimulate both additional research and effective conservation action.

ACKNOWLEDGMENTS

G. A. Bencke provided valuable insight regarding several topics during the initial phases of manuscript preparation. We are also grateful to I. Boldrini, P. M. A. Ferreira, H. Hasenack, R. B. Setubal, and E. Vélez for sharing their expertise and information concerning grassland regional status, conservation strategies, and other topics. We thank R. Clay for his comments on Paraguayan grasslands. We also thank G. Ritchison, P. Vickery, M. Codesido, and two anonymous reviewers for comments and suggestions that significantly improved our manuscript. ABA's work was funded by the Rufford Small Grants Foundation, the Wildlife Conservation Society, Cleveland Metroparks Zoo, the Neotropical Grassland Conservancy and the

Instituto de Conservación Neotropical. JPI's work is funded by CONICET, Universidad Nacional Mar del Plata, Beca "Conservar la Argentina" (Aves Argentinas). RAD is funded by Universidade Católica de Pelotas, SAVE Brasil, and CAPES. CSF is supported by Fundação Grupo Boticário de Proteção a Natureza, Neotropical Grassland Conservancy, CNPq, Igré- Associação Sócio Ambientalista, SAVE Brasil, and PUCRS. ASDG is funded by CONICET—Universidad de Buenos Aires and Aves Argentinas, and supported by Alparamis S.A. and Conservation Land Trust.

LITERATURE CITED

- AGUILERA, M., D. STEINACKER, M. DEMARÍA, AND A. AVILA. 1998. Estados y transiciones de los pastizales de *Sorghastrum pellitum* del área mediana central de San Luis, Argentina. *Ecotrópicos* 11: 107–120.
- AIZEN, M. A., L. A. GARIBALDI, AND M. DONDO. 2009. Soybean expansion and agriculture diversity in Argentina. *Ecología Austral* 19: 45–54.
- ALFARO, M., A. AZPIROZ, T. RABAU, AND M. ABREU. 2008. Distribution, relative abundance, and habitat use of four species of Neotropical shorebirds in Uruguay. *Ornitología Neotropical* 19 (Suppl): 461–472.
- ALLAN, D. G., J. A. HARRISON, R. A. NAVARRO, B. W. VAN WILGEN, AND M. W. THOMPSON. 1997. Impact of commercial afforestation on bird populations in Mpumalanga Province, South Africa—insights from bird-atlas data. *Biological Conservation* 79: 173–185.
- APREMAVI [online]. 2012. Corredor Rio Pelotas. <<http://www.apremavi.org.br>> (16 March 2012).
- ARBALLO, E., AND J. C. GAMBAROTTA. 1987. Registro de la Tijereta de las Pajas para el Uruguay. *Nuestras Aves* 5: 16–17.
- ARETA, J. I. 2008. Entre Ríos Seed-eater (*Sporophila zelichi*): a species that never was. *Journal of Field Ornithology* 79: 352–363.
- ASKINS, R. A., F. CHÁVEZ-RAMÍREZ, B. C. DALE, C. A. HAAS, J. R. HERKERT, F. KNOFF, AND P. D. VICKERY. 2007. Conservation of grassland birds in North America: understanding ecological processes in different regions. *Ornithological Monographs* 64: 1–46.
- AZPIROZ, A. B. 2000. Biología y conservación del Dragón (*Xanthopsar flavus*, Icteridae) en la Reserva de Biosfera Bañados del Este. PROBIDES, Rocha, Uruguay.
- . 2005. Conservation of Pampas Meadowlark (*Sturnella defilippii*) in Uruguay. *Cotinga* 23: 71–73.
- AZPIROZ, A. B., AND J. G. BLAKE. 2009. Avian assemblages in altered and natural grasslands in the Northern Campos of Uruguay. *Condor* 111: 21–35.
- , AND F. RILLA MANTA. 2007. Biodiversity and conservation in the Bañados del Este, southeastern Uruguay. In: Proceedings of the waterbirds around the world conference (G. C. Boere, C. A. Galbraith, AND D. A. Stroud, eds.), pp. 186–187. The Stationary Office, Edinburgh, UK.
- BALDI, G., J. P. GUERSHMAN, AND J. M. PARUELO. 2006. Characterizing fragmentation in temperate South America grasslands. *Agriculture, Ecosystems and Environment* 116: 197–208.
- , M. D. NOSETTO, M. R. ARAGÓN, F. AVERSA, J. M. PARUELO, AND E. G. JOBBÁGY. 2008. Long-term satellite NDVI datasets: Evaluating their ability to detect ecosystem functional changes in South America. *Sensors* 8: 5397–5425.
- , AND J. M. PARUELO. 2008. Land-use and land cover dynamics in South American temperate grasslands. *Ecology and Society* 13: 6.
- BARRIOS, L., AND A. RODRIGUEZ. 2004. Behavioural and environmental correlates of soaring-bird mortality at on-shore turbines. *Journal of Applied Ecology* 41: 72–81.
- BEHLING, H., V. D. P. PILLAR, AND S. G. BAUERMANN. 2005. Late Quaternary grassland (Campos), gallery forest, fire and climate dynamics, studied by pollen, charcoal and multivariate analysis of the São Francisco de Assis core in western Rio Grande do Sul (southern Brazil). *Review of Palaeobotany and Palynology* 133: 235–248.
- BEISSINGER, S. R., AND M. I. WESTPHAL. 1998. On the use of demographic models of population viability in endangered species management. *Journal of Wildlife Management* 62: 821–841.
- BELLIS, L. M., M. B. MARTELLA, AND J. L. NAVARRO. 2004a. Habitat use by wild and captive-reared Greater Rheas in agricultural landscapes. *Oryx* 38: 304–310.
- , ———, AND ———. 2004b. Home range of Greater and Lesser rhea in Argentina: relevance to conservation. *Biodiversity and Conservation* 13: 2589–2598.
- BELTON, W. 1994. Aves do Rio Grande do Sul, distribuição e biologia. Unisinos, São Leopoldo, Brazil.
- BENCKE, G. A. 2001. Lista de referência das aves do Rio Grande do Sul. Porto Alegre, Fundação Zoobotânica do Rio Grande do Sul, Brazil.
- , R. A. DIAS, L. BUGONI, C. E. AGNE, C. S. FONTANA, G. N. MAURÍCIO, AND D. B. MACHADO. 2010. Revisão e atualização da lista das aves do Rio Grande do Sul, Brasil. *Iheringia* 100: 519–556.
- , C. S. FONTANA, R. A. DIAS, G. N. MAURÍCIO, AND J. K. F. MÄHLER, Jr. 2003. Aves. In: Livro vermelho da fauna ameaçada de extinção no Rio Grande do Sul (C. S. Fontana, G. A. Bencke, AND R. E. dos Reis, eds.), pp. 189–479. Edipucrs, Porto Alegre, Brazil.
- BERNARDOS, J. N., AND M. E. ZACCAGNINI [online]. 2008. Evaluación del riesgo de toxicidad aguda para aves por uso de insecticidas en arroceras. In: Primer taller para la Conservación de Aves Playeras Migratorias en Arroceras del Cono Sur (V. M. de la Balze AND D. E. Blanco, eds.). Wetlands International, Buenos Aires, Argentina. <<http://lac.wetlands.org>> (12 December 2011)
- , ———, P. MINEAU, J. DECARRE, AND R. DE CARLI. 2007. Calculadora de riesgo ecotoxicológico para aves: sistema soporte de decisiones para el control de plagas con criterios ambientales. Versión 2.0. INTA, Buenos Aires, Argentina.
- BERTHRONG, S. T., C. W. SCHADT, G. PIÑEIRO, AND R. B. JACKSON. 2009. Afforestation alters the composition of functional genes in soil and biogeochemical

- processes in South American grasslands. *Applied and Environmental Microbiology* 75: 6240–6248.
- BILENCA, D., AND F. MIÑARRO. 2004. Identificación de Áreas Valiosas de Pastizal (AVPs) en las Pampas y Campos de Argentina, Uruguay y sur de Brasil. Fundación Vida Silvestre, Buenos Aires, Argentina.
- BIRDLIFE INTERNATIONAL [online]. 2012. Threatened birds of the world. Species factsheets. <http://www.birdlife.org> (20 March 2012)
- BLANCO, D. E., R. BANCHS, AND P. CANEVARI. 1993. Critical sites for the Eskimo Curlew (*Numenius borealis*), and other Nearctic grassland shorebirds in Argentina and Uruguay. U.S. Fish and Wildlife Service, Wetlands for the Americas, Buenos Aires, Argentina.
- , R. B. LANCTOT, J. P. ISACCH, AND V. A. GILL. 2004. Pastizales templados del Sur de América del Sur como hábitat de aves playeras migratorias. *Ornitología Neotropical* 15: 159–167.
- , B. LÓPEZ-LANÚS, R. A. DIAS, A. AZPIROZ, AND F. RILLA. 2006. Uso de arrozceras por chorlos y.
- , S. M. ZALBA, C. J. BERENQUER, G. PUGNALI, AND D. H. RODRÍGUEZ-GOÑI. 2003. Estado y conservación del cauquén colorado *Chloephaga rubidiceps* Sclater (Aves, Anatidae) en su zona de invernada (Provincia de Buenos Aires, Argentina). *Revista Chilena de Historia Natural* 76: 47–55.
- BOLDRINI, I. I., L. EGGERS, L. A. MENTZ, S. T. S. MIOTTO, N. I. MATZENBACHER, H. M. LONGHI-WAGNER, R. TREVISAN, A. A. SCHNEIDER, AND R. B. SETÚBAL. 2009. Flora. In: Biodiversidade dos Campos do Planalto das Araucárias (I. I. Boldrini, ed.), pp. 40–94. Ministério do Meio Ambiente, Brasília, Brazil.
- BRANDÃO, T., R. TREVISAN, AND T. BOTH. 2007. Unidades de conservação e os campos do sul do Rio Grande do Sul. *Revista Brasileira de Biociências* 5: 843–845.
- BRISTOT, A. 2001. Planalto das Araucárias – um ecossistema em perigo de extinção? *Agoecologia e Desenvolvimento Rural Sustentável* 2: 24–31.
- BROCKERHOFF, E. G., H. JACTEL, J. A. PARROTTA, C. P. QUINE, AND J. SAYER. 2008. Plantation forests and biodiversity: oxymoron or opportunity? *Biodiversity and Conservation* 17: 925–951.
- BROOK, B. W., J. J. O'GRADY, A. P. CHAPMAN, M. A. BURGMAN, H. R. AKÇAKAYA, AND R. FRANKHAM. 2000. Predictive accuracy of Population Viability Analysis in conservation biology. *Nature* 404: 385–387.
- BUCHER, E. H., AND M. NORES. 1988. Present status of birds in steppes and savannas of northern and central Argentina. In: *Ecology and conservation of grassland birds* (P. D. Goriup, ed.), pp. 71–80. International Council for Bird Preservation, Cambridge, UK.
- BURGOS, J. J., AND A. VIDAL. 1951. Los climas de la República Argentina según la nueva clasificación de Thornthwaite. *Revista Meteoros* 1: 3–32.
- BUTCHER, G. S., M. R. FULLER, L. S. McALLISTER, AND P. H. GEISSLER. 1990. An evaluation of the Christmas Bird Count for monitoring population trends of selected species. *Wildlife Society Bulletin* 18: 129–134.
- CAMARGO, O. A. (org.). 2002. Atlas eólico: Rio Grande do Sul. Secretaria de Energia, Minas e Comunicações, SEMC, Porto Alegre, Brazil.
- CANAVELLI, S. B., M. J. BECHARD, B. WOODBRIDGE, M. N. KOCHERT, J. J. MACEDA, AND M. E. ZACCAGNINI. 2003. Habitat use by Swainson's Hawks on their austral wintering grounds in Argentina. *Journal of Raptor Research* 37: 125–134.
- , M. E. ZACCAGNINI, J. TORRESIN, N. C. CALAMARI, M. P. DUCOMMUN, AND P. CAPLLONCH. 2004. Monitoreo extensivo de aves en el centro-sur de Entre Ríos. In: *Temas de la biodiversidad del litoral fluvial argentino* (F. G. Aceñolaza, ed.), pp. 349–362. Instituto Superior de Correlación Geológica, San Miguel de Tucumán, Argentina.
- CARDONI, D. A. 2011. Efectos de disturbios antrópicos en el uso de hábitat de aves de marismas: consideraciones para un uso sustentable. Ph.D. dissertation, Universidad Nacional de Mar del Plata, Mar del Plata, Argentina.
- , J. P. ISACCH, M. E. FANJUL, M. ESCAPA, AND O. IRIBARNE. 2011. Relationship between anthropogenic sewage discharge, marsh structure and bird assemblages in an SW Atlantic saltmarsh. *Marine Environmental Research* 71: 122–130.
- CARTES, J. L., AND R. P. CLAY. 2009. Paraguay. In: *Important Bird Areas Americas—Priority sites for biodiversity conservation* (C. Devenish, D. F. Díaz Fernández, R. P. Clay, I. Davidson, AND I. Yépez Zabala, eds.), pp. 297–306. BirdLife Conservation Series 16, BirdLife International, Quito, Ecuador.
- CASAÑAS, H.E., I. ROESLER, AND J. KLAVINS. 2007. Historia natural y distribución de la Cachirla Trinadora (*Anthus chacoensis*). *Hornero* 22: 59–63.
- CLAY, R. P., H. DEL CASTILLO, AND J. DE EGEA. 2008. Paraguay: contextos eco-regionales, geográficos y socioeconómicos. In: *Áreas importantes para la conservación de las aves del Paraguay* (Guyra Paraguay, ed.), pp. 31–44. Guyra Paraguay/BirdLife International, Asunción, Paraguay.
- CHEBEZ, J. C. 2008. Los que se van. Fauna argentina amenazada. Tomo 2. Albatros, Buenos Aires, Argentina.
- CODESIDO, M., AND R. M. FRAGA. 2009. Distributions of threatened grassland passerines of Paraguay, Argentina and Uruguay, with new locality records and notes on their natural history and habitat. *Ornitología Neotropical* 20: 585–595.
- , C. GONZÁLEZ-FISCHER, AND D. BILENCA. 2011. Distributional changes of landbird species in agroecosystems of central Argentina. *Condor* 113: 266–273.
- , ———, AND ———. 2012. Agricultural land-use, avian nesting and rarity in the Pampas of central Argentina. *Emu* 112: 46–54.
- COLLAR, N. J., AND P. ANDREW. 1988. *Birds to watch: the ICBP world checklist of threatened birds*. ICBP and IUCN, Cambridge, UK.
- , M. J. CROSBY, AND A. J. STATTERSFIELD. 1994. *Birds to watch 2: the world list of threatened birds*. BirdLife International, Cambridge, UK.
- , L. P. GONZAGA, N. KREBBE, A. MADROÑO NIETO, L. G. NARANJO, T. A. PARKER III, AND D. C. WEGE. 1992. Threatened birds of the Americas: the

- ICPB/IUCN Red Data Book. International Council for Bird Preservation, Cambridge, UK.
- COMPARATORE, V. M., M. M. MARTÍNEZ, A. I. VASALLO, O. M. BARG, AND J. P. ISACCH. 1996. Abundancia y relaciones con el hábitat de aves y mamíferos en pastizales de *Paspalum quadrifarium* (paja colorada) manejados con fuego (Provincia de Buenos Aires, Argentina). *Interciencia* 21: 228–237.
- COMPARATORE, V., AND C. YAGUEDDÚ. 2007. Diet of the Greater Rhea (*Rhea americana*) in an agroecosystem of the Flooding Pampa, Argentina. *Ornitología Neotropical* 18: 187–194.
- CORDEIRO, J. L. P., AND H. HASENACK. 2009. Cobertura vegetal atual do Rio Grande do Sul. In: Campos Sulinos: conservação e uso sustentável (V. P. Pillar, S. C. Müller, Z. M. S. Castilhos, AND A. V. Á. Jacques, eds.), pp. 285–299. Ministério do Meio Ambiente, Brasília, Brazil.
- CRAWSHAW, D., M. DALL'AGNOL, J. L. P. CORDEIRO, AND H. HASENACK. 2007. Caracterização dos campos sul-rio-grandenses: uma perspectiva da ecologia da paisagem. *Boletim Gaúcho de Geografia* 33:233–252.
- DE EGEE, J., M. PEÑA-CHOCARRO, C. ESPADA, AND S. KNAPP. 2012. Checklist of vascular plants of the Department of Ñeembucú, Paraguay. *PhytoKeys* 9: 15–179.
- DEL CASTILLO, H., AND R. P. CLAY. 2004. Annotated checklist of the birds of Paraguay. Asociación Guyra Paraguay, Asuncion, Paraguay.
- DEMARÍA, M. R., I. AGUADO SUÁREZ, AND D. F. STEIMAKER. 2008. Reemplazo y fragmentación de pastizales pampeanos semiáridos en San Luis, Argentina. *Ecología Austral* 18: 55–70.
- , W. J. MCSHEA, K. KOY, AND N. MACEIRA. 2003. Pampas deer conservation with respect to habitat loss and protected area considerations in San Luis, Argentina. *Biological Conservation* 115: 121–130.
- DEVELEY, P. F., AND J. M. GOERCK. 2009. Brazil. In: Important Bird Areas Americas—Priority sites for biodiversity conservation (C. Devenish, D. F. Díaz Fernández, R. P. Clay, I. Davidson, AND I. Yépez Zabala, eds.), pp. 99–112. BirdLife International, Quito, Ecuador.
- DEVENISH, C., D. F. DÍAZ FERNÁNDEZ, R. P. CLAY, I. DAVIDSON, AND I. YÉPEZ ZABALA (eds.). 2009. Important Bird Areas Americas. Priority sites for biodiversity conservation. BirdLife International, Quito, Ecuador.
- DIAS, R. A., AND G. N. MAURICIO. 2002. Natural history notes and conservation of a Saffron-cowled Blackbird *Xanthopsar flavus* population in the southern coastal plain of Rio Grande do Sul, Brazil. *Bird Conservation International* 12: 225–268.
- DI GIACOMO, A. S. 2005b. Áreas importantes para la conservación de las aves en la Argentina. Sitios prioritarios para la conservación de la biodiversidad. Aves Argentinas / Asociación Ornitológica del Plata, Buenos Aires, Argentina.
- , AND A. G. DI GIACOMO. 2004. Extinción, historia natural y conservación de las poblaciones del Yetapá de Collar (*Alectrurus risora*) en la Argentina. *Ornitología Neotropical* 15: 1–14.
- , ———, AND J. R. CONTRERAS. 2005. Status and conservation of the Bobolink (*Dolichonyx oryzivorus*) in Argentina. In: Bird conservation implementation and integration in the Americas (T. Rich, AND C. J. Ralph, eds.), pp. 519–524. General Technical Report PSW-GTR-191, Pacific Southwest Research Station, U.S. Forest Service, Albany, CA.
- , ———, AND J. C. REBORDA. 2011b. Male and female reproductive success in a threatened polygynous species: the Strange-Tailed Tyrant (*Alectrurus risora*). *Condor* 103: 619–628.
- DI GIACOMO, A. S., AND S. KRAPOVICKAS. 2001. Afforestation threatens Argentina's grasslands. *World Birdwatch* 23: 24–25.
- , AND A. F. PARERA. 2008. 20 áreas prioritarias para la conservación de las aves migratorias neárticas en los pastizales del cono sur de Sudamérica. *Aves Argentinas / Asociación Ornitológica del Plata, Buenos Aires, Argentina*.
- , P. VICKERY, H. CASAÑAS, O. SPITZNAGEL, C. OSTROSKY, S. KRAPOVICKAS, AND A. BOSSO. 2010. Landscape associations of globally threatened grassland birds in the Aguapey River Important Bird Area, Corrientes, Argentina. *Bird Conservation International* 20: 62–73.
- DI GIACOMO, A. G. 2005a. Aves de la Reserva El Bagual. In: Historia natural y paisaje de la Reserva El Bagual, Provincia de Formosa, Argentina. Inventario de la fauna de vertebrados y de la flora vascular de un área protegida del Chaco Húmedo (A. G. Di Giacomo, AND S. F. Krapovickas, eds.), pp. 201–465. Aves Argentinas / Asociación Ornitológica del Plata, Buenos Aires, Argentina.
- , A. S. DI GIACOMO, AND J. C. REBORDA. 2011a. Effects of grassland burning on reproductive success of globally threatened Strange-tailed Tyrants *Alectrurus risora*. *Bird Conservation International* 21: 411–422.
- Dinama/Mvotma. 1999. Propuesta de estrategia nacional para la conservación y uso sostenible de la diversidad biológica del Uruguay. Proyecto URU/96/G 31, FMAM, PNUD, Montevideo, Uruguay.
- DONALD, P. F., R. E. GREEN, AND M. F. HEATH. 2001. Agricultural intensification and the collapse of Europe's farmland bird populations. *Proceedings of the Royal Society B* 268: 25–29.
- , F. J. SANDERSON, I. J. BURFIELD, AND F. P. J. VAN BOMMEL. 2006. Further evidence of continent-wide impacts of agricultural intensification on European farmland birds 1990–2000. *Agriculture, Ecosystems and Environment* 116: 189–196.
- ECHEVERRÍA, A. I., A. I. VASSALLO, AND J. P. ISACCH. 2006. Experimental analysis of novelty responses in a bird assemblage inhabiting a suburban marsh. *Canadian Journal of Zoology* 84: 974–980.
- FAUBA [online]. 2011. Estiman usos de la tierra en la Provincia de Buenos Aires. <http://agro.fauga.info/node/206#_ftn1> (22 December 2011).
- FERNÁNDEZ, G., G. POSSE, V. FERRETTI, AND F. M. GABELLI. 2003. Bird-habitat relationship for the declining Pampas Meadowlark populations in the southern Pampas grasslands. *Biological Conservation* 115: 139–148.

- FILLOY, J., AND M. I. BELLOCO. 2007. Patterns of bird abundance along the agricultural gradient of the Pampean region. *Agriculture, Ecosystem and Environment* 120: 291–298.
- , ZURITA, G.A., CORBELL, J.M., AND M. I. BELLOCO. 2010. On the similarity among bird communities: testing the influence of distance and land use. *Acta Oecologica* 36: 333–338.
- FISCHER, J., AND D. B. LINDENMAYER. 2007. Landscape modification and habitat fragmentation: a synthesis. *Global Ecology and Biogeography* 16: 265–280.
- FISHER, R. J., AND S. K. DAVIS. 2010. From Wiens to Robel: a review of grassland-bird habitat selection. *Journal of Wildlife Management* 74: 265–73.
- FONTANA, C. S. 1994. História natural de *Heteroxolmis dominicana* (Vieillot, 1823) (Aves, Tyrannidae) com ênfase na relação ecológica com *Xanthopsar flavus* (Gmelin, 1788) (Aves Icteridae) no nordeste do Rio Grande do Sul. M.S. thesis. Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil.
- , C. E. ROVEDDER, M. REPENNING, AND M. L. GONÇALVES. 2008. Estado atual do conhecimento e conservação da avifauna dos Campos de Cima da Serra do sul do Brasil, Rio Grande do Sul e Santa Catarina. *Revista Brasileira de Ornitologia* 16: 281–307.
- , M. REPENNING, AND C. E. ROVEDDER. 2009. Fauna Terrestre: Aves. In: *Biodiversidade dos campos do Planalto das Araucárias* (I. I. Boldrini, ed.), pp. 159–207. Ministério do Meio Ambiente, Brasília, Brazil.
- FRAGA, R. M. 2003. Distribution, natural history and conservation of the Black-and-White Monjita (*Heteroxolmis dominicana*) in Argentina, a species vulnerable to extinction. *Ornitologia Neotropical* 14: 1–12.
- , H. CASAÑAS, AND G. PUGNALI. 1998. Natural history and conservation of the endangered Saffron-cowled Blackbird *Xanthopsar flavus* in Argentina. *Bird Conservation International* 8: 255–267.
- FUHLENDORF, S. D., AND D. M. ENGLE. 2001. Restoring heterogeneity on rangelands: ecosystem management based on evolutionary grazing patterns. *BioScience* 51: 625–632.
- GABELLI, F. M., G. J. FERNÁNDEZ, V. FERRETTI, G. POSSE, E. COCONIER, H. J. GAVIEIRO, P. E. LLAMBÍAS, P. I. PELÁEZ, M. L. VALLÉS, AND P. L. TUBARO. 2004. Range contraction in the Pampas Meadowlark *Sturnella defilippii* in the southern pampas grasslands of Argentina. *Oryx* 38: 164–170.
- GHIONE, S., AND D. MARTINO. 2008. Biodiversidad. Informe del estado del ambiente. In: *GEO Uruguay 2008* (S. Ghione AND D. Martino, eds.), pp. 178–239. Gráfica Mosca, Montevideo, Uruguay.
- GIBSON, E. 1885. Notes on the birds of Paysandú, Republic of Uruguay. *Ibis* 5: 275–283.
- GIORDANO, P. F., J. L. NAVARRO, AND M. B. MARTELLA. 2009. Building large-scale spatially explicit models to predict the distribution of suitable habitat patches for the Greater Rhea (*Rhea americana*), a near-threatened species. *Biological Conservation* 143: 357–365.
- GOLDSTEIN, M. I., B. WOODBRIDGE, M. E. ZACCAGNINI, S. G. CANAVELLI, AND A. LANUSSE. 1996. An assessment of mortality of Swainson's Hawks in wintering grounds in Argentina. *Journal of Raptor Research* 30: 106–107.
- GOLDSTEIN, M. I., T. E. LACHER, JR, B. WOODBRIDGE, M. J. BECHARD, S. B. CANAVELLI, M. E. ZACCAGNINI, G. P. COBB, E. J. SCOLLON, R. TRIBOLET, AND M. J. HOOPER. 1999a. Monocrotophos-induced mass mortality of Swainson's Hawks in Argentina, 1995–96. *Ecotoxicology* 8: 201–214.
- , ———, M. E. ZACCAGNINI, M. L. PARKER, AND M. J. HOOPER. 1999b. Monitoring and assessment of Swainson's Hawks in Argentina following restrictions on monocrotophos use, 1996–97. *Ecotoxicology* 8: 215–224.
- GORIUP, P. D. 1988. Ecology and conservation of grassland birds. International Council for Bird Preservation, Cambridge, UK.
- GORLERI, M. C. 2005. Caracterización climática del Chaco Húmedo. In: *Historia natural y paisaje de la Reserva El Bagual, Formosa, Argentina. Inventario de la fauna de vertebrados y de la flora vascular de un área del Chaco Húmedo* (A. G. Di Giacomo AND S. F. Krapovickas, eds.), pp. 13–25. Aves Argentinas/Asociación Ornitológica del Plata, Buenos Aires, Argentina.
- Guyra Paraguay. 2005. Atlas de las aves del Paraguay. Asunción, Paraguay.
- HASSE, G. 2006. Eucalipto: histórias de um imigrante vegetal. JÁ Editores, Porto Alegre, Brazil.
- HEDRICK, P. W. 2005. Large variance in reproductive success and the Ne/N ratio. *Evolution* 59: 1596–1599.
- HENWOOD, W. D. 2010. Toward a strategy for the conservation and protection of the world's temperate grasslands. *Great Plains Research* 20: 121–134.
- HERRERA, L., V. COMPARATORE, AND P. LATERRA. 2004. Habitat relations of *Rhea americana* in an agroecosystem of Buenos Aires Province, Argentina. *Biological Conservation* 119: 363–369.
- HERRERA, L. P., P. LATERRA, N. MACEIRA, K. D. ZELAYA, AND G. A. MARTÍNEZ. 2009. Fragmentation status of tall-tussock grassland relicts in the Flooding Pampa, Argentina. *Rangeland Ecology and Management* 62: 73–82.
- HOEKSTRA, J. M., T. M. BOUCHER, T. H. RICKETTS, AND C. ROBERTS. 2005. Confronting a biome crisis: global disparities of habitat loss and protection. *Ecology Letters* 8: 23–29.
- HOOPER, M. J., P. MINEAU, M. E. ZACCAGNINI, G. W. WINEGRAD, AND B. WOODBRIDGE. 1999. Monocrotophos and the Swainson's Hawk. *Pesticide Outlook* 10: 97–102.
- , ———, ———, AND ———. 2003. Pesticides and international migratory bird conservation. In: *Handbook of ecotoxicology* (D. J. Hoffman, B. A. Rattner, G. A. Burton, Jr., AND J. Cairns, Jr., eds.), pp. 737–753. Lewis Publishers, Boca Raton, FL.
- HUDSON, W. H. 1920. *Birds of La Plata*. J. M. Dent and Sons, Toronto, Canada.
- HÜFFNER, A., AND B.C. ENGEL (eds.) 2011. *Grandes e pequenas centrais hidrelétricas na bacia do Rio Uruguai: guias para ONGs e movimentos sociais. Amigos da Terra – NatBrasil*. Porto Alegre, Brazil.
- IBAMA. 2009. O uso sustentável da reserva legal: diretrizes para o manejo da pecuária nos campos do

- bioma Pampa. Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis/RS, Porto Alegre, Brazil.
- ICMBio. *In press*. Plano de ação nacional para a conservação dos passeriformes ameaçados dos Campos Sulinos e Espinilho. Instituto Chico Mendes de Conservação da Biodiversidade, Brasília, Brazil.
- IGNIS [online]. 2010. Lista de espécies ameaçadas de extinção em Santa Catarina. <<http://www.ignis.org.br>> (5 July 2011).
- IHERING, H. VON. 1899. As aves do Estado do Rio Grande do Sul. In: Anuário do Estado do Rio Grande do Sul para o ano de 1900, pp. 113–154. Gundlach and Krahe, Porto Alegre, Brazil.
- ISACCH, J. P. 2008. Implementing the Biosphere Reserve concept: the case of Parque Atlántico Mar Chiquito Biosphere Reserve from Argentina. *Biodiversity and Conservation* 17: 1799–1804.
- , M. S. BÓ V. COMPARATORE, L. P. HERRERA, R. J. VARGAS, AND M. M. MARTÍNEZ. 2001. Las aves de los pastizales costeros del SE de la provincia de Buenos Aires. In: Reserva de Biosfera Mar Chiquita: Características físicas, biológicas y ecológicas (O. Iribarne, ed.), pp. 269–286. Editorial Martín, Mar del Plata, Argentina.
- , M. S. BÓ, N. O. MACEIRA, M. R. DEMARÍA, AND S. PELUC. 2003. Composition and seasonal changes of the bird community of natural grasslands in the western Pampa of Argentina. *Journal of Field Ornithology* 74: 59–65.
- , AND D. A. CARDONI. 2011. Different grazing strategies are necessary to conserve endangered grassland birds in short and tall salty grasslands of the flooding Pampas. *Condor* 113: 724–734.
- , S. HOLZ, L. RICCI, AND M. M. MARTÍNEZ. 2004. Post-fire vegetation change and bird use of a salt marsh in coastal Argentina. *Wetlands* 24: 235–243.
- , N. O. MACEIRA, M. S. BÓ, M. R. DEMARÍA, AND S. PELUC. 2005. Bird-habitat relationship in semiarid natural grasslands and exotic pastures in the west pampas of Argentina. *Journal of Arid Environment* 62: 267–283.
- , AND M. M. MARTÍNEZ. 2001. Estacionalidad y relaciones con la estructura del hábitat de la comunidad de aves de pastizales de paja colorada (*Paspalum quadrifarium*) manejados con fuego en la provincia de Buenos Aires, Argentina. *Ornitología Neotropical* 12: 345–354.
- , AND ———. 2003a. Temporal variation in abundance and population status of nonbreeding Nearctic and Patagonian shorebirds in the flooding pampa grasslands of Argentina. *Journal of Field Ornithology* 74: 233–242.
- , AND ———. 2003b. Habitat use by nonbreeding shorebirds in flooding pampa grasslands of Argentina. *Waterbirds* 26: 494–500.
- JOBBÁGY, E. G., AND R. B. JACKSON. 2004. Groundwater use and salinization with grassland afforestation. *Global Change Biology* 10: 1299–1312.
- JOHNSON, M. D., T. W. SHERRY, R. T. HOLMES, AND P. P. MARRA. 2006. Measuring habitat quality for a migratory songbird wintering in natural and agricultural areas. *Conservation Biology* 20: 1433–1444.
- JOHNSON, R. J., J. A. JEDLICKA, J. E. QUINN, AND J. R. BRANDLE. 2011. Global perspectives on birds in agricultural landscapes. In: *Issues in agroecology: present status and future prospectus*, volume 1, Integrating agriculture, conservation and ecotourism: examples from the field (W. B. Campbell and S. L. Ortíz, eds.), pp. 55–140. Springer, London, UK.
- JOSEPH, L. 1997. Towards a broader view of Neotropical migrants: consequences of a re-examination of austral migration. *Ornitología Neotropical* 8: 31–36.
- KLEIN, R. M. 1978. Flora ilustrada catarinense: mapa fitogeográfico do Estado de Santa Catarina. Herbário Barbosa Rodrigues, Itajaí, Brazil.
- KRAUSMAN, R. P. 1999. Some basic principles of habitat use, grazing behavior of livestock and wildlife. *Idaho Forest, Wildlife and Range Experiment Station Bulletin* 70: 85–90.
- LANGTOT, R. B., D. E. BLANCO, R. A. DIAS, J. P. ISACCH, V. A. GILL, J. B. ALMEIDA, K. DELHEY, P. PETRACCI, G. A. BENCKE, AND R. A. BALBUENO. 2002. Conservation status of the Buff-breasted Sandpiper: historic and contemporary distribution and abundance in South America. *Wilson Bulletin* 114: 44–72.
- , ———, M. OESTERHELD, R. A. BALBUENO, J. P. GUERSCHMAN, AND G. PIÑEIRO. 2004. Assessing habitat availability and use by Buff-breasted Sandpipers (*Tryngites subruficollis*) wintering in South America. *Ornitología Neotropical* 15: 367–376.
- LANGSTON, R. H. W., AND J. D. PULLAN. 2003. Windfarms and birds: an analysis of the effects of windfarms on birds, and guidance on environmental assessment criteria and site selection issues. *Birdlife International, RSPB, Sandy, UK*.
- LANTSCHNER, M., V. RUSCH, AND C. PEYROU. 2008. Bird assemblages in pine plantations replacing native ecosystems in NW Patagonia. *Biodiversity and Conservation* 17: 969–989.
- LATERA, P., O. R. VIGNOLIO, L. G. HIDALGO, O. N. FERNÁNDEZ, M. A. CAUHEPE, AND N. O. MACEIRA. 1998. Dinámica de pajonales de paja colorada (*Paspalum* spp.) manejados con fuego y pastoreo en la Pampa Deprimida Argentina. *Ecotrópicos* 11: 141–149.
- LEITE, P. F. 2002. Contribuição ao conhecimento fitoecológico do sul do Brasil. *Ciência & Ambiente* 24: 51–73.
- LEÓN, R. J. C., AND S. E. BURKART. 1998. El pastizal de la Pampa Deprimida: estados alternativos. *Ecotrópicos* 11: 121–130.
- LEVEAU, L., AND C. LEVEAU. 2004. Riqueza y abundancia de aves en agroecosistemas pampeanos durante el período post-reproductivo. *Ornitología Neotropical* 15: 371–380.
- LIPSEY, M. K., AND P. A. R. HOCKEY. 2010. Do ecological networks in South African commercial forests benefit grassland birds? A case study of a pine plantation in KwaZulu-Natal. *Agriculture, Ecosystems and Environment* 137: 133–142.
- LOPES, L. E., G. B. MALACCO, E. F. ALTEFF, M. FERREIRA DE VASCONCELOS, D. HOFFMANN AND L. F. SILVEIRA. 2010. Range extensions and conservation of some threatened or little known Brazilian grassland birds. *Bird Conservation International* 20:84–94.

- LÓPEZ-LANÚS, B., P. GRILLI, E. COCONIER, A. DI GIACOMO, AND R. BANCHS. 2008. Categorización de las aves de la Argentina según su estado de conservación. Informe de Aves Argentinas / AOP y Secretaría de Ambiente y Desarrollo Sustentable, Buenos Aires, Argentina.
- , AND G. MARINO. 2010. Ecología del Charlatán y su estado actual en la Provincia de Santa Fe, Argentina. *Aves Argentinas/Asociación Ornitológica del Plata*, Buenos Aires, Argentina.
- MACHADO, É., AND L. F. SILVEIRA. 2011. Plumage variability and taxonomy of the Capped Seedeater *Sporophila bouvreuil* (Aves: Passeriformes: Emberizidae). *Zootaxa* 2781: 49–62.
- MADSEN, J., R. MATUS, O. BLANK, L. BENEGAS, G. MATEAZZI, AND D. E. BLANCO. 2003. Population status of the Ruddy-headed Goose (*Chloephaga rubidiceps*) in Tierra del Fuego and mainland Patagonia (Chile and Argentina). *Ornitología Neotropical* 14: 15–28.
- MANNING, A. D., D. B. LINDENMAYER, AND H. A. NIX. 2004. Continua and umwelt: novel perspectives on viewing landscapes. *Oikos* 104: 621–628.
- MARINO, G. D. 2008. Buenas prácticas ganaderas para conservar la vida silvestre de las Pampas. Una guía para optimizar la producción y conservar la biodiversidad de los pastizales de la Bahía Samborombón y la cuenca del Río Salado. *Aves Argentinas*, Buenos Aires, Argentina.
- MARTELLA, M. B., J. L. NAVARRO GONNET, AND S. A. MONGE. 1996. Diet of Greater Rheas in an agroecosystem of central Argentina. *Journal of Wildlife Management* 60: 586–592.
- MARTINO, D., AND M. METHOL. 2008. Cambios en el uso de la tierra. In: *GEO Uruguay 2008* (S. Ghione AND D. Martino, eds.), pp. 56–117. Gráfica Mosca, Montevideo, Uruguay.
- MATURO, H. M., L. J. OAKLEY, AND D. E. PRADO. 2005. Vegetación y posición fitogeográfica de la Reserva El Bagual. In: *Historia natural y paisaje de la Reserva El Bagual*, Formosa, Argentina. Inventario de la fauna de vertebrados y de la flora vascular de un área del Chaco Húmedo (A. G. Di Giacomo AND S. F. Krapovickas, eds.), pp. 59–73. *Aves Argentinas / Asociación Ornitológica del Plata*, Buenos Aires, Argentina.
- MEDAN, D., J. P. TORRETTA, K. HODARA, E. B. DE LA FUENTE, AND N. H. MONTALDO. 2011. Effects of agriculture expansion and intensification on the vertebrate and invertebrate diversity in the Pampas of Argentina. *Biodiversity and Conservation* 20: 3077–3100.
- METZGER, J. P., T. M. LEWINSOHN, C. A. JOLY, L. M. VERDADE, L. A. MARTINELLI, AND R. R. RODRIGUES. 2010. Brazilian law: full speed in reverse? *Science* 329: 276–277.
- MMA. 2007. Áreas prioritárias para a conservação, uso sustentável e repartição de benefícios da biodiversidade brasileira: atualização—Portaria no. 9. Ministério do Meio Ambiente, Brasília, Brazil.
- MORALES, C., H. DEL CASTILLO, S. CENTRÓN, AND F. PALACIOS. 2010. El contexto ambiental de los cultivos de arroz en Paraguay. In: *Arroz un Negocio Creciente* (A. Friedman AND B. Wei, eds.), pp. 49–52. USAID/Paraguay Vende, Asunción, Paraguay.
- MORRIS, W. F., AND D. F. DOAK. 2002. Quantitative conservation biology: the theory and practice of Population Viability Analysis. Sinauer Associates, Sunderland, MA.
- MYERS, J. P., AND L. P. MYERS. 1979. Shorebirds of coastal Buenos Aires Province, Argentina. *Ibis* 121: 186–200.
- NAROSKY, T., AND A. DI GIACOMO. 1993. Las aves de la Provincia de Buenos Aires: distribución y estatus. *Asociación Ornitológica del Plata*, Vázquez Mazzini Editores, Buenos Aires, Argentina.
- NETTO, C. G. A. M. 2009. O futuro dos campos: possibilidades econômicas de continuidade da bovinocultura de corte no Rio Grande do Sul. In: *Campos Sulinos: conservação e uso sustentável* (V. P. Pillar, S. C. Müller, Z. M. S. Castilhos, AND A. V. Á. Jacques, eds.), pp. 380–390. Ministério do Meio Ambiente, Brasília, Brazil.
- NUNNEY, L. 1993. The influence of mating system and overlapping generations on effective population size. *Evolution* 47: 1329–1341.
- 1996. The influence of variation in female fecundity on effective population size. *Biological Journal of the Linnean Society* 59: 411–425.
- OLSON, D. M., E. DINERSTEIN, E. D. WIKRAMANAYAKE, N. D. BURGESS, G. V. N. POWELL, E. C. UNDERWOOD, J. A. D'AMICO, I. ITOUA, H. E. STRAND, J. C. MORRISON, C. J. LOUCKS, T. F. ALLNUTT, T. H. RICKETTS, Y. KURA, J. F. LAMOREUX, W. W. WETTENGEL, P. HEDAO, AND K. R. KASSEM. 2001. Terrestrial ecoregions of the world: a new map of life on Earth. *BioScience* 51: 933–938.
- OVERBECK, G. E., S. C. MÜLLER, A. FIDELIS, J. PFADENHAUER, V. D. PILLAR, C. C. BLANCO, I. I. BOLDRINI, R. BOTH, AND E. D. FORNECK. 2007. Brazil's neglected biome: the South Brazilian Campos. Perspectives in Plant Ecology, Evolution and Systematics 9: 101–116.
- PARUELO, J. M., J. P. GUERSCHMAN, G. BALDI, AND C. DI BELLA. 2004. La estimación de la superficie agrícola. Antecedentes y una propuesta metodológica. *Inter-ciencia* 29: 421–427.
- PARUELO, J. M., J. P. GUERSCHMAN, G. PIÑEIRO, E. G. JOBBÁGY, S. R. VERÓN, G. BALDI, AND S. BAEZA. 2006. Cambios en el uso de la tierra en Argentina y Uruguay: marcos conceptuales para su análisis. *Agrociencias* 10: 47–61.
- PARUELO, J. M., J. P. GUERSCHMAN, AND S. R. VERÓN. 2005. Expansión agrícola y cambios en el uso del suelo. *Ciencia Hoy* 15: 14–23.
- PEDRANA, J., J. P. ISACCH, AND M. S. BÓ. 2008. Habitat relationships of diurnal raptors at local and landscape scales in southern temperate grasslands of Argentina. *Emu* 108: 301–310.
- PEREIRA, J. A., R. D. QUINTANA, AND S. MONGE. 2003. Diets of Plains Vizcacha, Greater Rhea and cattle in Argentina. *Journal of Range Management* 56: 13–20.
- PETRY, M. V., AND L. KRÜGER. 2010. Frequent use of burned grasslands by the vulnerable Saffron-cowled Blackbird *Xanthopsar flavus*: implications for the

- conservation of the species. *Journal of Ornithology* 151: 599–605.
- PILLAR, V. D., AND F. L. QUADROS. 1997. Grassland-forest boundaries in southern Brazil. *Coenosis* 12: 119–126.
- PILLAR, V. D., AND E. VÉLEZ. 2010. Extinção dos Campos Sulinos em unidades de conservação: um fenômeno natural ou um problema ético? *Natureza & Conservação* 8: 84–88.
- PINHEIRO, R. T., AND G. LOPEZ. 1999. Abundancia del tinamú manchado (*Nothura maculosa*) y del tinamú alirrojo (*Rhynchotus rufescens*) en un área cinegética del Río Grande Do Sul (Brasil). *Ornitología Neotropical* 10: 35–41.
- PREVEDELLO, J. A., AND M. V. VIEIRA. 2010. Does the type of matrix matter? A quantitative review of the evidence. *Biodiversity and Conservation* 19: 1205–1223.
- PRIETO, A. 2000. Vegetational history of the late glacial-Holocene transition in the grasslands of eastern Argentina. *Paleogeography, Paleoclimatology, Paleocology* 157: 167–188.
- PRUGH, L. R., K. E. HODGES, A. R. E. SINCLAIR, AND J. S. BRASHARES. 2008. Effect of habitat area and isolation on fragmented animal populations. *Proceedings of the National Academy of Sciences USA* 105: 20770–20775.
- REINO, L., M. PORTO, R. MORGADO, F. CARVALHO, A. MIRA, AND P. BEJA. 2010. Does afforestation increase bird nest predation risk in surrounding farmland? *Forest Ecology and Management* 260: 1359–1366.
- REMSEN, J. V., JR., C. D. CADENA, A. JARAMILLO, M. NORES, J. F. PACHECO, J. PÉREZ-EMÁN, M. B. ROBBINS, F. G. STILES, D. F. STOTZ, AND K. J. ZIMMER [online]. 2012. A classification of the bird species of South America. American Ornithologists' Union. <<http://www.museum.lsu.edu/Remsen/SACCBaseline.html>> (21 March 2012)
- REPENNING, M. 2012. História natural de uma população migratória de *Sporophila* aff. *plumbea* (Aves, Emberizidae) do sul do Brasil. M.S. thesis, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil.
- REPENNING, M., C. E. ROVEDDER, AND C. S. FONTANA. 2010. Distribuição e biologia de aves nos campos de altitude do sul do Planalto Meridional Brasileiro. *Revista Brasileira de Ornitologia* 18: 283–306.
- RIVERA-MILÁN, F. F., M. E. ZACCAGNINI, AND S. B. CANAVELLI. 2004. Field trials of line-transect surveys of bird carcasses in agro-ecosystems of Argentina's Pampas region. *Wildlife Society Bulletin* 32: 1219–1228.
- ROVEDDER, C. E. 2011. História natural de *Sporophila melanogaster* (Pelzeln 1870) (Aves: Emberizidae) com ênfase em sua biologia reprodutiva. M. S. thesis, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil.
- RUMBOLL, M. 1975. El Cauquén de Cabeza Colorada (*Chloephaga rubidiceps*): una nota de alarma. *El Hornero* 11: 315–316.
- RUMBOLL, M. 1979. El estado actual de *Chloephaga rubidiceps*. *Acta Zoologica Lilloana* 34: 153–154.
- SALA, O. E., M. OESTERHELD, R. J. C. LEÓN, AND A. SORIANO. 1986. Grazing effects upon plant community structure in subhumid grasslands of Argentina. *Vegetatio* 67: 27–32.
- SANTOS, R. E. F., AND P. SCHERER-NETO. 2009. Aves de campos e várzeas. In: Planos de Ação para Espécies de Aves Ameaçadas no Paraná (IAP/Projeto Paraná Biodiversidade, ed.). Instituto Ambiental do Paraná, Curitiba, Brazil.
- SARASOLA, J. H., M. A. GALMES, AND M. Á. SANTILLÁN. 2007. Ecology and conservation of Swainson's Hawk (*Buteo swainsoni*) in Argentina. *Hornero* 22: 173–184.
- SARASOLA, J. H., AND J. J. NEGRO. 2006. Role of alien tree stands on current distribution and social behavior of Swainson's Hawks *Buteo swainsoni* in the Argentine pampas. *Journal of Biogeography* 33: 1096–1101.
- SARASOLA, J. H., J. J. NEGRO, V. SALVADOR, AND J. J. MACEDA. 2005. Hailstorms as a cause of mass mortality of Swainson's Hawks in their wintering grounds. *Journal of Wildlife Diseases* 41: 643–646.
- SCHRAG, A. M., M. E. ZACCAGNINI, N. CALAMARI, AND S. CANAVELLI. 2009. Climate and land-use influences on avifauna in central Argentina: broad-scale patterns and implications of agricultural conversion for biodiversity. *Agriculture, Ecosystem and Environment* 132: 135–142.
- SICK, H. 1997. *Ornitologia Brasileira*. Nova Fronteira, Rio de Janeiro, Brazil.
- SILVEIRA, L., AND J. ALONSO. 2009. Runoff modifications due to the conversion of natural grasslands to forests in a large basin in Uruguay. *Hydrological Processes* 23: 320–329.
- SORIANO, A., R. J. C. LEÓN, O. E. SALA, S. LAVADO, V. A. DEREGIBUS, M. A. CAUHEPÉ, O. A. SCAGLIA, A. C. A. VELÁSQUEZ, AND J. H. LEMCOFF. 1991. Río de la Plata Grasslands. In: *Ecosystems of the world 8A, natural grasslands, introduction and Western Hemisphere* (R. T. Coupland, ed.), pp. 367–407. Elsevier, New York, NY.
- STOTZ, D. E., J. W. FITZPATRICK, T. A. PARKER III, AND D. K. MOSKOVITS. 1996. Neotropical birds: ecology and conservation. University of Chicago Press, Chicago, IL.
- STRAUBE, F., C. A. URBEN-FILHO, AND D. KAJIWARA. 2004. Aves. In: *Livro vermelho da fauna ameaçada no Estado do Paraná* (S. B. Mikich AND R. S. Bérnils, eds.), pp. 143–496. Instituto Ambiental do Paraná, Curitiba, Brazil.
- STRUM, K. M., M. J. HOOPER, K. A. JOHNSON, R. B. LANCTOT, M. E. ZACCAGNINI, AND B. K. SANDERCOCK. 2010. Exposure of migratory shorebirds to cholinesterase-inhibiting contaminants in the Western Hemisphere. *Condor* 112: 15–28.
- THOMPSON, J. J. 2004. Tinamous and agriculture: lessons learned from the Galliformes. *Ornitología Neotropical* 15: 301–307.
- THOMPSON, J. J., AND J. P. CARROLL. 2009. Habitat use and survival of the Spotted Tinamou (*Nothura maculosa*) in agroecosystems in the Province of Buenos Aires, Argentina. In: *Proceedings of Gamebird 2006. Quail VI and Perdix XII* (S. B. Cederbaum,

- B. C. Faircloth, T. M. Terhune, J. J. Thompson, AND J. P. Carroll, eds.), pp. 111–119. Warnell School of Forestry and Natural Resources, University of Georgia, Athens, GA.
- TUBARO, P. L., AND F. M. GABELLI. 1999. The decline of the Pampas Meadowlark: difficulties of applying IUCN criteria to Neotropical grassland birds. *Studies in Avian Biology* 19: 250–257.
- VÉLEZ, E., L. CHOMENKO, W. SCHAFFER, AND M. MADEIRA. 2009. Um panorama sobre as iniciativas de conservação dos Campos Sulinos. In: *Campos Sulinos: conservação e uso sustentável* (V. P. Pillar, S. C. Müller, Z. M. S. Castilhos, AND A. V. Á. Jacques, eds.), pp. 356–379. Ministério do Meio Ambiente, Brasília, Brazil.
- VICKERY, P. D., D. E. BLANCO, AND B. LÓPEZ-LANÚS. 2010. Conservation plan for the Upland Sandpiper (*Bartramia longicauda*), version 1.1. Manomet Center for Conservation Sciences, Manomet, MA.
- VICKERY, P. D., H. E. CASAÑAS, AND A. S. DI GIACOMO. 2003. Effects of altitude on the distribution of Nearctic and resident grassland birds in Córdoba province, Argentina. *Journal of Field Ornithology* 74: 172–178.
- VICKERY, P. D., M. L. HUNTER, JR., AND J. V. WELLS. 1992. Is density an indicator of breeding success? *Auk* 109: 706–710.
- VICKERY, P. D., P. L. TUBARO, J. M. C. DA SILVA, B. G. PETERJOHN, J. R. HERKERT, AND R. B. CAVALCANTI. 1999. Conservation of grassland birds in the Western Hemisphere. *Studies in Avian Biology* 19: 2–26.
- VIGLIZZO, E. F., F. LECTORA, A. J. PORDOMINGO, J. N. BERNARDOS, Z. E. ROBERTO, AND H. DEL VALLE. 2001. Ecological lessons and applications from one century of low external-input farming in the pampas of Argentina. *Agriculture, Ecosystem and Environment* 83: 65–81.
- VIGLIZZO, E. F., A. J. PORDOMINGO, D. BUSCHIAZZO, AND M. G. CASTRO. 2005. A methodological approach to assess cross-scale relations and interactions in agricultural ecosystems of Argentina. *Ecosystems* 8: 546–558.
- WIED, M. [Prinzen zu]. 1831. Beiträge zur Naturgeschichte von Brasilien. III(2). Landes-Industrie-Comptoirs, Weimar, Germany.
- WORLD RESOURCES 2000–2001. People and ecosystems: the fraying web of life. World Resources Institute, Washington, D.C.
- WOODBIDGE, B., K. K. FINLEY, AND S. T. SEAGER. 1995. An investigation of the Swainson's Hawk in Argentina. *Journal of Raptor Research* 29: 202–204.
- ZACCAGNINI, M. E. 2004. ¿Porqué monitoreo ecotoxicológico de diversidad de aves en sistemas productivos? In: *INTA expone 2004. Volumen III* (E. Larrea, ed.), pp. 69–89. INTA, Buenos Aires, Argentina.
- ZACCAGNINI, M. E., J. N. BERNARDOS, AND P. MINEAU. 2005. Calculadora de riesgo ecotoxicológico para aves, versión 1.0. and 1.1. INTA, Buenos Aires, Argentina.
- ZALBA, S. M., AND N. C. COZZANI. 2004. The impact of feral horses on grassland bird communities in Argentina. *Animal Conservation* 7: 35–44.

APPENDIX 1

Bird species associated with grasslands in Southeastern South America, with type of grassland and other habitats used, preferred grass height, conservation status (Birdlife International 2012), and migratory status (following Narosky and Di Giacomo 1993, Belton 1994, Stotz et al. 1996, Joseph 1997, Sick 1997).

Family/Species	Common name	Grassland use ^a	Other habitats ^b	Grass height ^c	Conservation status ^d	Migratory status ^e
Rheidae						
<i>Rhea americana</i>	Greater Rhea	2	Agr	Broad	NT	RES
Tinamidae						
<i>Rhynchotus rufescens</i>	Red-winged Tinamou	2	Sav	Broad	LC	RES
<i>Nothura darwinii</i>	Darwin's Nothura	2	Agr, Sav	Broad	LC	RES
<i>N. maculosa</i>	Spotted Nothura	2	Agr	Broad	LC	RES
Anatidae						
<i>Chloephaga picta</i>	Upland Goose	2	Agr	Short	LC	SACT
<i>C. poliocephala</i>	Ashy-headed Goose	2	Agr	Short	LC	SACT
<i>C. rubidiceps</i>	Ruddy-headed Goose	2	Agr	Short	LC	SACT
Ardeidae						
<i>Bubulcus ibis</i>	Cattle Egret	2	Agr, Wet	Broad	LC	RES
Threskiornithidae						
<i>Theristicus caudatus</i>	Buff-necked Ibis	2	Agr, Sav, Wet	Short	LC	RES
<i>T. melanopsis</i>	Black-faced Ibis	2	Agr	Short	LC	SACT
Accipitridae						
<i>Elanus leucurus</i>	White-tailed Kite	3	Agr, Sav	Broad	LC	RES
<i>Circus cinereus</i>	Cinereous Harrier	2	Agr, Wet	Broad	LC	RES
<i>C. buffoni</i>	Long-winged Harrier	2	Agr, Wet	Broad	LC	RES
<i>Buteogallus meridionalis</i>	Savanna Hawk	3	Agr, Sav	Broad	LC	RES
<i>B. coronatus</i>	Crowned Eagle	3	Sav	Broad	EN	RES
<i>Geranoaetus albicaudatus</i>	White-tailed Hawk	2	Sav	Broad	LC	RES
<i>G. melanoleucus</i>	Black-chested Buzzard-Eagle	2	Sav	Broad	LC	RES
<i>Buteo swainsoni</i>	Swainson's Hawk	2	Agr	Broad	LC	PNW
Charadriidae						
<i>Vanellus chilensis</i>	Southern Lapwing	2	Agr, Wet	Short	LC	RES
<i>Pluvialis dominica</i>	American Golden-Plover	2	Agr, Wet	Short	LC	PNW
<i>Charadrius modestus</i>	Rufous-chested Dotterel	2	Wet	Short	LC	SACT
<i>Oreopholus ruficollis</i>	Tawny-throated Dotterel	2	Agr	Short	LC	SACT
Scolopacidae						
<i>Numenius borealis</i>	Eskimo Curlew	1	–	Short	CR	PNW
<i>Bartramia longicauda</i>	Upland Sandpiper	2	Agr	Broad	LC	PNW
<i>Tryngites subruficollis</i>	Buff-breasted Sandpiper	2	Agr, Wet	Short	NT	PNW
Thinocoridae						
<i>Thinocorus rumicivorus</i>	Least Seedsnipe	2	Agr	Short	LC	SACT
Strigidae						
<i>Athene cunicularia</i>	Burrowing Owl	2	Agr	Short	LC	RES
<i>Asio flammeus</i>	Short-eared Owl	2	Agr, Wet	Broad	LC	RES
Caprimulgidae						
<i>Chordeiles nacunda</i>	Nacunda Nighthawk	2	Agr, Wet	Broad	LC	SATT
<i>Caprimulgus longirostris</i>	Band-winged Nightjar	3	Sav	Broad	LC	SACT
<i>Eleothreptus anomalus</i>	Sickle-winged Nightjar	2	Wet	Tall	NT	RES
Trochilidae						
<i>Polytmus guainumbi</i>	White-tailed Goldenthrout	3	Sav	Broad	LC	RES
Picidae						
<i>Colaptes campestris</i>	Campo Flicker	3	Agr, Sav	Broad	LC	RES
Cariamidae						
<i>Cariama cristata</i>	Red-legged Seriema	2	Sav	Broad	LC	RES
Falconidae						
<i>Caracara plancus</i>	Southern Caracara	2	Agr, Sav, Wet	Broad	LC	RES
<i>Milvago chimango</i>	Chimango Caracara	2	Agr, Sav, Wet	Broad	LC	RES
<i>Falco sparverius</i>	American Kestrel	3	Agr, Sav	Broad	LC	RES
<i>F. femoralis</i>	Applomado Falcon	2	Agr, Sav	Broad	LC	RES

Continued

APPENDIX—Continued

Family/Species	Common name	Grassland use ^a	Other habitats ^b	Grass height ^c	Conservation status ^d	Migratory status ^e
Furnariidae						
<i>Geositta cucicularia</i>	Common Miner	2	Wet	Short	LC	RES
<i>Furnarius rufus</i>	Rufous Hornero	3	Agr, Sav	Broad	LC	RES
<i>Cinclodes pabsti</i>	Long-tailed Cinclodes	1	—	Short	NT	RES
<i>C. fuscus</i>	Buff-winged Cinclodes	3	Wet	Short	LC	SACT
<i>Phacellodomus striaticollis</i>	Freckle-breasted Thornbird	3	Sav, Wet	Tall	LC	RES
<i>Anumbius annumbi</i>	Firewood-gatherer	3	Sav	Broad	LC	RES
<i>Asthenes hudsoni</i>	Hudson's Canastero	2	Wet	Broad	NT	RES
<i>A. modesta</i>	Cordilleran Canastero	1	—	Short	LC	RES
<i>Cranioleuca sulphurifera</i>	Sulphur-throated Spinetail	3	Wet	Tall	LC	RES
<i>Spartonoica maluroides</i>	Bay-capped Wren-Spinetail	2	Wet	Tall	NT	RES
Tyrannidae						
<i>Polystictus pectoralis</i>	Bearded Tachuri	1	—	Tall	NT	SATT
<i>Pseudocolaptes flaviventris</i>	Warbling Doradito	3	Wet	Tall	LC	RES
<i>Culicivora caudacuta</i>	Sharp-tailed Tyrant	2	Wet	Tall	VU	RES
<i>Lessonia rufa</i>	Austral Negrito	2	Agr, Wet	Short	LC	SACT
<i>Knipolegus lophotes</i>	Crested Black-Tyrant	2	Sav	Broad	LC	RES
<i>Hymenops perspicillatus</i>	Spectacled Tyrant	2	Wet	Broad	LC	RES
<i>Agriornis montanus</i>	Black-billed Shrike-Tyrant	1	—	Short	LC	RES
<i>Xolmis cinereus</i>	Gray Monjita	2	Sav	Broad	LC	RES
<i>X. irupero</i>	White Monjita	3	Agr, Sav	Broad	LC	RES
<i>X. dominicanus</i>	Black-and-white Monjita	2	Wet	Broad	VU	RES
<i>Neoxolmis rufiventris</i>	Chocolate-vented Tyrant	2	Agr	Short	LC	SACT
<i>Gubernetes yetapa</i>	Streamer-tailed Tyrant	2	Wet	Tall	LC	RES
<i>Alectrurus tricolor</i>	Cock-tailed Tyrant	1	—	Tall	VU	RES
<i>A. risora</i>	Strange-tailed Tyrant	2	Wet	Tall	VU	RES
<i>Machetornis rixosa</i>	Cattle Tyrant	3	Agr, Sav	Broad	LC	RES
<i>Tyrannus savana</i>	Fork-tailed Flycatcher	3	Agr, Sav	Broad	LC	SATT
Hirundinidae						
<i>Alopochelidon fucata</i>	Tawny-headed Swallow	2	Wet	Broad	LC	RES
<i>Progne tapera</i>	Brown-chested Martin	2	Agr, Sav	Broad	LC	SATT
<i>Tachycineta leucorhoa</i>	White-rumped Swallow	3	Agr, Sav, Wet	Broad	LC	RES
<i>T. meyeri</i>	Chilean Swallow	3	Agr, Wet	Broad	LC	SACT
<i>Hirundo rustica</i>	Barn Swallow	3	Agr, Wet	Broad	LC	PNW*
<i>Petrochelidon pyrrhonota</i>	Cliff Swallow	3	Agr, Sav	Broad	LC	PNW*
Troglodytidae						
<i>Cistothorus platensis</i>	Sedge Wren	2	Wet	Tall	LC	RES
Mimidae						
<i>Mimus saturninus</i>	Chalk-browed Mockingbird	3	Agr, Sav	Broad	LC	RES
Motacillidae						
<i>Anthus lutescens</i>	Yellowish Pipit	2	Agr, Wet	Short	LC	RES
<i>A. furcatus</i>	Short-billed Pipit	2	Agr	Short	LC	RES
<i>A. chacoensis</i>	Pampas Pipit	2	Agr	Broad	LC	RES
<i>A. correndera</i>	Correndera Pipit	2	Agr, Wet	Broad	LC	RES
<i>A. nattereri</i>	Ochre-breasted Pipit	1	—	Broad	VU	RES
<i>A. bellmayri</i>	Hellmayr's Pipit	1	—	Broad	LC	RES
Thraupidae						
<i>Donacospiza albifrons</i>	Long-tailed Reed Finch	2	Sav, Wet	Broad	LC	RES
<i>Poospiza nigrorufa</i>	Black-and-rufous Warbling-Finch	3	Wet	Tall	LC	RES
<i>Sicalis citrina</i>	Stripe-tailed Yellow-Finch	2	Sav	Short	LC	RES
<i>S. lebruni</i>	Patagonian Yellow-Finch	1	—	Broad	LC	RES
<i>S. luteola</i>	Grassland Yellow-Finch	2	Agr, Wet	Broad	LC	RES
<i>Emberizoides herbicola</i>	Wedge-tailed Grass-Finch	2	Sav	Tall	LC	RES
<i>E. ypiranganus</i>	Lesser Grass-Finch	2	Wet	Tall	LC	RES
<i>Embernagra platensis</i>	Great Pampa-Finch	2	Wet	Broad	LC	RES
<i>Volatinia jacarina</i>	Blue-black Grassquit	2	Agr, Wet	Tall	LC	SATT
<i>Sporophila aff. plumbea</i>	Plumbeous Seedeater	1	—	Tall	LC	SATT

Continued

APPENDIX—Continued

Family/Species	Common name	Grassland use ^a	Other habitats ^b	Grass height ^c	Conservation status ^d	Migratory status ^e
<i>S. collaris</i>	Rusty-collared Seedeater	3	Wet	Tall	LC	RES
<i>S. bouvreuil</i>	Capped Seedeater	2	Wet	Tall	LC	SATT
<i>S. hypoxantha</i>	Tawny-bellied Seedeater	2	Wet	Tall	LC	SATT
<i>S. ruficollis</i>	Dark-throated Seedeater	2	Wet	Tall	NT	SATT
<i>S. palustris</i>	Marsh Seedeater	2	Wet	Tall	EN	SATT
<i>S. hypochroma</i>	Rufous-rumped Seedeater	2	Wet	Tall	NT	SATT
<i>S. cinnamomea</i>	Chestnut Seedeater	2	Wet	Tall	VU	SATT
<i>S. melanogaster</i>	Black-bellied Seedeater	2	Wet	Tall	NT	SATT
<i>Catamenia analis</i>	Band-tailed Seedeater	1	–	Broad	LC	RES
<i>Coryphaspiza melanotis</i>	Black-masked Finch	2	Sav	Broad	VU	RES
Emberizidae						
<i>Ammodramus humeralis</i>	Grassland Sparrow	2	Agr	Broad	LC	RES
Icteridae						
<i>Agelasticus thilius</i>	Yellow-winged Blackbird	3	Wet	Broad	LC	RES
<i>Xanthopsar flavus</i>	Saffron-cowled Blackbird	2	Agr, Wet	Broad	VU	RES
<i>Pseudoleistes guirahuro</i>	Yellow-rumped Marshbird	2	Agr, Wet	Broad	LC	RES
<i>P. virescens</i>	Brown-and-yellow Marshbird	2	Agr, Wet	Broad	LC	RES
<i>Molothrus rufoaxillaris</i>	Screaming Cowbird	3	Sav	Broad	LC	RES
<i>M. bonariensis</i>	Shiny Cowbird	3	Agr, Sav	Broad	LC	RES
<i>Sturnella superciliaris</i>	White-browed Blackbird	2	Agr	Broad	LC	RES
<i>S. defilippii</i>	Pampas Meadowlark	1	–	Broad	VU	RES
<i>S. loyca</i>	Long-tailed Meadowlark	2	Agr, Sav	Broad	LC	RES
<i>Dolichonyx oryzivorus</i>	Bobolink	3	Agr, Wet	Tall	LC	PNW

^aGrassland-use categories are: (1) species restricted solely to grassland habitats (i.e., do not use alternative habitats throughout the SESA grasslands), (2) species that make extensive use of grassland habitats, but also use other habitats in the SESA grasslands, and (3) species that make extensive use of grassland habitats only in certain subregions of the SESA grasslands.

^bAlternative habitats used are: Agr: row crop habitats, Sav: shrub and savanna habitats, and Wet: wetland habitats; hyphens denote species that do not use alternative habitats in the SESA grasslands.

^cAssociation with homogeneous short (Short) and tall (Tall) grass; the Broad category indicates species that are less selective in terms of grassland height and can make broader use of grassland structure.

^dCR: Critically Endangered, EN: Endangered, VU: Vulnerable, NT: Near Threatened, and LC: Least Concern.

^eMigration systems (sensu Joseph 1997): RES: Resident, PNW: Pan New World Migration System, SACT: South American Cool-Temperate Migration System, and SATT: South American Temperate-Tropical Migration System; * = breeding populations recently established within SESA.