


# Quantification of the effect of host patch configuration on the abundance of *Bemisia tabaci* in central Argentina: a multimodel inference approach

## Research Paper

Mariano P. Grilli<sup>1,2</sup> , Marina Bruno<sup>3</sup> and Romina Fachinetti<sup>1</sup>

**Cite this article:** Grilli MP, Bruno M, Fachinetti R (2022). Quantification of the effect of host patch configuration on the abundance of *Bemisia tabaci* in central Argentina: a multimodel inference approach. *Bulletin of Entomological Research* 1–9. <https://doi.org/10.1017/S0007485322000311>

Received: 13 December 2021

Revised: 31 March 2022

Accepted: 12 May 2022

**Keywords:**

Bemisia tabaci; host patch configuration; spatial distribution; Multimodal inference

**Author for correspondence:**

Mariano P. Grilli,

Email: [mariano.grilli@unc.edu.ar](mailto:mariano.grilli@unc.edu.ar)

<sup>1</sup>Centro de Relevamiento y Evaluación de Recursos Agrícolas y Naturales (CREAN-IMBIV), CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina; <sup>2</sup>Cátedra de Bioestadística I y II, FCEfyN, Universidad Nacional de Córdoba, Córdoba, Argentina and <sup>3</sup>Cátedra de Microbiología, Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba, Córdoba, Argentina

**Abstract**

*Bemisia tabaci* is a complex of species, which is considered the most common and important pest of a wide range of crops belonging to many different botanical families. In Argentina, this species is recognized as a vector of geminiviruses, and Middle East-Asia Minor 1, Mediterranean, New World and New World 2 have been found to coexist in the same area. Landscape elements, like habitat patch area and isolation, define the habitat configuration and have a direct effect on insect populations between and within host patches. In this paper, we analyse the effect of potato patch configuration on the distribution and abundance of *B. tabaci*. Potato patches were identified using Landsat TM5 and TM7 images, and a supervised classification was performed to quantify the spatial distribution of the patches in the whole study area. Potato patch metrics were estimated using Fragstats 4.4. Generalized linear mixed models were employed to analyse the relationship between whiteflies and landscape configuration, through a multimodel inference approach, finding that *B. tabaci* abundance and landscape metrics were very variable. After a multimodel selection process, we found that perimeter-to-area ratio and Euclidean distance between patches were the variables that best explained whitefly abundance in potato patches. Implications of these findings are discussed.

**Introduction**

*Bemisia tabaci* is a cryptic species complex (De Barro *et al.*, 2011), considered the most common and important pest of a wide range of crops belonging to many different botanical families, due to its wide distribution and the serious damage it causes to host plants (Brown *et al.*, 1995; Oliveira *et al.*, 2001). It is a polyphagous and invasive species, colonizing more than 1000 different plant species and causing important direct losses by feeding or acting as a vector for more than 300 plant viruses (Navas-Castillo *et al.*, 2011). *B. tabaci* is one of the most damaging pests in tropical and subtropical regions (Byrne and Bellows, 1991), affecting the yield of various agricultural and horticultural crops (Cahill *et al.*, 1996). The broad range of hosts of *B. tabaci* for feeding and/or reproducing include horticultural crops such as potato (*S. tuberosum* L.), sweet potato (*Ipomoea batatas* L.), eggplant (*Solanum melongena* L.), and other horticultural extensive crops (Brown *et al.*, 1995).

In South America, the *B. tabaci* species complex is one of the few whitefly species damaging crops, directly by feeding, or by virus transmission (Krause-Sakate *et al.*, 2020). Direct damage occurs by phloem feeding and the excretion of the honeydew on leaves and fruits, which serves as a substrate for the growth of sooty mould that covers the surface, interfering with photosynthesis and causing losses in plant productivity (Kanakala and Ghanim, 2015). However, the most important damage caused by whiteflies to agriculture is virus transmission, particularly by the *B. tabaci* species complex, which can transmit viruses of the genera Begomovirus, Carlavirus, Crinivirus, Ipomovirus, Torradovirus, and Ploverovirus (Navas-Castillo *et al.*, 2011). In Argentina, since 1986, *B. tabaci* is recognized as the vector of a geminivirus of the Bean golden mosaic virus group on soybean (Ploper *et al.*, 1989).

Among the different members of the complex, two of the cryptic species, the putative species Middle East-Asia Minor 1 (MEAM1) and Mediterranean (usually referred to in the literature as B and Q biotypes, respectively) are known to be the most invasive species of the complex (Perring, 2001). In many areas, MEAM1 and Mediterranean are associated with the displacement of local whitefly species (Liu *et al.*, 2007). The invasion of MEAM1

and the Mediterranean in many regions of the world led to epidemics of plant disease caused by begomovirus transmitted through *B. tabaci* (Hogenhout *et al.*, 2008). In Argentina, MEAM1 was first detected on horticultural crops, weeds, and cotton (Viscarret *et al.*, 2003), and the putative Mediterranean species was detected in sweet pepper and melon (Grille *et al.*, 2011). Coexisting with these two invading species, recent studies showed two indigenous species, one that belongs to the New World putative species, and a second one referred to as New World 2 (Alemandri *et al.*, 2015). MEAM1 has been able to displace the New World putative species (Perring and Symmes, 2006), but the presence and persistence of New World 2 in areas invaded by MEAM1 suggest that this species retains the capacity to resist displacement or interbreeding (Alemandri *et al.*, 2012).

The spatial pattern of vegetation affects the distribution and abundance of herbivorous insects within agroecosystems (Grilli and Bruno, 2007; Grilli and Fachinetti, 2017). Habitat patch area, habitat patch isolation, and characteristics of the landscape surrounding the patch define the habitat configuration and affect direct or indirectly the insect population between and within patches (Grilli and Fachinetti, 2019). However, the role of habitat amount and configuration on species occurrence and abundance is one of the major focuses of research in ecology and biogeography (Saura, 2021). There is intense debate on the relative importance of habitat quantity (total area of habitat) and the spatial configuration of that habitat (the spatial arrangement of habitat) for biodiversity patterns and persistence (Fahrig, 2013; Fahrig *et al.*, 2015; Haddad *et al.*, 2015; Martin, 2018). The most widely accepted conceptual model, and the prevailing consensus among ecologists, has been that both habitat quantity and habitat configuration (e.g., fragmentation) are important and should be considered in conservation management (e.g., Haddad *et al.*, 2015).

The spatial configuration of habitat components (e.g., fragmentation and isolation of habitat types) is often important because individuals moving between patches are key to accessing resources in different habitats (Thies *et al.*, 2003). Furthermore, the effects of the landscape configuration on individual species may depend upon the extent to which a species uses one or multiple habitat types within a landscape (Ewers and Didham, 2006). The relationship between host patches and insect density within the patch is very variable and depends on the species and its life history. Steffan-Dewenter and Tscharnkte (2000) showed that population densities of monophagous species increased, but oligophagous and polyphagous species decreased, with habitat area. In the case of *B. tabaci*, their abundance and growth in a crop field is affected by many factors, such as the intercropping distance within the field, and the spatial arrangement of hosts around the crop field (Macfadyen *et al.*, 2018).

In Córdoba, the main area of potato production is placed in the central area of the province. Because of the climate of the area, there are two seasons, an early season by the end of winter when seed potato is produced, and a late season during autumn when consumption potato is produced. In this area, all the produced potatoes are of the same variety, *S. tuberosum* L. var. Spunta (Huarte and Capezio, 2013), and in particular during autumn, potato is the only crop present in the field.

Research on the distribution of *B. tabaci* has been limited in Argentina, despite its status as a pest increasing in recent years

(Alemandri *et al.*, 2012, 2014, 2015). In this paper, we analyse the effect of host patch configuration on the distribution and abundance of *B. tabaci* on consumption potato fields during autumn in central Argentina.

## Materials and methods

### Study area

The study was conducted in potato plots (*S. tuberosum* L.) in the central area of Córdoba province, Argentina (fig. 1). In this area, potato is planted by the end of the summer and harvested in autumn. The study area is one of the most important production areas provincially and nationally (Quattrini, 2005).

### Insect sampling

Adult individuals of *B. tabaci* were collected using yellow sticky traps, ten of which were placed within each potato patch (focal patches). The traps were made from a yellow cylinder supporting plastic film coated with adhesive and were placed at 1 m above the ground level. The plastic film was replaced with a clean one every 15 days during the sampling periods in two consecutive years. The films were taken to the laboratory, where *B. tabaci* were identified. Adults and nymphs of *B. tabaci* were identified in the fields and in the traps by their morphology according to Caballero (1996).

### Host patch assessment


For the sampling, we selected a total of 15 potato patches each year. The spatial position of each host patch was established using a GPS, identified on scenes path/row 229/82 on Landsat TM5 and Landsat ETM7 images. The images were georeferenced to the latitude/longitude reference system and atmospheric and radiometrically corrected.

### Host patch identification by supervised classification

A supervised classification was employed to determine land use, based on spectral brightness for six spectral bands in the visible and reflected infrared regions of the electromagnetic spectrum for each crop. To identify potato patches, four classes of land cover were considered in the analysis: bare soil, potato (host patches), woodlands, and pastures and weeds. Training site areas were digitized and signatures were created describing each informational class. Images were classified using Fisher's linear discriminant classifier (Landgrebe, 2003). Finally, accuracy was assessed by generating a random set of locations for verifying the true land cover type. An error matrix was applied *a posteriori* to compare the classes obtained with the real classes found in the field, and to obtain the kappa index of agreement for each class (Congalton and Green, 2019). All the image processing was performed using Terrset software (Eastman, 2020). Once the potato patches were identified, patch metrics of these were obtained from each scene with Fragstats 4.2 software (Mcgarigal and Ene, 2015).

### Landscape metrics

Considering the average dispersal distance of *B. tabaci* as reviewed by Byrne (1999), a 2000 m diameter area around each insect



**Figure 1.** Study area showing the distribution of the potato plots sampled in the central area of Córdoba, Argentina.

sampling site was extracted from each classified image to estimate host patch metrics. A total of four patch metrics representing the three basic patch configuration properties was obtained for each focal potato patch: two patch size-related metrics: Patch Area and Patch Perimeter; one patch shape complexity metric: Patch

Perimeter-to-area ratio; and a patch isolation metric: Patch Euclidean Nearest Neighbour Distance.

**Patch area (AREA)**

The area in hectares of the focal patch containing the sticky traps.

**Table 1.** Main *a priori* hypotheses (and predictions) about the effects of patch metric covariates on the abundance of *B. tabaci* in potato patches

No.	Hypothesis	Predictions	Independent variable involved
0	There is no effect of host patch configuration on the abundance of <i>B. tabaci</i>	The abundance of <i>B. tabaci</i> in a potato patch varies randomly across all the host patches	NULL
1	Patch size has a positive effect on the abundance of <i>B. tabaci</i> in a particular patch	An increase of the focal potato patch area will increase the abundance of adult individuals within the patch	AREA
2	Patch isolation has a negative effect on the abundance of <i>B. tabaci</i> in a particular patch	An increase of the Euclidean distance to the nearest neighbour from a focal potato patch to other potato patches will decrease the abundance of adult individuals within the focal patch	ENN
3	Patch edge has a positive effect on the abundance of <i>B. tabaci</i> in a particular patch	An increase of the focal potato patch perimeter will increase the abundance of adult individuals within the patch	PERIMETER
4	Patch shape complexity has a positive effect on the abundance of <i>B. tabaci</i> in a particular patch	An increase of the perimeter-to-area ratio will increase the abundance of adult individuals within the patch	PARA

### Patch perimeter (PERIMETER)

The perimeter in metres of the potato focal patch from which samples were collected.

### Perimeter-to-area ratio (PARA)

The ratio of the patch perimeter (m) to its area (m<sup>2</sup>).

### Euclidean nearest neighbour distance (ENN)

The distance (m) to the nearest neighbouring patch of the same type, based on shortest edge-to-edge distance.

### Data analysis

First, we recorded whitefly abundance per trap in every host patch during the study period. Generalized linear mixed models (GLMMs) were employed to analyse the relationships between the numbers of whiteflies collected in each potato field and the set of preselected patch metrics as independent variables. A multi-model inference approach (Grueber *et al.*, 2011) was employed to classify models and select the best model of the set. As sampling was longitudinal throughout the sampling period, sampling dates were set as a random effect. A set of GLMMs was pre-defined representing specific versions of the hypotheses proposed, based on the four patch configuration metrics (table 1). GLMMs were fitted using maximum likelihood and the relative performance of each model, evaluated with Akaike's information criteria (AICc) (Burnham and Anderson, 2002). Negative binomial was preferred over the Poisson distribution, based on the AICc of the models. The set of models to compare included a null model representing the null hypothesis of random variation of *B. tabaci* abundance in host patches. Models were fitted using lme4 and MASS packages in R 4.0.4 (R Core Team, 2019). Inference was based on the full results of each model set's analysis, and specifically on the covariate structure of better- vs. worse-performing models, as assessed by AICc (with lower scores signalling a better compromise between model fit and model complexity) (Burnham and Anderson, 2002). Once the best model was identified, the relationship between *B. tabaci* abundance and patch metric was estimated. Confidence intervals for each independent variable included in the model and the variance inflation factor (VIF) were also estimated to test the autocorrelation of the

independent variables. Finally, to clearly represent the effect of the selected variables on the full model, effects plots were performed using the package Effects from R, following the method described by Fox (2003).

### Results

*B. tabaci* abundance was variable between patches, with a maximum mean value of 389 ( $\pm 117$ ) individuals in plot 4 and a minimum mean value of 20 ( $\pm 1$ ) individuals collected in plot 11 (fig. 2).

Classification of land use by Fisher's linear discriminant classifier proved to be very precise. The error matrix accounted for 0.84 overall kappa index of the land use classification for potato, with a kappa index of agreement of 0.97.

### Host patch metrics

Focal potato patch metrics were also very variable in the study area, with areas (AREA) that ranged from 2.07 to 180.3 ha, focal patch perimeters (PERIMETER) ranging from 780 to 39,780 m, and perimeter-to-area ratios of focal patches (PARA) ranging from 115.07 to 571.43.

Euclidean nearest neighbour distance (ENN) of the focal patch to other potato patches also varied greatly, ranging from 60 to 480 m (table 2).

### Multimodel inference of *B. tabaci* abundance in potato patches

A 'confidence set' or 'credibility set' of models that were the most realistically likely to be the best approximating model was produced for each host patch variable. We ranked all the models from the best downwards and proceeded down the list until the cumulative Akaike's weight. This was done to explore which variables or combination of variables best explained the abundance of *B. tabaci* in the potato field. The results indicated that the best (most parsimonious) model that accounted for 99% of the cumulative weight was model 7 ( $M_7$ ), which includes perimeter-to-area ratio (PARA) and Euclidean distance to the nearest neighbour of the same class (ENN) as independent variables. This means that there is a 99% chance that  $M_7$  is the best approximating model describing the data, given the candidate set of models considered (table 3). We found that the two variables included in  $M_7$  were




Figure 2. Mean number of *B. tabaci* collected in each sampled potato plot.

significant (Wald chi-square test). A confidence interval (95%) was estimated. The full model showed that both independent variables, perimeter-to-area ratio (PARA) of potato plots and Euclidean distance to the nearest neighbour (ENN), have a negative relationship with *B. tabaci* abundance and show no autocorrelation between them (table 4). Numerically this means that patches with a PARA of 115 have almost three times more whiteflies than patches with a PARA of 571 (fig. 3a). In the case of the isolation of the potato patch, as represented by ENN, patches with a separation of 60 m from the nearest potato patch have 1.5 times more whiteflies than patches separated by 480 m from other potato patches (fig. 3b).

**Discussion**

To understand and explain which factors cause spatial variation in insect pest abundance, it is necessary to first understand the

Table 2. Mean, standard error (SE) and range values of the selected potato patch metrics in the study area

Variable	Mean	SE	Min	Max
AREA	46.49	2.52	2.07	180.36
PERIMETER	8434.09	471.52	780	39,780
PARA	241.28	5.67	115.07	571.43
ENN	206.98	6.33	60	480

different processes occurring at multiple spatial scales (Andersson et al., 2013), of which agricultural landscapes are good examples.

The study area was in a highly productive agricultural department of Córdoba province, with a developed gravity irrigation network. Potato (*S. tuberosum*) is one of the most important crops cultivated in the department (Sánchez and Barberis, 2013). Whitefly density was variable between potato plots.

Table 3. Multimodel inference selection of all the candidate models for the abundance of *B. tabaci* in potato fields

Model	Selected variables	AICc	Delta_AICc	AICcWt
M <sub>7</sub>	PARA ENN	4748.16	0	0.99
M <sub>3</sub>	PARA	4757.69	9.53	0.01
M <sub>6</sub>	PERIMETER ENN	4789.22	41.06	0
M <sub>2</sub>	PERIMETER	4805.69	57.53	0
M <sub>4</sub>	ENN	4806.5	58.34	0
M <sub>5</sub>	AREA ENN	4807.59	59.43	0
M <sub>0</sub>	NULL	4815.2	67.04	0
M <sub>1</sub>	AREA	4817.13	68.97	0

AICc, Akaike's information criterion; Delta\_AICc, difference between the best AICc model and the next AICc model; AICcWt, Akaike's weight.

**Table 4.** GLMM of the best model explaining the abundance of *B. tabaci* in each sampling plot based on the two proposed metrics

Model	AICc	Variables included	Coefficient	VIF	95% confidence interval
M <sub>7</sub>	4748.16	PARA	-0.0024	1	-0.0030/-0.0019
		ENN	-0.0009	1	-0.0015/-0.0004

AICc, Akaike's information criteria; VIF, variance inflation factor.


Values of VIF close to 1 indicate that there is no correlation among the predictor variables.

The method employed to determine the land use and the presence and distribution of host patches (potato plots) proved to be a very accurate and reliable tool for establishing the spatio-temporal

distribution of landscape suitable for the pest. Potato patches in the study area varied in size, shape, and spatial distribution. This variability was reflected other landscape metrics, e.g., small plots had higher perimeter-to-area ratios than large plots, and those with elongated shapes or irregular perimeters had higher perimeter-to-area ratios than plots of the same area with compact shapes and unbroken perimeters (Helzer and Jelinski, 1999). Cultivated patches in this area depend on the size of the farm and the farmer's decision about what to plant at each moment. This situation makes the distribution of crop patches very variable in space and time.

The multimodel inference method proved to be useful as a tool for model selection and to establish the role of the different covariates analysed (Burnham and Anderson, 1998, 2002; Anderson *et al.*, 2000). Our results clearly indicate that the abundance of adult individuals in a particular potato patch was mostly affected by the shape of the patch and the distance to other patches in the nearby area. The model shows a negative relationship between perimeter-to-area ratio and the abundance of dispersing individuals. Perimeter-to-area ratio (PARA) is a metric that quantifies the relationship between the patch size and its perimeter. In this case, a negative relationship between PARA and the abundance of whiteflies flying within the patch means that large patches, i.e., patches with lower PARA, will have more whiteflies than small patches, i.e., patches with higher PARA values. The same reasoning is valid for patches with a complex shape: more irregular potato patches will have higher PARA values, while regular patches will have lower PARA values. This means that large regular patches will have more flying whiteflies than small irregular ones.

Different insect species respond differently to host patch size (Bowers and Matter, 1997; Connor *et al.*, 2000; Bender *et al.*, 2003). The most traditional approach to the effect of host patch size on insect pests was formulated by Root (1973). This hypothesis predicts that specialist herbivores would reach higher densities in larger patches, which is considered an important guiding principle when studying the relationship between patch size and the population density of insect pests. The mechanism is based on the idea that there will be larger emigration rates out of smaller patches and larger immigration rates into larger patches (Hambäck and Englund, 2005). Since the hypothesis was formulated, many studies have quantified the relationship between patch size and population density for different organisms but the results have been very variable (Bowers and Matter, 1997; Connor *et al.*, 2000; Bender *et al.*, 2003). Some studies have found strong positive relationships between animal density and patch size, while others showed negative or no relationship at all (Bach, 1988). Hambäck and Englund (2005) found that any study on the relation between animal density and patch size should start from a thorough understanding of species dispersal and host search mode. They concluded, as did Bukovinsky



**Figure 3.** Effect display of the covariates included in the best model (M<sub>7</sub>). Dependent variable is in logarithmic scale. (a) Effect of PARA on *B. tabaci* abundance in potato plots. (b) Effect of ENN on *B. tabaci* abundance in potato plots. Grey area indicates the confidence interval of the model. For details see Fox and Hong (2009).

*et al.* (2005), that providing information on a species search mode can greatly improve the predictability of a movement-based hypothesis for understanding density–area relations (Hambäck and Englund, 2005).

On the other hand, the negative relationship between PARA and flying individuals within the patch may also be explained as an effect not only of the size of the patch, but of its shape. As we previously mentioned, large regular patches have lower PARA values, and low PARA values are related to higher *B. tabaci* abundance. And the opposite is also true; small irregular patches have higher PARA values, and higher PARA means patch values are related to lower amounts of flying *B. tabaci* within the patch. There are two key aspects of landscape boundaries that are predicted to influence the movement of insects: shape and contrast (Stamps *et al.*, 1987; Forman, 1995). Among the various descriptors of patch shape, patch perimeter-to-area ratio has received the most attention regarding its effect on animal movement into and out of the patches. The general theory states that, as the perimeter-to-area ratio increases, emigration also increases (Turchin, 1998). When the patch perimeter-to-area ratio is high, boundaries are more easily encountered by individuals living within the patch, increasing emigration from those patches. And this seems to be the case in our results. In other words, patch shape will affect insect movements between habitat patches (Stamps *et al.*, 1987). We can conclude that the negative relationship between potato patch perimeter-to-area ratio and whitefly density within host patches is a result of the way whiteflies disperse and search for new host patches (Byrne and Bellows, 1991).

The colonization of new host patches by herbivorous insects is mainly driven by the species' dispersal ability. Whiteflies are considered to be weak fliers, unable to actively fly more than 100 m (Byrne and Bellows, 1991). This is why the availability of source habitats in the nearby landscape is a key factor for the successful colonization of new habitat patches and population build-up for herbivorous insects. For example, the cabbage whitefly (*Aleyrodes proletella*) can colonize oilseed rape fields up to 1000 m away (Ludwig *et al.*, 2018). Nevertheless, the role of more distant source habitats is not clear. Byrne *et al.* (1996) found that *B. tabaci* had a bimodal dispersal pattern, with peaks at 100 and 2000 m from the dispersal source. They concluded that these peaks in dispersal of *B. tabaci* were the result of two morphs that differ in their specific ability to fly. During migration flights, they are attracted by ultraviolet or skylight, ignoring host cues (Döring, 2014), but afterwards, host finding is similar to the behaviour during trivial flights, characterized by attraction to green and yellow surfaces (Blackmer *et al.*, 1994), followed by a response to olfactory cues (Döring, 2014), and finally evaluating host suitability by probing the plant after landing (Noldus *et al.*, 1986). It is evident that host patch geometry plays a central role, as observed in our results. The dispersal behaviour described by Byrne *et al.* (1996) is also the most obvious explanation of the role of potato patches in the surrounding landscape in the process of colonizing the local patch. We found that the mean Euclidean distance of potato patches within the study area was 207 m, clearly within the dispersal range described by Byrne *et al.* (1996). Based on our models, we were able to quantify the effect of PARA and ENN on *B. tabaci* abundance in a particular patch.

Potato is the only green crop during autumn in the study area. During this period *B. tabaci* has no other alternative than moving from one potato field to another. All the other classified land covers are of non-host: bare soil, woodlands that are mainly of

*Vachellia caven*, *Prosopis alba*, and *Lithraea molleoides*, pastures and weeds. Based on this, to study the response of *B. tabaci* to host patch metrics and landscape configuration is valid, as no alternative hosts are present in the field during this time.

We concluded that there is a strong effect of the shape of the patch, with almost three times more whiteflies within the range of perimeter-to-area ratio observed, followed by the effect of the separation of the patches. The results obtained in this paper are a clear indication that effective management of whiteflies at a field scale would be viable only if addressed with an area-wide management approach.

**Conflict of interest.** The authors declare that they have no conflict of interests. This article does not contain any studies with human participants or animals performed by any of the authors.

## References

- Alemandri V, De Barro PJ, Bejerman N, Arguello-Caro EB, Dumon AD, Mattio MF, Rodriguez SM and Truol G (2012) Species within the *Bemisia tabaci* (Hemiptera: Aleyrodidae) complex in soybean and bean crops in Argentina. *Journal of Economic Entomology* **105**, 48–53.
- Alemandri V, Martino JA, Di Feo L and Truol G (2014) Indigenous and introduced species of the *Bemisia tabaci* complex in sweet potato crops from Argentina. *Agriscientia* **31**, 103–107.
- Alemandri V, Vaghi MCG, Dumón AD, Argüello CEB, Mattio MF, García MS, López LPM and Truol G (2015) Three members of the *Bemisia tabaci* (Hemiptera: Aleyrodidae) cryptic species complex occur sympatrically in Argentine horticultural crops. *Journal of Economic Entomology* **108**, 405–413.
- Anderson DR, Burnham KP and Thompson WL (2000) Null hypothesis testing: problems, prevalence, and an alternative. *Journal of Wildlife Management* **64**, 912–923.
- Andersson P, Löfstedt C and Hambäck PA (2013) Insect density–plant density relationships: a modified view of insect responses to resource concentrations. *Oecologia* **173**, 1333–1344.
- Bach CE (1988) Effects of host plant patch size on herbivore density: underlying mechanisms. *Ecology* **69**, 1103–1117.
- Bender DJ, Tischendorf L and Fahrig L (2003) Using patch isolation metrics to predict animal movement in binary landscapes. *Landscape Ecology* **18**, 17–39.
- Blackmer JL, Byrne DN and Tu Z (1994) Behavioral, morphological, and physiological traits associated with migratory *Bemisia tabaci* (Homoptera: Aleyrodidae). *Journal of Insect Behavior* **8**, 251–267.
- Bowers MA and Matter SF (1997) Landscape ecology of mammals: relationships between density and patch-size. *Journal of Mammalogy* **78**, 999–1013.
- Brown JK, Frohlich DR and Rosell RC (1995) The sweetpotato or silverleaf whiteflies: biotypes of *Bemisia tabaci* or a species complex? *Annual Review of Entomology* **40**, 511–534.
- Bukovinsky T, Potting RPJ, Clough Y, van Lenteren JC and Vet LEM (2005) The role of pre- and post-alighting detection mechanisms in the responses to patch size by specialist herbivores. *Oikos* **109**, 435–446.
- Burnham KP and Anderson DR (1998) *Model Selection and Multimodel Inference*. Berlin: Springer.
- Burnham KP and Anderson DR (2002) *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*, 2nd Edn. Berlin: Springer.
- Byrne DN (1999) Migration and dispersal by the sweet potato whitefly, *Bemisia tabaci*. *Agricultural and Forest Meteorology* **97**, 309–316.
- Byrne DN and Bellows TR (1991) Whitefly biology. *Annual Review of Entomology* **36**, 431–475.
- Byrne DN, Rathman RJ, Orum TV and Palumbo JC (1996) Localized migration and dispersal by the sweet potato whitefly, *Bemisia tabaci*. *Oecologia* **105**, 320–328.
- Caballero R (1996) Identificación de moscas blancas. In Hilje L (ed.), *Metodología para el estudio y manejo de moscas blancas y geminivirus*. Turrialba, Cartago, Costa Rica: CATIE, pp. 1–10.

- Cahill M, Denholm I, Ross G, Gorman K and Johnston D** (1996) Relationship between bioassay data and the simulated field performance of insecticides against susceptible and resistant adult *Bemisia tabaci* (Homoptera: Aleyrodidae). *Bulletin of Entomological Research* **86**, 109–116.
- Congalton RG and Green K** (2019) *Assessing the Accuracy of Remotely Sensed Data: Principles and Practices*, Third Edition. Boca Raton, FL, USA: CRC Press.
- Connor EF, Courtney AC and Yoder JM** (2000) Individuals–area relationships: the relationship between animal population density and area. *Ecology* **81**, 734–748.
- De Barro PJ, Liu SS, Boykin LM and Dinsdale AB** (2011) *Bemisia tabaci*: a statement of species status. *Annual Review of Entomology* **56**, 1–19.
- Döring TF** (2014) How aphids find their host plants, and how they don't. *Annals of Applied Biology* **165**, 3–26.
- Eastman R** (2020) *TerrSet 2020 – Geospatial Monitoring and Modeling System Manual*. Worcester, MA: Clark Laboratories, Clark University, 391 p.
- Ewers RM and Didham R** (2006) Continuous response functions for quantifying the strength of edge effects. *Journal of Applied Ecology* **43**, 527–536.
- Fahrig L** (2013) Rethinking patch size and isolation effects: the habitat amount hypothesis. *Journal of Biogeography* **40**, 1649–163.
- Fahrig L, Girard J, Duro D, Pasher J, Smith A, Javorek S, King D, Freemark Lidsay K, Mitchel S and Tischendorf L** (2015) Farmlands with smaller crop fields have higher within-field biodiversity. *Agriculture Ecosystem and Environment* **200**, 219–234.
- Forman RTT** (1995) *Land Mosaics: The Ecology of Landscapes and Regions*. Cambridge, UK: Cambridge University Press.
- Fox J** (2003) Effect displays in R for generalised linear models. *Journal of Statistical Software* **8**, 1–27. Available at <http://www.jstatsoft.org/v08/i15/>.
- Fox J and Hong J** (2009) Effect displays in R for Multinomial and Proportional-Odds Logit Models: Extensions to the effects Package. *Journal of Statistical Software* **32**, 1–24.
- Grille G, Gauthier N, Buenahora J, Basso C and Bonato O** (2011) First report of the Q biotype of *Bemisia tabaci* in Argentina and Uruguay. *Phytoparasitica* **39**, 235–238.
- Grilli MP and Bruno MA** (2007) Regional abundance of a planthopper pest: the effect of host patch area and configuration. *Entomologia Experimentalis et Applicata* **122**, 133–143.
- Grilli MP and Fachinetti R** (2017) The role of sex and mating status in the expansion process of *Arhopalus rusticus* (Coleoptera: Cerambycidae) – an exotic cerambycid in Argentina. *Environmental Entomology* **46**, 714–721.
- Grilli MP and Fachinetti R** (2019) Can forest pattern affect the distribution and abundance of *Arhopalus rusticus* (Coleoptera: Cerambycidae)? A landscape perspective in central Argentina. *International Journal of Pest Management* **65**, 268–275.
- Grueber CE, Nakagawa S, Laws RJ and Jamieson IG** (2011) Multimodel inference in ecology and evolution: challenges and solutions. *Journal of Evolutionary Biology* **24**, 699–711.
- Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, Holt RD, Lovejoy TE, Sexton JO, Austin MP, Collins CD, Cook WM, Damschen EI, Ewers RM, Foster BL, Jenkins CN, King AJ, Laurance WF, Levey DJ, Margules CR, Melbourne BA, Nicholls AO, Orrock JL, Song DX and Townshend JR** (2015) Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* **1**, e1500052.
- Hambäck PA and Englund G** (2005) Patch area, population density and the scaling of migration rates: the resource concentration hypothesis revisited. *Ecology Letters* **8**, 1057–1065.
- Helzer CJ and Jelinski DE** (1999) The relative importance of patch area and perimeter-area ratio to grassland breeding birds. *Ecological Applications* **9**, 1448–1458.
- Hogenhout SA, Ammar ED, Whitfield AE and Redinbaugh MG** (2008) Insect vector interactions with persistently transmitted viruses. *Annual Review of Phytopathology* **46**, 327–359.
- Huarte M and Capezio S** (2013) Cultivo de Papa. Unidad Integrada Balcarce INTA FCA UNMdP.CA. Available at [https://inta.gob.ar/sites/default/files/script-tmp-inta-huarte\\_capezio\\_papa2013.pdf](https://inta.gob.ar/sites/default/files/script-tmp-inta-huarte_capezio_papa2013.pdf).
- Kanakala S and Ghanim M** (2015) Advances in the genomics of the whitefly *Bemisia tabaci*: an insect pest and a virus vector. In *Short Views on Insect Genomics and Proteomics*, vol. 4. Cham, Switzerland: Springer International Publishing Switzerland, pp. 19–40. ISBN 978-3-319-24242-2.
- Krause-Sakate R, Watanabe LFM, Gorayeb ES, da Silva FB, Alvarez D, Bello VH, Nogueira AM, de Marchi BR, Vicentin E, Ribeiro-Junior MR, Marubayashi J, Rojas-Bertini CA, Muller C, Oliveira de Freitas Bueno RC, Rosales M, Ghanim M and Agenor Pavan M** (2020) Population dynamics of whiteflies and associated viruses in South America: research progress and perspectives. *Insects* **11**, 847.
- Landgrebe DA** (2003) *Signal Theory Methods in Multispectral Remote Sensing*. Hoboken, NY: Wiley, p. 528.
- Liu SS, De Barro PJ, Xu J, Luan JB, Zang LS, Ruan YM and Wan FH** (2007) Asymmetric mating interactions drive widespread invasion and displacement in a whitefly. *Science (New York, N.Y.)* **318**, 1769–1772.
- Ludwig M, Schlinkert H and Meyhofer R** (2018) Wind-modulated landscape effects on colonization of Brussels sprouts by insect pests and their syrphid antagonists. *Agricultural and Forest Entomology* **20**, 141–149.
- Macfadyen S, Paull C, Boykin LM, De Barro P, Maruthi MN, Otim M, Kalyebi A, Vassão DG, Sseruwagi P, Tay WT, Delatte H, Seguni Z, Colvin J and Omongo CA** (2018) Cassava whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) in East African farming landscapes: a review of the factors determining abundance. *Bulletin of Entomological Research* **20**, 1–18.
- Martin CA** (2018) An early synthesis of the habitat amount hypothesis. *Landscape Ecology* **33**, 1831–1835.
- McGarigal K and Ene E** (2015) Fragstats 4.2. A spatial pattern analysis program for categorical maps.
- Navas-Castillo J, Fiallo-Olive' E and Sanchez-Campos S** (2011) Emerging virus diseases transmitted by whiteflies. *Annual Review of Phytopathology* **49**, 219–248.
- Noldus LPJJ, Rumei X and van Lenteren JC** (1986) The parasite–host relationship between *Encarsia formosa* Gahan (Hymenoptera, Aphelinidae) and *Trialetrodes vaporariorum* (Westwood) (Homoptera, Aleyrodidae). XIX. Feeding-site selection by the greenhouse whitefly. *Journal of Applied Entomology* **101**, 492–507.
- Oliveira MRV, Henneberry TJ and Anderson P** (2001) History, current status, and collaborative research projects for *Bemisia tabaci*. *Crop Protection* **20**, 709–723.
- Perring TM** (2001) The *Bemisia tabaci* species complex. *Crop Protection* **20**, 725–737.
- Perring TM and Symmes EJ** (2006) Courtship behavior of *Bemisia argentifolii* (Hemiptera: Aleyrodidae) and whitefly mate recognition. *Annals of the Entomological Society of America* **99**, 598–606.
- Ploper LD, Laguna IG, Truol G and Rodríguez Pardini P** (1989) Infección doble con virus del mosaico de la soja (SMV) y un virus de partículas isométricas en cultivos de soja en la Provincia de Salta, Argentina. In Pascale AJ (ed.), *Proceedings World Soybean Research Conference IV*. Buenos Aires, Argentina: Editora S.R.L., pp. 2105–2111.
- Quattrini MM** (2005) *Cultivo de papa, en Olericultura, tomo II*. Córdoba: Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba, pp. 102–119.
- R Core Team** (2019) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Available at <http://www.R-project.org/>.
- Root RB** (1973) Organization of a plant–arthropod association in simple and diverse habitats: the fauna of collards (*Brassica oleracea*). *Ecological Monographs* **43**, 95–124.
- Sánchez C and Barberis NA** (2013) *Caracterización del territorio Centro de la provincia de Córdoba*. Córdoba, Argentina: Ediciones INTA. Estación Experimental Agropecuaria Manfredi.
- Saura S** (2021) The habitat amount hypothesis implies negative effects of habitat fragmentation on species richness. *Journal of Biogeography* **48**, 11–22.
- Stamps JA, Buechner M and Krishnan V** (1987) The effects of edge permeability and habitat geometry on emigration from patches of habitat. *American Naturalist* **129**, 533–552.
- Steffan-Dewenter I and Tscharrntke T** (2000) Butterfly community structure in fragmented habitats. *Ecology Letters* **3**, 449–456.
- Thies C, Steffan-Dewenter I and Tscharrntke T** (2003) Effects of landscape context on herbivory and parasitism at different spatial scales. *Oikos* **101**, 18–25.



**Turchin P** (1998) *Quantitative Analysis of Movement*. Sunderland, MA, USA: Sinauer Associates.

**Viscarret MM, Torres-Jerez II, Agostini de Manero E, Lopez SN, Botto EE and Brown JK** (2003) Mitochondrial DNA evidence for a distinct New

World group of *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) indigenous to Argentina and Bolivia, and presence of the Old World B Biotype Argentina. *Annals of the Entomological Society of America* **96**, 65–72.