



A global horizon scan of issues impacting marine and coastal biodiversity conservation

James E. Herbert-Read ^{1,37} ✉, Ann Thornton ^{2,37} ✉, Diva J. Amon ^{3,4}, Silvana N. R. Birchenough ⁵, Isabelle M. Côté ⁶, Maria P. Dias ^{7,8}, Brendan J. Godley⁹, Sally A. Keith ¹⁰, Emma McKinley ¹¹, Lloyd S. Peck ¹², Ricardo Calado¹³, Omar Defeo ¹⁴, Steven Degraer ¹⁵, Emma L. Johnston ¹⁶, Hermanni Kaartokallio¹⁷, Peter I. Macreadie ¹⁸, Anna Metaxas ¹⁹, Agnes W. N. Muthumbi²⁰, David O. Obura ^{21,22}, David M. Paterson²³, Alberto R. Piola ^{24,25}, Anthony J. Richardson ^{26,27}, Irene R. Schloss ^{28,29,30}, Paul V. R. Snelgrove ³¹, Bryce D. Stewart³², Paul M. Thompson ³³, Gordon J. Watson ³⁴, Thomas A. Worthington ², Moriaki Yasuhara ³⁵ and William J. Sutherland^{2,36}

The biodiversity of marine and coastal habitats is experiencing unprecedented change. While there are well-known drivers of these changes, such as overexploitation, climate change and pollution, there are also relatively unknown emerging issues that are poorly understood or recognized that have potentially positive or negative impacts on marine and coastal ecosystems. In this inaugural Marine and Coastal Horizon Scan, we brought together 30 scientists, policymakers and practitioners with transdisciplinary expertise in marine and coastal systems to identify new issues that are likely to have a significant impact on the functioning and conservation of marine and coastal biodiversity over the next 5–10 years. Based on a modified Delphi voting process, the final 15 issues presented were distilled from a list of 75 submitted by participants at the start of the process. These issues are grouped into three categories: ecosystem impacts, for example the impact of wildfires and the effect of poleward migration on equatorial biodiversity; resource exploitation, including an increase in the trade of fish swim bladders and increased exploitation of marine collagens; and new technologies, such as soft robotics and new biodegradable products. Our early identification of these issues and their potential impacts on marine and coastal biodiversity will support scientists, conservationists, resource managers and policymakers to address the challenges facing marine ecosystems.

The fifteenth Conference of the Parties (COP) to the United Nations Convention on Biological Diversity will conclude negotiations on a global biodiversity framework in late-2022 that will aim to slow and reverse the loss of biodiversity and establish goals for positive outcomes by 2050¹. Currently recognized drivers of declines in marine and coastal ecosystems include overexploitation of resources (for example, fishes, oil and gas), expansion of anthropogenic activities leading to cumulative impacts on the marine and coastal environment (for example, habitat loss, introduction of contaminants and pollution) and effects of climate change (for example, ocean warming, freshening and acidification). Within these broad categories, marine and coastal ecosystems face a wide range of emerging issues that are poorly recognized or understood, each having the potential to impact biodiversity. Researchers, conservation practitioners and marine resource managers must identify, understand and raise awareness of these relatively ‘unknown’ issues to catalyse further research into their underlying processes and impacts. Moreover, informing the public and policymakers of these issues can mitigate potentially negative impacts through precautionary principles before those effects become realized: horizon scans provide a platform to do this.

Horizon scans bring together experts from diverse disciplines to discuss issues that are (1) likely to have a positive or negative impact on biodiversity and conservation within the coming years and (2) not well known to the public or wider scientific community or face a substantial ‘step-change’ in their importance or application².

Horizon scans are an effective approach for pre-emptively identifying issues facing global conservation³. Indeed, marine issues previously identified through this approach include microplastics⁴, invasive lionfish⁴ and electric pulse trawling⁵. To date, however, no horizon scan of this type has focused solely on issues related to marine and coastal biodiversity, although a scan on coastal shorebirds in 2012 identified potential threats to coastal ecosystems⁶. This horizon scan aims to benefit our ocean and human society by stimulating research and policy development that will underpin appropriate scientific advice on prevention, mitigation, management and conservation approaches in marine and coastal ecosystems.

Results

We present the final 15 issues below in thematic groups identified post-scoring, rather than rank order (Fig. 1).

Ecosystem impacts. *Wildfire impacts on coastal and marine ecosystems.* The frequency and severity of wildfires are increasing with climate change⁷. Since 2017, there have been fires of unprecedented scale and duration in Australia, Brazil, Portugal, Russia and along the Pacific coast of North America. In addition to threatening human life and releasing stored carbon, wildfires release aerosols, particles and large volumes of materials containing soluble forms of nutrients including nitrogen, phosphorus and trace metals such as copper, lead and iron. Winds and rains can transport these materials over long distances to reach coastal and marine ecosystems.

Australian wildfires, for example, triggered widespread phytoplankton blooms in the Southern Ocean⁸ along with fish and invertebrate kills in estuaries⁹. Predicting the magnitude and effects of these acute inputs is difficult because they vary with the size and duration of wildfires, the burning vegetation type, rainfall patterns, riparian vegetation buffers, dispersal by aerosols and currents, seasonal timing and nutrient limitation in the recipient ecosystem. Wildfires might therefore lead to beneficial, albeit temporary, increases in primary productivity, produce no effect or have deleterious consequences, such as the mortality of benthic invertebrates, including corals, from sedimentation, coastal darkening (see below), eutrophication or algal blooms¹⁰.

Coastal darkening. Coastal ecosystems depend on the penetration of light for primary production by planktonic and attached algae and seagrass. However, climate change and human activities increase light attenuation through changes in dissolved materials modifying water colour and suspended particles. Increased precipitation, storms, permafrost thawing and coastal erosion have led to the 'browning' of freshwater ecosystems by elevated organic carbon, iron and particles, all of which are eventually discharged into the ocean¹¹. Coastal eutrophication leading to algal blooms compounds this darkening by further blocking light penetration. Additionally, land-use change, dredging and bottom fishing can increase seafloor disturbance, resuspending sediments and increasing turbidity. Such changes could affect ocean chemistry, including photochemical degradation of dissolved organic carbon and generation of toxic chemicals. At moderate intensities, limited spatial scales and during heatwaves, coastal darkening may have some positive impacts such as limiting coral bleaching on shallow reefs¹² but, at high intensities and prolonged spatial and temporal extents, lower light-regimes can contribute to cumulative stressor effects thereby profoundly altering ecosystems. This darkening may result in shifts in species composition, distribution, behaviour and phenology, as well as declines in coastal habitats and their functions (for example, carbon sequestration)¹³.

Increased toxicity of metal pollution due to ocean acidification. Concerns about metal toxicity in the marine environment are increasing as we learn more about the complex interactions between metals and global climate change¹⁴. Despite tight regulation of pollutants and remediation efforts in some countries, the high persistence of metals in contaminated sediments results in the ongoing remobilization of existing metal pollutants by storms, trawling and coastal development, augmented by continuing release of additional contaminants into coastal waters, particularly in urban and industrial areas across the globe¹⁴. Ocean acidification increases the bioavailability, uptake and toxicity of metals in seawater and sediments, with direct toxicity effects on some marine organisms¹⁵. Not all biogeochemical changes will result in increased toxicity; in pelagic and deep-sea ecosystems, where trace metals are often deficient, increasing acidity may increase bioavailability and, in shallow waters, stimulate productivity for non-calcifying phytoplankton¹⁶. However, increased uptake of metals in wild-caught and farmed bivalves linked to ocean acidification could also affect human health, especially given that these species provide 25% of the world's seafood. The combined effects of ocean acidification and metals could not only increase the levels of contamination in these organisms but could also impact their populations in the future¹⁴.

Equatorial marine communities are becoming depauperate due to climate migration. Climate change is causing ocean warming, resulting in a poleward shift of existing thermal zones. In response, species are tracking the changing ocean environmental conditions globally, with range shifts moving five times faster than on land¹⁷. In mid-latitudes and higher latitudes, as some species move away from

current distribution ranges, other species from warmer regions can replace them¹⁸. However, the hottest climatic zones already host the most thermally tolerant species, which cannot be replaced due to their geographical position. Thus, climate change reduces equatorial species richness and has caused the formerly unimodal latitudinal diversity gradient in many communities to now become bimodal. This bimodality (dip in equatorial diversity) is projected to increase within the next 100 years if carbon dioxide emissions are not reduced¹⁹. The ecological consequences of this decline in equatorial zones are unclear, especially when combined with impacts of increasing human extraction and pollution²⁰. Nevertheless, emerging ecological communities in equatorial systems are likely to have reduced resilience and capacity to support ecosystem services and human livelihoods.

Effects of altered nutritional content of fish due to climate change. Essential fatty acids (EFAs) are critical to maintaining human and animal health and fish consumption provides the primary source of EFAs for billions of people. In aquatic ecosystems, phytoplankton synthesize EFAs, such as docosahexaenoic acid (DHA)²¹, with pelagic fishes then consuming phytoplankton. However, concentrations of EFAs in fishes vary, with generally higher concentrations of omega-3 fatty acids in slower-growing species from colder waters²². Ongoing effects of climate change are impacting the production of EFAs by phytoplankton, with warming waters predicted to reduce the availability of DHA by about 10–58% by 2100²³; a 27.8% reduction in available DHA is associated with a 2.5°C rise in water temperature²¹. Combined with geographical range shifts in response to environmental change affecting the abundance and distribution of fishes, this could lead to a reduction in sufficient quantities of EFAs for fishes, particularly in the tropics²⁴. Changes to EFA production by phytoplankton in response to climate change, as shown for Antarctic waters²⁵, could have cascading effects on the nutrient content of species further up the food web, with consequences for marine predators and human health²⁶.

Resource exploitation. *The untapped potential of marine collagens and their impacts on marine ecosystems.* Collagens are structural proteins increasingly used in cosmetics, pharmaceuticals, nutraceuticals and biomedical applications. Growing demand for collagen has fuelled recent efforts to find new sources that avoid religious constraints and alleviate risks associated with disease transmission from conventional bovine and porcine sources²⁷. The search for alternative sources has revealed an untapped opportunity in marine organisms, such as from fisheries bycatch²⁸. However, this new source may discourage efforts to reduce the capture of non-target species. Sponges and jellyfish offer a premium source of marine collagens. While the commercial-scale harvesting of sponges is unlikely to be widely sustainable, there may be some opportunity in sponge aquaculture and jellyfish harvesting, especially in areas where nuisance jellyfish species bloom regularly (for example, Mediterranean and Japan Seas). The use of sharks and other cartilaginous fish to supply marine collagens is of concern given the unprecedented pressure on these species. However, the use of coproducts derived from the fish-processing industry (for example, skin, bones and trims) offers a more sustainable approach to marine collagen production and could actively contribute to the blue bio-economy agenda and foster circularity²⁹.

Impacts of expanding trade for fish swim bladders on target and non-target species. In addition to better-known luxury dried seafoods, such as shark fins, abalone and sea cucumbers, there is an increasing demand for fish swim bladders, also known as fish maw³⁰. This demand may trigger an expansion of unsustainable harvests of target fish populations, with additional impacts on marine biodiversity through bycatch^{30,31}. The fish swim-bladder trade has gained




Fig. 1 | The 15 horizon issues presented in thematic groups: ecosystem impacts, resource exploitation and new technologies. Numbers refer to the order presented in this article, rather than final ranking. Image of brine pool courtesy of the NOAA Office of Ocean Exploration and Research, Gulf of Mexico 2014. Image of biodegradable bag courtesy of Katie Dunkley.

a high profile because the overexploitation of totoaba (*Totoaba macdonaldi*) has driven both the target population and the vaquita (*Phocoena sinus*) (which is bycaught in the Gulf of Mexico fishery) to near extinction³². By 2018, totoaba swim bladders were being sold for US\$46,000 kg⁻¹. This extremely lucrative trade disrupts efforts to encourage sustainable fisheries. However, increased demand on the totoaba was itself caused by overexploitation over the last century of the closely related traditional species of choice, the Chinese bahaba (*Bahaba taipingensis*). We now risk both repeating this pattern and increasing its scale of impact, where depletion of a target species causes markets to switch to species across broader taxonomic and biogeographical ranges³¹. Not only does this cascading effect threaten other croakers and target species, such as catfish and pufferfish but maw nets set in more diverse marine habitats are likely to create bycatch of sharks, rays, turtles and other species of conservation concern.

Impacts of fishing for mesopelagic species on the biological ocean carbon pump. Growing concerns about food security have generated interest in harvesting largely unexploited mesopelagic fishes that live at depths of 200–1,000 m (ref. ³³). Small lanternfishes (Myctophidae) dominate this potentially 10 billion ton community, exceeding the mass of all other marine fishes combined³⁴ and spanning millions of square kilometres of the open ocean. Mesopelagic fish are generally unsuitable for human consumption but could potentially provide fishmeal for aquaculture³⁴ or be used for fertilizers. Although we know little of their biology, their diel vertical migration transfers carbon, obtained by feeding in surface waters at night, to deeper waters during the day across many hundreds and even thousands of

metres depth where it is released by excretion, egestion and death. This globally important carbon transport pathway contributes to the biological pump³⁵ and sequesters carbon to the deep sea³⁶. Recent estimates put the contribution of all fishes to the biological ocean pump at 16.1% (\pm s.d. 13%) (ref. ³⁷). The potential large-scale removal of mesopelagic fishes could disrupt a major pathway of carbon transport into the ocean depths.

Extraction of lithium from deep-sea brine pools. Global groups, such as the Deep-Ocean Stewardship Initiative, emphasize increasing concern about the ecosystem impacts from deep-sea resource extraction³⁸. The demand for batteries, including for electric vehicles, will probably lead to a demand for lithium that is more than five times its current level by 2030³⁹. While concentrations are relatively low in seawater, some deep-sea brines and cold seeps offer higher concentrations of lithium. Furthermore, new technologies, such as solid-state electrolyte membranes, can enrich the concentration of lithium from seawater sources by 43,000 times, increasing the energy efficiency and profitability of lithium extraction from the sea³⁹. These factors could divert extraction of lithium resources away from terrestrial to marine mining, with the potential for significant impacts to localized deep-sea brine ecosystems. These brine pools probably host many endemic and genetically distinct species that are largely undiscovered or awaiting formal description. Moreover, the extremophilic species in these environments offer potential sources of marine genetic resources that could be used in new biomedical applications including pharmaceuticals, industrial agents and biomaterials⁴⁰. These concerns point to the need to better quantify and




Fig. 2 | Stepwise process used to identify, score and present the 15 horizon issues likely to impact marine and coastal biodiversity conservation in the next 5–10 years. Left and right columns show the process for the first and second rounds of scoring, respectively.

monitor biodiversity in these extreme environments to establish baselines and aid management.

New technologies. *Colocation of marine activities.* Climate change, energy needs and food security have moved to the top of global policy agendas⁴¹. Increasing energy needs, alongside the demands of fisheries and transport infrastructure, have led to the proposal of collocated and multifunctional structures to deliver economic benefits, optimize spatial planning and minimize the environmental impacts of marine activities⁴². These designs often bring technical, social, economic and environmental challenges. Some studies have begun to explore these multipurpose projects (for example, offshore windfarms collocated with aquaculture developments and/or Marine Protected Areas) and how to adapt these concepts to ensure they are 'fit for purpose', economically viable and reliable. However, environmental and ecosystem assessment, management and regulatory frameworks for collocated and multi-use structures need to be established to prevent

these activities from compounding rather than mitigating the environmental impacts from climate change⁴³.

Floating marine cities. In April 2019, the UN-HABITAT programme convened a meeting of scientists, architects, designers and entrepreneurs to discuss how floating cities might be a solution to urban challenges such as climate change and lack of housing associated with a rising human population (<https://unhabitat.org/roundtable-on-floating-cities-at-unhq-calls-for-innovation-to-benefit-all>). The concept of floating marine cities—hubs of floating structures placed at sea—was born in the middle of the twentieth century and updated designs now aim to translate this vision into reality⁴⁴. Oceanic locations provide benefits from wave and tidal renewable energy and food production supported by hydroponic agriculture⁴⁵. Modular designs also offer greater flexibility than traditional static terrestrial cities, whereby accommodation and facilities could be incorporated or removed in response to changes in population or specific events. The cost of construction in harsh offshore environ-




Fig. 3 | Median rank of each issue versus proportion of issues participants had previously heard of. a, Round 1. Each point represents an individual issue. For all issue titles, see Supplementary Table 1. Issues in dark blue were retained for the second round. Issues that were ranked higher were generally those that participants had not heard of (Spearman rank correlation = 0.38, $P < 0.001$). **b**, Round 2. Scores as in round 1. For titles of the second round of 32 issues, see Supplementary Table 2. The 15 final issues (marked in red) achieved the top ranks (horizontal dashed line) and had only been heard of by 50% of participants (vertical dashed line). Red circles, squares and triangles denote issues relating to ecosystem impacts, resource exploitation and new technologies, respectively. The two grey issues marked with crosses were discounted during final discussions because participants could not identify the horizon component of these issues.

ments, rather than technology, currently limits the development of marine cities and potential designs will need to consider the consequences of more frequent and extreme climate events. Although the artificial hard substrates created for these floating cities could act as stepping stones, facilitating species movement in response to climate change⁴⁶, this could also increase the spread of invasive species. Finally, the development of offshore living will raise issues in relation to governance and land ownership that must be addressed for marine cities to be viable⁴⁷.

Trace-element contamination compounded by the global transition to green technologies. The persistent environmental impacts of metal and metalloid trace-element contamination in coastal sediments are now increasing after a long decline⁴⁸. However, the complex sources of contamination challenge their management. The acceleration of the global transition to green technologies, including electric vehicles, will increase demand for batteries by over 10% annually in the coming years⁴⁹. Electric vehicle batteries currently depend almost exclusively on lithium-ion chemistries, with potential trace-element emissions across their life cycle from raw material extraction to recycling or end-of-life disposal. Few jurisdictions treat lithium-ion batteries as harmful waste, enabling landfill disposal with minimal recycling⁴⁹. Cobalt and nickel are the primary ecotoxic elements in next-generation lithium-ion batteries⁵⁰, although there is a drive to develop a cobalt-free alternative likely to contain higher nickel content⁵⁰. Some battery binder and electrolyte chemicals are toxic to aquatic life or form persistent organic pollutants during incomplete burning. Increasing pollution from battery production, recycling and disposal in the next decade could substantially increase the potentially toxic trace-element contamination in marine and coastal systems worldwide.

New underwater tracking systems to study non-surfacing marine animals. The use of tracking data in science and conservation has grown exponentially in recent decades. Most trajectory data collected on marine species to date, however, has been restricted to

large and near-surface species, limited by the size of the devices and reliance on radio signals that do not propagate well underwater. New battery-free technology based on acoustic telemetry, named ‘underwater backscatter localization’ (UBL), may allow high-accuracy (<1 m) tracking of animals travelling at any depth and over large distances⁵¹. Still in the early stages of development, UBL technology has significant potential to help fill knowledge gaps in the distribution and spatial ecology of small, non-surfacing marine species, as well as the early life-history stages of many species⁵², over the next decades. However, the potential negative impacts of this methodology on the behaviour of animals are still to be determined. Ultimately, UBL may inform spatial management both in coastal and offshore regions, as well as in the high seas and address a currently biased perspective of how marine animals use ocean space, which is largely based on near-surface or aerial marine megafauna (for example, ref. ⁵³).

Soft robotics for marine research. The application and utility of soft robotics in marine environments is expected to accelerate in the next decade. Soft robotics, using compliant materials inspired by living organisms, could eventually offer increased flexibility at depth because they do not face the same constraints as rigid robots that need pressurized systems to function⁵⁴. This technology could increase our ability to monitor and map the deep sea, with both positive and negative consequences for deep-sea fauna. Soft-grab robots could facilitate collection of delicate samples for biodiversity monitoring but, without careful management, could also add pollutants and waste to these previously unexplored and poorly understood environments⁵⁵. With advancing technology, potential deployment of swarms of small robots could collect basic environmental data to facilitate mapping of the seabed. Currently limited by power supply, energy-harvesting modules are in development that enable soft robots to ‘swallow’ organic material and convert it into power⁵⁶, although this could result in inadvertently harvesting rare deep-sea organisms. Soft robots themselves may also be ingested by predatory species mistaking them for prey. Deployment of soft robotics will require careful monitoring of both its benefits and risks to marine biodiversity.

The effects of new biodegradable materials in the marine environment. Mounting public pressure to address marine plastic pollution has prompted the replacement of some fossil fuel-based plastics with bio-based biodegradable polymers. This consumer pressure is creating an economic incentive to adopt such products rapidly and some companies are promoting their environmental benefits without rigorous toxicity testing and/or life-cycle assessments. Materials such as polybutylene succinate (PBS), polylactic acid (PLA) or cellulose and starch-based materials may become marine litter and cause harmful effects akin to conventional plastics⁵⁷. The long-term and large-scale effect of the use of biodegradable polymers in products (for example, clothing) and the unintended release of byproducts, such as microfibres, into the environment remain unknown. However, some natural microfibres have greater toxicity than plastic microfibres when consumed by aquatic invertebrates⁵⁸. Jurisdictions should enact and enforce suitable regulations to require the individual assessment of all new materials intended to biodegrade in a full range of marine environmental conditions. In addition, testing should include studies on the toxicity of major transition chemicals created during the breakdown process⁵⁹, ideally considering the different trophic levels of marine food webs.

Discussion

This scan identified three categories of horizon issues: impacts on, and alterations to, ecosystems; changes to resource use and extraction; and the emergence of technologies. While some of the issues discussed, such as improved monitoring of species (underwa-

ter tracking and soft robotics) and more sustainable resource use (marine collagens), may have some positive outcomes for marine and coastal biodiversity, most identified issues are expected to have substantial negative impacts if not managed or mitigated appropriately. This imbalance highlights the considerable emerging pressures facing marine ecosystems that are often a byproduct of human activities.

Four issues identified in this scan related to ongoing large-scale (hundreds to many thousands of square kilometres) alterations to marine ecosystems (wildfires, coastal darkening, depauperate equatorial communities and altered nutritional fish content), either through the impacts of global climate change or other human activities. There are already clear impacts of climate change, for example, on stores of blue carbon (for example, ref. ⁶⁰) and small-scale fisheries (for example, ref. ⁶¹) but the identification of these issues highlights the need for global action that reverses such trends. The United Nations Decade of Ocean Science for Sustainable Development (2021–2030) is now underway, aligning with other decadal policy priorities, including the Sustainable Development Goals (<https://sdgs.un.org/>), the 2030 targets for biodiversity to be agreed in 2022, the conclusion of the ongoing negotiations on biodiversity beyond national jurisdictions (BBNJ) (<https://www.un.org/bbnj/>), the UN Conference on Biodiversity (COP15) (<https://www.unep.org/events/conference/un-biodiversity-conference-cop-15>) and the UN Climate Change Conference 2021 (COP26) (<https://ukcop26.org/>). While some campaigns to allocate 30% of the ocean to Marine Protected Areas by 2030 are prominently aired⁶², the unintended future consequences of such protection and how to monitor and manage these areas, remain unclear^{63–65}.

Another set of issues related to anticipated increases in marine resource use and extraction (swim bladders, marine collagens, lithium extraction and mesopelagic fisheries). The complex issue of mitigating the impacts on marine conservation and biodiversity of exploiting and using newly discovered resources must consider public perceptions of the ocean^{66,67}, market forces and the sustainable blue economy^{68,69}.

The final set of issues related to new technological advancements, with many offering more sustainable opportunities, albeit some having potentially unintended negative consequences on marine and coastal biodiversity. For example, trace-element contamination from green technologies and harmful effects of biodegradable products highlights the need to assess the step-changes in impacts from their increased use and avoid the paradox of technologies designed to mitigate the damaging effects of climate change on biodiversity themselves damaging biodiversity. Indeed, the impacts on marine and coastal biodiversity from emerging technologies currently in development (such as underwater tracking or soft robotics) need to be assessed before deployment at scale.

There are limitations to any horizon scanning process that aims to identify global issues and a different group of experts may have identified a different set of issues. By inviting participants from a range of subject backgrounds and global regions and asking them to canvass their network of colleagues and collaborators, we aimed to identify as broad a set of issues as possible. We acknowledge, however, that only about one-quarter of the participants were from non-academic organizations, which may have skewed the submitted issues and how they were voted on. However, others³ reported no significant correlation between participants' areas of research expertise and the top issues selected in the horizon scan conducted in 2009. Therefore, horizon scans do not necessarily simply represent issues that reflect the expertise of participants. We also sought to achieve diversity by inviting participants from 22 countries and actively seeking representatives from the global south. However, the final panel of 30 participants spanned only 11 countries, most in the global north. We were forced by the COVID-19 pandemic to hold the scan online and while we hoped that this would enable

participants to engage from around the world alleviating broader global inequalities in science⁶³, digital inequality was in fact enhanced during the pandemic⁷⁰. Our experience highlights the need for other mechanisms that can promote global representation in these scans.

This Marine and Coastal Horizon Scan seeks to raise awareness of issues that may impact marine and coastal biodiversity conservation in the next 5–10 years. Our aim is to bring these issues to the attention of scientists, policymakers, practitioners and the wider community, either directly, through social networks or the mainstream media. Whilst it is almost impossible to determine whether issues gained prominence as a direct result of a horizon scan, some issues featured in previous scans have seen growth in reporting and awareness. Others³ found that 71% of topics identified in the Horizon Scan in 2009 had seen an increase in their importance over the next 10 years. Issues such as microplastics and invasive lionfish had received increased research and investment from scientists, funders, managers and policymakers to understand their impacts and the horizon scans may have helped motivate this increase. Horizon scans, therefore, should primarily act as signposts, putting focus onto particular issues and providing support for researchers and practitioners to seek investment in these areas.

Whilst recognizing that marine and coastal environments are complex social-ecological systems, the role of governance, policy and litigation on all areas of marine science needs to be developed, as it is yet to be established to the same extent as in terrestrial ecosystems⁷¹. Indeed, tackling many of the issues presented in this scan will require an understanding of the human dimensions relating to these issues, through fields of research including but not limited to ocean literacy^{72,73}, social justice, equity⁷⁴ and human health⁷⁵. Importantly, however, horizon scanning has proved an efficient tool in identifying issues that have subsequently come to the forefront of public knowledge and policy decisions, while also helping to focus future research. The scale of the issues facing marine and coastal areas emphasizes the need to identify and prioritize, at an early stage, those issues specifically facing marine ecosystems, especially within this UN Decade of Ocean Science for Sustainable Development.

Methods

Identification of issues. In March 2021, we brought together a core team of 11 participants from a broad range of marine and coastal disciplines. The core team suggested names of individuals outside their subject area who were also invited to participate in the horizon scan. To ensure we included as many different subject areas as possible within marine and coastal conservation, we selected one individual from each discipline. Our panel of experts comprised 30 (37% female) marine and coastal scientists, policymakers and practitioners (27% from non-academic institutions), with cross-disciplinary expertise in ecology (including tropical, temperate, polar and deep-sea ecosystems), palaeoecology, conservation, oceanography, climate change, ecotoxicology, technology, engineering and marine social sciences (including governance, blue economy and ocean literacy). Participants were invited from 22 countries across six continents, resulting in a final panel of 30 experts from 11 countries (Europe $n = 17$ (including the three organizers); North America and Caribbean $n = 4$; South America $n = 3$; Australasia $n = 3$; Asia $n = 1$; Africa $n = 2$). All experts co-authored this paper.

To reduce the potential for bias in the identification of suitable issues, each participant was invited to consult their own network and required to submit two to five issues that they considered new and likely to have a positive or negative impact on marine and coastal biodiversity conservation in the next 5–10 years (Supplementary Information text describes instructions given to participants). Each issue was described in paragraphs of ~200 words (plus references). Due to the COVID-19 pandemic, participants relied mainly on virtual meetings and online communication using email, social-media platforms, online conferences and networking events. Through these channels ~680 people were canvassed by the participants, counting all direct in-person or online discussions as individual contacts but treating social-media posts or generic emails as a single contact. This process resulted in a long list of 75 issues that were considered in the first round of scoring (see Supplementary Table 1 for the full list of initially submitted issues).

Round 1 scoring. The initial list of proposed issues was then shortened through a scoring process. We used a modified Delphi-style⁷⁶ voting process, which has

been consistently applied in horizon scans since 2009 (refs. 477) (see Fig. 2 for the stepwise process). This process ensured that consideration and selection of issues remained repeatable, transparent and inclusive. Panel members were asked to confidentially and independently score the long list of 75 issues from 1 (low) to 1,000 (high) on the basis of the following criteria:

- Whether the issue is new (with 'new' issues scoring higher) or is a well-known issue likely to exhibit a significant step-change in impact
- Whether the issue is likely to be important and impactful over the next 5–10 years
- Whether the issue specifically impacts marine and coastal biodiversity

Participants were also asked whether they had heard of the issue or not. 'Voter fatigue' can result in issues at the end of a lengthy list not receiving the same consideration as those at the beginning⁶. We counteracted this potential bias by randomly assigning participants to one of three differently ordered long-lists of issues. Participants' scores were converted to ranks (1–75). We had aimed to retain the top 30 issues with the highest median ranks for the second round of assessment at the workshop but kept 31 issues because two issues achieved equal median ranks. In addition, we identified one issue that had been incorrectly grouped with three others and presented this as a separate issue. The subsequent online workshop to discuss this shortlist, therefore, considered the top-ranked 32 issues (Fig. 3a) (see Supplementary Table 2 for the full list).

Workshop and round 2 scoring. Before the workshop, each participant was assigned up to four of the 32 issues to research in more detail and contribute further information to the discussion. We convened a one-day workshop online in September 2021. The geographic spread of participants meant that time zones spanned 17 h. Despite these constraints, discussions remained detailed, focused, varied and lively. In addition, participants made use of the chat function on the platform to add notes, links to articles and comments to the discussion. After discussing each issue, participants re-scored the topic (1–1,000, low to high) based on novelty and the issue's importance for, and probable impact on, marine and coastal biodiversity (3 participants out of 30 did not score all issues and therefore their scores were discounted). At the end of the selection process, scores were again converted to ranks and collated. Highest-ranked issues were then discussed by correspondence focusing on the same three criteria as outlined above, after which the top 15 horizon issues were selected (Fig. 3b).

Reporting summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The datasets generated during and/or analysed during the current study are available from figshare <https://doi.org/10.6084/m9.figshare.19703485.v1>. Source data are provided with this paper.

Received: 12 November 2021; Accepted: 24 May 2022;

Published online: 07 July 2022

References

- Diaz, S. et al. Set ambitious goals for biodiversity and sustainability. *Science* **370**, 411–413 (2020).
- Sutherland, W. J. & Woodroof, H. J. The need for environmental horizon scanning. *Trends Ecol. Evol.* **24**, 523–527 (2009).
- Sutherland, W. J. et al. Ten years on: a review of the first global conservation horizon scan. *Trends Ecol. Evol.* **34**, 139–153 (2019).
- Sutherland, W. J. et al. A horizon scan of global conservation issues for 2010. *Trends Ecol. Evol.* **25**, 1–7 (2010).
- Sutherland, W. J. et al. A horizon scan of global conservation issues for 2016. *Trends Ecol. Evol.* **31**, 44–53 (2016).
- Sutherland, W. J. et al. A horizon scanning assessment of current and potential future threats facing migratory shorebirds. *Ibis* **154**, 663–679 (2012).
- Bowman, D. M. J. S. et al. Vegetation fires in the Anthropocene. *Nat. Rev. Earth Environ.* **1**, 500–515 (2020).
- Tang, W. et al. Widespread phytoplankton blooms triggered by 2019–2020 Australian wildfires. *Nature* **597**, 370–375 (2021).
- Silva, L. G. M. et al. Mortality events resulting from Australia's catastrophic fires threaten aquatic biota. *Glob. Change Biol.* **26**, 5345–5350 (2020).
- Abram, N. J., Gagan, M. K., McCulloch, M. T., Chappell, J. & Hantoro, W. S. Coral reef death during the 1997 Indian Ocean Dipole linked to Indonesian wildfires. *Science* **301**, 952–955 (2003).
- Solomon, C. T. et al. Ecosystem consequences of changing inputs of terrestrial dissolved organic matter to lakes: current knowledge and future challenges. *Ecosystems* **18**, 376–389 (2015).
- Sully, S. & van Woerk, R. Turbid reefs moderate coral bleaching under climate related temperature stress. *Glob. Change Biol.* **26**, 1367–1373 (2021).
- Blain, C. O., Hansen, S. C. & Shears, N. T. Coastal darkening substantially limits the contribution of kelp to coastal carbon cycles. *Glob. Change Biol.* **27**, 5547–5563 (2021).
- Stewart, B. D. et al. Metal pollution as a potential threat to shell strength and survival in marine bivalves. *Sci. Total Environ.* **755**, 143019 (2021).
- Roberts, D. A. et al. Ocean acidification increases the toxicity of contaminated sediments. *Glob. Change Biol.* **19**, 340–351 (2013).
- Hauton, C. et al. Identifying toxic impact of metals potentially released during deep-sea mining—a synthesis of the challenges to quantifying risk. *Front. Mar. Sci.* **4**, 368 (2017).
- Chaudhary, C. et al. Global warming is causing a more pronounced dip in marine species richness around the equator. *Proc. Natl Acad. Sci. USA* **118**, e2015094118 (2021).
- Burrows, M. T. et al. Geographical limits to species-range shifts are suggested by climate velocity. *Nature* **507**, 492–495 (2014).
- Yasuhara, M. et al. Past and future decline of tropical pelagic biodiversity. *Proc. Natl Acad. Sci. USA* **117**, 12891–12896 (2020).
- Pandolfi, J. M. et al. Are U.S. coral reefs on the slippery slope to slime? *Science* **307**, 1725–1726 (2005).
- Hixson, S. M. & Arts, M. T. Climate warming is predicted to reduce omega-3, long-chain, polyunsaturated fatty acid production in phytoplankton. *Glob. Change Biol.* **22**, 2744–2755 (2016).
- Hicks, C. C. et al. Harnessing global fisheries to tackle micronutrient deficiencies. *Nature* **574**, 95–98 (2019).
- Colombo, S. M. et al. Projected declines in global DHA availability for human consumption as a result of global warming. *Ambio* **49**, 865–880 (2020).
- Lam, V. W. et al. Climate change, tropical fisheries and prospects for sustainable development. *Nat. Rev. Earth Environ.* **1**, 440–454 (2020).
- Antacl, J. C. et al. Increase in unsaturated fatty acids in Antarctic phytoplankton under ocean warming and glacial melting scenarios. *Sci. Total Environ.* **790**, 147879 (2021).
- Maire, E. et al. Micronutrient supply from global marine fisheries under climate change and overfishing. *Curr. Biol.* **18**, 4132–4138 (2021).
- Lim, Y. S., Ok, Y. J., Hwang, S. Y., Kwak, J. Y. & Yoon, S. Marine collagen as a promising biomaterial for biomedical applications. *Mar. Drugs* **17**, 467 (2019).
- Xu, N. et al. Marine-derived collagen as biomaterials for human health. *Front. Nutr.* **8**, 702108 (2021).
- Vieira, H., Leal, M. C. & Calado, R. Fifty shades of blue: how blue biotechnology is shaping the bioeconomy. *Trends Biotechnol.* **38**, 940–943 (2020).
- Ben-Hasan, A. et al. China's fish maw demand and its implications for fisheries in source countries. *Mar. Policy* **132**, 104696 (2021).
- Sadovy de Mitcheson, Y., To, A. W. L., Wong, N. W., Kwan, H. Y. & Bud, W. S. Emerging from the murk: threats, challenges and opportunities for the global swim bladder trade. *Rev. Fish. Biol. Fish.* **29**, 809–835 (2019).
- Brownell, R. L. Jr et al. Bycatch in gillnet fisheries threatens critically endangered small cetaceans and other aquatic megafauna. *Endang. Species Res.* **40**, 285–296 (2019).
- Webb, T. J., Vanden Berghe, E. & O'Dor, R. K. Biodiversity's big wet secret: the global distribution of marine biological records reveals chronic under-exploration of the deep pelagic ocean. *PLoS ONE* **5**, e10223 (2010).
- St. John, M. A. et al. A dark hole in our understanding of marine ecosystems and their services: perspectives from the mesopelagic community. *Front. Mar. Sci.* **3**, 31 (2016).
- Thomsen, L. et al. The oceanic biological pump: rapid carbon transfer to depth at continental margins during winter. *Sci. Rep.* **7**, 10763 (2017).
- Roberts, C. M., Hawkins, J. P., Hindle, K., Wilson, R. W. & O'Leary, B. C. *Entering the Twilight Zone: The Ecological Role and Importance of Mesopelagic Fishes* (Blue Marine Foundation, 2020).
- Cavan, E. L., Laurenceau-Cornec, E. C., Bressac, M. & Boyd, P. W. Exploring the ecology of the mesopelagic biological pump. *Prog. Oceanogr.* **176**, 102125 (2019).
- Levin, L. A. et al. Climate change considerations are fundamental to management of deep-sea resource extraction. *Glob. Change Biol.* **26**, 4664–4678 (2020).
- Li, Z. et al. Continuous electrical pumping membrane process for seawater lithium mining. *Energy Environ. Sci.* **14**, 3152–3159 (2021).
- Lin, M., Gai, Y., Guo, X., Hou, Y. & Zeng, R. Properties and applications of extremozymes from deep-sea extremophilic microorganisms: a mini review. *Mar. Drugs* **17**, 656 (2019).
- Mbow, C. et al. in *IPCC Special Report on Climate Change and Land* (eds Shukla, P. R. et al.) 437–550 (IPCC, 2019).
- Christie, N., Smyth, K., Barnes, R. & Elliott, M. Co-location of activities and designations: a means of solving or creating problems in marine spatial planning? *Mar. Pol.* **43**, 254–261 (2014).
- Mayer-Pinto, M., Dafforn, K. A. & Johnston, E. L. A decision framework for coastal infrastructure to optimize biotic resistance and resilience in a changing climate. *BioScience* **69**, 833–843 (2019).
- Wang, C. M. & Wang, B. T. in *ICSCEA 2019* (eds Reddy, J. N. et al.) 3–29 (Springer, 2020).
- Ross, C. T. F. & McCullough, R. R. Conceptual design of a floating island city. *J. Ocean Technol.* **5**, 120–121 (2010).

46. Dong, Y.-w, Huang, X.-w, Wang, W., Li, Y. & Wang, J. The marine 'great wall' of China: local- and broad-scale ecological impacts of coastal infrastructure on intertidal macrobenthic communities. *Divers. Distrib.* **22**, 731–744 (2016).
47. Flikkema, M. M. B., Lin, F.-Y., van der Plank, P. P. J., Koning, J. & Waals, O. Legal issues for artificial floating islands. *Front. Mar. Sci.* **8**, 619462 (2021).
48. Richir, J., Bray, S., McAleese, T. & Watson, G. J. Three decades of trace element sediment contamination: the mining of governmental databases and the need to address hidden sources for clean and healthy seas. *Environ. Int.* **149**, 106362 (2021).
49. Zhao, Y. et al. A review on battery market trends, second-life reuse, and recycling. *Sustain. Chem.* **2**, 167–205 (2021).
50. Li, W., Lee, S. & Manthiram, A. High-Nickel NMA: a cobalt-free alternative to NMC and NCA cathodes for lithium-ion batteries. *Adv. Mater.* **32**, 2002718 (2020).
51. Ghaffarivardavagh, R., Afzal, S. S., Rodriguez, O. & Adib, F. in *SIGCOMM '20 Proc. 19th ACM Workshop on Hot Topics in Networks* 125–131 (Association for Computing Machinery, 2020).
52. Hazen, E. L. et al. Ontogeny in marine tagging and tracking science: technologies and data gaps. *Mar. Ecol. Prog. Ser.* **457**, 221–240 (2012).
53. Davies, T. E. et al. Tracking data and the conservation of the high seas: opportunities and challenges. *J. Appl. Ecol.* **58**, 2703–2710 (2021).
54. Aracri, S. et al. Soft robots for ocean exploration and offshore operations: a perspective. *Soft Robot.* <https://doi.org/10.1089/soro.2020.0011> (2021).
55. Li, G. et al. Self-powered soft robot in the Mariana Trench. *Nature* **591**, 66–71 (2021).
56. Philamore, H., Ieropoulos, I., Stinchcombe, A. & Rossiter, J. Toward energetically autonomous foraging soft robots. *Soft Robot.* **3**, 186–197 (2016).
57. Manfra, L. et al. Biodegradable polymers: a real opportunity to solve marine plastic pollution? *J. Hazard. Mater.* **416**, 125763 (2021).
58. Kim, D., Kim, H. & An, Y. J. Effects of synthetic and natural microfibers on *Daphnia magna*: are they dependent on microfiber type? *Aquat. Toxicol.* **240**, 105968 (2021).
59. Degli-Innocenti, F., Bellia, G., Tosin, M., Kapanen, A. & Itävaara, M. Detection of toxicity released by biodegradable plastics after composting in activated vermiculite. *Polym. Degrad. Stab.* **73**, 101–106 (2001).
60. Macreadie, P. I. et al. The future of blue carbon science. *Nat. Commun.* **10**, 3998 (2019).
61. Short, R. E. et al. Harnessing the diversity of small-scale actors is key to the future of aquatic food systems. *Nat. Food* **2**, 733–741 (2021).
62. Watson, J. E. M. et al. Set a global target for ecosystems. *Nature* **578**, 360–362 (2020).
63. Obura, D. O. et al. Integrate biodiversity targets from local to global levels. *Science* **373**, 746 (2021).
64. Barnes, M. D., Glew, L., Wyborn, C. & Craigie, I. D. Prevent perverse outcomes from global protected area policy. *Nat. Ecol. Evol.* **2**, 759–762 (2018).
65. Grorud-Colvert, K. et al. The MPA Guide: a framework to achieve global goals for the ocean. *Science* **373**, eabf0861 (2021).
66. Jefferson, R. L., McKinley, E., Griffin, H., Nimmo, A. & Fletcher, S. Public perceptions of the ocean: lessons for marine conservation from a global research review. *Front. Mar. Sci.* **8**, 711245 (2021).
67. Potts, T., Pita, C., O'Higgins, T. & Mee, L. Who cares? European attitudes towards marine and coastal environments. *Mar. Pol.* **72**, 59–66 (2016).
68. Bennett, N. J. et al. Towards a sustainable and equitable blue economy. *Nat. Sustain.* **2**, 991–993 (2019).
69. Jouffray, J.-B., Blasiak, R., Norström, A. V., Österblom, H. & Nyström, M. The blue acceleration: the trajectory of human expansion into the ocean. *One Earth* **2**, 43–54 (2020).
70. Zheng, Y. & Walsham, G. Inequality of what? An intersectional approach to digital inequality under Covid-19. *Inf. Organ.* **31**, 100341 (2021).
71. Blythe, J. L., Armitage, D., Bennett, N. J., Silver, J. J. & Song, A. M. The politics of ocean governance transformations. *Front. Mar. Sci.* **8**, 634718 (2021).
72. Brennan, C., Ashley, M. & Molloy, O. A system dynamics approach to increasing ocean literacy. *Front. Mar. Sci.* **6**, 360 (2019).
73. Stoll-Kleemann, S. Feasible options for behavior change toward more effective ocean literacy: a systematic review. *Front. Mar. Sci.* **6**, 273 (2019).
74. Bennett, N. J. et al. Advancing social equity in and through marine conservation. *Front. Mar. Sci.* **8**, 711538 (2021).
75. Short, R. E. et al. Review of the evidence for oceans and human health relationships in Europe: a systematic map. *Environ. Int.* **146**, 106275 (2021).
76. Mukherjee, N. et al. The Delphi technique in ecology and biological conservation: applications and guidelines. *Methods Ecol. Evol.* **6**, 1097–1109 (2015).
77. Sutherland, W. J. et al. A 2021 horizon scan of emerging global biological conservation issues. *Trends Ecol. Evol.* **36**, 87–97 (2021).

Acknowledgements

This Marine and Coastal Horizon Scan was funded by Oceankind. S.N.R.B. is supported by EcoStar (DM048) and Cefas (My time). R.C. acknowledges FCT/MCTES for the financial support to CESAM (UIDP/50017/2020, UIDB/50017/2020, LA/P/0094/2020) through national funds. O.D. is supported by CSIC Uruguay and Inter-American Institute for Global Change Research. J.E.H.-R. is supported by the Whitten Lectureship in Marine Biology. S.A.K. is supported by a Natural Environment Research Council grant (NE/S00050X/1). P.I.M. is supported by an Australian Research Council Discovery Grant (DP200100575). D.M.P. is supported by the Marine Alliance for Science and Technology for Scotland (MASTS). A.R.P. is supported by the Inter-American Institute for Global Change Research. W.J.S. is funded by Arcadia. A.T. is supported by Oceankind. M.Y. is supported by the Deep Ocean Stewardship Initiative and bioDISCOVERY. We are grateful to everyone who submitted ideas to the exercise and the following who are not authors but who suggested a topic that made the final list: R. Brown (colocation of marine activities), N. Graham and C. Hicks (altered nutritional content of fish), A. Thornton (soft robotics), A. Vincent (fish swim bladders) and T. Webb (mesopelagic fisheries).

Author contributions

J.E.H.-R. and A.T. contributed equally to the manuscript. J.E.H.-R., A.T. and W.J.S. devised, organized and led the Marine and Coastal Horizon Scan. D.J.A., S.N.R.B., I.M.C., M.P.D., B.J.G., S.A.K., E.M. and L.S.P. formed the core team and are listed alphabetically in the author list. All other authors, R.C., O.D., S.D., E.L.J., H.K., P.I.M., A.M., A.W.N.M., D.O.O., D.M.P., A.R.P., A.J.R., I.R.S., P.V.R.S., B.D.S., P.M.T., G.J.W., T.A.W. and M.Y. are listed alphabetically. All authors contributed to and participated in the process and all were involved in writing and editing the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41559-022-01812-0>.

Correspondence and requests for materials should be addressed to James E. Herbert-Read or Ann Thornton.

Peer review information *Nature Ecology & Evolution* thanks Camille Mellin, Prue Addison and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2022

¹Department of Zoology, University of Cambridge, Cambridge, UK. ²Conservation Science Group, Department of Zoology, Cambridge University, Cambridge, UK. ³SpeSeas, D'Abadie, Trinidad and Tobago. ⁴Marine Science Institute, University of California, Santa Barbara, CA, USA. ⁵The Centre for Environment, Fisheries and Aquaculture Science (Cefas), Lowestoft, UK. ⁶Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada. ⁷Centre for Ecology, Evolution and Environmental Changes (cE3c), Department of Animal Biology, Faculdade de Ciências da Universidade de Lisboa, Lisbon, Portugal. ⁸BirdLife International, The David Attenborough Building, Cambridge, UK. ⁹Centre for Ecology and Conservation, University of Exeter, Penryn, UK. ¹⁰Lancaster Environment Centre, Lancaster University, Lancaster, UK. ¹¹School of Earth and Environmental Sciences, Cardiff University, Cardiff, UK. ¹²British Antarctic Survey, Natural Environment Research Council, Cambridge, UK. ¹³ECOMARE, CESAM—Centre for Environmental and Marine Studies, Department of Biology, University of Aveiro, Santiago University Campus, Aveiro, Portugal. ¹⁴Laboratory of Marine Sciences (UNDECIMAR), Faculty of Sciences, University of the Republic, Montevideo, Uruguay. ¹⁵Royal Belgian Institute of Natural Sciences, Operational Directorate Natural Environment, Marine Ecology and Management, Brussels, Belgium. ¹⁶School of Biological, Earth, and Environmental Sciences, University of New South Wales, Sydney, New South Wales, Australia. ¹⁷Finnish Environment Institute, Helsinki, Finland. ¹⁸Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Burwood Campus, Burwood, Victoria, Australia. ¹⁹Department of Oceanography, Dalhousie University, Halifax, Nova Scotia, Canada. ²⁰Department of Biology, University of Nairobi, Nairobi, Kenya. ²¹Coastal Oceans Research and Development in the Indian Ocean, Mombasa, Kenya. ²²School of Biological Sciences, University of Queensland, St Lucia, Brisbane, Queensland, Australia. ²³Scottish Oceans Institute, School of Biology, University of St Andrews, St Andrews, UK. ²⁴Servicio de Hidrografía Naval, Buenos Aires, Argentina. ²⁵Instituto Franco-Argentino sobre Estudios de Clima y sus Impactos, CONICET/CNRS, Universidad de Buenos Aires, Buenos Aires, Argentina. ²⁶School of Mathematics and Physics, The University of Queensland, St Lucia, Brisbane, Queensland, Australia. ²⁷Commonwealth Scientific and Industrial Research Organisation (CSIRO) Oceans and Atmosphere, Queensland Biosciences Precinct, St Lucia, Brisbane, Queensland, Australia. ²⁸Instituto Antártico Argentino, Buenos Aires, Argentina. ²⁹Centro Austral de Investigaciones Científicas (CADIC-CONICET), Ushuaia, Argentina. ³⁰Universidad Nacional de Tierra del Fuego, Antártida e Islas del Atlántico Sur, Ushuaia, Argentina. ³¹Department of Ocean Sciences and Biology Department, Memorial University, St John's, Newfoundland and Labrador, Canada. ³²Department of Environment and Geography, University of York, York, UK. ³³Lighthouse Field Station, School of Biological Sciences, University of Aberdeen, Cromarty, UK. ³⁴Institute of Marine Sciences, School of Biological Sciences, University of Portsmouth, Portsmouth, UK. ³⁵School of Biological Sciences, Area of Ecology and Biodiversity, Swire Institute of Marine Science, Institute for Climate and Carbon Neutrality, Musketeers Foundation Institute of Data Science, and State Key Laboratory of Marine Pollution, The University of Hong Kong, Kadoorie Biological Sciences Building, Hong Kong, China. ³⁶Biosecurity Research Initiative at St Catharine's (BioRISC), St Catharine's College, University of Cambridge, Cambridge, UK. ³⁷These authors contributed equally: James E. Herbert-Read, Ann Thornton. [✉]e-mail: jh2223@cam.ac.uk; at915@cam.ac.uk

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection

Data analysis

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

A data availability statement is included. The datasets generated during and/or analysed during the current study are available from Figshare <https://doi.org/10.6084/m9.figshare.19703485.v1>.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	Horizon scan exercise involving trans-disciplinary team of 30 global experts in marine and coastal ecosystems.
Research sample	30 participants scored issues throughout the exercise.
Sampling strategy	Score-sheets (Excel) were sent to all participants to complete and return.
Data collection	Score-sheets from participants were submitted, confidentially, to AT.
Timing and spatial scale	March - June 2021: Issues submitted. June 2021: Round 1 scoring carried out. July 2021: 32 top-ranked 32 issues sent out to participants. September 2021: Round 2 - online workshop and discussion. Scores submitted to AT and top-ranked issues revealed.
Data exclusions	Three participants did not score all issues in round 2, and therefore their scores were discounted.
Reproducibility	We used a modified Delphi technique ensuring a repeatable, transparent scoring system.
Randomization	In round 1, participants were assigned one of three differently ordered score-sheets to complete. This reduced the potential for 'voter-fatigue' when issues at the end of a long list don't receive the same consideration as those at the beginning.
Blinding	All scores were submitted to AT confidentially. Only AT was aware of individual scores.
Did the study involve field work?	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involvement
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Human research participants
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern

Methods

n/a	Involvement
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging