

Short Communication

On the identity of "*Dendrilla membranosa*" (Porifera, Dendroceratida) *sensu* Burton, and the specimens collected in Argentina, SW Atlantic Ocean

Laura Schejter¹ , Pilar Ríos² , Javier Cristobo²  & Rob Van Soest³ 

¹Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP)
Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)
Mar del Plata, Argentina

²Instituto Español de Oceanografía, Centro Oceanográfico de Gijón (IEO-CSIC), Gijón, España

³Naturalis Biodiversity Center (NBC), Leiden, The Netherlands

Corresponding author: Laura Schejter (schejter@inidep.edu.ar)

ABSTRACT. There has been confusion regarding the specimens identified as "*Dendrilla membranosa* (Pallas, 1766)" in Antarctic and subantarctic waters, considering that the original description corresponded to specimens from the Indian Ocean. In this study, we clarified the identification of the specimens collected in Argentinian waters, SW Atlantic Ocean, that should be identified as *Dendrilla antarctica* Topsent, 1905, updating its distributional range and considering the new records reported in the present study.

Keywords: *Dendrilla antarctica*; sponges; distribution; Burdwood bank; Tierra del Fuego

The sponge *Dendrilla membranosa* (Pallas, 1766) was originally described as *Spongia membranosa* based on a specimen collected in the Indian Ocean with no precise definition of the type locality. The specimen was likely collected along the way from Amsterdam to Indonesia, with stops at South Africa, Seychelles, India, and possibly China (Van Soest, *comm. pers.*). There is no physical type of this species, only an illustration by Seba (1759: Plate 95, Fig. 3) indicated by Pallas as the type (Pallas 1766: 398, Fig. 1a) from specimens of his collection, which has been dispersed over various museums and have not been reliably retrieved. Although Seba's illustration and Pallas' description are sufficient for typification under ICZN Art. 12.2.1 and 72.5.6, the properties of *S. membranosa* remain unclear (cf. Wiedenmayer 1989, Bergquist 1996). Further, the identity of it as a species of *Dendrilla* Von Lendenfeld, 1883 is not based on verifiable physical type material, and the taxonomic view of the species in this genus have been limited by inconsistency (Bergquist & Cook 2002).

Centuries later, Burton (1929) reported *D. membranosa* specimens from Antarctica and proposed the synonymy of *D. membranosa* with *D. antarctica* Topsent, 1905. Afterward, he registered this species again from Antarctica, Burdwood Bank, South Georgias, and the Malvinas (Falkland) Islands (Burton 1932, 1934a). However, in 1934, Burton also published a report on sponges from the Great Barrier Reef Expedition, performed in 1928-1929 (Burton 1934a), and also registered specimens under the name of "*Dendrilla membranosa*" collected in the North Australian waters. Remarkably, Burton did not describe any specimens he assigned to *D. membranosa*, merely listing the various names and specimens he considered synonymous. For the Antarctic material, he relied on Topsent's description of *D. antarctica*; for the tropical Indo-West Pacific material, he relied on Ridley's *Aplysina membranosa* and *A. pallasi*.

Bergquist (1980) created the genus *Dictyodendrilla*, and proposed to include the species *D. membranosa*, as




Figure 1. a) *Dendrilla membranosa* (from Seba 1759, Plate 95 Fig. 3), b) *Dendrilla antarctica* (from Topsent, 1908), c-e) *D. antarctica* from Burdwood Bank and Tierra del Fuego (present study).

she distilled from Burton's (1934a) descriptive notes that his specimens from the Great Barrier Reef were reticulated. Later, Bergquist (1996) argued that there is no way we can establish whether Pallas' specimen was reticulated or not, so she proposed that Pallas' material be reassigned as *D. membranosa*, and the tropical

specimens of Burton be kept in *Dictyodendrilla*. This transfer needed Burton's tropical material to have a new name, for which she proposed *Dendrilla pallasii* (Ridley, 1884) (originally *Aplysina*). In the Bergquist (1996) publication, she wrote "*Dendrilla membranosa sensu* Burton, 1932", but this was probably an

unfortunate misspelling because it makes sense that the discussed specimens were those collected during the Great Barrier Reef Expedition (Burton 1934a), not those referenced from Antarctic waters (Burton 1932, 1934b). Although Bergquist (1996) proposed that the tropical *D. membranosa sensu* Burton should be renamed *Dictyodendrilla pallasii* (Ridley, 1884), the evidence she presented for this proposal lacks substance, as Ridley's material was not redescribed. To solve the identity and name for the tropical "*Dendrilla membranosa*" is beyond this present short communication, although it should be addressed soon using integrative systematics. It is still considered a valid species; however, it is listed as inaccurate for the Antarctic region by World Register of Marine Species (WoRMS) (De Voogd et al. 2022).

Returning to colder waters, *D. antarctica* was extensively described by Topsent (1905) for Wandel, Wiencke, and Anvers Islands (Antarctica), from shallow waters of 20-40 m, and illustrated later in Topsent (1908) (Fig. 1b). This is a common and often abundant species distributed in coastal Antarctic waters, reported and studied from a taxonomic, biological, ecological, and chemical perspective by Hentschel (1914), Vacelet (1958), Desqueyroux-Faúndez (1989), Koutsouveli et al. (2018), Sacristán-Soriano et al. (2020) and Bory et al. (2020), among others. Desqueyroux-Faúndez (1989) and Goodwin et al. (2019) also noted the confusion generated by Burton (1929, 1932, 1934b) with *D. membranosa* when they were studying Antarctic specimens of *Dendrilla*, but this information was overlooked in their extensive descriptive works.

Between April 22 and May 12, 2017, an Argentinian Expedition onboard the research vessel "Puerto Deseado" was undertaken. This expedition aimed to acquire knowledge on oceanography, geology, biology, ecology, and the conservation value of the Marine Protected Area Namuncurá/ Burdwood Bank (created in 2013, Law 26875) and nearby regions (Falabella et al. 2017). Samples of the benthic community were collected using bottom trawls to characterize the bottom landscape and to create a species inventory of this understudied region where fragile, vulnerable, and also many undescribed species inhabited (e.g. Schejter et al. 2016, 2017, 2020, López-Gappa et al. 2018, Pérez & Cordeiro 2020). During sponge sampling procedures, specimens that fit the description of the genus *Dendrilla* were recorded (Figs. 1c-e). They were collected at stations 24 (54°19.9'S, 59°53.7'W; 97 m depth; Marine Protected Area Namuncurá-Burdwood Bank; deposited at MACN-In 43803) and 40 (58°S,

67°01'W; 49 m; Tierra del Fuego coastal waters; deposited at MACN-In-43804) of the "Puerto Deseado" 2017 Expedition to Burdwood Bank. It was the first time (after Burton's studies) that specimens of *Dendrilla* were collected in the SW Atlantic Ocean, outside Antarctic waters. The collected specimens are flabellate to irregularly branched. Some were beige, others bright yellow in life, while the preserved specimens were beige or became violet to nearly black. They have a smooth conulose surface. The skeleton was composed of spongin fibers visible at the terminal parts, protruding in the conules (Figs. 1c-e). Additionally, new bioactive compounds were recently discovered from specimens collected at Tierra del Fuego during the expedition mentioned above (Prieto et al. 2022).

According to the literature, only "*Dendrilla membranosa*" was previously reported among the Dendroceratida and Darwinellidae in the studied area (Burton 1934b, López-Gappa & Landoni 2005). It is widely accepted that many Antarctic species could reach Argentinian subantarctic waters in their distributions (i.e. Bertolino et al. 2007, Figuerola et al. 2014, Schejter & Bremec 2015). Moreover, according to Leiva et al. (2019), *D. antarctica*, was found to have high levels of gene flow and the potential for long-distance larval dispersal. Our specimens very much fit the description of *D. antarctica* provided by Topsent (1905, 1908), but also that of "*Dendrilla membranosa*" provided by Burton (1929, 1932, 1934a). Given the previous confusion mentioned above and based on our experience and the arguments discussed, we propose that the specimens identified as *D. membranosa sensu* Burton (1929, 1932, 1934b) in the SW Atlantic Ocean, outside Antarctica, should be considered to belong to *D. antarctica*. In accordance with the observations made by Desqueyroux-Faúndez (1989) and Goodwin et al. (2019) for Antarctic specimens. In this way, the distribution range of *D. antarctica* should be updated. It would comprise not only the Antarctic domain (Antarctica, Kerguelen Islands, South Georgias Islands) but also Tierra del Fuego (Argentina) (data from the present study), Burdwood Bank (data from the present study), and Malvinas (Falkland) Islands. Wrong identification of a species may produce a cascade-like process, with consequences at multiple levels (ecology, biogeography, diversity, conservation) (Bortolus 2008, Thomson et al. 2018); hence, molecular and integrative systematics should be encouraged. Of the 10 valid species under the genus *Dendrilla* (De Voogd et al. 2022), only *D. antarctica* is distributed in the cold waters of Antarctica and the southern South American waters. The other eight species (besides *D. membra-*

nosa) are distributed in Australian waters (*D. cactos* (Selenka, 1867), *D. cruor* (Carter, 1886), *D. rosea* Von Lendenfeld, 1883), New Zealand, and New Caledonia (*D. rosea*), West Pacific Ocean (*D. lendenfeldi* Hentschel, 1912, *D. mertoni* Hentschel, 1912), Maldives (*D. cactos*), Mediterranean Sea (*D. acantha* Vacelet, 1958, *D. cirsioides* Topsent, 1893), North Atlantic Ocean (*D. acantha*) and the Caribbean Sea (*D. camera* (de Laubenfels, 1936)) (De Voogd et al. 2022).

ACKNOWLEDGMENTS

Laura Schejter wishes to thank members of the scientific staff that helped during sampling procedures onboard the RV "Puerto Deseado," particularly Dr. G. Lovrich, General Scientific Coordinator of the activities developed in the MPA Namuncurá/Burdwood bank. We also thank the suggestions and comments of the anonymous reviewers that contributed to improving this note's final version. The expedition was supported by the Marine Protected Area Namuncurá/Burdwood bank Law 26875/2013, and the study was partially supported by PICT 2019-4233 to LS. MPA Namuncurá Contribution N°65 and INIDEP Contribution N°2269. Voucher specimens were deposited in the Museo Argentino de Ciencias Naturales "Bernardino Rivadavia" (MACN-In), Argentina.

REFERENCES

Bergquist, P.R. 1980. A revision of the supraspecific classification of the orders Dictyoceratida, Dendroceratida and Verongida (class Demospongiae). *New Zealand Journal of Zoology*, 7: 443-503.

Bergquist, P.R. 1996. The marine fauna of New Zealand: Porifera: Demospongiae. Part 5. Dendroceratida and Halisarca. *New Zealand Oceanographic Institute Memoir*, 107: 1-53.

Bergquist, P.R. & Cook, S.D.C. 2002. Family Darwinellidae Merejkowsky, 1879. In: Hooper, J.N.A. & Van Soest, R.W.M. (Eds.). *Systema Porifera. A guide to the classification of sponges*. Kluwer Academic/Plenum, New York, 2: 1068-1071.

Bertolino, M., Schejter, L., Calcinai, B., Cerrano, C. & Bremec, C. 2007. Sponges from a submarine canyon of the Argentine Sea. In: Custódio, M.R., Lôbo-Hajdu, G., Hajdu, E. & Muricy, G. (Eds.). *Porifera research: biodiversity, innovation and sustainability*. Museu Nacional, Rio de Janeiro, pp. 189-201.

Bortolus, A. 2008. Error cascades in the biological sciences: the unwanted consequences of using bad taxonomy in ecology. *Ambio*, 37: 114-118.

Bory, A., Shilling, A.J., Allen, J., Azhari, A., Roth, A., Shaw, L.N., et al. 2020. Bioactivity of spongian diterpenoid scaffolds from the Antarctic sponge *Dendrilla antarctica*. *Marine Drugs*, 18: 327. doi: 10.3390/md18060327

Burton, M. 1929. Porifera. Part II. Antarctic sponges. *British Antarctic ('Terra Nova') Expedition, 1910. Natural History Report, London, Zoology*, 6: 393-458.

Burton, M. 1932. Sponges. *Discovery Reports*, 6: 237-392.

Burton, M. 1934a. Sponges. *Scientific reports of the Great Barrier Reef Expedition 1928-29*, 4: 513-621.

Burton, M. 1934b. Sponges. In: *Further zoological results of the Swedish Antarctic Expedition 1901-03 under the direction of Dr. Otto Nordenskjöld*. Norstedt & Söner, Stockholm, pp. 1-58.

Desqueyroux-Faúndez, R. 1989. Demospongiae (Porifera) del litoral chileno antártico. *Serie Científica INACH*, 39: 97-158.

De Voogd, N.J., Alvarez, B., Boury-Esnault, N., Carballo, J.L., Cárdenas, P., Díaz, M.-C., et al. 2022. World Porifera database. [<https://www.marinespecies.org/porifera>]. Reviewed: May 23, 2022.

Falabella, V., Caille, G., Campagna, C., Krapovickas, S., Lovrich, G., Moreno, D., et al. 2017. Área Marina Protegida Namuncurá - Banco Burdwood. In: Falabella, V. (Ed.). *Contribuciones para la línea de base y el plan de manejo*. Jefatura de Gabinete de Ministros, Buenos Aires.

Figuerola, B., Gordon, D.P., Polonio, V., Cristobo, J. & Avila, C. 2014. Cheilostome bryozoan diversity from the southwest Atlantic region: is Antarctica really isolated? *Journal of Sea Research*, 85: 1-17.

Goodwin, C., Berman, J. & Hendry, K.R. 2019. Demosponges from the sublittoral and shallow-circalittoral (<24 m depth) Antarctic Peninsula with a description of four new species and notes on *in situ* identification characteristics. *Zootaxa*, 4658: 461-508. doi: 10.11646/zootaxa.4658.3.3

Hentschel, E. 1914. Monaxone Kieselschwämme und Hornschwämme der Deutschen Südpolar-Expedition 1901-1903. *Deutsche Südpolar-Expedition*, 15: 35-141.

Koutsouveli, V., Taboada, S., Moles, J., Cristobo, J., Ríos, P., Bertran, A., et al. 2018. Insights into the reproduction of some Antarctic dendroceratid, poecilosclerid, and haplosclerid demosponges. *Plos One*, 13: e0192267. doi: 10.1371/journal.pone.0192267

Leiva, C., Taboada, S., Kenny, N.J., Combosch, D., Giribet, G., Jombart, T. & Riesgo, A. 2019. Population

- substructure and signals of divergent adaptive selection despite admixture in the sponge *Dendrilla antarctica* from shallow waters surrounding the Antarctic Peninsula. *Molecular Ecology*, 28: 3151-3170. doi: 10.1111/mec.15135
- López-Gappa, J. & Landoni, N.A. 2005. Biodiversity of Porifera in the Southwest Atlantic between 35°S and 56°S. *Revista del Museo Argentino de Ciencias Naturales*, 7: 191-219. doi: 10.22179/REVMACN.7.328
- López-Gappa, J., Liuzzi, M.G. & Zelaya, D.G. 2018. A new genus and species of cheilostome bryozoan associated with hermit crabs in the subantarctic Southwest Atlantic. *Polar Biology*, 41: 733-41. doi: 10.1007/s00300-017-2234-9
- Pallas, P.S. 1766. *Elenchus zoophytorum sistens generum adumbrationes generaliores et specierum cognitarum succinctas descriptiones, cum selectis auctorum synonymis*. Franciscum Varrentrapp, Hagae.
- Pérez, C.D. & Cordeiro, R.T. 2020. *Ideogorgia laurae*, an uncommon new octocoral species (Alcyonacea: Keroididae) from a newly established Marine Protected Area at Burdwood Bank, Argentina. *Polar Biology*, 43: 63-69. doi: 10.1007/s00300-019-02604-2
- Prieto, I., Pérez, M., García, M., Blunstein, G., Schejter, L. & Palermo, J. 2022. Antifouling diterpenoids from the sponge *Dendrilla antarctica*. *Chemistry and Biodiversity*, 19: e202100618. doi: 10.1002/cbdv.202100618
- Sacristán-Soriano, O., Pérez-Criado, N. & Avila, C. 2020. Host species determines symbiotic community composition in Antarctic sponges (Porifera: Demospongiae). *Frontiers in Marine Science*, 7: 1-11. doi: 10.3389/fmars.2020.00474
- Schejter, L. & Bremec, C. 2015. First record and range extension of the Antarctic coral *Flabellum (Flabellum) impensum* in Argentinean coastal waters. *Marine Biodiversity Records*, 8: e104. doi: 10.1017/S1755267215000858
- Schejter, L., Bertolino, M. & Calcinai, B. 2017. Description of *Antho (Plocamia) bremecae* sp. nov. and checklist of Microcionidae (Demospongiae: Poecilosclerida) from Burdwood Bank and neighboring areas, SW Atlantic Ocean. *Zootaxa*, 4312: 580-594. doi: 10.11646/zootaxa.4312.3.11
- Schejter, L., Genzano, G., Gaitán, E., Pérez, C.D. & Bremec, C.S. 2020. Benthic communities in the Southwest Atlantic Ocean: conservation value of animal forests at the Burdwood Bank slope. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 30: 426-439. doi: 10.1002/aqc.3265
- Schejter, L., Rimondino, C., Chiesa, I., Díaz de Astarloa, J.M., Doti, B.L., Elías, R., et al. 2016. Namuncurá Marine Protected Area: an oceanic hot spot of benthic biodiversity at Burdwood Bank, Argentina. *Polar Biology*, 39: 2373-2386. doi: 10.1007/s00300-016-1913-2
- Seba, A. 1759. *Locupletissimi rerum naturalium thesauri accurata descriptio, et iconibus artificiosissimis expressio, per universam physices historiam: opus, cui, in hoc rerum genere, nullum par exstitit / Description exacte des principales curiosités naturelles du magnifique cabinet d'Albert Seba*. 4 vols. Apud Janssonio-Waesbergios & J. Wetstenium & Gul. Smith, 1734-1765, Amstelaedami.
- Thomson, S.A., Pyle, R.L., Ahyong, S.T., Alonso-Zarazaga, M., Ammirati, J., Araya, J.F., et al. 2018. Taxonomy based on science is necessary for global conservation. *Plos Biology*, 16: e2005075. doi: 10.1371/journal.pbio.2005075
- Topsent, E. 1905. Notes sur les éponges recueillies par le Français dans l'Antarctique. Description d'une *Dendrilla* nouvelle. *Bulletin du Muséum National d'Histoire Naturelle*, 11: 502-505.
- Topsent, E. 1908. Spongiaires. Expédition antarctique française (1903-1905) commandée par le Dr. Jean Charcot (Paris), 4: 1-37.
- Vacelet, J. 1958. *Dendrilla acantha* nov. sp. nouvelle éponge cornée Méditerranéenne. Remarques sur les genres *Dendrilla* Lendenfeld et *Megalopastas* Dendy. *Recueil des Travaux de la Station Marine d'Endoume*, 14: 143-147.
- Von Lendenfeld, R. 1883. Über Coelenteraten der Südsee. II. Mittheilung. Neue Aplysinidae. *Zeitschrift für Wissenschaftliche Zoologie*, 38: 234-313.
- Wiedenmayer, F. 1989. Demospongiae (Porifera) from northern Bas Strait, southern Australia. *Memoirs of the Museum of Victoria*, 50: 1-242. doi: 10.24199/j.mmv.1989.50.01