

1 REVIEW ARTICLE

2
3 **Beyond transcription: compelling open questions in plant RNA**
4 **biology**

5
6 **Pablo A. Manavella^{1,*}, Micaela A. Godoy Herz², Alberto R. Kornblihtt², Reed**
7 **Sorenson³, Leslie E. Sieburth³, Kentaro Nakaminami⁴, Motoaki Seki^{4,5,6}, Yiliang**
8 **Ding⁷, Qianwen Sun^{8,9}, Hunseung Kang¹⁰, Federico D. Ariel¹, Martin Crespi^{11,12},**
9 **Axel Giudicatti¹, Qiang Cai¹³, Hailing Jin¹⁴, Xiaoqi Feng⁷, Yijun Qi^{15,16}, Craig S.**
10 **Pikaard¹⁷**

11
12 #All authors contributed equally to this article and are listed in the order in which their questions appear.

13 *Author for correspondence: pablomanavella@ial.santafe-conicet.gov.ar

14
15 **Affiliations**

16 1. Instituto de Agrobiotecnología del Litoral (CONICET-UNL), Cátedra de Biología Celular y Molecular,
17 Facultad de Bioquímica y Ciencias Biológicas, Universidad Nacional del Litoral, 3000 Santa Fe,
18 Argentina.

19 2. Universidad de Buenos Aires (UBA), Facultad de Ciencias Exactas y Naturales, Departamento de
20 Fisiología, Biología Molecular y Celular and CONICET-UBA, Instituto de Fisiología, Biología Molecular y
21 Neurociencias (IFIBYNE), (C1428EHA), Buenos Aires, Argentina. (ark@fbmc.fcen.uba.ar)

22 3. School of Biological Sciences, University of Utah 84112. (sieburth@biology.utah.edu)

23 4. Center for Sustainable Resource Science, RIKEN, Kanagawa, 230-0045, Japan
24 (motoaki.seki@riken.jp)

25 5. Cluster for Pioneering Research, RIKEN, Saitama, 351-0198, Japan

26 6. Kihara Institute for Biological Research, Yokohama City University, Kanagawa, 244-0813, Japan.

27 7. Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park, Norwich,
28 NR4 7UH, United Kingdom. (Yiliang.Ding@jic.ac.uk) (Xiaoqi.Feng@jic.ac.uk)

29 8. Center for Plant Biology, School of Life Sciences, Tsinghua University, Beijing 100084, China.
30 (sunqianwen@mail.tsinghua.edu.cn)

31 9. Tsinghua-Peking Center for Life Sciences, Beijing 100084, China.

32 10. Department of Applied Biology, College of Agriculture and Life Sciences, Chonnam National
33 University, Gwangju 61186, Korea. (hskang@jnu.ac.kr)

34 11. Institute of Plant Sciences Paris Saclay IPS2, CNRS, INRA, Université Evry, Université Paris-Saclay,
35 Bâtiment 630, 91405 Orsay, France. (martin.crespi@universite-paris-saclay.fr)

36 12. Institute of Plant Sciences Paris-Saclay IPS2, Université de Paris, Bâtiment 630, 91405 Orsay,
37 France.

38 13. State Key Laboratory of Hybrid Rice, College of Life Science, Wuhan University, Wuhan 430072,
39 China.

40 14. Department of Microbiology and Plant Pathology and Center for Plant Cell Biology, Institute for
41 Integrative Genome Biology, University of California, Riverside, CA 92507, USA. (hailingj@ucr.edu)

42 15. Center for Plant Biology, School of Life Sciences, Tsinghua University, Beijing 100084, China.
43 (qiyijun@tsinghua.edu.cn)

44 16. Tsinghua-Peking Center for Life Sciences, Beijing 100084, China

45 17. Howard Hughes Medical Institute, Department of Biology and Department of Molecular and Cellular
46 Biochemistry, Indiana University, Bloomington, IN 47405. (cpikaard@indiana.edu)

1
2 **Short title: Open questions in plant RNA biology**
3
4

5 **Abstract**

6 The study of RNAs has become one of the most influential research fields in
7 contemporary biology and biomedicine. In the last few years, new sequencing
8 technologies have produced an explosion of new and exciting discoveries in the field
9 but have also given rise to many open questions. Defining these questions, together
10 with old, long-standing gaps in our knowledge, is the spirit of this article. The breadth of
11 topics within RNA biology research is vast, and every aspect of the biology of these
12 molecules contains countless exciting open questions. Here, we asked twelve groups to
13 discuss their most compelling question among some plant RNA biology topics. The
14 following vignettes cover RNA alternative splicing; RNA dynamics; RNA translation;
15 RNA structures; R-loops; epitranscriptomics; long noncoding RNAs; small RNA
16 production and their functions in crops; small RNAs during gametogenesis and in cross-
17 kingdom RNA interference; and RNA-directed DNA methylation. In each section, we will
18 present the current state-of-the-art in plant RNA biology research before asking the
19 questions that will surely motivate future discoveries in the field. We hope this article will
20 spark a debate about the future perspective on RNA biology and provoke novel
21 reflections in the reader.

22
23 **Introduction**

24 (Written by Pablo A. Manavella)

25 In all living organisms, DNA is the molecule storing all genetic information, while RNA
26 carries this data to the ribosomes to be translated into proteins. While DNA is
27 omnipresent in our imagination, making star appearances in movies, TV shows, and
28 books, the contribution of RNA to life is less recognized by society. However, as a
29 consequence of the recent COVID pandemic, the word "RNA" has reached most people
30 on the planet as they learned about the RNA-based genome of the virus and the
31 therapeutic use of RNA vaccines. Thus, the concept of 'information flow', that is the
32 decoding of DNA to protein using an RNA intermediate, has suddenly become the
33 center of attention and conversations. What remains largely unknown to the general
34 audience is that the advent of sequencing technologies has made it clear that RNA is
35 not only a coding molecule but also has various other functions, mostly in the form of
36 cellular non-coding RNA transcripts.

37 The study of RNAs has emerged as a particularly important research field in
38 contemporary biology, especially in plant biology, where these molecules execute many
39 actions during development and response to the environment. Advances in sequencing
40 technologies have allowed the global analysis of RNA modifications, the resolution of
41 RNA secondary structures, the mapping of epigenetic modifications, the identification of

1 RNA-edited sequences, and the discovery of novel classes of RNAs resulting in a
2 revolution in molecular biology that is just starting.

3 In this article, we gathered twelve experts in different aspects of plant RNA biology to
4 discuss some of the most compelling open questions in the field. Each section
5 discusses long-standing open questions of the field as well as questions that have only
6 begun to emerge from breakthrough discoveries. We hope this article helps stimulate
7 the community and sparks new ideas and research projects that will expand the
8 frontiers of RNA biology knowledge in plants.

9 10 **How does light control RNA alternative splicing in plants?**

11 (Written by Micaela Godoy Herz and Alberto Kornblihtt)

12 Plants rely on light as their main source of energy, but light also regulates many
13 developmental and physiological responses during the plant life cycle (Arsovski et al.,
14 2012). Furthermore, light signals induce a massive reprogramming of gene expression
15 in plants (Tognacca et al., 2020). Alternative splicing produces multiple mRNA variants
16 from a single locus. Splicing and alternative splicing are coupled with transcription, and
17 factors that regulate transcription also affect alternative splicing (Kornblihtt et al., 2013).

18 Our laboratory showed how light regulates plant alternative splicing through the
19 chloroplast (Petrillo et al., 2014). Light and dark conditions affect alternative splicing of a
20 subset of *Arabidopsis* (*Arabidopsis thaliana*) genes preferentially encoding proteins
21 involved in RNA processing. This effect requires functional chloroplasts: treatment of
22 *Arabidopsis* seedlings with drugs that impair the chloroplast photosynthetic transport
23 chain inhibit the effect of light on alternative splicing. Moreover, the effect of light is also
24 observed in roots when communication with leaves –the photosynthetic tissue– is not
25 interrupted (Petrillo et al., 2014). Light, sensed by the chloroplast, indeed triggers a
26 retrograde signal that regulates alternative splicing not only in leaves, but also in roots.

27 How does light cause splicing responses in roots? In a recent work, Riegler and
28 collaborators investigated this shoot-to-root signaling: they showed that alternative
29 splicing responses in roots are not directly caused by light, but are instead most likely
30 triggered by sugars. The kinase TARGET OF RAPAMYCIN (TOR) plays a key role in
31 this signaling pathway. Sugars activate the TOR pathway and act as mobile signals to
32 coordinate alternative splicing responses throughout the plant (Riegler et al., 2021).

33 These results afforded us a better understanding of how mobile signals regulate
34 alternative splicing throughout the entire plant in response to light. One remaining
35 outstanding open question is what happens in the nucleus: that is, what are the
36 mechanisms involved in this regulation of alternative splicing in plants?

37 We performed different experiments to address the role of transcription elongation and
38 determined that the light control of alternative splicing responds to a kinetic coupling
39 mechanism (Godoy Herz et al., 2019). Briefly, the kinetic coupling model explains how
40 changes in RNA Polymerase II (Pol II) elongation rate influence alternative splicing.

1 Each splice site consists of a consensus sequence that is recognized by spliceosomal
2 components, although 'strong' splice sites (those that are close to the consensus
3 sequence) are more efficiently recognized than 'weak' splice sites, which are
4 suboptimal. In the example illustrated in Figure 1, there is an alternative splicing event
5 with two 3' splice sites: a weak upstream 3' splice site, and a strong downstream splice
6 site. If Pol II elongation rate is fast, both sites are presented to the splicing machinery at
7 the same time, and the strong 3' splice site is recognized by the splicing machinery
8 more efficiently, resulting in exon skipping. However, if Pol II transcription rate is slow,
9 the splicing machinery will recognize the upstream, weaker, 3' splice site first, and
10 afterwards the strong 3' splice site, which leads to exon inclusion (Godoy Herz et al.,
11 2019). We showed by different experimental approaches that light promoted
12 transcription elongation in Arabidopsis, while Pol II elongation was slower in darkness.
13 Furthermore, the light control of alternative splicing and elongation was abolished in
14 plants lacking function for TRANSCRIPTION FACTOR II S (TFIIS) in a previous report
15 (Dolata et al., 2015): These TFIIS mutant plants did not respond to light signaling on a
16 group of alternative splicing events. This result demonstrated that coupling between
17 transcription and splicing is important for a whole organism to respond to environmental
18 cues (Figure 1).

19 Plant lines with higher Pol II transcription activity were recently generated by introducing
20 point mutations in NRPB2, the second largest subunit of Pol II. As a result, an
21 accelerated Pol II elongation rate increased the polymerase signal in gene bodies,
22 which appeared to modulate alternative splicing choices (Leng et al., 2020)..

23 Even though our knowledge of alternative splicing in plants has grown significantly in
24 the last decade, many important open questions remain. It has been shown that, in
25 response to light, sugars activate the TOR pathway, which in turn regulates alternative
26 splicing. But how does TOR regulate alternative splicing in the nucleus? In the
27 chloroplast, the exact nature of the chloroplast retrograde signal that regulates
28 alternative splicing remain unknown, although it may be triggered by the oxidation state
29 of the plastoquinone pool connecting both photosystems (Petrillo et al., 2014).

30 Moving forward, inside the nucleus, how light promotes Pol II elongation is unknown:
31 what makes Pol II elongation faster in the light, and slower in darkness? There are
32 many possible mechanisms that might explain how the chloroplast regulates
33 transcription elongation. Furthermore, the role of chromatin modifications on the
34 regulation of alternative splicing in plants remains an interesting field to investigate.
35 Previous studies in mammalian systems have shown that histone post-translational
36 modifications play a key role in the regulation of alternative splicing decisions. Treating
37 cell cultures with drugs that open chromatin structure promoted changes in alternative
38 splicing by facilitating Pol II elongation and exon skipping (Schor et al., 2009). By
39 contrast, cell differentiation results in an increase in intragenic silencing chromatin
40 marks that raised the rate of higher exon inclusion (Schor et al., 2013). In our work,

1 histone acetylation mimics the effect of light on alternative splicing, but light does not
2 affect the levels of this histone modification (Godoy Herz et al., 2019). Future
3 experiments will be needed to address the role of chromatin structure in splicing
4 regulation in plants.

5 Moreover, coupling between transcription elongation and alternative splicing may also
6 act in response to other environmental stimuli, like temperature. A recent work shows
7 that the TFIIIS elongation factor is required for thermal adaptation in *Arabidopsis*
8 (Szadeczky-Kardoss et al., 2022). Furthermore, analyses of plant native elongation
9 transcript sequencing (plaNET-seq) experiments in response to cold showed changes in
10 Pol II promoter-proximal stalling and at the 3' end of genes (Kindgren et al., 2020).

11 Finally, it would be interesting to study if these mechanisms of gene expression
12 regulation are also conserved in other plants and other photosynthetic organisms like
13 algae. Future work from different groups will be needed to address these questions.
14

15 **Acknowledgments**

16 We would like to thank M. Guillermina Kubaczka.
17

18 **Funding**

19 This work was supported by the ANPCyT (Agencia Nacional de Promoción Científica y
20 Tecnológica, Argentina, PICT 2019-862) and the Lounsbery Foundation.
21
22

23 **The invisible world of RNA dynamics**

24 (Written by Reed Sorenson and Leslie E. Sieburth)

25 Transcriptomics has transformed our understanding of molecular responses to signals.
26 The abundance of many mRNAs can be robustly upregulated or downregulated, and
27 many regulated genes bring about changes in development or physiology. Indeed,
28 measurements of RNA abundance are so ingrained in our thinking that changes in RNA
29 levels are frequently referred to as *gene expression* or *transcriptional responses*.
30 However, alongside regulatory events that lead to changes in mRNA levels, there lurks
31 the largely unseen layer of mRNA decay rate regulation. In addition to RNAs with
32 modified rates of decay and changes in their abundance, rates of decay can also be
33 modified for mRNAs whose abundances are held steady. This largely invisible dynamic
34 regulation is just beginning to be investigated, and so there are numerous unanswered
35 questions, including why decay rates are modified independently of changes in
36 abundance, how this modulation occurs, and whether this regulation has implications for
37 mRNAs that do show changes in their abundance.

38 RNA abundances are influenced by both synthesis (transcription) and decay, and the
39 rate of RNA turnover is called flux (Figure 2A). Wide variations in flux have been
40 observed in all deep RNA decay analyses, but whether flux rates affect RNA

1 abundances and/or regulation is still an open question. A special case of mRNA flux
2 regulation occurs when both the transcription and decay rates of an mRNA are
3 modulated, and yet the mRNA abundance does not change. This phenomenon is called
4 'RNA buffering' because transcription and decay rates are balanced to maintain steady
5 abundances (Figure 2B). RNA buffering has been documented in Arabidopsis, but we
6 are at the very beginning of understanding all aspects of this process, including both
7 how and why some mRNAs become buffered.

8 The system where RNA buffering is best understood is budding yeast (*Saccharomyces*
9 *cerevisiae*). A mysterious observation led to its discovery: mutants with defects in either
10 RNA decay or transcription were found to maintain normal mRNA abundances. It turned
11 out that the initial defect, in e.g. RNA decay, was accompanied by a compensatory
12 change (e.g. in transcription). That is, normal abundances of mRNAs in many
13 transcription and RNA decay mutants were maintained by precisely balanced changes
14 through RNA buffering (Haimovich et al., 2013; Sun et al., 2013a; Timmers and Tora,
15 2018; Hartenian and Glaunsinger, 2019). Because most mRNA decay occurs in the
16 cytoplasm, while transcription takes place in the nucleus, RNA buffering requires not
17 just precise regulation, but also communication between the nucleus and the cytoplasm.
18 Mechanisms underlying this regulation are still emerging and somewhat controversial,
19 but studies in yeast have revealed RNA decay proteins relocating to the nucleus and
20 displaying novel functions. For example, Sun and colleagues showed that the yeast
21 5'→3' EXORIBONUCLEASE 1 (XRN1) moves from the cytoplasm to the nucleus, where
22 it binds DNA and influences transcription of buffered RNAs (Sun et al., 2013a).
23 Similarly, upon nuclear RNA exosome dysfunction, RNA buffering was activated by
24 global attenuation of transcription via stabilization of the mRNA encoding HISTONE
25 SIRTUIN DEACETYLASE (HST3) (Bryll and Peterson, 2022). RNA buffering has also
26 been observed in *Drosophila* (*Drosophila melanogaster*), where it was used for gene
27 dosage compensation (Faucillion et al., 2022).

28 It was a similarly mysterious observation that led us to discover RNA buffering in
29 Arabidopsis (Sorenson et al., 2018). Cytoplasmic mRNA decay initiates through
30 deadenylation, and decay in the 3'→5' direction can be catalyzed by either the RNA
31 exosome or SUPPRESSOR OF VARICOSE (SOV)/ DIS3-LIKE EXONUCLEASE 2
32 (DIS3L2), while decay in the 5'→3' direction is initiated by decapping followed by
33 exoribonucleolytic digestion by XRN4 (Labno et al., 2016). Because the popular wild-
34 type accession Columbia-0 (Col-0) harbors a *sov* loss-of-function mutation (Zhang et
35 al., 2010) possible functions of this decay pathway were mysterious. To understand why
36 *sov* mutants did not show an abnormal phenotype, and also identify mRNA substrates
37 of decapping and SOV, we compared genome-wide RNA decay rates for wild type and
38 RNA decay mutants. In *varicose* (*vcs*) mutants (which lack mRNA decapping) the
39 expectation that mRNA decapping substrates would decay more slowly was largely
40 observed. The most common decay pattern (seen in >7,000 RNAs) was half-lives that

1 were longer in *vcs*, and longer still in *vcs sov* double mutants, indicating that these
2 RNAs were typically degraded by VCS, but upon loss of VCS, SOV provided back-up.
3 However, many mRNAs in *sov* mutants showed a surprising shift to shorter half-lives.
4 Moreover, mRNA decapping (via VCS) was required to sustain these shorter half-lives.
5 This unusual decay rate shift had no significant effect on RNA abundances, indicative of
6 RNA buffering and explaining the lack of phenotypic consequences in *sov* mutants in
7 Col-0. Data suggestive of RNA buffering was also identified in an Arabidopsis study of
8 cold response (Arae et al., 2017). We do not know whether plants use a mechanism
9 similar to that of yeast for RNA buffering; however there are no reports of XRN4 being
10 found in the nucleus (suggesting that RNA buffering in Arabidopsis might differ
11 mechanistically from yeast), and the shifting of SOV substrates to decapping via VCS
12 has not been described previously.

13 Conventional views of gene expression place all the action on those mRNAs whose
14 abundances are altered. However, RNA buffering turns this conventional view on its
15 head by demonstrating that many mRNAs with stable unchanging abundances also
16 undergo complex regulation. And this observation leads to an even bigger open
17 question: how are some mRNAs buffered so that their abundances do not vary, while
18 others appear to be able to freely increase or decrease in abundance? If only specific
19 mRNAs are buffered, perhaps they share a common sequence motif, e.g. for a
20 regulatory RNA-binding protein. Alternatively, buffering of RNA abundances may be a
21 default state, and some sort of licensing might be required to allow mRNAs to undergo
22 changes in abundance (Figure 2B). What distinguishes RNAs to be buffered from those
23 licensed to undergo alterations in their abundance? Attractive candidates can be found
24 in the expanding world of RNA modifications, from differing caps to covalent
25 modifications such as N⁶-methyladenosine (m⁶A) or structure (Kwok et al., 2013; Mauer
26 et al., 2017; Anderson et al., 2018; Reichel et al., 2019; Wang et al., 2019; Zhang et al.,
27 2019a). Addressing these many open questions will require much deeper understanding
28 of RNA kinetics.

29

30 **Funding**

31 This work was supported by the National Science Foundation (NSF MCB-1616779).

32

33

34 **Stabilization of mRNA and translational regulation by stress granules in response** 35 **to environmental conditions**

36 (Written by Kentaro Nakaminami and Motoaki Seki)

37 Current technologies used to analyze gene expression have enabled a high-level of
38 resolution on the expression of thousands of genes. Advancements in proteomic
39 technologies have also greatly improved the comprehensive analysis of proteins. Thus,
40 it has become possible to analyze plant physiology and metabolism in great detail using

1 various analytical methods. Collectively, these studies have empirically indicated that
2 gene and protein abundance patterns are not always identical based on the results of
3 multiomics analyses. Major factors contributing to the observed differences between
4 mRNA and protein patterns are post-transcriptional regulation of mRNA and
5 translational regulation of proteins. Both mechanisms fine-tune which mRNAs are
6 translated into proteins to regulate the physiology and metabolism of living organisms.
7 Various events occur between mRNA transcription and translation via the activity of
8 RNA-binding proteins (RBPs) that determine whether proteins are synthesized (Burjoski
9 and Reddy, 2021). These events begin with quality control of transcribed mRNAs,
10 followed by degradation of unnecessary or aberrant mRNAs, or protein translation.
11 Additionally, mRNAs can be temporarily stored via a stabilization system for subsequent
12 activation in response to environmental changes and other stimuli. Although mRNA
13 abundance is affected by the balance between transcription and degradation, the
14 amount of protein is not always proportional to mRNA abundance, and is affected by
15 post-transcriptional regulatory mechanisms such as the speed of translation and
16 translational inhibition. mRNA degradation is regulated by mRNA-protein (mRNP)
17 complexes called processing bodies (PBs); translation is carried out by ribosome
18 complexes (poly-ribosomes or polysomes), while mRNA stabilization or storage occurs
19 in stress granules (SGs) (Chantarachot and Bailey-Serres, 2018; Maruri-Lopez et al.,
20 2021). These granules are not organelles but rather membraneless RNA granules
21 formed via liquid-liquid phase separation (LLPS) (Emenecker et al., 2020). They have
22 been reported to be present in both animals and plants and are becoming a growing
23 research focus. This section of the present review discusses the nature of SGs, which
24 are responsible for the mechanisms of translational regulation, and how mRNAs are
25 stabilized and stored in these bodies.

26 Plants suppress mRNA translation when they are subjected to severe stress
27 (Merchante et al., 2017). This strategy reduces energy expenditure under stress
28 conditions, as only essential proteins are synthesized. Since translation requires
29 considerable energy, reducing energy requirements during stress contributes to
30 increased survival rates. Importantly, active but selective translation must operate
31 during stress response in plants since essential proteins are still translated. The
32 temporary storage of mRNA in SGs during stress conditions can be rapidly reversed,
33 with mRNAs being released in a translationally active form (polysomes) as plants
34 recover from stress conditions (Kosmacz et al., 2019). The mechanisms responsible for
35 determining target selectivity and translation timing by mRNP complexes, however,
36 have not been clearly elucidated.

37 SG complexes that form in the cytoplasm during stress are conserved in eukaryotes
38 (Maruri-Lopez et al., 2021). SG formation in plants is triggered by a variety of stresses,
39 including high temperature, hypoxia, high salinity, and darkness (Chantarachot and
40 Bailey-Serres, 2018; Hamada et al., 2018). An SG is composed of translationally

1 arrested mRNAs and proteins related to the initiation of translation, such as translation
2 initiation factors, small subunits of ribosomal RNA (rRNA), poly(A)-binding proteins, as
3 well as regulatory RBPs that inhibit translation. Recently, hundreds of proteins have
4 been characterized as SG components by combining immunoprecipitation (IP) and
5 genome-wide mRNA-binding interactome capture methods with proteomic analyses
6 (Chantarachot and Bailey-Serres, 2018; Kosmacz et al., 2019; Maronedze et al., 2019;
7 Gutierrez-Beltran et al., 2021; Maruri-Lopez et al., 2021). These results have suggested
8 that SGs are formed not only upon heat and hypoxia stresses, but also by drought
9 stress, resulting in translational repression. The components discovered in these
10 studies were not revealed based on their homology to SG components in animals and
11 yeasts as in previous studies, but rather were directly identified by the indicated
12 methodologies as components of SGs. Although many SG components have been
13 isolated with this approach, proteins within SGs also include translation-promoting
14 proteins such as translation initiation factors, and not all are related to translation
15 inhibition. It is necessary to consider the components and functions of SGs, including
16 spatiotemporal factors such as the dynamics of SG formation/dissociation, timing and
17 localization. SGs suppress translation and protect transcribed mRNAs from degradation
18 by temporarily storing selected mRNAs. SGs can be disassembled during stress
19 recovery and the stored mRNAs then become accessible for immediate translation. This
20 rapid reactivation is believed to be a response to environmental changes. Previous
21 studies have identified SG-regulated target mRNAs by analyzing RBPs present in SGs.
22 The identification of untranslated target mRNAs stored in SG has provided information
23 on the translational control or selective translation mechanism that occurs in response
24 to stress. In previous studies using hypoxic and heat stress samples, various direct
25 target mRNA identifications have been performed with multiomics analyses such as
26 RNA immunoprecipitation followed by sequencing (RIP-seq) analysis, transcriptome
27 analysis, translatoome and mRNA degradation rate analysis (Sorenson and Bailey-
28 Serres, 2014; Nguyen et al., 2016; Tian et al., 2022; Zhu et al., 2022). Although many
29 mRNA targets that are thought to be regulated by SGs have been revealed, their
30 subsequent fates, such as when translation-inhibited mRNAs are finally translated into
31 proteins, still remains unclear at this time.

32 While multiomics analyses, such as a RIP analysis combined with translatoome and
33 polysomal analyses, have enabled the identification of SG components and direct target
34 mRNAs that are SG-regulated, there are still many open questions that are await
35 clarification. For example, OLIGOURIDYLATE BINDING PROTEIN 1b (UBP1b), an SG
36 component, localizes to the nucleus under non-stress conditions. UBP1b-stress
37 granules (UBP1b-SG) are induced to form in the cytoplasm in response to heat stress,
38 and candidate target mRNAs for UBP1b have been identified. UBP1b is present in both
39 the nucleus and cytoplasm, but its precise location where it exerts its mRNA
40 stabilization role remains unclear (Figure 3). In addition, PBs and SGs co-localize,

1 suggesting that their constituent components might interact (Hamada et al., 2018).
2 Although many mRNAs have been described as SG targets, not all will be
3 translationally inhibited. It is plausible that some targets might be degraded by PBs
4 rather than stabilized by SGs; the underlying mechanism of recruitment of target
5 mRNAs remains to be elucidated. Future studies should elucidate how targeted mRNAs
6 are exactly regulated by SGs. Clarifying the mechanism(s) of selective translation will
7 be a major step forward in understanding stress responses in plants.
8

9 **Acknowledgments**

10 We would like to acknowledge Journal Doctors Inc. for critical reading of the manuscript.
11

12 **Funding**

13 This work was supported by grants from RIKEN, Grants-in-Aid for Scientific Research
14 provided by the Ministry of Education, Culture, Sports, Science and Technology
15 (Innovative Areas 18H04705) to MS, and grants from RIKEN, Next Generation
16 Acceleration Research Program to KN.
17

19 **The pervasive function of RNA structure in plant growth and development**

20 (Written by Yiliang Ding)

21 Plant growth and development is a continuous process starting with embryogenesis,
22 and the formation of the embryonic root and shoot, followed by organogenesis of
23 diverse organs such as roots, leaves, branches, and flowers. Plants rely on gene
24 expression regulation to achieve specific cell differentiation and elongation to form
25 different organs. This extremely high coordination of gene expression at both temporal
26 and spatial levels requires diverse regulatory mechanisms to achieve evolutionary
27 fitness. Furthermore, plants have evolved to adapt to wide-ranging environmental
28 conditions, acquiring highly dynamic regulation of gene expression in response to
29 different environmental factors. In addition to gene sequence content, RNA structure is
30 another important property of genes that can dynamically regulate gene expression at
31 the post-transcriptional level (Zhang and Ding, 2021).

32 Recent advances in RNA structure studies have enabled unprecedented opportunities
33 to determine the functional importance of RNA structure across varied aspects of plant
34 growth and development. For instance, the antisense long noncoding RNA (lncRNA),
35 *COOLAIR*, folds into a complex RNA structure (Hawkes et al., 2016) that was
36 suggested to suppress transcription of the key flowering gene, *FLOWERING LOCUS C*
37 (*FLC*) and promote flowering following vernalization. Another key regulator of plant
38 vascular development, *JULGI* (*JUL*), was shown to limit phloem differentiation through
39 its direct interaction with an RNA tertiary structure motif, RNA G-quadruplex, on the 5'
40 untranslated regions (5' UTRs) of *SUPPRESSOR OF MAX2 1-LIKE4/5* (*SMXL4/5*)

1 mRNA to suppress their translation (Cho et al., 2018). Other studies have shown that
2 RNA G-quadruplex affects plant root growth and development (Foley et al., 2017;
3 Zhang et al., 2019b; Yang et al., 2020a). Extensive studies have indicated that RNA
4 structural conformations change in response to temperature (Su et al., 2018; Chung et
5 al., 2020), light (Gawronski et al., 2021), salinity stress (Kramer et al., 2020; Tack et al.,
6 2020), and phosphate starvation (Reis et al., 2021). These changes subsequently alter
7 gene expression at the post-transcriptional level such as translation and RNA
8 degradation (Su et al., 2018; Chung et al., 2020; Kramer et al., 2020; Tack et al., 2020;
9 Gawronski et al., 2021; Reis et al., 2021). These recent studies have focused on either
10 identifying a specific RNA structural element on a specific transcript, or determining
11 global associations between RNA structure features and corresponding molecular
12 functions, and further support the growing evidence that highlights the importance of
13 RNA structure across diverse aspects of plant growth and development.

14 Since every mRNA is capable of folding into a particular RNA structure, this question
15 has stimulated interest to explore the pervasive role of RNA structure in individual
16 genes to gain a more comprehensive understanding of RNA structure-mediated
17 regulation in plant growth and development. To achieve this in-depth understanding, the
18 strategies employed for studying RNA structure functionality need to reach a new level.
19 A promising approach may be the capability of achieving specific cell type resolution. In
20 plants, although stem cells are pluripotent, their cellular trajectories are limited in scope
21 because the identity of any given cell depends on its position relative to its neighbors.
22 For instance, root growth starts from sets of stem cell initials in the quiescent center
23 (QC), which generate continuous parallel files of epidermal cells that divide in a
24 transverse, anticlinal orientation. Cells then divide in the meristematic zone before
25 starting to elongate into the differentiation zone of the mature root. After division, cells
26 remain in the same position and belong to the same lineage (Costa, 2016).
27 Interestingly, all these cell types share the same genetic information encoded in their
28 DNA, but with diverse cellular conditions. The folding status of RNA structure is highly
29 dependent on cellular conditions such as ion concentrations and interacting proteins
30 (Zhang and Ding, 2021). Thus, it is likely that RNAs may fold differently to specify gene
31 function in different cell types, resulting in unique cell identities (Figure 4). Future
32 research could focus on dissecting the extent of RNA structure diversities across
33 individual cell types. Indeed, the development of single-cell RNA structure profiling will
34 advance our understanding of RNA structural dynamics in plant cells.

35 Another future perspective could be to elucidate how RNA structures serve as
36 environmental sensors. During growth, plants are constantly challenged by fluctuating
37 environmental conditions such as biotic and abiotic stresses. Other abiotic stresses
38 such as flooding and drought are likely to affect the folding status of RNA structures due
39 to changing molecular concentrations in the cells (Zhang and Ding, 2021). During
40 pathogen infection, many metabolites are significantly altered that may also influence

1 RNA folding (Zhang and Ding, 2021). Additional research could focus on dissecting the
2 detailed mechanisms of RNA structure-mediated stress responses including
3 comprehensive assessments of different stresses, or different degrees and duration of
4 stress. Finally, it may be possible to assess the evolution of RNA structures across the
5 plant kingdom. Previous studies have illustrated the evidence of evolutionary selection
6 of certain RNA structure motifs (Yang et al., 2020a) and distinguished RNA structure
7 features in specific species (Deng et al., 2018; Yang et al., 2021). Studies of
8 evolutionary RNA structures may shed novel insight into understanding nucleotide
9 diversities in noncoding regions and at synonymous codon positions. Extension of RNA
10 structure studies with a phylogenomic perspective may provide an evolutionary
11 perspective on RNA structure functionality.

12 With the rise of transcriptome-wide RNA structuromes, large volumes of RNA structure
13 data now provide the necessary scope for deep learning applications with the potential
14 for translating fundamental knowledge into RNA structure-based molecular design. For
15 instance, from transcriptome-wide RNA structure and RNA stability data, we can now
16 learn and predict what kind of RNA structure features are responsible for RNA stability.
17 It may be possible to customize these RNA structure features into more or less stable
18 RNAs of interest. Where RNA structure acts as a post-transcriptional regulator, directly
19 affecting protein production, RNA structure-guided molecular design may offer the
20 potential for new avenues in synthetic biology.

21 Recent technological advances have significantly pushed the discovery of RNA
22 structure functionalities forward. Further innovations in PacBio and Nanopore
23 technologies to study RNA structures may offer more accurate RNA structure
24 information at single-base resolution. These upcoming developments will invigorate
25 RNA structure views to individual RNA structure conformations rather than the familiar
26 bulk conformation. RNA structures may be a type of hidden “codon” embedded in every
27 gene that facilitates the complexity of gene expression regulation. The rapid growth of
28 RNA structure research may ultimately reveal the regulatory power of RNA structures in
29 every aspect of plant growth and development.

30

31 **Acknowledgements**

32 Thanks to members of the Ding lab, Prof. Huakun Zhang and Miss Wenqing Sun for
33 their inspiring discussions. My apologies to those colleagues whose work could not be
34 cited due to length restrictions.

35

36 **Funding**

37 This work was supported by the United Kingdom Biotechnology and Biological Sciences
38 Research Council (BBSRC: BBS/E/J/000PR9788), the European Commission Horizon
39 2020 European Research Council (ERC) Starting Grant [680324] and the associated
40 Proof of Concept Grant [966855].

3 Sensing, regulation, and functions of R-loops in plants

4 (Written by Qianwen Sun)

5 The R-loop, a three-stranded chromatin structure comprising one single-stranded DNA
6 molecule and one RNA:DNA hybrid duplex, is widely distributed in the genome, with
7 essential roles in multiple cellular and disease processes (Garcia-Muse and Aguilera,
8 2019; Brickner et al., 2022; Petermann et al., 2022). Recent advances in genome-wide
9 detection methods have broadened our understanding of the distribution and dynamic
10 patterns of R-loops (Xu et al., 2022b). R-loops are involved in many biological
11 processes related to genome regulation, including transcription, replication, DNA
12 damage and repair, and chromatin organization (Zhou et al., 2022a). The biological
13 study of R-loops in plants began in 2013 when we discovered that an R-loop formed on
14 the promoter of the antisense lncRNA *COOLAIR* and affected the expression of *FLC*
15 (Sun et al., 2013b). In 2017, following the development of ssDRIP-seq (single-strand
16 DNA ligation-based library construction after DNA:RNA hybrid immunoprecipitation,
17 followed by sequencing), the localization of R-loops in the Arabidopsis genome was
18 revealed (Xu et al., 2017). The R-loop profiles of other plants have since been disclosed
19 (Figure 5). Through analysis of genome-wide data (Xu et al., 2017; Xu et al., 2020b),
20 some unique features of R-loop distribution in the nuclear genomes of plants have
21 emerged, prompting intriguing research directions in plant R-loop biology.

22 While analyzing the genome-wide distribution of R-loops, we identified a unique group
23 of R-loops formed by antisense lncRNAs near transcription start sites (TSS) named
24 asTSS_R-loops, (Xu et al., 2017). Similar patterns of R-loop distribution have also been
25 observed in other plant species, such as rice (*Oryza sativa*) and maize (*Zea mays*)
26 (Figure 5, and summarized in (Zhou et al., 2022a)). These conserved patterns raise
27 several questions: what are the functions of these asTSS_R-loops; how are they
28 transcribed, and is the transcriptional initiation of the antisense lncRNAs specific to
29 particular physiological or pathological responses? Another notable finding was the
30 prevalence of transfer RNA (tRNA)-promoted sense R-loops throughout the genome
31 (Xu et al., 2017). We discovered that these intragenic R-loops orchestrated
32 transcriptional interference between Pol II and Pol III, thus regulating the expression of
33 oxidative-responsive genes (Liu and Sun, 2021). Surprisingly, a large proportion of R-
34 loops is located in constitutive pericentromeric heterochromatin and overlaps with
35 H3K9me2 and H3K27me1 heterochromatic marks in Arabidopsis (Xu et al., 2017). This
36 observation raises intriguing questions about the functions of R-loops in
37 heterochromatin formation and organization in plants.

38 R-loops play important roles in cellular reprogramming in mammals (Li et al., 2020b;
39 Yan et al., 2020). During the lifecycle of Arabidopsis, R-loops showed a range of
40 dynamic changes during generational switches (such as flowering and germination) and

1 during recovery from long-term heat-stress treatment (Xu et al., 2020b). During the
2 transition from vegetative growth to flowering, R-loop formation decreased dramatically,
3 whereas from flower development to germination, R-loop formation gradually increased,
4 suggesting that the global reprogramming of R-loops also occurs in Arabidopsis. These
5 dramatic changes in R-loop formation likely co-occur with other events of genome
6 regulation, such as DNA replication and transcriptional reprogramming. It will be
7 important to explore the biological functions and regulatory mechanisms of R-loop
8 reprogramming during key developmental transitions in plants. Conversely, R-loops
9 likely function in transcriptional reprogramming during physiological and pathological
10 processes. Interestingly, Moore *et al.* proposed a model of R-loop-mediated
11 transcriptional reprogramming during plant defense responses (Moore et al., 2011),
12 although experimental evidence is still lacking.

13 Most R-loops form and function in *cis*. However, *trans*-formed R-loops may also play
14 important roles in plants. For example, the lncRNA *APOLO* promoted *trans*-R-loop
15 formation and altered chromatin loop conformation (Ariel et al., 2020). Current detection
16 methods cannot provide information about whether R-loops form in *cis* or in *trans*,
17 underscoring the need to develop a high-throughput technique for distinguishing *cis*- or
18 *trans*-R-loops globally. Moreover, it would be useful to alter the levels of R-loops at
19 specific genomic loci (Liu and Sun, 2021), but there is currently no efficient way to
20 modulate an entire group of R-loops (such as asTSS_R-loops) jointly. asTSS_R-loops
21 were recently proposed to promote co-transcriptional micro RNA (miRNA) processing
22 (Gonzalo et al., 2022). However, the lack of tools for modulating R-loops makes it
23 challenging to study the functions of particular groups of R-loops with similar distribution
24 patterns in the genome. Alternatively, identifying the specific regulators of a particular
25 group of R-loops could help solve this problem.

26 To date, several R-loop modulators have been identified in plants (Zhou et al., 2022a).
27 Among these, the evolutionarily conserved RNase H1 proteins specifically remove the
28 RNA moiety in RNA:DNA hybrids, thus resolving R-loops efficiently. The Arabidopsis
29 genome encodes three RNase H1 proteins: AtRNH1A, AtRNH1B, and AtRNH1C (Yang
30 et al., 2017). While AtRNH1B and AtRNH1C are involved in stabilizing the genome
31 integrity of semi-autonomous organelles (mitochondria and chloroplasts) (Yang et al.,
32 2017; Cheng et al., 2021; Wang et al., 2021b), the biological function of nucleus-
33 localized AtRNH1A is still unclear. The biological functions of RNase H1 proteins and
34 other R-loop regulators in different plant species also need to be further explored.

35 Organisms must integrate and coordinate the activities of different tissues and cell
36 types. Precisely analyzing the genome-wide patterns of R-loops from ultra-low input
37 samples is difficult using current methods (Zhou et al., 2022a). It is imperative to
38 establish ultralow-input (or even single-cell) R-loop profiling techniques to systematically
39 explore the functions of R-loops in critical genomic events in complex tissues. Such
40 tools could be powerful for dissecting R-loop distribution and dynamics in specific cell

1 types and during specific differentiation programs, such as double fertilization in plants.
2 As the topological state of the genome could affect R-loop formation, it would be useful
3 to develop tools to quantitatively measure topological conformation and explore how the
4 3D genome organization influences R-loop formation. Advanced computational
5 predictions of R-loops genome-wide could complement experimental approaches for
6 species with available genome sequence information.

7 Chloroplasts and mitochondria are semi-autonomous organelles of endosymbiotic origin
8 with their own genetic materials. In the face of complex external environmental
9 conditions and internal growth and developmental factors, how these organelles
10 maintain their genome stability has long been unclear. We recently discovered that R-
11 loops act as regulatory centers in determining the stability of organelle genomes (Figure
12 5). R-loops play both positive and negative roles in maintaining the stability of the
13 organellar genome, which not only causes genome instability by modulating head-on
14 transcription-replication conflicts (Yang et al., 2017; Yang et al., 2020b) but also
15 promotes DNA damage repair (Cheng et al., 2021; Wang et al., 2021b). However, our
16 knowledge about how R-loop levels are sensed and adjusted to maintain normal
17 organellar function is still in its infancy. For example, in cells lacking mitochondrion-
18 localized AtRNH1B, chloroplast-localized AtRNH1C sensed high R-loop levels and
19 relocalized to mitochondria via an unknown mechanism (Figure 5). This observation
20 raises the following intriguing question: Do plants sense R-loops during chloroplast-
21 mitochondria communication? Furthermore, how do plants coordinate and adjust R-loop
22 levels inside and between cells, and how is this process managed in response to
23 physiological and pathological processes?

24 **Acknowledgements**

25 I greatly appreciate my current and previous lab members and collaborators for their
26 efforts and contributions to the R-loop biology studies. I thank Dr. Jincong Zhou for the
27 assistant of Figure 5 illustration. I apologize to the colleagues whose work could not be
28 cited due to the length limitation.

29 **Funding**

30 The Sun Lab is supported by grants from the National Natural Science Foundation of
31 China (grant nos. 91740105, 91940306, and 31822028) and Tsinghua-Peking Center
32 for Life Sciences.
33
34
35
36

37 **Epitranscriptomic mRNA modification: a potent regulatory mechanism in plant 38 development and stress responses**

39 (Written by Hunseung Kang)

1 Epitranscriptomic RNA modifications, which are analogous to epigenetic regulation that
2 involves DNA methylation and histone modifications, are emerging as a new layer of
3 gene regulation. These modifications play a pivotal role in fine-tuning plant development
4 and fitness to changing environmental cues. At least 160 mRNA modifications have
5 been identified to date, among which N⁶-methyladenosine (m⁶A), N¹-methyladenosine
6 (m¹A), and 5-methylcytidine (m⁵C) are common and abundant internal modifications
7 observed in coding RNAs; m⁶A is the most prevalent internal modification in eukaryotic
8 mRNAs (Boccaletto et al., 2018). Methyltransferases (referred to as “writers”),
9 demethylases (referred to as “erasers”), and RNA-binding proteins (referred to as
10 “readers”) are cellular components responsible for the installation, removal, and
11 interpretation of m⁶A marks, respectively (Figure 6). Recent transcriptome-wide m⁶A
12 mapping, as well as the identification and characterization of m⁶A writers, readers, and
13 erasers in *Arabidopsis* and model crops, have enhanced our understanding of the
14 dynamics, distribution, regulatory mechanisms, and biological functions of m⁶A
15 methylation in plant development and stress responses.

16 Transcriptome-wide analyses of m⁶A methylation patterns in plants have led to the
17 identification of an RR(m⁶A)CH (R = A/G; H = A/C/U) motif found in all eukaryotes (Luo
18 et al., 2014; Duan et al., 2017; Hu et al., 2021) and a URU(m⁶A)Y (Y = C/U) motif
19 unique to plants (Arribas-Hernandez et al., 2018; Arribas-Hernandez et al., 2021a;
20 Arribas-Hernandez et al., 2021b; Hu et al., 2021; Hu et al., 2022). The presence of the
21 plant-specific m⁶A motif, as well as the common m⁶A motif conserved across all
22 eukaryotes, suggests that m⁶A modifications exert multifaceted functions in plants. The
23 m⁶A writers responsible for these modifications include METHYLTRANSFERASE A
24 (MTA), MTB, FKBP12-interacting protein 37 (FIP37), VIRILIZER (VIR), the E3 ubiquitin
25 ligase HAKAI, and FIONA1 (FIO1) (Ruzicka et al., 2017); reviewed in (Hu et al., 2019;
26 Xu et al., 2022a). The three m⁶A erasers, AlkB homolog 2 (ALKBH2), ALKBH9B, and
27 ALKBH10B, have been confirmed as m⁶A demethylases (Duan et al., 2017; Martinez-
28 Perez et al., 2017; Zhou et al., 2019) (Figure 6). YT521-B homology (YTH)-domain
29 proteins have been characterized as m⁶A readers that recognize m⁶A marks and affect
30 the stability, translation, nucleus-to-cytoplasm movement, and alternative
31 polyadenylation of m⁶A-modified transcripts (Arribas-Hernandez et al., 2018; Scutenaire
32 et al., 2018; Wei et al., 2018; Hu et al., 2019; Arribas-Hernandez et al., 2021a; Arribas-
33 Hernandez et al., 2021b; Song et al., 2021; Hou et al., 2022) (Figure 6). Dynamic and
34 reversible m⁶A methylation play vital roles in embryogenesis, morphogenesis, trichome
35 morphology, root development, and fruit ripening (Ruzicka et al., 2017; Arribas-
36 Hernandez et al., 2018; Hu et al., 2019; Zhou et al., 2019; Hu et al., 2022) (Figure 6).
37 Accumulating evidence has highlighted the pivotal roles of m⁶A modifications in plant
38 growth and development. However, several questions, including the mechanism by
39 which m⁶A is added to, or removed from, mRNA transcripts in a growth stage-

1 dependent manner and differentially regulates the abundance of transcripts crucial for
2 plant development, remain unanswered.

3 Mapping and characterization of mRNA modifications in plant stress responses are
4 currently at the nascent stage. Bioinformatics analyses revealed that the expression
5 levels of m⁶A writers, erasers, and readers change differentially in response to diverse
6 stresses (Hu et al., 2019), suggesting a vital role for m⁶A methylation in plant stress
7 responses. Recent molecular evidence has established a link between mRNA
8 modifications and transcript levels involved in plant stress responses (Hou et al., 2021;
9 Hu et al., 2021; Hou et al., 2022). Notably, m⁶A modifications play crucial roles in plant
10 responses to diverse stresses, including salt, drought, and nutrient (nitrate) starvation,
11 by affecting mRNA stability, alternative polyadenylation, and translation efficiency of
12 stress-responsive genes (Hou et al., 2021; Hu et al., 2021; Hou et al., 2022). However,
13 the precise mechanism underlying RNA modification-mediated gene regulation during
14 stress adaptation requires further investigation. Therefore, the crucial aspects that
15 remain unexplored are the mechanisms by which RNA modification patterns vary under
16 specific stress conditions and the association of these modifications with stress-induced
17 alterations in transcript and protein levels.

18 Most studies conducted thus far have focused on the cellular components responsible
19 for RNA methylation and their roles in the nucleus and cytoplasm. Chloroplast and
20 mitochondrial RNAs are highly m⁶A-methylated, accounting for 98–100% and 86–90%
21 of the transcripts in chloroplasts and mitochondria, respectively (Luo et al., 2014; Wang
22 et al., 2017b). Therefore, RNA methylation might likely exert crucial roles in plant
23 organelles. However, the nature and identity of writers, erasers, and readers in
24 chloroplasts and mitochondria, except m⁴C and m₂⁶A rRNA writers in chloroplasts, are
25 largely unknown (reviewed in (Manduzio and Kang, 2021). Analysis of chloroplast
26 proteomes by liquid chromatography-tandem mass spectrometry and prediction of
27 organelle-localized proteins have revealed that the m⁶A writer components MTA, MTB,
28 and FIP37 found in plant nuclei were also possibly localized in chloroplasts and
29 mitochondria and several putative S-adenosyl methionine (SAM)-dependent
30 methyltransferase proteins are present in the chloroplasts of Arabidopsis (reviewed in
31 (Manduzio and Kang, 2021). Further verification of the methyltransferase activity of
32 these putative writer proteins, as well as the previously unknown erasers or readers in
33 chloroplasts and mitochondria, will help elucidate the significance of RNA modifications
34 in plant organelles.

35 Rapid progress in transcriptome-wide mapping and the identification of writers, readers,
36 and erasers have unraveled the regulatory roles of m⁶A modification in plant
37 development and stress responses. Nonetheless, many challenges remain in mapping
38 m⁶A modifications at single-base resolution using recently advanced sequencing
39 methods, including Nanopore direct transcriptome deep sequencing (RNA-seq),
40 MAZTER-seq, m⁶A-REF-seq (m⁶A-sensitive RNA-Endoribonuclease-Facilitated

1 sequencing), and miCLIP-seq (m⁶A individual-nucleotide-resolution cross-linking and
2 immunoprecipitation combined with high-throughput sequencing). Furthermore,
3 characterizing novel cellular components of writers, readers, and erasers in crops will
4 help firmly establish the molecular link between m⁶A, crop productivity, and stress
5 adaptation. Recent findings have suggested that m⁶A is associated with LLPS, which
6 expands the repertoire of regulatory mechanisms crucial for cellular responses to
7 developmental and environmental cues (Scutenaire et al., 2018; Ries et al., 2019; Song
8 et al., 2021). Integrating these molecular insights to the regulatory roles of m⁶A
9 modification with novel genome-editing technologies, including A-to-G base editing to
10 modify potential m⁶A sites and clustered regularly interspaced short palindromic repeat
11 (CRISPR)/CRISPR-associated nuclease 13 (Cas13)-based targeted RNA methylation
12 (Liu et al., 2019; Li et al., 2020a), will greatly facilitate epitranscriptomics research and
13 lead to the development of a potential strategy for breeding stress-tolerant crops via
14 precisely engineered RNA modifications. Further exploration of this field is warranted,
15 and we anticipate exciting discoveries in the near future.

16

17 **Acknowledgments**

18 I would like to acknowledge the seminal contributions of those colleagues whose work
19 could not be cited owing to length restrictions.

20

21 **Funding**

22 This work was supported by grants from the Mid-career Researcher Program through
23 the National Research Foundation of Korea, funded by the Ministry of Science, ICT and
24 Future Planning (NRF-2021R1A2C1004187), Republic of Korea, and the New Breeding
25 Technologies Development Program (PJ01652401), Rural Development Administration,
26 Republic of Korea.

27

28

29 **Decoding the grammar of plant long noncoding RNAs**

30 (Written by Federico D. Ariel and Martin Crespi)

31 The inspection of the presence and combination of domains within a protein is generally
32 a good starting point to infer its potential molecular action. This information is then
33 complemented with subcellular localization studies, biochemical characterization,
34 analysis of expression patterns of the encoding gene across multicellular organisms and
35 genetic approaches to propose a biological role of the given gene in plants. By contrast,
36 the comprehensive functional characterization of lncRNAs (Wierzbicki et al., 2021) is a
37 challenging task that should take into account (i) their promiscuous or specific
38 interaction with other molecules based on their sequence and/or structure; (ii) their
39 redundancy with other unrelated transcripts; (iii) their subcellular localization; (iv) their
40 role within molecular regulatory networks; and (v) an eventual RNA biological activity

1 (Figure 7). In the last 15 years, thousands of lncRNAs have been annotated from a
2 growing number of plant species, although their functional characterization lags behind,
3 thus severely hindering the differentiation between transcriptional noise and biologically
4 relevant noncoding transcripts. Identifying general molecular features linking specific
5 lncRNAs with their targets have uncovered certain mechanisms. For instance, target
6 mimicry of miRNAs (RNA molecules acting as decoy of miRNAs blocking their activity)
7 was demonstrated for *INDUCED BY PHOSPHATE STARVATION 1 (IPS1)* and could
8 be later predicted *in silico* for other lncRNAs across species (Franco-Zorrilla et al.,
9 2007). However, for the large majority of lncRNAs acting through other molecular
10 mechanisms, there is no evident features to define their targets in order to dissect the
11 molecular basis governing their action in plants.

12 Upon extensive annotation of lncRNAs across species, future screenings for biological
13 functions, likely based on systematic CRISPR-derived approaches, may empower the
14 selection of novel relevant lncRNAs for in-depth molecular characterization. In addition,
15 integration of lncRNA expression patterns from transcriptomic data of multiple wild-type
16 plants, mutants, and natural accessions in response to environmental and
17 developmental cues will position the lncRNA of interest within particular regulatory
18 networks driving plant development and/or adaptation to the environment.

19 Specific lncRNAs have been shown to interact with protein partners in ribonucleoprotein
20 (RNP) complexes (modulating their stability, subcellular localization, or their activity),
21 DNA (forming RNA-DNA duplexes known as R-loops), or other transcripts (such as
22 antisense RNAs, forming paired RNA regions triggering mRNA degradation or
23 promoting translation) (Lucero et al., 2021). Future research to generalize these
24 interactions may include global identification of RNAs forming R-loops or interacting with
25 specific RNP complexes involved in splicing modulation (Rigo et al., 2020) or the
26 translational machinery (Bazin et al., 2017). Another emerging mechanism is the
27 interaction of lncRNAs with chromatin-related proteins linked to epigenetic regulations
28 such as LIKE HETEROCHROMATIN PROTEIN 1 (LHP1), CURLY LEAF (CLF),
29 MEIOTIC F-BOX (MOF), ARABIDOPSIS TRITHORAX-LIKE PROTEIN1 (ATX1) or
30 WD40 REPEAT 5A (WDR5a) (Fonouni-Farde et al., 2021) although their binding
31 specificity remains uncertain, or even with transcription factors (e.g. WRKY42, (Moison
32 et al., 2021)). In addition, the identification of nascent RNAs (Kindgren et al., 2020) as
33 well as chromatin-associated lncRNAs based on chromatin isolation and high
34 throughput sequencing techniques will further contribute to creating a matrix of lncRNA
35 features underlying their function. Altogether, mapping lncRNA interactions with DNA,
36 chromatin, and proteins involved in a wide range of mechanisms in model and crop
37 plants should set the stage for a comprehensive classification of lncRNAs enabling the
38 search of singularities and commonalities behind the functions of noncoding transcripts.
39 The identification of protein partners and lncRNA-interacting nucleic acids using
40 biotinylated probes for the purification of lncRNA-containing complexes followed by

1 mass spectrometry or DNA sequencing is an initial key goal to define the lncRNA
2 interactome, despite the potential artifacts linked to these approaches (Machyna and
3 Simon, 2018). Alongside the genome-wide identification of lncRNAs participating in
4 alternative RNP complexes, the detailed characterization of selected lncRNA actions on
5 these complexes remains essential to better understand the diversity of regulatory
6 mechanisms involving noncoding transcription.

7 Another major question in lncRNA biology and biochemistry concerns transcript
8 structure (Zhu et al., 2021). Secondary and tertiary structures of RNAs are very likely
9 determinant features for their dynamic interaction with proteins and other partners.
10 Considering that plants cannot modulate their body temperature, the structure of
11 lncRNAs may serve as potential versatile molecules acting as thermosensors in order to
12 rapidly adjust epigenomic features and alternative splicing, two major processes
13 affected by ambient temperature (John et al., 2021; Perrella et al., 2022). A growing
14 number of prediction tools based on classical and machine learning approaches have
15 shed light on this field (Bugnon et al., 2022), although the biochemical characterization
16 of individual or groups of plant lncRNAs is just starting. In general, genome-wide
17 approaches for the mapping of double-stranded RNAs (dsRNAs) or chemical
18 degradation profiles to reconstruct transcript structures fail to deliver enough data about
19 low-abundance lncRNAs. However, *in vitro* transcription of selected lncRNAs followed
20 by biochemical approaches ignores the enormous collection of epitranscriptomic
21 modifications as well as their *in vivo* interaction with partner molecules, which are likely
22 to affect RNA structure (Miller et al., 2022).

23 Similar to the study of metazoan lncRNAs, cell biology techniques, notably single
24 molecule RNA (smRNA) fluorescence *in situ* hybridization (FISH) (Duncan et al., 2017),
25 can contribute to our understanding of the mechanisms involving specific lncRNAs. As a
26 complement to subcellular fractionation studies followed by high-throughput
27 sequencing, smRNA FISH may not only indicate whether a given lncRNA accumulates
28 in the nucleus or the cytoplasm, but also reveal its distribution in “speckles”, or
29 localization in specific loci, in subcellular compartments, in non-membranous organelles
30 or particles. However, the technical difficulties related to the presence of cell wall
31 barriers in plant tissues prevents the accessibility of fluorescent oligonucleotide probes,
32 thus delaying the massive use of this approach by most plant RNA biology groups, in
33 comparison to labs working on mammalian cell culture models.

34 The fields of plant RNA biology and biochemistry will need to integrate cell biology, RNP
35 proteomics, genomic and genetic approaches to unveil the function and evolution of the
36 noncoding transcriptome, in particular during differentiation and environmental stress
37 responses. Evolutionary analysis at a global level (e.g. involving synteny) of lncRNAs
38 exhibiting common features (e.g. integration into specific RNPs), together with the in-
39 depth characterization of specific leading cases, will achieve a better understanding of
40 the structures and sequences (likely very short) setting the specificity rules of their

1 interaction with partner molecules. As the RNA interactome ultimately determines their
2 function, these integrated approaches will hopefully help us uncover the grammar of
3 plant lncRNAs.

4 5 **ACKNOWLEDGEMENTS**

6 FA and MC are part of the International Associated Laboratory NOCOSYM (CNRS).
7
8

9 **Emerging and long-standing questions about miRNA biogenesis in plants**

10 (Written by Axel Giudicatti and Pablo A. Manavella)

11 From the point of view of RNA biology, miRNAs are exciting molecules. Not only do
12 mature miRNAs target other RNA molecules to block their translation or trigger their
13 degradation, but their precursors undergo nearly all the regulatory features described in
14 this article. For instance, many *MIRNA* genes contain introns that affect the processing
15 of the primary transcripts (pri-miRNAs) (Stepien et al., 2017); the ribonucleotides of
16 miRNA precursors can be modified or edited to change their regulatory outcome
17 (Mingardi et al., 2018; Bhat et al., 2020); pri-miRNA secondary structure fluctuations
18 define miRNA biogenesis (Wang et al., 2018b; Re et al., 2019); even aTSS_R-loops
19 were recently shown to promote co-transcriptional processing of miRNAs (Gonzalo et
20 al., 2022). These features make miRNAs a unique entity where many aspects of RNA
21 biology converge. Even after more than 20 years of research, all these aspects of
22 miRNA biology present unresolved questions and intriguing gaps in our knowledge. For
23 instance, although we know that there is a crosstalk between splicing and pri-miRNA
24 processing (Stepien et al., 2017), it is unclear how these two processes interact. The
25 transcription and processing of pri-miRNAs is coupled (Fang et al., 2015; Gonzalo et al.,
26 2022). This observation opens the possibility that the crosstalk between pri-miRNA
27 processing and splicing only exists for miRNAs processed co-transcriptionally where
28 both machineries, the spliceosome and microprocessor, meet. In this scenario, it is
29 unclear whether the miRNA processing factors and splicing factors act cooperatively or
30 simply interfere entropically with each other over the nascent pri-miRNAs during
31 maturation. Advances in RNA sequencing technologies, especially of nascent RNAs,
32 will help our understanding of how these two processes are connected (Figure 8C).

33 On its own, the discovery of coupling between transcription and miRNA processing,
34 initially suggested by the ground-breaking work of Fang *et al* (Fang et al., 2015) and
35 further confirmed in 2022 (Gonzalo et al., 2022), opened many exciting new avenues of
36 inquiry. The recruitment of the microprocessor to *MIRNA* loci is a well-reported
37 phenomenon (Fang et al., 2015; Cambiagno et al., 2021). However, how the
38 microprocessor specifically recognizes these loci over any other Pol II-transcribed
39 region remains an enigma. Still, the association of the microprocessor to *MIRNAS*
40 requires the presence of the pri-miRNA transcript (Fang et al., 2015). Thus, it is possible

1 that the Dicing complex recognizes the stem-loop structure within pri-miRNA transcripts,
2 thereby giving specificity to the system. Co-transcriptional miRNA processing appeared
3 favored in those loci containing aTSS_R-loops (Gonzalo et al., 2022). These three-
4 stranded chromatin structures may also provide an initial signal promoting the
5 recruitment of the microprocessor to these loci, although their functions in this process
6 are still merely hypothetical. Nevertheless, this result raises the possibility that the
7 three-stranded hybrid is the platform upon which the microprocessor is built. It will be
8 interesting to study whether any of the proteins proposed to link the microprocessor to
9 chromatin have affinity for R-loops, either for the single-stranded DNA or the RNA/DNA
10 hybrid (Figure 8B). The assembly of the processing complex also presents a
11 challenging, but very relevant, problem to solve; which is the hierarchical order of
12 recruitment of the microprocessor components to *MIRNA* loci? Another compelling
13 question raised from the discovery of the processing of nascent pri-miRNAs is whether
14 co-transcriptionally processed miRNAs have distinct functions. In this sense, it was
15 recently shown that the protein HASTY (HST) is required for both the assembly of the
16 microprocessor at *MIRNA* loci and to promote the non-cell-autonomous function of
17 miRNAs (Brioudes et al., 2021; Cambiagno et al., 2021). It is therefore possible that
18 miRNAs processed during transcription take a particular road that makes them mobile
19 molecules (Figure 8E). Perhaps this pool of miRNAs somehow avoids loading into
20 ARGONAUTE 1 (AGO1), an event proposed to lock miRNAs inside the cell, preventing
21 their movement (Devers et al., 2020; Fan et al., 2022; Voinnet, 2022). It is curious that
22 the precise mechanisms of miRNA movement between cells and whether such
23 movement is chaperoned, still remains unknown. In fact, this question is probably one of
24 the longest-standing questions in the field.

25 Among the four DICER-Like (DCL) enzymes in *Arabidopsis*, DCL1 is the main actor in
26 miRNA processing, due to its nuclear localization and preference to process imperfect
27 stem-loop folded RNAs. Within the pri-miRNA stem-loop, DCL1 recognizes structural
28 features that guide processing to release a unique miRNA duplex (Bologna et al., 2013;
29 Manavella et al., 2019). It was recently shown that the folding of pri-miRNAs can be
30 altered, consequently modifying processing efficiency (Wang et al., 2018b). In addition,
31 nucleotides at pri-miRNAs can be modified and even edited, although the influence that
32 these events have over the miRNA processing were not demonstrated in plants
33 (Mingardi et al., 2018; Bhat et al., 2020). The role of RNA editing, modification, and re-
34 folding in miRNA processing are just emerging as important regulatory mechanisms and
35 deserve our attention. It is expected, as plants are nonthermogenic organisms, that the
36 secondary structure of plant pri-miRNAs will fluctuate with ambient temperature, likely
37 affecting their processing. Thus, it can be envisioned that some miRNAs may even act
38 as thermosensors (Figure 8A). Although we do have some evidence that temperature
39 changes how miRNAs are processed (Re et al., 2019), much more needs to be done on
40 this subject.

1 D-bodies are one of the most intriguing elements in the miRNA pathway. These discrete
2 membraneless nuclear speckles are the typical localization of many fluorescently-
3 tagged miRNA biogenesis proteins (Fang and Spector, 2007). The localization of these
4 proteins led to the proposal that D-bodies are the center of miRNA processing in plants.
5 A recent study showed that D-bodies arise through SERRATE (SE)-mediated phase
6 separation (Xie et al., 2021). Disruption of SE phase separation, and thus D-body
7 formation, by deleting the N-terminal intrinsically disordered region (IDR) of SE reduces
8 miRNA accumulation, supporting the idea that D-bodies are sites of pri-miRNA
9 processing. The role of D-bodies in miRNA processing is also supported by several
10 studies showing a correlation between D-body formation and miRNA production.
11 Intriguingly, other studies have shown that the disappearance of D-bodies does not
12 affect the ability of the cell to produce miRNAs (discussed by (Mencia et al., 2022)).
13 This observation suggests that D-bodies are not the sole place of miRNA processing
14 and raises the possibility that compensatory mechanisms act to offset the reduction of
15 miRNAs caused by the loss of D-bodies. While we now know that miRNA can be
16 processed co-transcriptionally (Fang et al., 2015; Cambiagno et al., 2021; Gonzalo et
17 al., 2022), many pri-miRNAs are partially or entirely processed in the nucleoplasm, likely
18 in D-bodies (Gonzalo et al., 2022). Thus, a balance between these two processing sites
19 may buffer any fluctuation in processing and maintain stable levels of miRNAs. Given
20 the current data, it is hard to simply categorize D-bodies as the only place where miRNA
21 biogenesis occurs. It is also possible that D-bodies are not unique entities but rather a
22 collection of small micro-reactors of different compositions and functions (Figure 8D).
23 This idea goes along with the finding that despite localizing to D-bodies, some miRNA
24 factors do not co-localize with each other (Tomassi et al., 2020). We previously
25 discussed several possible scenarios for D-body functions (Mencia et al., 2022).
26 Defining the nature and role of D-bodies and their crosstalk with co-transcriptional
27 processing is at the frontier of miRNA research, although it is a technically challenging
28 goal. Future studies applying state-of-the-art *in vivo* immunostaining and biochemical
29 approaches will certainly surprise us with new discoveries about these nuclear
30 speckles.

31 These are only a few of the many open questions regarding how miRNAs are produced
32 and do not even consider the equally large number of questions we have regarding how
33 miRNAs act once loaded into the RNA-induced silencing complex (RISC) and how
34 these molecules are stabilized or degraded when necessary.

35

36 **Acknowledgments**

37 Thanks to all those whose important contributions inspired and promote the miRNA
38 research and that could not be cited here due to the short format of this article.

39

40

1 **Funding**

2 Work on miRNA biogenesis in the Manavella group is supported by ANPCyT (Agencia
3 Nacional de Promoción Científica y Tecnológica, Argentina) and UNL (Universidad
4 Nacional del Litoral).

7 **Cross-kingdom RNAi**

8 (Written by Qiang Cai and Hailing Jin)

9 Over the years, studies on extracellular RNAs, including small RNAs (sRNAs), have
10 focused mostly on their movement between cells and tissues within an organism (Liu
11 and Chen, 2018; Huang et al., 2019). Naturally occurring sRNA trafficking across
12 organismal boundaries between interacting organisms that induces gene silencing in
13 the counter party, a biological phenomenon named cross-kingdom/species RNA
14 interference (RNAi), was first described in plant-fungal interactions (Weiberg et al.,
15 2013). During infection, the gray mold fungal pathogen *Botrytis cinerea* delivers its
16 sRNAs, called sRNA effectors, into plant cells and hijacks the plant RNAi machinery to
17 silence those host genes that are involved in plant immunity (Weiberg et al., 2013).
18 Similar phenomena were later observed in many plant pathogens and parasites. For
19 example, sRNAs from the fungal pathogens *Verticillium dahlia* (causing verticillium wilt
20 of cotton [*Gossypium hirsutum*]) and *Puccinia striiformis* (causing stripe rust of wheat
21 [*Triticum aestivum*]) can move into their plant host and silence plant defense genes
22 (Wang et al., 2016; Wang et al., 2017a). Similarly, oomycete pathogens, such as
23 *Hyaloperonospora arabidopsidis* (causing downy mildew of Arabidopsis) (Dunker et al.,
24 2020), also utilize cross-kingdom RNAi to achieve aggressive infection. Furthermore,
25 miRNAs from parasitic plant dodders (*Cuscuta campestris*) act as cross-species
26 regulators of gene expression in their plant hosts, suggesting that mobile sRNAs act as
27 virulence factors during parasitism (Shahid et al., 2018). Cross-kingdom RNAi is not
28 limited to pathogenic interactions but also exists in symbiotic interacting systems. A
29 recently discovered fungal miRNA from the beneficial ectomycorrhizal fungus *Pisolithus*
30 *microcarpus* enters *Eucalyptus grandis* root cells and stabilizes the symbiotic interaction
31 by silencing several nucleotide-binding (NB)-ARC domain-containing proteins from the
32 host (Wong-Bajracharya et al., 2022). Even for prokaryotic pathogens that do not have
33 a canonical RNAi pathway, rhizobial tRNA-derived short RNAs act as functional sRNAs
34 moving into plant cells to silence nodulation-related target genes (Ren et al., 2019).
35 Most strikingly, the molecular mechanism underlying cross-kingdom RNAi is also
36 conserved. The sRNAs from the fungal pathogens *B. cinerea* and *V. dahlia*, the
37 oomycete pathogen *H. arabidopsidis*, and the rhizobium were all found to be loaded into
38 the plant host AGO1 to silence host target genes (Weiberg et al., 2013; Wang et al.,
39 2016; Ren et al., 2019; Dunker et al., 2020).

1 Recent studies have shown that cross-kingdom RNAi is bidirectional, and many plant
2 species can also transport endogenous sRNAs into their interacting pathogens (Cai et
3 al., 2021; Liu et al., 2021). For example, Arabidopsis plants send miRNAs, phased
4 secondary small interfering RNAs (phasiRNAs), and other endogenous short interfering
5 RNAs (siRNAs) into interacting *B. cinerea* cells (Cai et al., 2018). These transported
6 host sRNAs can silence *B. cinerea* virulence-related genes, many of which are involved
7 in fungal vesicle-trafficking pathways (Cai et al., 2018). Cross-kingdom sRNA trafficking
8 from host plants into pathogens was also observed in other plant-fungal systems, such
9 as cotton-*V. dahliae* and wheat-*F. graminearum* interaction systems (Cai et al., 2021).
10 It has been demonstrated that plant sRNAs are transported into fungal cells mainly by
11 extracellular vesicles (EVs) (Cai et al., 2018). EVs are heterogeneous membrane-
12 encapsulated structures that transport different RNA and protein cargoes between cells
13 (Mathieu et al., 2019). EVs play an important role in sRNA trafficking between cells and
14 tissues in both animal and plant systems (Cai et al., 2019; Mathieu et al., 2019). Like
15 animal cells, a heterogeneous population of EVs exists in plants (Cai et al., 2019;
16 Huang et al., 2021b). In Arabidopsis, a distinct class of EVs, called tetraspanin (TET)-
17 positive exosomes, are responsible for secretion and transport of functional sRNAs and
18 play a significant role in cross-kingdom RNAi and plant-microbial interactions (Cai et al.,
19 2018; He et al., 2021).
20 How specific plant sRNAs are selectively loaded into EVs has long remained poorly
21 understood. A recent study identified a list of EV-localized RNA-binding proteins,
22 including AGO1, DEAD-box RNA helicases (RH11, RH37, and RH52), and ANNEXIN 1
23 and 2 (ANN1 and ANN2) (He et al., 2021). AGO1, RH11, and RH37 were shown to
24 selectively bind to a set of sRNAs that are found in EVs, and contribute to selective
25 sRNA loading into EVs, mostly TET-positive exosomes, whereas ANN1/2 bind to RNAs
26 non-specifically. The level of sRNAs is reduced in EVs isolated from *ann1 ann2*
27 mutants, which indicates that ANN1/2 are involved in sRNA stabilization in EVs,
28 although they do not contribute to selective sRNA loading (He et al., 2021).
29 Research on cross-kingdom RNAi and sRNA trafficking is still in its infancy, and
30 increasing studies demonstrate that mobile RNA molecules are important regulatory
31 elements of the interaction between hosts and interacting organisms (Huang et al.,
32 2019). Bidirectional cross-kingdom RNAi has been developed during the co-
33 evolutionary arms race between hosts and pathogens, which has become a widespread
34 molecular regulatory mechanism in plant-microbial interaction and plays a significant
35 role in host immunity and pathogen virulence (Huang et al., 2019). Current studies show
36 that EVs are essential in transporting sRNAs from the plant hosts to pathogens (Cai et
37 al., 2021). EV-mediated sRNA transport has evolved in both plant and animal systems,
38 suggesting that it is likely a conserved mechanism for cell-to-cell communication. The
39 current understanding of cross-kingdom RNA transport is just the tip of the iceberg.
40 Many questions remaining to be answered in this field are: i) Can pathogen sRNAs act

1 as effector molecules, and can plants sense them as pathogen-associated molecular
2 patterns (PAMPs)? ii) Do plant EVs also transport other classes of RNAs, i.e., mRNAs
3 and lncRNAs, into pathogen cells to inhibit virulence? iii) Are there other mechanisms
4 by which plant RNAs are selectively loaded into EVs? iv) Do fungal pathogens also
5 utilize EVs to deliver RNAs into host plants? v) Besides EVs, do other pathways
6 contribute to cross-kingdom RNA transport? A better understanding of RNA
7 communications between interacting organisms will contribute to the development of
8 new strategies for disease control and crop protection, such as EV-based sRNA
9 fungicides.

10 **Funding**

11 Work in the Dr. Qiang Cai's laboratory was supported by grants from the National
12 Natural Science Foundation of China (32070288).
13
14
15

16 **Why does germline development require specialized small RNAs?**

17 (Written by Xiaoqi Feng)

18 Small interfering RNAs (siRNAs) move between cells and exert regulatory functions
19 during plant and animal development (Chen and Rechavi, 2022). Specialized,
20 somatically produced siRNAs play essential roles during plant germline development.
21 Similarly, a special army of siRNAs operates in the animal germline, called Piwi-
22 interacting sRNAs (piRNAs) (Ozata et al., 2019). A central question arising from these
23 reproductive-cell-specific siRNAs is why such specificity? What is intrinsic about sexual
24 reproduction that requires specialized siRNAs? This is arguably one of the most exciting
25 questions in RNA and reproductive biology. As these siRNAs have diverse, pleiotropic
26 roles during reproductive development, investigation of multiple eukaryotic lineages is
27 necessary to resolve this question.

28 Pioneering evidence of soma-germ siRNA movement in plants came from Arabidopsis
29 pollen where 21-nt siRNAs associate with derepressed transposable elements (TEs) in
30 the sperm companion cell, the vegetative cell (Slotkin et al., 2009) (VC; Figure 9).
31 These siRNAs, but not the TE transcripts, accumulate in the sperm cell, suggesting that
32 VC siRNAs can move into the sperm to reinforce TE silencing (Figure 9). Such TE
33 reactivation (and hence associated siRNAs) is likely confined to gamete companion
34 cells, as it is largely driven by a DNA demethylase, DEMETER (DME) (He et al., 2019),
35 whose encoding gene is specifically expressed in companion cells (Feng et al., 2013).
36 Indeed, TEs that are demethylated by DME in the VC are hypermethylated in sperm
37 (where *DME* is not expressed) in a DME-dependent manner (Ibarra et al., 2012). *DME*
38 is also expressed in the egg companion cell, the central cell, and likely activates siRNAs
39 moving into the egg (Ibarra et al., 2012; Feng et al., 2013) (Figure 9).

1 Since the above-mentioned study by Slotkin *et al.*, it has become clear that somatic
2 cells surrounding the germline produce distinct populations of siRNAs. An example is a
3 variant form of the small RNA-directed DNA methylation pathway (RdDM) in meiocyte
4 nurse cells (the tapetum; Figure 9). RdDM methylates TEs using 24-nt siRNAs
5 transcribed by RNA Polymerase IV (Pol IV), which is recruited by putative chromatin
6 remodelers, CLASSY1-4 (CLSY1-4). Somatic tissues mainly express *CLSY1* and
7 *CLSY2*, and their proteins recruit RdDM to thousands of repeats. In tapetal cells and
8 ovules, *CLSY3* is expressed at much higher levels than *CLSY1/2*, leading to a distinct
9 24-nt siRNA profile with the vast majority of siRNAs coming from a few hundred loci
10 (Long *et al.*, 2021; Zhou *et al.*, 2022b).

11 Although these germline siRNAs were discovered due to their roles in TE silencing,
12 increasing evidence links them to gene regulation, for example during pollen
13 development in *Capsella* (Wang *et al.*, 2020). 24-nt siRNAs produced by tapetal cells
14 methylate genes with similar but not identical sequences in male meiocytes (Walker *et al.*,
15 2018; Long *et al.*, 2021) (Figure 9), thereby regulating the splicing of a meiotic gene
16 and facilitating meiosis (Walker *et al.*, 2018; Long *et al.*, 2021). As the TE-silencing and
17 gene regulatory functions of germline siRNAs go hand in hand, it is tantalizing but
18 difficult to tease apart which is the primary function, if such a distinction is possible.

19 Compounding the complexity, germline siRNA biogenesis varies among plant species.
20 Although *Arabidopsis* meiotic 24-nt siRNAs are produced by Pol IV and RdDM, similarly
21 abundant 24-nt phased secondary siRNAs (phasiRNAs) in maize and rice tapetal cells
22 are produced by cleavage of noncoding Pol II transcripts by a miRNA (Liu *et al.*, 2020).
23 Monocot anther wall cells also accumulate an earlier wave of 21-nt phasiRNAs. Both
24 21-nt and 24-nt phasiRNAs have been proposed to move into meiotic cells and are
25 important for male fertility, especially under certain environmental conditions, although it
26 is still unclear why (Liu *et al.*, 2020; Zhou *et al.*, 2022c).

27 Another challenge is to elucidate the link between siRNA-mediated gene regulation and
28 germ cell differentiation. The most well-understood example is the differentiation of
29 female meiocytes, called megaspore mother cells (MMCs). Normally, only one
30 subepidermal (L2) cell adopts MMC fate and undergoes meiosis in each ovule (Figure
31 9). Multiple MMCs differentiate in mutants of RdDM or 21-22 nt *trans*-acting siRNA
32 (tasiRNA) pathways (Olmedo-Monfil *et al.*, 2010; Su *et al.*, 2020). Key components of
33 both pathways are specifically expressed in apical epidermal (L1) cells, suggesting that
34 these L1-produced siRNAs are essential for suppressing MMC fate in L2 cells (Figure
35 9). Importantly, causal links were made between L1-produced tasiRNAs, the repression
36 of *AUXIN RESPONSE FACTOR 3* (*ARF3*) in L2 cells, and the suppression of MMC fate
37 (Su *et al.*, 2017; Su *et al.*, 2020). However, this is unlikely the sole regulatory
38 mechanism for MMC differentiation, as mutations of other epigenetic pathways, such as
39 METHYLTRANSFERASE 1 (*MET1*)-mediated DNA methylation maintenance (Li *et al.*,
40 2017), also cause a similar supernumerary MMC phenotype. An indirect mechanism is

1 also plausible, e.g. failure of epigenetic silencing interferes with MMC meiosis or
2 function, which activates neighboring cells to adopt MMC fate as a compensating
3 mechanism.

4 A converging feature of germline siRNAs is their non-cell-autonomy, which raises the
5 question of why germ cells do not produce the siRNAs themselves, but instead rely on
6 neighboring companion/nurse cells. Many ideas have arisen: perhaps siRNA
7 biosynthesis exposes certain risks as it generally involves transcription of TEs, or nurse
8 cells might afford to sensitize their chromatin environment to unfurl their genome and
9 reveal potentially hazardous TEs (Feng et al., 2013), or maybe it is a question of why
10 not, as nurse cells are already geared to provide a wide range of nutrients and other
11 molecules to germ cells. These are exciting concepts ripe for exploration.

12 For Arabidopsis tapetal siRNAs, non-cell-autonomy may allow more precise control of
13 germline transcriptional regulation. Canonical RdDM is self-reinforcing, as DNA
14 methylation promotes the generation of methylation-inducing siRNAs by recruiting Pol
15 IV. The methylation arm of RdDM is tuned more aggressively in meiocytes to target
16 broader sequences, which allows the targeting of genes and fast-evolving TEs (Long et
17 al., 2021). However, given the self-reinforcing nature of RdDM, this broad-targeting
18 ability needs to be tightly controlled to prevent the long-term establishment of RdDM at
19 inappropriate genomic regions. Such control is achievable by cellular
20 compartmentalization: 24-nt siRNA biogenesis is confined to the tapetum, whereas
21 broad-targeting competence is restricted to male meiocytes (Long et al., 2021).

22 Understanding how germline siRNAs move between cells remains technically
23 challenging. Plasmodesmata provide symplastic connections between daughter cells
24 and are known to prevail in several scenarios of germline siRNA movement (Liu et al.,
25 2020; Long et al., 2021). However, in which form(s) and how does the silencing signal
26 move (Chen and Rechavi, 2022)? Furthermore, one cannot exclude the possibility of an
27 apoplasmic transport mechanism (reviewed before in the context of cross-kingdom
28 RNAi), warranting further investigation.

29 Finally, germline siRNAs undoubtedly have functions beyond those in germ cells.
30 siRNAs in sperm can act as quantitative measures of paternal genome dosage, whose
31 imbalance with maternal dosage causes seed abortion (Wang et al., 2018a). Similarly,
32 encountering gamete siRNAs in the zygote could, in theory, assess the compatibility of
33 parental genomes, leading to hybridization barriers (Bourc'his and Voinnet, 2010).
34 Although debated, endosperm siRNAs have also been proposed to move into the
35 embryo, where they may exert a transgenerational effect (Bourc'his and Voinnet, 2010).
36 siRNA pathways are known to be environmentally sensitive and malleable. Thus,
37 germline siRNAs might be inherited by the next generation to facilitate memory of the
38 environment and regulate the development of the offspring accordingly. The
39 transgenerational effect of siRNAs (if any) remains an exciting area for future
40 investigation.

Acknowledgements

My apologies to colleagues whose work could not be cited due to length restrictions.

Funding

Research in the Feng lab is supported by the Biotechnology and Biological Sciences Research Council (BBS0096201 and BBP0135111) and the European Research Council ('SexMeth' 804981).

The roles of small RNAs in the regulation of agronomic traits of crops

(Written by Yijun Qi)

Our knowledge of the biogenesis, action mode and biological roles of small RNAs has mostly been obtained from studies in Arabidopsis. However, findings in Arabidopsis cannot always be reasonably extrapolated to crops. Studies in crops, despite still being limited, have revealed that small RNAs play unexpected roles, particularly in the regulation of traits of agronomic significance.

Dozens of miRNAs have been shown to regulate crop development, metabolism and stress responses. For instance, miR156, one of the most conserved miRNAs among plant species, regulates juvenile to adult transition in Arabidopsis (Wang et al., 2009; Wu et al., 2009a). However, in rice, miR156 not only helps shape plant architecture but also regulates grain development and filling (Jiao et al., 2010). The conserved miR396, which in Arabidopsis regulates plant development, targets and regulates the transcription factor gene *HaWRKY6* in sunflower (*Helianthus annuus*) during heat response (Giacomelli et al., 2012). There are many species-specific miRNAs in crops. For example, miR528, a monocot-specific miRNA, targets a number of genes involved in a variety of developmental processes or biotic and abiotic stress responses (Chen et al., 2019). How conserved miRNAs gain more regulatory functions and how species-specific miRNAs have been acquired by certain crops remain to be fully elucidated. Dissection of diversified roles of miRNAs in crops will greatly improve our understanding of the range of miRNA-mediated regulation.

In addition to canonical 21-nt miRNAs, there is a distinct class of 24-nt long miRNAs, referred to as lmiRNAs, in rice (Wu et al., 2009b). lmiRNAs regulate transcription via directing DNA methylation at target sites (Wu et al., 2010). It remains unclear how prevalent lmiRNAs are among crops. lmiRNAs that have been functionally characterized were all found to regulate rice biotic stress responses (Jiang et al., 2020a; Zhou et al., 2020; Campo et al., 2021). This result raises the question as to whether lmiRNAs evolved for plant stress responses and adaptation to environmental changes. Systematic identification of lmiRNAs and their target genes in different crops will be necessary for a better understanding of lmiRNA evolution and function.

1 Twenty-four-nt siRNAs are produced mainly from TEs and direct DNA methylation at
2 target loci through RdDM. While Arabidopsis mutants lacking RdDM do not show
3 obvious phenotypes, rice RdDM mutants have pleiotropic alterations, including
4 dwarfism, an increase in rice tillering and a reduction in rice panicle size (Wei et al.,
5 2014; Xu et al., 2020a). In maize, loss of 24-nt siRNAs leads to dwarfism, altered leaf
6 polarity and development of feminized tassels (Alleman et al., 2006). These findings
7 indicate that 24-nt siRNAs are important regulators of agronomic traits in crops. The
8 more prevailing regulatory role of 24-nt siRNAs in rice and maize could be explained by
9 the fact that TEs are very abundant and dispersed in euchromatic regions in these
10 plants, which greatly increases the likelihood that RdDM at TEs regulates nearby genes.
11 Indeed, increased tillering in rice RdDM mutants is attributed to loss of RdDM at
12 miniature inverted-repeat transposable elements (MITEs) near *MIR156d/j* and *D14*,
13 which control rice tillering (Xu et al., 2020a). Interestingly, it has recently been shown
14 that 24-nt siRNA can direct DNA methylation at imperfectly matched targets in
15 Arabidopsis and cabbage (*Brassica rapa*) (Fei et al., 2021; Long et al., 2021; Burgess et
16 al., 2022), which may greatly increase the range and complexity of RdDM-mediated
17 gene regulation. For most 24-nt siRNAs, their tissue-specific expression, their targets,
18 and the effects of their loss remain unknown.

19 PhasiRNAs, secondary siRNAs that are produced following miRNA-directed target
20 mRNA cleavage, can be 21 or 24 nt in length, depending on the miRNA trigger.
21 PhasiRNAs are the predominant type of small RNAs in anthers in monocots, suggesting
22 that they play a pivotal role in crop reproduction. Supporting this notion, loss of 21-nt
23 phasiRNAs, or their activity, in rice leads to pollen sterility (Jiang et al., 2020b), and
24 overproduction of 21-nt phasiRNAs at the *Pms1* locus results in photoperiod-sensitive
25 male sterility, which allows the establishment of a two-line system for hybrid rice
26 breeding (Fan et al., 2016). 21-nt phasiRNAs were found to facilitate the progression of
27 meiosis by directing target mRNA cleavage (Jiang et al., 2020b; Zhang et al., 2020). As
28 these targets are regulated for successful meiosis, investigation of their functions could
29 be a shortcut to discovering genes and mechanisms important for crop reproduction.
30 Loss of 24-nt phasiRNAs causes reduced pollen fertility and seed-setting rate in rice
31 and temperature-sensitive male sterility in maize. There is some evidence supporting
32 the idea that 24-nt phasiRNAs direct DNA methylation in *cis* (Zhang et al., 2021).
33 Whether they can direct DNA methylation in *trans* and whether DNA methylation, if
34 established, can be passed to next generation and regulates grain development remain
35 to be explored.

36 tRNA-derived small RNAs (tsRNAs) and rRNA-derived small RNAs (rsRNAs) are two
37 classes of small RNAs that have recently been identified. Whereas we still have limited
38 information about the expression profile, modes of action, and biological roles of
39 rsRNAs in plants, tsRNAs have been profiled in Arabidopsis (Ma et al., 2021). tsRNA
40 levels appear to undergo dynamic changes in response to abiotic and biotic stresses. A

1 19-nt 5' tsRNA produced from tRNA-Ala regulates anti-fungal defense in Arabidopsis
2 (Gu et al., 2022). tsRNAs have not been well characterized in crops and their functions
3 remain to be revealed. It will be also interesting to investigate whether they are widely
4 involved in stress responses in crops.

5 Because many agronomic traits are controlled by small RNAs, manipulation of small
6 RNA-mediated gene regulation has emerged as an important strategy to achieving
7 desired agronomic traits. Unlike overexpressing or knocking out a gene, manipulation of
8 small RNA activity allows us to fine-tune or precisely control the expression of a gene.
9 Such changes in gene expression can be more physiologically relevant and may
10 overcome side effects induced by all-or-nothing approaches. Thus, this offers a great
11 new strategy to improve agronomic traits in crops.

12 **Acknowledgments**

13 My apology to colleagues whose work could not be cited due to length restrictions
14

15 **Funding**

16 Research in the Qi lab is supported by National Natural Science Foundation of China
17 (Grant No. 31788103).
18

19 **Open questions in the study of RNA-directed DNA methylation**

20 (Written by Craig S. Pikaard)

21 Eukaryotic cells protect themselves against TEs, viruses and other selfish genetic
22 elements using RNAi pathways dependent on siRNAs. In plants, siRNAs range in size
23 from 21-24 nt and mediate both post-transcriptional gene silencing (PTGS) and
24 transcriptional gene silencing. RdDM is an important aspect of transcriptional gene
25 silencing, involving siRNAs to bring about cytosine methylation of complementary DNA
26 sequences (Erdmann and Picard, 2020). Chemical modifications of histone proteins
27 also occur, in crosstalk with DNA methylation (Law and Jacobsen, 2010). Collectively,
28 DNA and histone modifications result in chromatin environments that suppress
29 promoter-dependent gene activation, but exactly how is not clear.

30 Most of what we know about RNA-dependent silencing in plants comes from studies of
31 Arabidopsis. At least two pathways contribute to RdDM: an initiation pathway that acts
32 on transcriptionally active transposons or invading viruses and a maintenance pathway
33 that perpetuates cytosine methylation at thousands of transposon loci throughout the
34 genome (Figure 10). The establishment pathway overlaps with the pathway for PTGS
35 (Nuthikattu et al., 2013) and begins with transposon, virus or transgene transcripts that
36 are somehow recognized as being different from other cellular RNAs (Hung and Slotkin,
37 2021), triggering their conversion into double-stranded RNA (dsRNA) by RNA-
38 DEPENDENT RNA POLYMERASE 6 (RDR6). The dsRNAs are then cut (diced) into 21-
39
40

1 or 22-nt siRNAs by the Dicer-like endonucleases DCL4 or DCL2 and loaded into an
2 Argonaute family protein, primarily AGO1 or AGO6 (Ariel and Manavella, 2021). siRNA-
3 AGO1 complexes can bind complementary target mRNAs to cause their destruction or
4 interfere with their translation, thus achieving PTGS. In parallel, 21-22-nt siRNAs bound
5 to AGO6 guide low-level cytosine methylation at complementary DNA sequences in
6 partnership with multisubunit RNA Polymerase V (Pol V) and the DNA
7 methyltransferase DOMAINS REARRANGED METHYLTRANSFERASE 2 (DRM2).
8 Low-level methylation is not sufficient for transcriptional gene silencing but serves as a
9 signal to recruit the machinery of the maintenance pathway, which accounts for the vast
10 majority of RdDM activity (Figure 10). This pathway involves RNA Polymerase IV (Pol
11 IV), RNA-DEPENDENT RNA POLYMERASE 2 (RDR2), DCL3, 24-nt siRNAs, AGO4,
12 Pol V, DRM2 and numerous helper activities implicated in Pol IV or Pol V recruitment or
13 chromatin modification and is dependent on 24-nt siRNAs (Figure 10).
14 The biogenesis of 24-nt siRNAs is understood in some detail, having been recapitulated
15 *in vitro* (Singh et al., 2019) using purified enzymes whose structures have recently been
16 resolved (Fukudome et al., 2021; Huang et al., 2021a; Wang et al., 2021a), yet
17 questions still remain. Pol IV acts first in the pathway, presumably initiating RNA
18 biosynthesis within the context of a melted DNA transcription bubble, as is the case for
19 other DNA-dependent RNA polymerases. However, Pol IV is unable to sustain
20 transcriptional elongation over more than ~12-16 nt into the double-stranded DNA
21 beyond the initiation bubble (Singh et al., 2019), for reasons that are not yet clear. This
22 behavior causes the polymerase to stall and then retreat, sliding backward along the
23 DNA template as the template and non-template strands reanneal (Fukudome et al.,
24 2021; Huang et al., 2021a), a phenomenon known as polymerase backtracking. As Pol
25 IV backtracks, the 3' end of its short (~30 nt) transcript becomes unpaired from the
26 template DNA strand and is extruded and becomes engaged by RDR2 (Huang et al.,
27 2021a), which uses the RNA as a template and initiates transcription 1-2 nt internal to
28 its 3' end (Fukudome et al., 2021). Whether the physical interactions of RDR2 with
29 specific Pol IV subunits stimulates Pol IV backtracking and disfavors Pol IV elongation
30 remains unclear, but is testable. Upon completing transcription of the Pol IV strand to
31 generate a dsRNA, RDR2 has an intrinsic terminal transferase activity that adds an
32 extra untemplated nucleotide to the 3' end of its transcript, and then RDR2 releases the
33 resulting dsRNA (Singh et al., 2019). Due to initiation by RDR2 internal to the 3' end of
34 the Pol IV transcript and its addition of an untemplated nucleotide to the 3' end of its
35 transcript, the resulting dsRNA has 3' overhangs of 1-2 nt at each end. These
36 overhangs, together with 5' nucleotide preferences, program alternative DCL3 dicing
37 reactions from either end of the dsRNAs, yielding siRNA duplexes that consist of a 24-nt
38 strand paired with a 23-nt strand or a pair of 24-nt strands (Loffer et al., 2022) (Figure
39 10). In the case of 24/23 duplexes, the 23-nt RNAs serve as so-called passenger
40 strands that help specify that the paired 24-nt strands are loaded into AGO4 to serve as

1 guide strands (Wang et al., 2022). The passenger strand is then sliced by AGO and
2 partially released. It is not clear how, or why, 24-nt siRNAs are specifically loaded as
3 guide strands given that 21-, 22-, 23- or 24-nt RNAs can be loaded into recombinant
4 AGO4 and guide slicing of target RNAs with similar efficiency (Wang et al., 2022). One
5 speculation is that a dsRNA-binding chaperone activity that can discriminate between 3'
6 overhangs of 1 or 2 nt orients the siRNA duplex such that the strand with the 2-nt
7 overhang is loaded into AGO4 as the guide strand. In the case of asymmetric 24/23
8 duplexes, the 24-nt strand would be oriented to become the guide whereas for
9 symmetrical 24/24 duplexes, with 2-nt overhangs at each end, guide strand choice
10 would presumably be random. Experiments are needed to test this hypothesis.

11 What happens following AGO4-siRNA loading is not clear. Early studies showed that
12 AGO4 localization at RdDM loci is dependent on Pol V transcription, that AGO4 can be
13 chemically crosslinked to Pol V transcripts (Wierzbicki et al., 2009) and that cytosine
14 methylation occurs where siRNAs overlap sites of Pol V occupancy (Wierzbicki et al.,
15 2012). Other studies have revealed that AGO4 can bind the C-terminal domain (CTD) of
16 the Pol V largest subunit and/or the Pol V-associated protein, SPT5L (Suppressor of Ty
17 insertion 5-like) (El-Shami et al., 2007; Bies-Etheve et al., 2009). Thus, AGO4-RNA and
18 AGO4-protein interactions are both likely to be important, but whether they occur
19 simultaneously or sequentially is unknown. And how does DNA methylation, and/or the
20 histone modifications that correlate with DNA methylation, ensue from these siRNA-
21 AGO4-Pol V interactions? There is co-immunoprecipitation evidence that DRM2 and
22 AGO4 can directly interact (Zhong et al., 2014), but RdDM has not yet been achieved *in*
23 *vitro*. Biochemical and structural studies that could reveal the spatial positions of the
24 proteins, RNAs and DNA strands when RdDM occurs would be break-through studies
25 for the field.

26 Other major unanswered questions pertain to how Pol IV and Pol V transcription is
27 initiated. Bacterial and archaeal multisubunit RNA polymerases, as well as eukaryotic
28 RNA polymerases I, II and III require DNA-binding transcription factors that recruit the
29 polymerase to promoters, melt the DNA in the vicinity of the start site and position the
30 polymerase to initiate transcription of one of the two DNA strands. However,
31 conventional transcription factors and promoters have not been implicated in Pol IV or
32 Pol V transcription. Instead, the evidence suggests that pre-existing chromatin
33 modifications serve as recruitment signals, with cytosine methylation in the CG context,
34 requiring MET1 and HISTONE DEACETYLASE 6 (HDA6) (Blevins et al., 2014), methyl
35 cytosine binding by SUPPRESSOR OF VARIATION 3-9 HOMOLOG PROTEIN 2/9
36 (SUVH2/9), or binding of methylated histone H3 lysine 9 (H3K9) by SAWADEE
37 HOMEODOMAIN HOMOLOG 1 (SHH1) implicated in Pol IV and/or Pol V recruitment
38 (Figure 10) (Erdmann and Picard, 2020). ATP-dependent DNA translocases are also
39 implicated, including the CLSY protein family in the case of Pol IV and DEFECTIVE IN
40 RNA-DIRECTED DNA METHYLATION 1 (DRD1) in the case of Pol V. However, there

1 is currently no biochemical evidence to suggest how promoter-independent DNA
2 melting, polymerase positioning or transcription initiation occurs for Pol IV or Pol V.
3 Once again, *in vitro* experiments with purified components will be needed to move from
4 knowing the list of proteins involved to knowing what they do and how they work.
5

6 7 REFERENCES

8 **Alleman, M., Sidorenko, L., McGinnis, K., Seshadri, V., Dorweiler, J.E., White, J.,**
9 **Sikkink, K., and Chandler, V.L.** (2006). An RNA-dependent RNA polymerase is
10 required for paramutation in maize. *Nature* **442**, 295-298.

11 **Anderson, S.J., Kramer, M.C., Gosai, S.J., Yu, X., Vandivier, L.E., Nelson, A.D.L.,**
12 **Anderson, Z.D., Beilstein, M.A., Fray, R.G., Lyons, E., and Gregory, B.D.** (2018).
13 N(6)-Methyladenosine Inhibits Local Ribonucleolytic Cleavage to Stabilize mRNAs in
14 Arabidopsis. *Cell Rep* **25**, 1146-1157 e1143.

15 **Arae, T., Isai, S., Sakai, A., Mineta, K., Yokota Hirai, M., Suzuki, Y., Kanaya, S.,**
16 **Yamaguchi, J., Naito, S., and Chiba, Y.** (2017). Co-ordinated Regulations of mRNA
17 Synthesis and Decay during Cold Acclimation in Arabidopsis Cells. *Plant Cell Physiol*
18 **58**, 1090-1102.

19 **Ariel, F., Lucero, L., Christ, A., Mammarella, M.F., Jegu, T., Veluchamy, A.,**
20 **Mariappan, K., Latrasse, D., Blein, T., Liu, C., Benhamed, M., and Crespi, M.**
21 (2020). R-Loop Mediated trans Action of the APOLO Long Noncoding RNA. *Mol Cell* **77**,
22 1055-1065 e1054.

23 **Ariel, F.D., and Manavella, P.A.** (2021). When junk DNA turns functional: transposon-
24 derived non-coding RNAs in plants. *J Exp Bot* **72**, 4132-4143.

25 **Arribas-Hernandez, L., Bressendorff, S., Hansen, M.H., Poulsen, C., Erdmann, S.,**
26 **and Brodersen, P.** (2018). An m(6)A-YTH Module Controls Developmental Timing and
27 Morphogenesis in Arabidopsis. *Plant Cell* **30**, 952-967.

28 **Arribas-Hernandez, L., Rennie, S., Koster, T., Porcelli, C., Lewinski, M., Staiger, D.,**
29 **Andersson, R., and Brodersen, P.** (2021a). Principles of mRNA targeting via the
30 Arabidopsis m(6)A-binding protein ECT2. *Elife* **10**.

31 **Arribas-Hernandez, L., Rennie, S., Schon, M., Porcelli, C., Enugutti, B.,**
32 **Andersson, R., Nodine, M.D., and Brodersen, P.** (2021b). The YTHDF proteins ECT2
33 and ECT3 bind largely overlapping target sets and influence target mRNA abundance,
34 not alternative polyadenylation. *Elife* **10**.

35 **Arsovski, A.A., Galstyan, A., Guseman, J.M., and Nemhauser, J.L.** (2012).
36 Photomorphogenesis. *Arabidopsis Book* **10**, e0147.

37 **Bazin, J., Baerenfaller, K., Gosai, S.J., Gregory, B.D., Crespi, M., and Bailey-**
38 **Serres, J.** (2017). Global analysis of ribosome-associated noncoding RNAs unveils new
39 modes of translational regulation. *Proc Natl Acad Sci U S A* **114**, E10018-E10027.

- 1 **Bhat, S.S., Bielewicz, D., Gulanicz, T., Bodi, Z., Yu, X., Anderson, S.J., Szewc, L.,**
2 **Bajczyk, M., Dolata, J., Grzelak, N., Smolinski, D.J., Gregory, B.D., Fray, R.G.,**
3 **Jarmolowski, A., and Szweykowska-Kulinska, Z.** (2020). mRNA adenosine
4 methylase (MTA) deposits m(6)A on pri-miRNAs to modulate miRNA biogenesis in
5 *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A* **117**, 21785-21795.
- 6 **Bies-Etheve, N., Pontier, D., Lahmy, S., Picart, C., Vega, D., Cooke, R., and**
7 **Lagrange, T.** (2009). RNA-directed DNA methylation requires an AGO4-interacting
8 member of the SPT5 elongation factor family. *EMBO Rep* **10**, 649-654.
- 9 **Blevins, T., Pontvianne, F., Cocklin, R., Podicheti, R., Chandrasekhara, C.,**
10 **Yerneni, S., Braun, C., Lee, B., Rusch, D., Mockaitis, K., Tang, H., and Pikaard,**
11 **C.S.** (2014). A two-step process for epigenetic inheritance in *Arabidopsis*. *Mol Cell* **54**,
12 30-42.
- 13 **Boccaletto, P., Machnicka, M.A., Purta, E., Piatkowski, P., Baginski, B., Wirecki,**
14 **T.K., de Crecy-Lagard, V., Ross, R., Limbach, P.A., Kotter, A., Helm, M., and**
15 **Bujnicki, J.M.** (2018). MODOMICS: a database of RNA modification pathways. 2017
16 update. *Nucleic Acids Res* **46**, D303-D307.
- 17 **Bologna, N.G., Schapire, A.L., and Palatnik, J.F.** (2013). Processing of plant
18 microRNA precursors. *Brief Funct Genomics* **12**, 37-45.
- 19 **Bourc'his, D., and Voinnet, O.** (2010). A small-RNA perspective on gametogenesis,
20 fertilization, and early zygotic development. *Science* **330**, 617-622.
- 21 **Brickner, J.R., Garzon, J.L., and Cimprich, K.A.** (2022). Walking a tightrope: The
22 complex balancing act of R-loops in genome stability. *Molecular cell*.
- 23 **Brioudes, F., Jay, F., Sarazin, A., Grentzinger, T., Devers, E.A., and Voinnet, O.**
24 (2021). HASTY, the *Arabidopsis* EXPORTIN5 ortholog, regulates cell-to-cell and
25 vascular microRNA movement. *EMBO J* **40**, e107455.
- 26 **Bryll, A.R., and Peterson, C.L.** (2022). Functional interaction between the RNA
27 exosome and the sirtuin deacetylase Hst3 maintains transcriptional homeostasis.
28 *Genes Dev* **36**, 17-22.
- 29 **Bugnon, L.A., Edera, A.A., Prochetto, S., Gerard, M., Raad, J., Fenoy, E., Rubiolo,**
30 **M., Chorostecki, U., Gabaldon, T., Ariel, F., Di Persia, L.E., Milone, D.H., and**
31 **Stegmayer, G.** (2022). Secondary structure prediction of long noncoding RNA: review
32 and experimental comparison of existing approaches. *Brief Bioinform* **23**.
- 33 **Burgess, D., Chow, H.T., Grover, J.W., Freeling, M., and Mosher, R.A.** (2022). Ovule
34 siRNAs methylate protein-coding genes in trans. *Plant Cell* **34**, 3647-3664.
- 35 **Burjoski, V., and Reddy, A.S.N.** (2021). The Landscape of RNA-Protein Interactions in
36 Plants: Approaches and Current Status. *Int J Mol Sci* **22**.
- 37 **Cai, Q., He, B., and Jin, H.** (2019). A safe ride in extracellular vesicles - small RNA
38 trafficking between plant hosts and pathogens. *Curr Opin Plant Biol* **52**, 140-148.

- 1 **Cai, Q., Qiao, L., Wang, M., He, B., Lin, F.M., Palmquist, J., Huang, S.D., and Jin, H.**
2 (2018). Plants send small RNAs in extracellular vesicles to fungal pathogen to silence
3 virulence genes. *Science* **360**, 1126-1129.
- 4 **Cai, Q., He, B., Wang, S., Fletcher, S., Niu, D., Mitter, N., Birch, P.R.J., and Jin, H.**
5 (2021). Message in a Bubble: Shuttling Small RNAs and Proteins Between Cells and
6 Interacting Organisms Using Extracellular Vesicles. *Annu Rev Plant Biol* **72**, 497-524.
- 7 **Cambiagno, D.A., Giudicatti, A.J., Arce, A.L., Gagliardi, D., Li, L., Yuan, W.,**
8 **Lundberg, D.S., Weigel, D., and Manavella, P.A.** (2021). HASTY modulates miRNA
9 biogenesis by linking pri-miRNA transcription and processing. *Mol Plant* **14**, 426-439.
- 10 **Campo, S., Sanchez-Sanuy, F., Camargo-Ramirez, R., Gomez-Ariza, J., Baldrich,**
11 **P., Campos-Soriano, L., Soto-Suarez, M., and San Segundo, B.** (2021). A novel
12 Transposable element-derived microRNA participates in plant immunity to rice blast
13 disease. *Plant Biotechnol J* **19**, 1798-1811.
- 14 **Chantarachot, T., and Bailey-Serres, J.** (2018). Polysomes, Stress Granules, and
15 Processing Bodies: A Dynamic Triumvirate Controlling Cytoplasmic mRNA Fate and
16 Function. *Plant Physiol* **176**, 254-269.
- 17 **Chen, C., Liu, Y., and Xia, R.** (2019). Jack of Many Trades: The Multifaceted Role of
18 miR528 in Monocots. *Mol Plant* **12**, 1044-1046.
- 19 **Chen, X., and Rechavi, O.** (2022). Plant and animal small RNA communications
20 between cells and organisms. *Nat Rev Mol Cell Biol* **23**, 185-203.
- 21 **Cheng, L., Wang, W., Yao, Y., and Sun, Q.** (2021). Mitochondrial RNase H1 activity
22 regulates R-loop homeostasis to maintain genome integrity and enable early
23 embryogenesis in Arabidopsis. *PLoS Biol* **19**, e3001357.
- 24 **Cho, H., Cho, H.S., Nam, H., Jo, H., Yoon, J., Park, C., Dang, T.V.T., Kim, E., Jeong,**
25 **J., Park, S., Wallner, E.S., Youn, H., Park, J., Jeon, J., Ryu, H., Greb, T., Choi, K.,**
26 **Lee, Y., Jang, S.K., Ban, C., and Hwang, I.** (2018). Translational control of phloem
27 development by RNA G-quadruplex-JULGI determines plant sink strength. *Nat Plants* **4**,
28 376-390.
- 29 **Chung, B.Y.W., Balcerowicz, M., Di Antonio, M., Jaeger, K.E., Geng, F., Franaszek,**
30 **K., Marriott, P., Brierley, I., Firth, A.E., and Wigge, P.A.** (2020). An RNA
31 thermoswitch regulates daytime growth in Arabidopsis. *Nat Plants* **6**, 522-532.
- 32 **Costa, S.** (2016). Cell identity: a matter of lineage and neighbours. *New Phytol* **210**,
33 1155-1158.
- 34 **Deng, H., Cheema, J., Zhang, H., Woolfenden, H., Norris, M., Liu, Z., Liu, Q., Yang,**
35 **X., Yang, M., Deng, X., Cao, X., and Ding, Y.** (2018). Rice In Vivo RNA Structurome
36 Reveals RNA Secondary Structure Conservation and Divergence in Plants. *Mol Plant*
37 **11**, 607-622.
- 38 **Devers, E.A., Brosnan, C.A., Sarazin, A., Albertini, D., Amsler, A.C., Brioudes, F.,**
39 **Jullien, P.E., Lim, P., Schott, G., and Voinnet, O.** (2020). Movement and differential

1 consumption of short interfering RNA duplexes underlie mobile RNA interference. *Nat*
2 *Plants* **6**, 789-799.

3 **Dolata, J., Guo, Y., Kolowerzo, A., Smolinski, D., Brzyzek, G., Jarmolowski, A., and**
4 **Swiezewski, S.** (2015). NTR1 is required for transcription elongation checkpoints at
5 alternative exons in Arabidopsis. *EMBO J* **34**, 544-558.

6 **Duan, H.C., Wei, L.H., Zhang, C., Wang, Y., Chen, L., Lu, Z., Chen, P.R., He, C., and**
7 **Jia, G.** (2017). ALKBH10B Is an RNA N(6)-Methyladenosine Demethylase Affecting
8 Arabidopsis Floral Transition. *Plant Cell* **29**, 2995-3011.

9 **Duncan, S., Olsson, T.S.G., Hartley, M., Dean, C., and Rosa, S.** (2017). Single
10 Molecule RNA FISH in Arabidopsis Root Cells. *Bio Protoc* **7**, e2240.

11 **Dunker, F., Trutzenberg, A., Rothenpieler, J.S., Kuhn, S., Prols, R., Schreiber, T.,**
12 **Tissier, A., Kemen, A., Kemen, E., Huckelhoven, R., and Weiberg, A.** (2020).
13 Oomycete small RNAs bind to the plant RNA-induced silencing complex for virulence.
14 *Elife* **9**.

15 **El-Shami, M., Pontier, D., Lahmy, S., Braun, L., Picart, C., Vega, D., Hakimi, M.A.,**
16 **Jacobsen, S.E., Cooke, R., and Lagrange, T.** (2007). Reiterated WG/GW motifs form
17 functionally and evolutionarily conserved ARGONAUTE-binding platforms in RNAi-
18 related components. *Genes Dev* **21**, 2539-2544.

19 **Emenecker, R.J., Holehouse, A.S., and Strader, L.C.** (2020). Emerging Roles for
20 Phase Separation in Plants. *Dev Cell* **55**, 69-83.

21 **Erdmann, R.M., and Picard, C.L.** (2020). RNA-directed DNA Methylation. *PLoS Genet*
22 **16**, e1009034.

23 **Fan, L., Zhang, C., Gao, B., Zhang, Y., Stewart, E., Jez, J., Nakajima, K., and Chen,**
24 **X.** (2022). Microtubules promote the non-cell autonomous action of microRNAs by
25 inhibiting their cytoplasmic loading onto ARGONAUTE1 in Arabidopsis. *Dev Cell* **57**,
26 995-1008 e1005.

27 **Fan, Y., Yang, J., Mathioni, S.M., Yu, J., Shen, J., Yang, X., Wang, L., Zhang, Q.,**
28 **Cai, Z., Xu, C., Li, X., Xiao, J., Meyers, B.C., and Zhang, Q.** (2016). PMS1T,
29 producing phased small-interfering RNAs, regulates photoperiod-sensitive male sterility
30 in rice. *Proc Natl Acad Sci U S A* **113**, 15144-15149.

31 **Fang, X., Cui, Y., Li, Y., and Qi, Y.** (2015). Transcription and processing of primary
32 microRNAs are coupled by Elongator complex in Arabidopsis. *Nat Plants* **1**, 15075.

33 **Fang, Y., and Spector, D.L.** (2007). Identification of nuclear dicing bodies containing
34 proteins for microRNA biogenesis in living Arabidopsis plants. *Curr Biol* **17**, 818-823.

35 **Faucillion, M.L., Johansson, A.M., and Larsson, J.** (2022). Modulation of RNA
36 stability regulates gene expression in two opposite ways: through buffering of RNA
37 levels upon global perturbations and by supporting adapted differential expression.
38 *Nucleic Acids Res* **50**, 4372-4388.

39 **Fei, Y., Nyiko, T., and Molnar, A.** (2021). Non-perfectly matching small RNAs can
40 induce stable and heritable epigenetic modifications and can be used as molecular

1 markers to trace the origin and fate of silencing RNAs. *Nucleic Acids Res* **49**, 1900-
2 1913.

3 **Feng, X., Zilberman, D., and Dickinson, H.** (2013). A conversation across
4 generations: soma-germ cell crosstalk in plants. *Dev Cell* **24**, 215-225.

5 **Foley, S.W., Gosai, S.J., Wang, D., Selamoglu, N., Sollitti, A.C., Koster, T., Steffen,**
6 **A., Lyons, E., Daldal, F., Garcia, B.A., Staiger, D., Deal, R.B., and Gregory, B.D.**
7 (2017). A Global View of RNA-Protein Interactions Identifies Post-transcriptional
8 Regulators of Root Hair Cell Fate. *Dev Cell* **41**, 204-220 e205.

9 **Fonouni-Farde, C., Ariel, F., and Crespi, M.** (2021). Plant Long Noncoding RNAs:
10 New Players in the Field of Post-Transcriptional Regulations. *Noncoding RNA* **7**.

11 **Franco-Zorrilla, J.M., Valli, A., Todesco, M., Mateos, I., Puga, M.I., Rubio-Somoza,**
12 **I., Leyva, A., Weigel, D., Garcia, J.A., and Paz-Ares, J.** (2007). Target mimicry
13 provides a new mechanism for regulation of microRNA activity. *Nat Genet* **39**, 1033-
14 1037.

15 **Fukudome, A., Singh, J., Mishra, V., Reddem, E., Martinez-Marquez, F., Wenzel, S.,**
16 **Yan, R., Shiozaki, M., Yu, Z., Wang, J.C., Takagi, Y., and Pikaard, C.S.** (2021).
17 Structure and RNA template requirements of Arabidopsis RNA-DEPENDENT RNA
18 POLYMERASE 2. *Proc Natl Acad Sci U S A* **118**.

19 **Garcia-Muse, T., and Aguilera, A.** (2019). R Loops: From Physiological to Pathological
20 Roles. *Cell* **179**, 604-618.

21 **Gawronski, P., Enroth, C., Kindgren, P., Marquardt, S., Karpinski, S., Leister, D.,**
22 **Jensen, P.E., Vinther, J., and Scharff, L.B.** (2021). Light-Dependent Translation
23 Change of Arabidopsis psbA Correlates with RNA Structure Alterations at the
24 Translation Initiation Region. *Cells* **10**.

25 **Giacomelli, J.I., Weigel, D., Chan, R.L., and Manavella, P.A.** (2012). Role of recently
26 evolved miRNA regulation of sunflower HaWRKY6 in response to temperature damage.
27 *New Phytol* **195**, 766-773.

28 **Godoy Herz, M.A., Kubaczka, M.G., Brzyzek, G., Servi, L., Krzyszton, M., Simpson,**
29 **C., Brown, J., Swiezewski, S., Petrillo, E., and Kornblihtt, A.R.** (2019). Light
30 Regulates Plant Alternative Splicing through the Control of Transcriptional Elongation.
31 *Mol Cell* **73**, 1066-1074 e1063.

32 **Gonzalo, L., Tossolini, I., Gulanicz, T., Cambiagno, D.A., Kasprowicz-Maluski, A.,**
33 **Smolinski, D.J., Mammarella, M.F., Ariel, F.D., Marquardt, S., Szweykowska-**
34 **Kulinska, Z., Jarmolowski, A., and Manavella, P.A.** (2022). R-loops at microRNA
35 encoding loci promote co-transcriptional processing of pri-miRNAs in plants. *Nat Plants*
36 **8**, 402-418.

37 **Gu, H., Lian, B., Yuan, Y., Kong, C., Li, Y., Liu, C., and Qi, Y.** (2022). A 5' tRNA-Ala-
38 derived small RNA regulates anti-fungal defense in plants. *Sci China Life Sci* **65**, 1-15.

39 **Gutierrez-Beltran, E., Elander, P.H., Dalman, K., Dayhoff, G.W., 2nd, Moschou,**
40 **P.N., Uversky, V.N., Crespo, J.L., and Bozhkov, P.V.** (2021). Tudor staphylococcal

1 nuclease is a docking platform for stress granule components and is essential for
2 SnRK1 activation in Arabidopsis. *EMBO J* **40**, e105043.

3 **Haimovich, G., Medina, D.A., Causse, S.Z., Garber, M., Millan-Zambrano, G.,**
4 **Barkai, O., Chavez, S., Perez-Ortin, J.E., Darzacq, X., and Choder, M.** (2013). Gene
5 expression is circular: factors for mRNA degradation also foster mRNA synthesis. *Cell*
6 **153**, 1000-1011.

7 **Hamada, T., Yako, M., Minegishi, M., Sato, M., Kamei, Y., Yanagawa, Y., Toyooka,**
8 **K., Watanabe, Y., and Hara-Nishimura, I.** (2018). Stress granule formation is induced
9 by a threshold temperature rather than a temperature difference in Arabidopsis. *J Cell*
10 *Sci* **131**.

11 **Hartenian, E., and Glaunsinger, B.A.** (2019). Feedback to the central dogma:
12 cytoplasmic mRNA decay and transcription are interdependent processes. *Crit Rev*
13 *Biochem Mol Biol* **54**, 385-398.

14 **Hawkes, E.J., Hennelly, S.P., Novikova, I.V., Irwin, J.A., Dean, C., and**
15 **Sanbonmatsu, K.Y.** (2016). COOLAIR Antisense RNAs Form Evolutionarily Conserved
16 Elaborate Secondary Structures. *Cell Rep* **16**, 3087-3096.

17 **He, B., Cai, Q., Qiao, L., Huang, C.Y., Wang, S., Miao, W., Ha, T., Wang, Y., and Jin,**
18 **H.** (2021). RNA-binding proteins contribute to small RNA loading in plant extracellular
19 vesicles. *Nat Plants* **7**, 342-352.

20 **He, S.B., Vickers, M., Zhang, J.Y., and Feng, X.Q.** (2019). Natural depletion of histone
21 H1 in sex cells causes DNA demethylation, heterochromatin decondensation and
22 transposon activation. *Elife* **8**, e42530.

23 **Hou, N., Li, C., He, J., Liu, Y., Yu, S., Malnoy, M., Mobeen Tahir, M., Xu, L., Ma, F.,**
24 **and Guan, Q.** (2022). MdMTA-mediated m(6) A modification enhances drought
25 tolerance by promoting mRNA stability and translation efficiency of genes involved in
26 lignin deposition and oxidative stress. *New Phytol* **234**, 1294-1314.

27 **Hou, Y., Sun, J., Wu, B., Gao, Y., Nie, H., Nie, Z., Quan, S., Wang, Y., Cao, X., and**
28 **Li, S.** (2021). CPSF30-L-mediated recognition of mRNA m(6)A modification controls
29 alternative polyadenylation of nitrate signaling-related gene transcripts in Arabidopsis.
30 *Mol Plant* **14**, 688-699.

31 **Hu, J., Manduzio, S., and Kang, H.** (2019). Epitranscriptomic RNA Methylation in Plant
32 Development and Abiotic Stress Responses. *Front Plant Sci* **10**, 500.

33 **Hu, J., Cai, J., Umme, A., Chen, Y., Xu, T., and Kang, H.** (2022). Unique features of
34 mRNA m6A methylomes during expansion of tomato (*Solanum lycopersicum*) fruits.
35 *Plant Physiol* **188**, 2215-2227.

36 **Hu, J., Cai, J., Park, S.J., Lee, K., Li, Y., Chen, Y., Yun, J.Y., Xu, T., and Kang, H.**
37 (2021). N(6) -Methyladenosine mRNA methylation is important for salt stress tolerance
38 in Arabidopsis. *Plant J* **106**, 1759-1775.

39 **Huang, C.Y., Wang, H., Hu, P., Hamby, R., and Jin, H.** (2019). Small RNAs - Big
40 Players in Plant-Microbe Interactions. *Cell Host Microbe* **26**, 173-182.

1 **Huang, K., Wu, X.X., Fang, C.L., Xu, Z.G., Zhang, H.W., Gao, J., Zhou, C.M., You,**
2 **L.L., Gu, Z.X., Mu, W.H., Feng, Y., Wang, J.W., and Zhang, Y.** (2021a). Pol IV and
3 RDR2: A two-RNA-polymerase machine that produces double-stranded RNA. *Science*
4 **374**, 1579-1586.

5 **Huang, Y., Wang, S., Cai, Q., and Jin, H.** (2021b). Effective methods for isolation and
6 purification of extracellular vesicles from plants. *J Integr Plant Biol* **63**, 2020-2030.

7 **Hung, Y.H., and Slotkin, R.K.** (2021). The initiation of RNA interference (RNAi) in
8 plants. *Curr Opin Plant Biol* **61**, 102014.

9 **Ibarra, C.A., Feng, X., Schoft, V.K., Hsieh, T.F., Uzawa, R., Rodrigues, J.A.,**
10 **Zemach, A., Chumak, N., Machlicova, A., Nishimura, T., Rojas, D., Fischer, R.L.,**
11 **Tamaru, H., and Zilberman, D.** (2012). Active DNA demethylation in plant companion
12 cells reinforces transposon methylation in gametes. *Science* **337**, 1360-1364.

13 **Jiang, G., Liu, D., Yin, D., Zhou, Z., Shi, Y., Li, C., Zhu, L., and Zhai, W.** (2020a). A
14 Rice NBS-ARC Gene Conferring Quantitative Resistance to Bacterial Blight Is
15 Regulated by a Pathogen Effector-Inducible miRNA. *Mol Plant* **13**, 1752-1767.

16 **Jiang, P., Lian, B., Liu, C., Fu, Z., Shen, Y., Cheng, Z., and Qi, Y.** (2020b). 21-nt
17 phasiRNAs direct target mRNA cleavage in rice male germ cells. *Nat Commun* **11**,
18 5191.

19 **Jiao, Y., Wang, Y., Xue, D., Wang, J., Yan, M., Liu, G., Dong, G., Zeng, D., Lu, Z.,**
20 **Zhu, X., Qian, Q., and Li, J.** (2010). Regulation of OsSPL14 by OsmiR156 defines
21 ideal plant architecture in rice. *Nat Genet* **42**, 541-544.

22 **John, S., Olas, J.J., and Mueller-Roeber, B.** (2021). Regulation of alternative splicing
23 in response to temperature variation in plants. *J Exp Bot* **72**, 6150-6163.

24 **Kindgren, P., Ivanov, M., and Marquardt, S.** (2020). Native elongation transcript
25 sequencing reveals temperature dependent dynamics of nascent RNAPII transcription
26 in Arabidopsis. *Nucleic Acids Res* **48**, 2332-2347.

27 **Kornblihtt, A.R., Schor, I.E., Allo, M., Dujardin, G., Petrillo, E., and Munoz, M.J.**
28 (2013). Alternative splicing: a pivotal step between eukaryotic transcription and
29 translation. *Nat Rev Mol Cell Biol* **14**, 153-165.

30 **Kosmacz, M., Gorka, M., Schmidt, S., Luzarowski, M., Moreno, J.C., Szlachetko, J.,**
31 **Leniak, E., Sokolowska, E.M., Sofroni, K., Schnittger, A., and Skirycz, A.** (2019).
32 Protein and metabolite composition of Arabidopsis stress granules. *New Phytol* **222**,
33 1420-1433.

34 **Kramer, M.C., Janssen, K.A., Palos, K., Nelson, A.D.L., Vandivier, L.E., Garcia,**
35 **B.A., Lyons, E., Beilstein, M.A., and Gregory, B.D.** (2020). N(6)-methyladenosine and
36 RNA secondary structure affect transcript stability and protein abundance during
37 systemic salt stress in Arabidopsis. *Plant Direct* **4**, e00239.

38 **Kwok, C.K., Ding, Y., Tang, Y., Assmann, S.M., and Bevilacqua, P.C.** (2013).
39 Determination of in vivo RNA structure in low-abundance transcripts. *Nat Commun* **4**,
40 2971.

- 1 **Labno, A., Tomecki, R., and Dziembowski, A.** (2016). Cytoplasmic RNA decay
2 pathways - Enzymes and mechanisms. *Biochim Biophys Acta* **1863**, 3125-3147.
- 3 **Law, J.A., and Jacobsen, S.E.** (2010). Establishing, maintaining and modifying DNA
4 methylation patterns in plants and animals. *Nat Rev Genet* **11**, 204-220.
- 5 **Leng, X., Ivanov, M., Kindgren, P., Malik, I., Thieffry, A., Brodersen, P., Sandelin,
6 A., Kaplan, C.D., and Marquardt, S.** (2020). Organismal benefits of transcription speed
7 control at gene boundaries. *EMBO Rep* **21**, e49315.
- 8 **Li, J., Chen, Z., Chen, F., Xie, G., Ling, Y., Peng, Y., Lin, Y., Luo, N., Chiang, C.M.,
9 and Wang, H.** (2020a). Targeted mRNA demethylation using an engineered dCas13b-
10 ALKBH5 fusion protein. *Nucleic Acids Res* **48**, 5684-5694.
- 11 **Li, L., Wu, W., Zhao, Y., and Zheng, B.** (2017). A reciprocal inhibition between ARID1
12 and MET1 in male and female gametes in Arabidopsis. *J Integr Plant Biol* **59**, 657-668.
- 13 **Li, Y., Song, Y., Xu, W., Li, Q., Wang, X., Li, K., Wang, J., Liu, Z., Velychko, S., Ye,
14 R., Xia, Q., Wang, L., Guo, R., Dong, X., Zheng, Z., Dai, Y., Li, H., Yao, M., Xue, Y.,
15 Scholer, H.R., Sun, Q., and Yao, H.** (2020b). R-loops coordinate with SOX2 in
16 regulating reprogramming to pluripotency. *Sci Adv* **6**, eaba0777.
- 17 **Liu, G., Kang, G., Wang, S., Huang, Y., and Cai, Q.** (2021). Extracellular Vesicles:
18 Emerging Players in Plant Defense Against Pathogens. *Front Plant Sci* **12**, 757925.
- 19 **Liu, K., and Sun, Q.** (2021). Intragenic tRNA-promoted R-loops orchestrate
20 transcription interference for plant oxidative stress responses. *Plant Cell* **33**, 3574-3591.
- 21 **Liu, L., and Chen, X.** (2018). Intercellular and systemic trafficking of RNAs in plants.
22 *Nat Plants* **4**, 869-878.
- 23 **Liu, X.M., Zhou, J., Mao, Y., Ji, Q., and Qian, S.B.** (2019). Programmable RNA N(6)-
24 methyladenosine editing by CRISPR-Cas9 conjugates. *Nat Chem Biol* **15**, 865-871.
- 25 **Liu, Y., Teng, C., Xia, R., and Meyers, B.C.** (2020). PhasiRNAs in Plants: Their
26 Biogenesis, Genic Sources, and Roles in Stress Responses, Development, and
27 Reproduction. *Plant Cell* **32**, 3059-3080.
- 28 **Loffer, A., Singh, J., Fukudome, A., Mishra, V., Wang, F., and Pikaard, C.S.** (2022).
29 A DCL3 dicing code within Pol IV-RDR2 transcripts diversifies the siRNA pool guiding
30 RNA-directed DNA methylation. *Elife* **11**.
- 31 **Long, J., Walker, J., She, W., Aldridge, B., Gao, H., Deans, S., Vickers, M., and
32 Feng, X.** (2021). Nurse cell--derived small RNAs define paternal epigenetic inheritance
33 in Arabidopsis. *Science* **373**.
- 34 **Lucero, L., Ferrero, L., Fonouni-Farde, C., and Ariel, F.** (2021). Functional
35 classification of plant long noncoding RNAs: a transcript is known by the company it
36 keeps. *New Phytol* **229**, 1251-1260.
- 37 **Luo, G.Z., MacQueen, A., Zheng, G., Duan, H., Dore, L.C., Lu, Z., Liu, J., Chen, K.,
38 Jia, G., Bergelson, J., and He, C.** (2014). Unique features of the m6A methylome in
39 Arabidopsis thaliana. *Nat Commun* **5**, 5630.

- 1 **Ma, X., Liu, C., Kong, X., Liu, J., Zhang, S., Liang, S., Luan, W., and Cao, X.** (2021).
2 Extensive profiling of the expressions of tRNAs and tRNA-derived fragments (tRFs)
3 reveals the complexities of tRNA and tRF populations in plants. *Sci China Life Sci* **64**,
4 495-511.
- 5 **Machyna, M., and Simon, M.D.** (2018). Catching RNAs on chromatin using
6 hybridization capture methods. *Brief Funct Genomics* **17**, 96-103.
- 7 **Manavella, P.A., Yang, S.W., and Palatnik, J.** (2019). Keep calm and carry on: miRNA
8 biogenesis under stress. *Plant J* **99**, 832-843.
- 9 **Manduzio, S., and Kang, H.** (2021). RNA methylation in chloroplasts or mitochondria in
10 plants. *RNA Biol* **18**, 2127-2135.
- 11 **Maronedze, C., Thomas, L., Lilley, K.S., and Gehring, C.** (2019). Drought Stress
12 Causes Specific Changes to the Spliceosome and Stress Granule Components. *Front*
13 *Mol Biosci* **6**, 163.
- 14 **Martinez-Perez, M., Aparicio, F., Lopez-Gresa, M.P., Belles, J.M., Sanchez-Navarro,**
15 **J.A., and Pallas, V.** (2017). Arabidopsis m(6)A demethylase activity modulates viral
16 infection of a plant virus and the m(6)A abundance in its genomic RNAs. *Proc Natl Acad*
17 *Sci U S A* **114**, 10755-10760.
- 18 **Maruri-Lopez, I., Figueroa, N.E., Hernandez-Sanchez, I.E., and Chodasiewicz, M.**
19 (2021). Plant Stress Granules: Trends and Beyond. *Front Plant Sci* **12**, 722643.
- 20 **Mathieu, M., Martin-Jaular, L., Lavieu, G., and Thery, C.** (2019). Specificities of
21 secretion and uptake of exosomes and other extracellular vesicles for cell-to-cell
22 communication. *Nat Cell Biol* **21**, 9-17.
- 23 **Matzke, M.A., and Mosher, R.A.** (2014). RNA-directed DNA methylation: an epigenetic
24 pathway of increasing complexity. *Nat Rev Genet* **15**, 394-408.
- 25 **Mauer, J., Luo, X., Blanjoie, A., Jiao, X., Grozhik, A.V., Patil, D.P., Linder, B.,**
26 **Pickering, B.F., Vasseur, J.J., Chen, Q., Gross, S.S., Elemento, O., Debart, F.,**
27 **Kiledjian, M., and Jaffrey, S.R.** (2017). Reversible methylation of m(6)Am in the 5' cap
28 controls mRNA stability. *Nature* **541**, 371-375.
- 29 **Mencia, R., Gonzalo, L., Tossolini, I., and Manavella, P.A.** (2022). Keeping up with
30 the miRNAs: current paradigms of the biogenesis pathway. *J Exp Bot*.
- 31 **Merchante, C., Stepanova, A.N., and Alonso, J.M.** (2017). Translation regulation in
32 plants: an interesting past, an exciting present and a promising future. *Plant J* **90**, 628-
33 653.
- 34 **Miller, H.E., Ilieva, M., Bishop, A.J.R., and Uchida, S.** (2022). Current Status of
35 Epitranscriptomic Marks Affecting lncRNA Structures and Functions. *Noncoding RNA* **8**.
- 36 **Mingardi, J., Musazzi, L., De Petro, G., and Barbon, A.** (2018). miRNA Editing: New
37 Insights into the Fast Control of Gene Expression in Health and Disease. *Mol Neurobiol*
38 **55**, 7717-7727.
- 39 **Moison, M., Pacheco, J.M., Lucero, L., Fonouni-Farde, C., Rodriguez-Melo, J.,**
40 **Mansilla, N., Christ, A., Bazin, J., Benhamed, M., Ibanez, F., Crespi, M., Estevez,**

1 **J.M., and Ariel, F.** (2021). The lncRNA APOLO interacts with the transcription factor
2 WRKY42 to trigger root hair cell expansion in response to cold. *Mol Plant* **14**, 937-948.

3 **Moore, J.W., Loake, G.J., and Spoel, S.H.** (2011). Transcription dynamics in plant
4 immunity. *Plant Cell* **23**, 2809-2820.

5 **Nguyen, C.C., Nakaminami, K., Matsui, A., Kobayashi, S., Kurihara, Y., Toyooka,**
6 **K., Tanaka, M., and Seki, M.** (2016). Oligouridylate Binding Protein 1b Plays an
7 Integral Role in Plant Heat Stress Tolerance. *Front Plant Sci* **7**, 853.

8 **Nuthikattu, S., McCue, A.D., Panda, K., Fultz, D., DeFraia, C., Thomas, E.N., and**
9 **Slotkin, R.K.** (2013). The initiation of epigenetic silencing of active transposable
10 elements is triggered by RDR6 and 21-22 nucleotide small interfering RNAs. *Plant*
11 *Physiol* **162**, 116-131.

12 **Olmedo-Monfil, V., Duran-Figueroa, N., Arteaga-Vazquez, M., Demesa-Arevalo, E.,**
13 **Autran, D., Grimanelli, D., Slotkin, R.K., Martienssen, R.A., and Vielle-Calzada, J.P.**
14 (2010). Control of female gamete formation by a small RNA pathway in *Arabidopsis*.
15 *Nature* **464**, 628-632.

16 **Ozata, D.M., Gainetdinov, I., Zoch, A., O'Carroll, D., and Zamore, P.D.** (2019). PIWI-
17 interacting RNAs: small RNAs with big functions. *Nat Rev Genet* **20**, 89-108.

18 **Perrella, G., Baurle, I., and van Zanten, M.** (2022). Epigenetic regulation of
19 thermomorphogenesis and heat stress tolerance. *New Phytol* **234**, 1144-1160.

20 **Petermann, E., Lan, L., and Zou, L.** (2022). Sources, resolution and physiological
21 relevance of R-loops and RNA-DNA hybrids. *Nat Rev Mol Cell Biol*.

22 **Petrillo, E., Godoy Herz, M.A., Fuchs, A., Reifer, D., Fuller, J., Yanovsky, M.J.,**
23 **Simpson, C., Brown, J.W., Barta, A., Kalyna, M., and Kornblihtt, A.R.** (2014). A
24 chloroplast retrograde signal regulates nuclear alternative splicing. *Science* **344**, 427-
25 430.

26 **Pikaard, C.S., Haag, J.R., Pontes, O.M., Blevins, T., and Cocklin, R.** (2012). A
27 transcription fork model for Pol IV and Pol V-dependent RNA-directed DNA methylation.
28 *Cold Spring Harb Symp Quant Biol* **77**, 205-212.

29 **Re, D.A., Lang, P.L.M., Yones, C., Arce, A.L., Stegmayer, G., Milone, D., and**
30 **Manavella, P.A.** (2019). Alternative use of miRNA-biogenesis co-factors in plants at low
31 temperatures. *Development* **146**.

32 **Reichel, M., Koster, T., and Staiger, D.** (2019). Marking RNA: m6A writers, readers,
33 and functions in *Arabidopsis*. *J Mol Cell Biol* **11**, 899-910.

34 **Reis, R.S., Deforges, J., Schmidt, R.R., Schippers, J.H.M., and Poirier, Y.** (2021).
35 An antisense noncoding RNA enhances translation via localized structural
36 rearrangements of its cognate mRNA. *Plant Cell* **33**, 1381-1397.

37 **Ren, B., Wang, X., Duan, J., and Ma, J.** (2019). Rhizobial tRNA-derived small RNAs
38 are signal molecules regulating plant nodulation. *Science* **365**, 919-922.

39 **Riegler, S., Servi, L., Scarpin, M.R., Godoy Herz, M.A., Kubaczka, M.G., Venhuizen,**
40 **P., Meyer, C., Brunkard, J.O., Kalyna, M., Barta, A., and Petrillo, E.** (2021). Light

1 regulates alternative splicing outcomes via the TOR kinase pathway. *Cell Rep* **36**,
2 109676.

3 **Ries, R.J., Zaccara, S., Klein, P., Olarerin-George, A., Namkoong, S., Pickering,**
4 **B.F., Patil, D.P., Kwak, H., Lee, J.H., and Jaffrey, S.R.** (2019). m(6)A enhances the
5 phase separation potential of mRNA. *Nature* **571**, 424-428.

6 **Rigo, R., Bazin, J., Romero-Barrios, N., Moison, M., Lucero, L., Christ, A.,**
7 **Benhamed, M., Blein, T., Huguet, S., Charon, C., Crespi, M., and Ariel, F.** (2020).
8 The Arabidopsis lncRNA ASCO modulates the transcriptome through interaction with
9 splicing factors. *EMBO Rep* **21**, e48977.

10 **Ruzicka, K., Zhang, M., Campilho, A., Bodi, Z., Kashif, M., Saleh, M., Eeckhout, D.,**
11 **El-Showk, S., Li, H., Zhong, S., De Jaeger, G., Mongan, N.P., Hejatko, J., Helariutta,**
12 **Y., and Fray, R.G.** (2017). Identification of factors required for m(6) A mRNA
13 methylation in Arabidopsis reveals a role for the conserved E3 ubiquitin ligase HAKAI.
14 *New Phytol* **215**, 157-172.

15 **Schor, I.E., Fiszbein, A., Petrillo, E., and Kornblihtt, A.R.** (2013). Intragenic
16 epigenetic changes modulate NCAM alternative splicing in neuronal differentiation.
17 *EMBO J* **32**, 2264-2274.

18 **Schor, I.E., Rascovan, N., Pelisch, F., Allo, M., and Kornblihtt, A.R.** (2009).
19 Neuronal cell depolarization induces intragenic chromatin modifications affecting NCAM
20 alternative splicing. *Proc Natl Acad Sci U S A* **106**, 4325-4330.

21 **Scutenaire, J., Deragon, J.M., Jean, V., Benhamed, M., Raynaud, C., Favory, J.J.,**
22 **Merret, R., and Bousquet-Antonelli, C.** (2018). The YTH Domain Protein ECT2 Is an
23 m(6)A Reader Required for Normal Trichome Branching in Arabidopsis. *Plant Cell* **30**,
24 986-1005.

25 **Shahid, S., Kim, G., Johnson, N.R., Wafula, E., Wang, F., Coruh, C., Bernal-**
26 **Galeano, V., Phifer, T., dePamphilis, C.W., Westwood, J.H., and Axtell, M.J.** (2018).
27 MicroRNAs from the parasitic plant *Cuscuta campestris* target host messenger RNAs.
28 *Nature* **553**, 82-85.

29 **Singh, J., Mishra, V., Wang, F., Huang, H.Y., and Pikaard, C.S.** (2019). Reaction
30 Mechanisms of Pol IV, RDR2, and DCL3 Drive RNA Channeling in the siRNA-Directed
31 DNA Methylation Pathway. *Mol Cell* **75**, 576-589 e575.

32 **Slotkin, R.K., Vaughn, M., Borges, F., Tanurdzic, M., Becker, J.D., Feijo, J.A., and**
33 **Martienssen, R.A.** (2009). Epigenetic reprogramming and small RNA silencing of
34 transposable elements in pollen. *Cell* **136**, 461-472.

35 **Song, P., Yang, J., Wang, C., Lu, Q., Shi, L., Tayier, S., and Jia, G.** (2021).
36 Arabidopsis N(6)-methyladenosine reader CPSF30-L recognizes FUE signals to control
37 polyadenylation site choice in liquid-like nuclear bodies. *Mol Plant* **14**, 571-587.

38 **Sorenson, R., and Bailey-Serres, J.** (2014). Selective mRNA sequestration by
39 OLIGOURIDYLATE-BINDING PROTEIN 1 contributes to translational control during
40 hypoxia in Arabidopsis. *Proc Natl Acad Sci U S A* **111**, 2373-2378.

- 1 **Sorenson, R.S., Deshotel, M.J., Johnson, K., Adler, F.R., and Sieburth, L.E.** (2018).
2 Arabidopsis mRNA decay landscape arises from specialized RNA decay substrates,
3 decapping-mediated feedback, and redundancy. *Proc Natl Acad Sci U S A* **115**, E1485-
4 E1494.
- 5 **Stepien, A., Knop, K., Dolata, J., Taube, M., Bajczyk, M., Barciszewska-Pacak, M.,**
6 **Pacak, A., Jarmolowski, A., and Szweykowska-Kulinska, Z.** (2017).
7 Posttranscriptional coordination of splicing and miRNA biogenesis in plants. *Wiley*
8 *Interdiscip Rev RNA* **8**.
- 9 **Su, Z., Tang, Y., Ritchey, L.E., Tack, D.C., Zhu, M., Bevilacqua, P.C., and Assmann,**
10 **S.M.** (2018). Genome-wide RNA structurome reprogramming by acute heat shock
11 globally regulates mRNA abundance. *Proc Natl Acad Sci U S A* **115**, 12170-12175.
- 12 **Su, Z., Wang, N., Hou, Z., Li, B., Li, D., Liu, Y., Cai, H., Qin, Y., and Chen, X.** (2020).
13 Regulation of Female Germline Specification via Small RNA Mobility in Arabidopsis.
14 *Plant Cell* **32**, 2842-2854.
- 15 **Su, Z., Zhao, L., Zhao, Y., Li, S., Won, S., Cai, H., Wang, L., Li, Z., Chen, P., Qin, Y.,**
16 **and Chen, X.** (2017). The THO Complex Non-Cell-Autonomously Represses Female
17 Germline Specification through the TAS3-ARF3 Module. *Curr Biol* **27**, 1597-1609
18 e1592.
- 19 **Sun, M., Schwalb, B., Pirkl, N., Maier, K.C., Schenk, A., Failmezger, H., Tresch, A.,**
20 **and Cramer, P.** (2013a). Global analysis of eukaryotic mRNA degradation reveals
21 Xrn1-dependent buffering of transcript levels. *Mol Cell* **52**, 52-62.
- 22 **Sun, Q., Csorba, T., Skourti-Stathaki, K., Proudfoot, N.J., and Dean, C.** (2013b). R-
23 loop stabilization represses antisense transcription at the Arabidopsis FLC locus.
24 *Science* **340**, 619-621.
- 25 **Szadeczky-Kardoss, I., Szaker, H.M., Verma, R., Darko, E., Pettko-Szandtner, A.,**
26 **Silhavy, D., and Csorba, T.** (2022). Elongation factor TFIIS is essential for heat stress
27 adaptation in plants. *Nucleic Acids Res* **50**, 1927-1950.
- 28 **Tack, D.C., Su, Z., Yu, Y., Bevilacqua, P.C., and Assmann, S.M.** (2020). Tissue-
29 specific changes in the RNA structurome mediate salinity response in Arabidopsis. *RNA*
30 **26**, 492-511.
- 31 **Tian, X., Qin, Z., Zhao, Y., Wen, J., Lan, T., Zhang, L., Wang, F., Qin, D., Yu, K.,**
32 **Zhao, A., Hu, Z., Yao, Y., Ni, Z., Sun, Q., De Smet, I., Peng, H., and Xin, M.** (2022).
33 Stress granule-associated TaMBF1c confers thermotolerance through regulating
34 specific mRNA translation in wheat (*Triticum aestivum*). *New Phytol* **233**, 1719-1731.
- 35 **Timmers, H.T.M., and Tora, L.** (2018). Transcript Buffering: A Balancing Act between
36 mRNA Synthesis and mRNA Degradation. *Mol Cell* **72**, 10-17.
- 37 **Tognacca, R.S., Kubaczka, M.G., Servi, L., Rodriguez, F.S., Godoy Herz, M.A., and**
38 **Petrillo, E.** (2020). Light in the transcription landscape: chromatin, RNA polymerase II
39 and splicing throughout Arabidopsis thaliana's life cycle. *Transcription* **11**, 117-133.

- 1 **Tomassi, A.H., Re, D.A., Romani, F., Cambiagno, D.A., Gonzalo, L., Moreno, J.E.,**
2 **Arce, A.L., and Manavella, P.A.** (2020). The Intrinsically Disordered Protein CARP9
3 Bridges HYL1 to AGO1 in the Nucleus to Promote MicroRNA Activity. *Plant Physiol* **184**,
4 316-329.
- 5 **Voinnet, O.** (2022). Revisiting small RNA movement in plants. *Nat Rev Mol Cell Biol* **23**,
6 163-164.
- 7 **Walker, J., Gao, H., Zhang, J., Aldridge, B., Vickers, M., Higgins, J.D., and Feng, X.**
8 (2018). Sexual-lineage-specific DNA methylation regulates meiosis in Arabidopsis. *Nat*
9 *Genet* **50**, 130-137.
- 10 **Wang, B., Sun, Y., Song, N., Zhao, M., Liu, R., Feng, H., Wang, X., and Kang, Z.**
11 (2017a). *Puccinia striiformis* f. sp. *tritici* microRNA-like RNA 1 (Pst-milR1), an important
12 pathogenicity factor of Pst, impairs wheat resistance to Pst by suppressing the wheat
13 pathogenesis-related 2 gene. *New Phytol* **215**, 338-350.
- 14 **Wang, F., Huang, H.-Y., Huang, J., Singh, J., and Pikaard, C.S.** (2022). Mechanisms
15 of AGO4 slicing-enhanced RNA-directed DNA methylation. *BioRxiv*.
- 16 **Wang, G., Jiang, H., Del Toro de Leon, G., Martinez, G., and Kohler, C.** (2018a).
17 Sequestration of a Transposon-Derived siRNA by a Target Mimic Imprinted Gene
18 Induces Postzygotic Reproductive Isolation in Arabidopsis. *Dev Cell* **46**, 696-705 e694.
- 19 **Wang, J.W., Czech, B., and Weigel, D.** (2009). miR156-regulated SPL transcription
20 factors define an endogenous flowering pathway in Arabidopsis thaliana. *Cell* **138**, 738-
21 749.
- 22 **Wang, M., Weiberg, A., Lin, F.M., Thomma, B.P., Huang, H.D., and Jin, H.** (2016).
23 Bidirectional cross-kingdom RNAi and fungal uptake of external RNAs confer plant
24 protection. *Nat Plants* **2**, 16151.
- 25 **Wang, Q., Xue, Y., Zhang, L., Zhong, Z., Feng, S., Wang, C., Xiao, L., Yang, Z.,**
26 **Harris, C.J., Wu, Z., Zhai, J., Yang, M., Li, S., Jacobsen, S.E., and Du, J.** (2021a).
27 Mechanism of siRNA production by a plant Dicer-RNA complex in dicing-competent
28 conformation. *Science* **374**, 1152-1157.
- 29 **Wang, W., Li, K., Yang, Z., Hou, Q., Zhao, W.W., and Sun, Q.** (2021b). RNase H1C
30 collaborates with ssDNA binding proteins WHY1/3 and recombinase RecA1 to fulfill the
31 DNA damage repair in Arabidopsis chloroplasts. *Nucleic Acids Res* **49**, 6771-6787.
- 32 **Wang, Y., Li, S., Zhao, Y., You, C., Le, B., Gong, Z., Mo, B., Xia, Y., and Chen, X.**
33 (2019). NAD(+)-capped RNAs are widespread in the Arabidopsis transcriptome and can
34 probably be translated. *Proc Natl Acad Sci U S A* **116**, 12094-12102.
- 35 **Wang, Z., Tang, K., Zhang, D., Wan, Y., Wen, Y., Lu, Q., and Wang, L.** (2017b). High-
36 throughput m6A-seq reveals RNA m6A methylation patterns in the chloroplast and
37 mitochondria transcriptomes of Arabidopsis thaliana. *PLoS One* **12**, e0185612.
- 38 **Wang, Z., Butel, N., Santos-Gonzalez, J., Borges, F., Yi, J., Martienssen, R.A.,**
39 **Martinez, G., and Kohler, C.** (2020). Polymerase IV Plays a Crucial Role in Pollen
40 Development in Capsella. *Plant Cell* **32**, 950-966.

- 1 **Wang, Z., Ma, Z., Castillo-Gonzalez, C., Sun, D., Li, Y., Yu, B., Zhao, B., Li, P., and**
2 **Zhang, X.** (2018b). SWI2/SNF2 ATPase CHR2 remodels pri-miRNAs via Serrate to
3 impede miRNA production. *Nature* **557**, 516-521.
- 4 **Wei, L., Gu, L., Song, X., Cui, X., Lu, Z., Zhou, M., Wang, L., Hu, F., Zhai, J.,**
5 **Meyers, B.C., and Cao, X.** (2014). Dicer-like 3 produces transposable element-
6 associated 24-nt siRNAs that control agricultural traits in rice. *Proc Natl Acad Sci U S A*
7 **111**, 3877-3882.
- 8 **Wei, L.H., Song, P., Wang, Y., Lu, Z., Tang, Q., Yu, Q., Xiao, Y., Zhang, X., Duan,**
9 **H.C., and Jia, G.** (2018). The m(6)A Reader ECT2 Controls Trichome Morphology by
10 Affecting mRNA Stability in Arabidopsis. *Plant Cell* **30**, 968-985.
- 11 **Weiberg, A., Wang, M., Lin, F.M., Zhao, H., Zhang, Z., Kaloshian, I., Huang, H.D.,**
12 **and Jin, H.** (2013). Fungal small RNAs suppress plant immunity by hijacking host RNA
13 interference pathways. *Science* **342**, 118-123.
- 14 **Wendte, J.M., and Pikaard, C.S.** (2017). The RNAs of RNA-directed DNA methylation.
15 *Biochim Biophys Acta Gene Regul Mech* **1860**, 140-148.
- 16 **Wierzbicki, A.T., Blevins, T., and Swiezewski, S.** (2021). Long Noncoding RNAs in
17 Plants. *Annu Rev Plant Biol* **72**, 245-271.
- 18 **Wierzbicki, A.T., Ream, T.S., Haag, J.R., and Pikaard, C.S.** (2009). RNA polymerase
19 V transcription guides ARGONAUTE4 to chromatin. *Nat Genet* **41**, 630-634.
- 20 **Wierzbicki, A.T., Cocklin, R., Mayampurath, A., Lister, R., Rowley, M.J., Gregory,**
21 **B.D., Ecker, J.R., Tang, H., and Pikaard, C.S.** (2012). Spatial and functional
22 relationships among Pol V-associated loci, Pol IV-dependent siRNAs, and cytosine
23 methylation in the Arabidopsis epigenome. *Genes Dev* **26**, 1825-1836.
- 24 **Wong-Bajracharya, J., Singan, V.R., Monti, R., Plett, K.L., Ng, V., Grigoriev, I.V.,**
25 **Martin, F.M., Anderson, I.C., and Plett, J.M.** (2022). The ectomycorrhizal fungus
26 *Pisolithus microcarpus* encodes a microRNA involved in cross-kingdom gene silencing
27 during symbiosis. *Proc Natl Acad Sci U S A* **119**.
- 28 **Wu, G., Park, M.Y., Conway, S.R., Wang, J.W., Weigel, D., and Poethig, R.S.**
29 (2009a). The sequential action of miR156 and miR172 regulates developmental timing
30 in Arabidopsis. *Cell* **138**, 750-759.
- 31 **Wu, L., Zhang, Q., Zhou, H., Ni, F., Wu, X., and Qi, Y.** (2009b). Rice MicroRNA
32 effector complexes and targets. *Plant Cell* **21**, 3421-3435.
- 33 **Wu, L., Zhou, H., Zhang, Q., Zhang, J., Ni, F., Liu, C., and Qi, Y.** (2010). DNA
34 methylation mediated by a microRNA pathway. *Mol Cell* **38**, 465-475.
- 35 **Xie, D., Chen, M., Niu, J., Wang, L., Li, Y., Fang, X., Li, P., and Qi, Y.** (2021). Phase
36 separation of SERRATE drives dicing body assembly and promotes miRNA processing
37 in Arabidopsis. *Nat Cell Biol* **23**, 32-39.
- 38 **Xu, L., Yuan, K., Yuan, M., Meng, X., Chen, M., Wu, J., Li, J., and Qi, Y.** (2020a).
39 Regulation of Rice Tillering by RNA-Directed DNA Methylation at Miniature Inverted-
40 Repeat Transposable Elements. *Mol Plant* **13**, 851-863.

- 1 **Xu, T., Wu, X., Wong, C.E., Fan, S., Zhang, Y., Zhang, S., Liang, Z., Yu, H., and**
2 **Shen, L.** (2022a). FIONA1-Mediated m(6) A Modification Regulates the Floral
3 Transition in Arabidopsis. *Adv Sci (Weinh)* **9**, e2103628.
- 4 **Xu, W., Li, K., Li, Q., Li, S., Zhou, J., and Sun, Q.** (2022b). Quantitative, Convenient,
5 and Efficient Genome-Wide R-Loop Profiling by ssDRIP-Seq in Multiple Organisms.
6 *Methods Mol Biol* **2528**, 445-464.
- 7 **Xu, W., Xu, H., Li, K., Fan, Y., Liu, Y., Yang, X., and Sun, Q.** (2017). The R-loop is a
8 common chromatin feature of the Arabidopsis genome. *Nat Plants* **3**, 704-714.
- 9 **Xu, W., Li, K., Li, S., Hou, Q., Zhang, Y., Liu, K., and Sun, Q.** (2020b). The R-Loop
10 Atlas of Arabidopsis Development and Responses to Environmental Stimuli. *Plant Cell*
11 **32**, 888-903.
- 12 **Yan, P., Liu, Z., Song, M., Wu, Z., Xu, W., Li, K., Ji, Q., Wang, S., Liu, X., Yan, K.,**
13 **Esteban, C.R., Ci, W., Belmonte, J.C.I., Xie, W., Ren, J., Zhang, W., Sun, Q., Qu, J.,**
14 **and Liu, G.H.** (2020). Genome-wide R-loop Landscapes during Cell Differentiation and
15 Reprogramming. *Cell Rep* **32**, 107870.
- 16 **Yang, X., Cheema, J., Zhang, Y., Deng, H., Duncan, S., Umar, M.I., Zhao, J., Liu, Q.,**
17 **Cao, X., Kwok, C.K., and Ding, Y.** (2020a). RNA G-quadruplex structures exist and
18 function in vivo in plants. *Genome Biol* **21**, 226.
- 19 **Yang, X., Yu, H., Sun, W., Ding, L., Li, J., Cheema, J., Ramirez-Gonzalez, R., Zhao,**
20 **X., Martin, A.C., Lu, F., Liu, B., Uauy, C., Ding, Y., and Zhang, H.** (2021). Wheat in
21 vivo RNA structure landscape reveals a prevalent role of RNA structure in modulating
22 translational subgenome expression asymmetry. *Genome Biol* **22**, 326.
- 23 **Yang, Z., Li, M., and Sun, Q.** (2020b). RHON1 Co-transcriptionally Resolves R-Loops
24 for Arabidopsis Chloroplast Genome Maintenance. *Cell Rep* **30**, 243-256 e245.
- 25 **Yang, Z., Hou, Q., Cheng, L., Xu, W., Hong, Y., Li, S., and Sun, Q.** (2017). RNase H1
26 Cooperates with DNA Gyrase to Restrict R-Loops and Maintain Genome Integrity in
27 Arabidopsis Chloroplasts. *Plant Cell* **29**, 2478-2497.
- 28 **Zhang, H., and Ding, Y.** (2021). Novel insights into the pervasive role of RNA structure
29 in post-transcriptional regulation of gene expression in plants. *Biochem Soc Trans* **49**,
30 1829-1839.
- 31 **Zhang, H., Zhong, H., Zhang, S., Shao, X., Ni, M., Cai, Z., Chen, X., and Xia, Y.**
32 (2019a). NAD tagSeq reveals that NAD(+)-capped RNAs are mostly produced from a
33 large number of protein-coding genes in Arabidopsis. *Proc Natl Acad Sci U S A* **116**,
34 12072-12077.
- 35 **Zhang, M., Ma, X., Wang, C., Li, Q., Meyers, B.C., Springer, N.M., and Walbot, V.**
36 (2021). CHH DNA methylation increases at 24-PHAS loci depend on 24-nt phased
37 small interfering RNAs in maize meiotic anthers. *New Phytol* **229**, 2984-2997.
- 38 **Zhang, W., Murphy, C., and Sieburth, L.E.** (2010). Conserved RNaseII domain protein
39 functions in cytoplasmic mRNA decay and suppresses Arabidopsis decapping mutant
40 phenotypes. *Proc Natl Acad Sci U S A* **107**, 15981-15985.

- 1 **Zhang, Y., Yang, M., Duncan, S., Yang, X., Abdelhamid, M.A.S., Huang, L., Zhang,**
2 **H., Benfey, P.N., Waller, Z.A.E., and Ding, Y.** (2019b). G-quadruplex structures trigger
3 RNA phase separation. *Nucleic Acids Res* **47**, 11746-11754.
- 4 **Zhang, Y.C., Lei, M.Q., Zhou, Y.F., Yang, Y.W., Lian, J.P., Yu, Y., Feng, Y.Z., Zhou,**
5 **K.R., He, R.R., He, H., Zhang, Z., Yang, J.H., and Chen, Y.Q.** (2020). Reproductive
6 phasiRNAs regulate reprogramming of gene expression and meiotic progression in rice.
7 *Nat Commun* **11**, 6031.
- 8 **Zhong, X., Du, J., Hale, C.J., Gallego-Bartolome, J., Feng, S., Vashisht, A.A.,**
9 **Chory, J., Wohlschlegel, J.A., Patel, D.J., and Jacobsen, S.E.** (2014). Molecular
10 mechanism of action of plant DRM de novo DNA methyltransferases. *Cell* **157**, 1050-
11 1060.
- 12 **Zhou, J., Zhang, W., and Sun, Q.** (2022a). R-loop: The new genome regulatory
13 element in plants. *J Integr Plant Biol*.
- 14 **Zhou, L., Tian, S., and Qin, G.** (2019). RNA methylomes reveal the m(6)A-mediated
15 regulation of DNA demethylase gene SIDML2 in tomato fruit ripening. *Genome Biol* **20**,
16 156.
- 17 **Zhou, M., Coruh, C., Xu, G., Martins, L.M., Bourbousse, C., Lambolez, A., and Law,**
18 **J.A.** (2022b). The CLASSY family controls tissue-specific DNA methylation patterns in
19 *Arabidopsis*. *Nat Commun* **13**, 244.
- 20 **Zhou, S.X., Zhu, Y., Wang, L.F., Zheng, Y.P., Chen, J.F., Li, T.T., Yang, X.M., Wang,**
21 **H., Li, X.P., Ma, X.C., Zhao, J.Q., Pu, M., Feng, H., Li, Y., Fan, J., Zhang, J.W.,**
22 **Huang, Y.Y., and Wang, W.M.** (2020). Osa-miR1873 fine-tunes rice immunity against
23 *Magnaporthe oryzae* and yield traits. *J Integr Plant Biol* **62**, 1213-1226.
- 24 **Zhou, X., Huang, K., Teng, C., Abdelgawad, A., Batish, M., Meyers, B.C., and**
25 **Walbot, V.** (2022c). 24-nt phasiRNAs move from tapetal to meiotic cells in maize
26 anthers. *New Phytol* **235**, 488-501.
- 27 **Zhu, J., Li, C., Peng, X., and Zhang, X.** (2021). RNA architecture influences plant
28 biology. *J Exp Bot* **72**, 4144-4160.
- 29 **Zhu, S., Gu, J., Yao, J., Li, Y., Zhang, Z., Xia, W., Wang, Z., Gui, X., Li, L., Li, D.,**
30 **Zhang, H., and Liu, C.** (2022). Liquid-liquid phase separation of RBGD2/4 is required
31 for heat stress resistance in *Arabidopsis*. *Dev Cell* **57**, 583-597 e586.

32
33
34

1 **Figure Legends**

2 **Figure 1. Light, sensed by the chloroplast, triggers a retrograde signal that**
3 **regulates alternative splicing in the nucleus.**

4 In the light, RNA polymerase II (Pol II) elongation rate is fast, resulting in exon skipping.
5 Leaf cells produce sugars that act as mobile signals to coordinate alternative splicing
6 responses throughout the whole plant, thus reaching root cells.

7
8 **Figure 2. RNA buffering is a flux-mediated regulatory mechanism that maintains**
9 **some mRNAs at a stable abundance.**

10 A. RNA abundance is influenced by the balance between RNA synthesis and
11 degradation. B. RNA flux describes the turnover rate of an mRNA, RNA buffering occurs
12 when the flux of an RNA shifts, but not its abundance.

13
14 **Figure 3. Hypothetical model of stress granule function.**

15 UBP1b localizes to the nucleus under non-stress conditions. UBP1b-stress granules
16 (SGs) are induced to form in the cytoplasm in response to heat stress. UBP1b-SGs
17 protect target mRNA from degradation during stress. Elucidation of the mechanism of
18 target mRNA recruitment and the timing of the translation of the protected mRNA will
19 provide critical information on the selective translation mechanisms utilized in plants in
20 response to stress.

21
22 **Figure 4. RNA structure may pervasively function in plant growth and**
23 **development.**

24 RNAs may fold into diverse RNA structures in different cell types and under different
25 environmental conditions. These dynamic and diverse RNA structures facilitate the
26 regulatory specificities of gene expression at post-transcriptional levels.

27
28 **Figure 5. R-loops in plant cells.**

29 Left, different distribution patterns of nuclear R-loops along the gene body in the
30 genomes of Arabidopsis, maize, and rice.

31 Right, chloroplast-localized AtRNH1C restricts RNA:DNA hybrid formation to release
32 head-on transcription-replication conflicts (TRCs) and to promote homologous
33 recombination (HR) repair in chloroplasts. Mitochondrion-localized AtRNH1B inhibits
34 homologous recombination at repeats in the mitochondrial genome by suppressing
35 RNA:DNA hybrid formation. In the absence of AtRNH1B, high levels of mitochondrial R-
36 loops stimulate the relocation of AtRNH1C to mitochondria.

37
38 **Figure 6. Regulatory roles of N⁶-methyladenosine (m⁶A) writers, erasers, and**
39 **readers in RNA metabolism, plant development, and stress responses.**

1 The cellular components responsible for installation, removal, and interpretation of m⁶A
 2 marks are methyltransferases (“writers”), demethylases (“erasers”), and RNA-binding
 3 proteins (“readers”), respectively. The m⁶A reader proteins YTH5, YTH9, and YTH13
 4 are also known as ECT4, ECT2, and ECT3, respectively. RNA methylation affects all
 5 aspects of RNA metabolism, including stability, export, intron splicing, and translational
 6 control, which are crucial for plant development and stress responses. Several potential
 7 m⁶A erasers and readers are yet to be identified.

8
 9 **Figure 7. Plant lncRNA grammar is determined by the transcript interactome.**

10 Multiple features contribute to the interaction of lncRNAs with DNA, protein partners or
 11 other RNA molecules. First, their expression pattern and their subcellular localization
 12 will restrict the range of potential partners. Second, the lncRNA interacting capacity
 13 depends on its sequence, post-transcriptional modifications, and secondary and tertiary
 14 structure adopted, which is, in turn, modulated by the interaction with partner molecules.
 15 Third, the resulting lncRNA interactome participates in the regulatory networks behind
 16 plant development and adaptation to the environment as all these factors can be
 17 responsive to environmental cues.

18
 19 **Figure 8. Unanswered questions of miRNA biogenesis.**

- 20 A. Can the alterations of processing efficiency caused by pri-miRNA refolding upon
 21 temperature change act as thermosensors during the plant response to heat?
 22 B. Can proteins specifically binding to the ssDNA or RNA/DNA strands of R-loops act as
 23 scaffold to recruit the microprocessor to *MIRNA* loci?
 24 C. How does the microprocessor and spliceosome interact?
 25 D. Can we define different D-bodies? And if so, can we establish the precise
 26 biochemistry within D-bodies during their maturation?
 27 E. Are co-transcriptionally processed miRNAs functionally different from their siblings
 28 produced post-transcriptionally, perhaps defining mobile miRNAs?

29
 30 **Figure 9. siRNA movement during male and female germline development in**
 31 **Arabidopsis.**

32 Here we use a more relaxed definition of the germline to indicate the cell lineage that
 33 undergoes meiosis and produces the gamete(s). The germline as strictly defined, is
 34 marked in red, ie. the generative and sperm cells (male) and the egg cell (female).
 35 Arrows mark the direction of the proposed siRNA movement. PMC, pollen mother cells;
 36 MMC, megaspore mother cells.

37
 38 **Figure 10. Establishment and maintenance of DNA methylation by RdDM**

39 21- and 22-nt siRNAs that are generated by DCL4 and DCL2 can bind to AGO1 to
 40 target mRNAs for post-transcriptional silencing (PTGS) or bind to AGO6 to initiate

1 RdDM in partnership with Pol V and DRM2. The latter enzymes are also key to the
2 major RdDM pathway that maintains silencing of thousands of loci and requires 24 -t
3 siRNAs that are generated by the Pol IV-RDR2 complex and DCL3 and loaded primarily
4 into AGO4. CG maintenance methylation, requiring MET1 and HDA6, is important for
5 both Pol IV and Pol V recruitment, and correlates with histone H3 lysine 9 dimethylation
6 (H3K9me2) among associated nucleosomes. Proteins that interact with these marks
7 and are implicated in Pol IV or Pol V transcriptional activity are indicated, as are histone
8 modifying enzymes involved in establishing repressive chromatin environments. The
9 figure is an update of the transcription fork model originally published in 2013 (Pikaard
10 et al., 2012), revised in 2017 (Wendte and Pikaard, 2017) and also adapted by other
11 authors (Matzke and Moshier, 2014).

12
13

ACCEPTED MANUSCRIPT




Figure 1
254x241 mm (x DPI)

1
2
3
4

1
2
3
4

5
6
7
8




Figure 2
83x39 mm (x DPI)






Figure 3
265x189 mm (x DPI)



1
2
3


Figure 4
290x204 mm (x DPI)



4
5
6

Figure 5
339x190 mm (x DPI)

1



2

3

4

5

Figure 6
158x98 mm (x DPI)

ACCEPTED MANUSCRIPT




Figure 7
185x136 mm (x DPI)

1
2
3
4




Figure 8
207x240 mm (x DPI)

1
2
3
4





Figure 9
106x131 mm (x DPI)

1
2
3
4

Initiation pathway:



Maintenance pathway:




Figure 10
254x190 mm (x DPI)

1
2
3