

## Journal Pre-proof

Improving the accuracy of beef cattle methane inventories in Latin America and Caribbean countries

Guilherme F.S. Congio, André Bannink, Olga L. Mayorga, João P.P. Rodrigues, Adeline Bougouin, Ermias Kebreab, Paulo C.F. Carvalho, Telma T. Berchielli, Maria E.Z. Mercadante, Sebastião C. Valadares-Filho, Ana L.C.C. Borges, Alexandre Berndt, Paulo H.M. Rodrigues, Juan C. Ku-Vera, Isabel C. Molina-Botero, Jacobo Arango, Ricardo A. Reis, Sandra L. Posada-Ochoa, Thierry R. Tomich, Octavio A. Castelán-Ortega, Marcos I. Marcondes, Carlos Gómez, Henrique M.N. Ribeiro-Filho, José I. Gere, Claudia Ariza-Nieto, Luis A. Giraldo, Horacio Gonda, María E. Cerón-Cucchi, Olegario Hernández, Patricia Ricci, Alexander N. Hristov



PII: S0048-9697(22)06227-1

DOI: <https://doi.org/10.1016/j.scitotenv.2022.159128>

Reference: STOTEN 159128

To appear in: *Science of the Total Environment*

Received date: 5 May 2022

Revised date: 18 September 2022

Accepted date: 26 September 2022

Please cite this article as: G.F.S. Congio, A. Bannink, O.L. Mayorga, et al., Improving the accuracy of beef cattle methane inventories in Latin America and Caribbean countries, *Science of the Total Environment* (2022), <https://doi.org/10.1016/j.scitotenv.2022.159128>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## Improving the accuracy of beef cattle methane inventories in Latin America and Caribbean countries

Guilherme F.S. Congio <sup>a,\*,1</sup>, André Bannink <sup>b</sup>, Olga L. Mayorga <sup>c</sup>, João P.P. Rodrigues <sup>d</sup>, Adeline Bougouin <sup>e</sup>, Ermias Kebreab <sup>e</sup>, Paulo C.F. Carvalho <sup>f</sup>, Telma T. Berchielli <sup>g</sup>, Maria E.Z. Mercadante <sup>h</sup>, Sebastião C. Valadares-Filho <sup>i</sup>, Ana L.C.C. Borges <sup>j</sup>, Alexandre Berndt <sup>k</sup>, Paulo H.M. Rodrigues <sup>l</sup>, Juan C. Ku-Vera <sup>m</sup>, Isabel C. Molina-Botero <sup>n</sup>, Jacobo Arango <sup>o</sup>, Ricardo A. Reis <sup>g</sup>, Sandra L. Posada-Ochoa <sup>p</sup>, Thierry R. Tomich <sup>q</sup>, Octavio A. Castelán-Ortega <sup>r</sup>, Marcos I. Marcondes <sup>s</sup>, Carlos Gómez <sup>n</sup>, Henrique M.N. Ribeiro-Filho <sup>t</sup>, José I. Gere <sup>u,v</sup>, Claudia Ariza-Nieto <sup>c</sup>, Luis A. Giraldo <sup>w</sup>, Horacio González <sup>x</sup>, María E. Cerón-Cucchi <sup>v,y</sup>, Olegario Hernández <sup>z</sup>, Patricia Ricci <sup>v,¶</sup>, Alexander N. Hristov <sup>‡</sup>

<sup>a</sup> Department of Animal Science, Luiz de Queiroz College of Agriculture, University of São Paulo, Piracicaba, SP 13418-900, Brazil

<sup>b</sup> Wageningen Livestock Research, Wageningen University & Research, Wageningen, AH 6700, The Netherlands

<sup>c</sup> Colombian Corporation for Agricultural Research, Tibaitatá, Bogotá D.C. 250047, Colombia

<sup>d</sup> Animal Science Institute, Department of Animal Production, Federal Rural University of Rio de Janeiro, Seropédica, RJ 23897-000, Brazil

<sup>e</sup> Department of Animal Science, University of California, Davis, CA 95618, USA

<sup>f</sup> Department of Forage Plants and Agrometeorology, Federal University of Rio Grande do Sul, Porto Alegre, RS 91501-970, Brazil

<sup>g</sup> Department of Animal Science, São Paulo State University, Jaboticabal, SP 14884-900, Brazil

<sup>h</sup> Institute of Animal Science, São Paulo Agribusiness Technology Agency, Sertãozinho, SP 14174-000, Brazil

<sup>i</sup> Department of Animal Science, Federal University of Viçosa, Viçosa, MG 36570-000, Brazil

<sup>j</sup> Department of Animal Science, Federal University of Minas Gerais, Belo Horizonte, MG 31270-901, Brazil

<sup>k</sup> Brazilian Agricultural Research Corporation, Embrapa Southeast Livestock, São Carlos, SP 13560-970, Brazil

<sup>l</sup> Department of Animal Nutrition and Production, Faculty of Veterinary Medicine and Animal Science, University of São Paulo, Pirassununga, SP 13635-900, Brazil

<sup>m</sup> Department of Animal Nutrition, Faculty of Veterinary Medicine and Animal Science, University of Yucatan, Mérida, Yucatán 97100, Mexico

<sup>n</sup> Department of Animal Husbandry, Faculty of Animal Science, National Agrarian University La Molina, Lima 15024, Peru

<sup>o</sup> International Center for Tropical Agriculture, Cali, Valle del Cauca 763537, Colombia

<sup>p</sup> Faculty of Agricultural Sciences, University of Antioquia, Medellín, Antioquia 050034, Colombia

<sup>q</sup> Brazilian Agricultural Research Corporation, Embrapa Dairy Cattle, Juiz de Fora, MG 36038-330, Brazil

<sup>r</sup> Faculty of Veterinary Medicine and Animal Science, Autonomous University of the State of Mexico, Toluca, Estado de México 50000, Mexico

<sup>s</sup> Department of Animal Sciences, Washington State University, Pullman, WA 99163, USA

<sup>t</sup> Department of Animal and Food Science, Santa Catarina State University, Lages, SC 88520-000, Brazil

<sup>u</sup> Regional Faculty of Buenos Aires, National Technological University, Buenos Aires C1179AAQ, Argentina

<sup>v</sup> National Scientific and Technical Research Council, Buenos Aires C1425FQB, Argentina

<sup>w</sup> Department of Animal Production, Faculty of Agricultural Sciences, National University of Colombia, Medellín, Antioquia 2037, Colombia

<sup>x</sup> Department of Animal Nutrition and Management, Faculty of Veterinary Medicine and Animal Science, Swedish University of Agricultural Sciences, Uppsala 75007, Sweden

<sup>y</sup> National Institute of Agricultural Technology, Institute of Pathobiology, Hurlingham C1686, Argentina

<sup>z</sup> National Institute of Agricultural Technology, Santiago del Estero G4200, Santiago del Estero, Argentina

<sup>¶</sup> National Institute of Agricultural Technology, Balcarce P1620, Argentina

<sup>‡</sup> Department of Animal Science, The Pennsylvania State University, University Park, PA 16802, USA

\* Corresponding author at: Department of Animal Science, Luiz de Queiroz College of Agriculture, University of São Paulo, Piracicaba 13418-900, SP, Brazil.

*E-mail address:* gcongio@gmail.com (G.F.S. Congio).

<sup>1</sup> Current address: Noble Research Institute LLC, 2510 Sam Noble Parkway, Ardmore, OK 73401, USA

## **Abstract**

On-farm methane (CH<sub>4</sub>) emissions need to be estimated accurately so that the mitigation effect of recommended practices can be accounted for. In the present study prediction equations for enteric CH<sub>4</sub> have been developed in lieu of expensive animal measurement approaches. Our objectives were to: (1) compile a dataset from individual beef cattle data for the Latin America and Caribbean (LAC) region; (2) determine main predictors of CH<sub>4</sub> emission variables; (3) develop and cross-validate prediction models according to dietary forage content (DFC); and (4) compare the predictive ability of these newly-developed models with extant equations reported in literature, including those currently used for CH<sub>4</sub>

inventories in LAC countries. After outlier's screening, 1100 beef cattle observations from 55 studies were kept in the final dataset (~ 50% of the original dataset). Mixed-effects models were fitted with a random effect of study. The whole dataset was split according to DFC into a subset for all-forage (DFC = 100%), high-forage ( $94\% \geq \text{DFC} \geq 54\%$ ), and low-forage ( $50\% \geq \text{DFC}$ ) diets. Feed intake and average daily gain (ADG) were the main predictors of  $\text{CH}_4$  emission ( $\text{g d}^{-1}$ ), whereas this was feeding level [dry matter intake (DMI) as % of body weight] for  $\text{CH}_4$  yield ( $\text{g kg}^{-1}$  DMI). The newly-developed models were more accurate than IPCC Tier 2 equations for all subsets. Simple and multiple regression models including ADG were accurate and a feasible option to predict  $\text{CH}_4$  emission when data on feed intake are not available. Methane yield was not well predicted by any extant equation in contrast to the newly-developed models. The present study delivered new models that may be alternatives for the IPCC Tier 2 equations to improve  $\text{CH}_4$  prediction for beef cattle in inventories of LAC countries based either on more or less readily available data.

**Keywords:** dietary nutrients; greenhouse gas; linear regression; livestock; methane conversion factor; model cross-validation

## 1. Introduction

Sixty per cent of all global methane ( $\text{CH}_4$ ) emissions are due to human-induced activities. Enteric  $\text{CH}_4$  fermentation and manure management represent 32% of global anthropogenic emissions, or 19.2% of total global  $\text{CH}_4$  emissions including non-anthropogenic sources (UNEP and CCAC, 2021). Developing countries contribute about 70% of livestock anthropogenic  $\text{CH}_4$  emissions globally of which 25% originates from Latin America and Caribbean (LAC) herds (UNEP and CCAC, 2021). The short lifetime of  $\text{CH}_4$  in the atmosphere, and its fast response to emission reduction, denotes that efforts to curb  $\text{CH}_4$  emissions will have a prompt global warming mitigation effect (Saunio et al., 2020). Therefore, reducing anthropogenic  $\text{CH}_4$  emissions in LAC countries, especially those from

enteric fermentation, may have an important role in the global endeavors to restrain temperature rise to 1.5-2.0 °C (Congio et al., 2021; Arndt et al., 2022).

Prediction equations have been widely developed in lieu of expensive and laborious animal CH<sub>4</sub> measurement approaches (Hristov et al., 2018). Previous studies, based mainly on data from developed countries, already indicated that diet- or region-specific models are more accurate than universal equations to predict CH<sub>4</sub> emission, which seem to be related in particular to differences in animal diets (Niu et al., 2018; Benaouda et al., 2019). When accurate, these equations would be convenient to estimate CH<sub>4</sub> emissions under a range of conditions, and next to optimize the allocation of resources for research programs on enteric CH<sub>4</sub>, which is particularly important for LAC developing countries (Congio et al., 2022a). Further, on-farm CH<sub>4</sub> emissions need to be estimated accurately so that mitigation of enteric CH<sub>4</sub> through better management practices is accounted for (Niu et al., 2018). Lastly, more accurate equations could aid LAC to improve the accuracy of their CH<sub>4</sub> inventories.

In 2018, beef cattle emitted 449 Mton of carbon dioxide equivalents due to ruminal CH<sub>4</sub> fermentation in the LAC region, or 39.5% of beef cattle emissions globally (FAOSTAT, 2020). Recent studies already developed CH<sub>4</sub> prediction models from ruminants in the LAC region. However, they were either focused only on dairy cattle (Congio et al., 2022a) and sheep (Congio et al., 2022b), or they used a limited database (Benaouda et al., 2020). The development of models specifically for beef cattle is still necessary due to specifics regarding livestock production systems (e.g., type and breed of animals) and diets (e.g., sources and levels of forage inclusion). Therefore, the objectives of the present study were to: (1) compile a comprehensive LAC dataset for individual beef cattle data; (2) determine main predictors of CH<sub>4</sub> emission variables; (3) develop and cross-validate prediction models according to different dietary inclusion levels of forage (i.e., the dietary forage content; DFC); and (4) compare the predictive ability of these newly-developed models with extant equations,

including those currently used in CH<sub>4</sub> inventories in LAC countries (IPCC 1997, 2006).

## 2. Material and methods

### 2.1. Dataset and data handling

The ‘Latin America Methane Project’ (LAMP) is a research initiative involving animal scientists from LAC countries studying *in vivo* CH<sub>4</sub> emission in ruminants and previous work was published by Congio et al. (2021, 2022a, 2022b). A comprehensive beef cattle dataset was compiled in LAMP which comprised 2194 individual beef cattle records from 67 studies (being 35 published and 32 unpublished). Studies performed from 2011 to 2021 by scientists from five LAC countries [Brazil,  $n = 1656$  (records) from 39 (studies); Mexico,  $n = 209$  from 11; Argentina,  $n = 178$  from 7; Colombia,  $n = 127$  from 9, Peru,  $n = 24$  from 1]. The database included CH<sub>4</sub> emission (g d<sup>-1</sup>) records along with corresponding dry matter intake (DMI; kg DM d<sup>-1</sup>), body weight (BW; kg), average daily gain (ADG; kg d<sup>-1</sup>), dietary forage content (DFC; % DM), crude protein (CP; % DM), ether extract (EE; % DM), neutral-detergent fiber (NDF; % DM), ash (% DM), and gross energy (GE; MJ kg<sup>-1</sup> DM) concentration. Where not available, dietary variables were calculated as follows. Dietary GE ( $n = 255$ ) according to Weiss and Tebbe (2019):  $\{GE = [(CP (\% DM) \times 0.056) + (EE (\% DM) \times 0.094)] + [(100 - CP - EE - ash (\% DM)) \times 0.042] \times 4.184\}$ . Publications from the LAC region were used to derive dietary NDF ( $n = 7$ ) and EE ( $n = 369$ ). Methane yield [CH<sub>4</sub> yield (g kg<sup>-1</sup> DMI) = CH<sub>4</sub> emission (g d<sup>-1</sup>) ÷ DMI (kg DM d<sup>-1</sup>)], gross energy intake [GEI (MJ d<sup>-1</sup>) = DMI (kg DM d<sup>-1</sup>) × dietary GE (MJ kg<sup>-1</sup> DM)], CH<sub>4</sub> conversion factor (Y<sub>m</sub>) [Y<sub>m</sub> (%) = CH<sub>4</sub> emission (g d<sup>-1</sup>) × 0.05565 ÷ GEI (MJ d<sup>-1</sup>) × 100], and feeding level (FL) [FL (%) = DMI (kg DM d<sup>-1</sup>) ÷ BW (kg) × 100], were computed for all observations.

Animal observations lacking CH<sub>4</sub> emission or DMI data were excluded from the dataset ( $n = 359$ ). Studies in which CH<sub>4</sub> emission was negatively correlated with DMI were also excluded (12 studies totaling 376 individual animal records) following discussion in a review



by Hristov et al. (2018). Additionally, records from treatments including anti-methanogenic additives or feed ingredients (*e.g.*, nitrate, tannins, lipid supplementation) were excluded ( $n = 163$ ) (Niu et al., 2018). Next, the identification of outliers was performed using the interquartile range approach (Zwillinger and Kokoska, 2000) considering a factor of 1.5 for extremes (van Lingen et al., 2019). It was performed based on all animal variables (Congio et al., 2022a). As a result, 1100 beef cattle observations from 55 studies were kept in the final dataset (~ 50% of the original dataset). This exclusion rate resembles that applied in previous meta-analyses for globally or regionally assembled databases where eventually also about 50% of the original records were retained (van Lingen et al., 2019; Congio et al., 2022b). The complete list of references of the refined dataset used in the current analysis is included in Supplementary Material.

## 2.2. Model development

Mixed-effects models were fitted using the lme4 procedure (Bates et al., 2015) of R statistical package (R Core Team 2020, version 4.0.2) according to:

$$Y_{ij} = \beta_0 + \beta_1 X_{ij1} + \beta_2 X_{ij2} + \dots + \beta_k X_{ijk} + S_i + \varepsilon_{ij}$$

where  $Y_{ij}$  denotes the  $j^{th}$  response variable of CH<sub>4</sub> emission (g d<sup>-1</sup>) or CH<sub>4</sub> yield (g kg<sup>-1</sup> DMI) from the  $i^{th}$  study;  $\beta_0$  denotes the fixed effect of intercept;  $X_{ij1}$  to  $X_{ijk}$  denote the fixed effects of predictor variables and  $\beta_1$  to  $\beta_k$  are the corresponding slopes;  $S_i$  and  $\varepsilon_{ij}$  are the random effect of study and residual error, respectively.

Models were developed following a sequential approach with predictor variables being incrementally added (van Lingen et al., 2019). Simple CH<sub>4</sub> emission models were based on DMI, GEI, or ADG. Then, multiple regression models assessed combinations of the above variables with NDF, EE, or CP separately. Finally, models were tested with a selection of dietary variables, a selection of DMI or GEI together with dietary variables, and a selection of all available variables, as well as a selection of all available variables except DMI or GEI

(Congio et al., 2022a). Methane yield models were developed considering ADG, FL, BW, GE, CP, EE, and NDF individually. Then, multiple CH<sub>4</sub> yield models were evaluated using a selection of dietary variables, and a selection of all available variables except DMI or GEI (as CH<sub>4</sub> yield is mathematically dependent on the latter variables). The backward multistep selection approach was used to select the most important variables to predict CH<sub>4</sub> emission and yield as described previously by van Lingen et al. (2019) and Congio et al. (2022a). The Bayesian information criterion was calculated for all fitted models, and those with the smallest values were chosen (James et al., 2014). The multi-collinearity among covariates in multiple regression models was verified considering the variance inflation factor (Zuur et al., 2010), and models were chosen when all covariates had a variance inflation factor lower than 3 (van Lingen et al., 2019).

### *2.3. Evaluation of developed models and extant equations*

A leave-one-out cross-validation was performed to evaluate the predictive ability of fitted prediction models at different levels (James et al., 2014). Studies were sequentially taken as the testing set for model evaluation, whereas those remained were considered as the training set for model fitting (van Lingen et al., 2019). Also, thirteen extant equations, including those from IPCC (1997, 2006) were evaluated (but not cross-validated). The most accurate equations from Ellis et al. (2007, 2009), Yan et al. (2009), Hristov et al. (2013), Ramin and Huhtanen (2013), Moraes et al. (2014), Charmley et al. (2016), Patra (2017), van Lingen et al. (2019), Benaouda et al. (2020), and Ribeiro et al. (2020) were selected to be evaluated considering the availability of predictor variables in the present dataset (equations are detailed in Table S4). To ensure independent evaluation, studies used to develop the extant equations that overlapped the current dataset were not included in the evaluations of those extant equations corresponding to the method adopted by van Lingen et al. (2019) and Congio et al. (2022a). Next to the use of the complete dataset, the dataset was split into

subsets according to DFC delivering subsets for data obtained with all-forage (DFC = 100%), high-forage ( $94\% \geq \text{DFC} \geq 54\%$ ), and low-forage ( $50\% \geq \text{DFC}$ ) diets, in order to derive separate prediction models for these subsets. These boundaries were defined to accommodate the wide range of beef cattle production systems from LAC region. A large proportion of beef cattle are raised in grazing systems with no supplement (*i.e.*, all-forage subset). The high-forage subset accommodates grazing animals receiving some level of concentrate and confined animals fed higher-forage diets, whereas the low-forage subset represents those are confined and fed higher levels of concentrate.

The all-forage subset comprised of mostly Angus (50%), Nellore (16%), and Brahman (10%) breeds. On the other hand, Nellore (35%), crosses including Zebu cattle (32%), Holstein (7%), Canchim (7%), Gyr (7%), Brahman (5%), and Brangus (5%) majorly composed the high-forage subset. Cattle fed low-forage diets were predominantly Nellore (72%), Holstein  $\times$  Charolais (7%), Canchim (7%), Holstein  $\times$  Zebu (7%). The main forage types for the all-forage subset were *Avena* spp. (33%), *Megathyrsus maximus* (32%), *Cynodon* spp. (9%), *Urochloa* spp. (9%), and *Pennisetum* spp. (6%), whereas corn silage (47%), *Urochloa* spp. (24%), and *M. maximus* (16%) were mostly represented in the high-forage subset. Corn silage (55%), fresh-cut sugar cane (16%), and *Urochloa* spp. (15%) were the main forages for the low-forage subset. Respiration chambers, GreenFeed<sup>TM</sup> (C-Lock Inc., Rapid City, SD), and the tracer gas sulfur hexafluoride (SF<sub>6</sub>) were used as CH<sub>4</sub> measurement techniques representing 8, 0 and 83%, 41, 10 and 44%, and 16, 41, and 44% of the data in the all-forage, high-forage and low-forage subsets, respectively. Further, DMI was mostly estimated using markers (81%) in the all-forage subset, whereas for the high-forage (80%) and low-forage (100%) subsets, the gravimetric technique was predominant.

Model performance was evaluated using a combination of statistic metrics including mean square prediction error (MSPE; Bibby and Toutenburg, 1977), mean bias (MB), slope

bias (SB), root MSPE (RMSPE), ratio between RMSPE and standard deviation of observed values (RSR; Moriasi et al., 2007), and concordance correlation coefficient (CCC; Lin, 1989). In the present analysis, we considered unsuitable models with  $RSR \geq 1.00$  (van Lingen et al., 2019). Equations used to calculate all above metrics are detailed in Supplementary Material.

### 3. Results and discussion

Several models to predict enteric CH<sub>4</sub> emissions from beef cattle have been published already (*e.g.*, Ellis et al., 2009; Charmley et al., 2016; van Lingen et al., 2019). However, these models were developed from small datasets, some used published treatment means data (*vs.* individual animal observations in the current study), and/or based on a restricted geographic region, which did not comprise data from LAC [*e.g.*, Ellis et al. (2007, 2009); Yan et al. (2009); Moraes et al. (2014); Charmley et al. (2016)]. Some studies included LAC data, but they either used limited databases (Benaouda et al., 2020) or included few data from LAC region (Escobar-Bahamondes et al., 2017). The ‘Global Network’ studies by Benaouda et al. (2019) and van Lingen et al. (2019) reported that regional equations are more accurate to predict CH<sub>4</sub> emissions, and that data from important beef-producing regions were essentially lacking. The current study is based on the largest database of individual animal data for beef cattle from the LAC region specifically including CH<sub>4</sub> measurements, and it therefore represents the greatest endeavour to date to establish accurate models for predicting CH<sub>4</sub> in beef cattle for this region.

#### 3.1. Dataset

Summary variable statistics for the complete dataset, and the all-forage, high-forage, and low-forage subsets are shown in Table 1. The complete dataset mainly constituted data for high-forage (47%) rather than all-forage (29.5%) and low-forage (23.5%) diets. The high- and low-forage subsets were composed only by data for growing cattle (100%), whereas all-

forage also included a small proportion of data for lactating beef cows (4%). Feed intake and BW increased as the dietary forage content of subsets decreased.

*Table 1*

Methane emission ranged from 38.2 from 364 g d<sup>-1</sup> in the current study, whereas Escobar-Bahamondes et al. (2017) and van Lingen et al. (2019) reported mean values ranging from 37.0 to 372 g d<sup>-1</sup> and from 37.4 to 322 g d<sup>-1</sup>. Methane yield ranged from 19.3 (low-forage) to 27.3 (all-forage) g kg<sup>-1</sup> DMI, whereas the Y<sub>m</sub> varied from 5.97 (low-forage) to 8.79% (all-forage) in the current study. van Lingen et al. (2019) and Escobar-Bahamondes et al. (2017), both using data predominantly from the EU and North America, reported CH<sub>4</sub> yield and Y<sub>m</sub> ranging from 20.0 to 22.1 g kg<sup>-1</sup> DMI and from 6.0 to 6.3%, respectively. Further, Benaouda et al. (2020), using a smaller dataset from LAC, reported mean values of CH<sub>4</sub> yield and Y<sub>m</sub> of 20.3 g kg<sup>-1</sup> DMI and 6.2%, respectively.

### *3.2. Methane emission*

Developed CH<sub>4</sub> emission equations and metrics of model performance are shown in Table 2. Average daily gain (Eqs. 1, 2, and 10), DMI (Eqs. 2 and 3), GEI (Eqs. 4, 5, 9, and 11), and dietary GE content (Eq. 11) had a positive relationship with CH<sub>4</sub> emission, whereas FL (Eq. 11) and dietary contents of EE (Eqs. 3, 5, and 11) and CP (Eqs. 9, 10, and 11) had a negative relationship. A negative relationship between CH<sub>4</sub> emission and dietary EE concentration is in line with studies by Moraes et al. (2014) and Escobar-Bahamondes et al. (2017). Increased dietary lipid supplies non-fermentable energy to the rumen, whereas suppresses archaeal activity, and may cause a shift in the rumen fermentation profile as well decrease NDF digestibility at higher inclusion levels (Hristov et al., 2013). Finally, dietary CP being negatively related with CH<sub>4</sub> emission agrees with Blaxter et al. (1971), which showed a decrease in CH<sub>4</sub> emission when diet CP increases.

Feed intake variables were the main predictors of CH<sub>4</sub> emission in line with previous

meta-analyses for beef cattle (van Lingen et al., 2019), dairy cattle (Niu et al., 2018), sheep (Congio et al., 2022b), and goats (Patra and Lalhriatpui, 2016). In the present meta-analysis, DMI was significantly and positively related to CH<sub>4</sub> emission with slopes varying from 19.2 to 19.6 g CH<sub>4</sub> kg<sup>-1</sup> DMI. This demonstrates that the greater the availability of substrate in the rumen, the greater the rate of formation CH<sub>4</sub> to be expected (Congio et al., 2022a). Furthermore, feed intake is a key covariate predicting CH<sub>4</sub> emissions because it is the consequence of certain dietary characteristics and animal nutritional requirements, both affecting the fermentative and digestive process (Charmley et al., 2016). Models based exclusively on feed chemical composition parameters did not perform well for the complete dataset as well as all subsets (results not shown), which is in line with Niu et al. (2018), reasserting also the key importance of DMI in determining CH<sub>4</sub> emission (Congio et al., 2022a).

Average daily gain (ADG) was positively related with CH<sub>4</sub> emission that agrees with previous results from Benaouda et al. (2020). This is probably due to the general positive relationship between feed intake and animal production variables (Hristov et al., 2005). Also, BW and CH<sub>4</sub> emission were positively corresponding to results of Moraes et al. (2014) and Benaouda et al. (2020). This is due to the proportionality between rumen volume and BW (Demment and Van Soest, 1985) as well as DMI. Heavier ruminants with higher requirements for maintenance energy are prone to consume extra feed and emit more CH<sub>4</sub> (Hristov et al., 2013).

Models including DMI, GEI, or ADG and dietary parameters (results not shown) were not more accurate than the one-variable models in the complete dataset and the high-forage subset. Nonetheless, the addition of EE (Eqs. 3 and 5) and CP (Eqs. 9 and 10) improved the model precision compared to those simple regressions in the all-forage and low-forage subsets, respectively. Models that included only dietary parameters were the least accurate

considering developed models in all subsets (results not shown).

*Table 2*

The ADG simple regression model (Eq. 1) was unique with a RSR < 1.00 (Table 2 and Fig. 1) in the complete dataset. For all-forage diets, the GEI model (Eq. 4) was slightly more accurate than the DMI model (Eq. 2), presenting smaller RSR (0.92 vs. 0.93) and RMSPE (31.8 vs. 32.2%), but both had low precision (MB ~ 15.0%). Inclusion of dietary EE content (Eqs. 3 and 5) improved the precision of these simple regression models (Table 2 and Fig. 2). The MB of those simple regressions including DMI (Eq. 2) or GEI (Eq. 4) averaged 15.92%, whereas the correction for dietary EE decreased the MB to an average of 6.97% (Eqs. 3 and 5). Both multiple regression models that allowed all variables (Eq. 6) and all variables except DMI (Eq. 7) for selection had the smallest RSR and RMSPE (Table 2), and included FL as well.

*Figs. 1 and 2*

Similar to the complete dataset, the ADG simple regression (Eq. 8) was the unique model that had RSR < 1.00 in the high-forage subset; however, it had low accuracy (SB = 11.67%) (Table 2 and Fig. 3). There were no simple regression models with RSR < 1.00 (results not shown) in the low-forage subset. Models including dietary CP content with GEI (Eq. 9) or ADG (Eq. 10) were associated with low RMSPE and good precision. The multiple regression model 11 selected GEI, FL, and dietary contents of GE, CP, and EE and had the best predictive accuracy (RMSPE = 18.9%) with negligible bias (MB = 0.13% and SB = 1.79%) (Table 2 and Fig. 4).

*Figs. 3 and 4*

Increasing complexity improved the predictive performance of CH<sub>4</sub> emission models only for the all-forage and low-forage subsets, but not for complete dataset and the high-forage subset. Usually, including dietary variables to feed intake simple models results in

more accurate models (Niu et al., 2018; Congio et al., 2022b). However, Ribeiro et al. (2020) and Congio et al. (2022a) also did not report more accurate equations comprising dietary variables using datasets containing LAC studies. The authors attributed those outcomes to a low variability of diets in terms of ingredients and nutrient contents (Ribeiro et al., 2020). Besides prediction potential, models with greater complexity might only be feasible in intensive beef cattle production systems. Extensive pasture-based systems are more common in LAC, however, and therefore simpler models based on easily available on-farm variables, such as ADG, probably are more practical to apply as discussed before by Congio et al. (2022a).

### *3.2.1. Performance of extant prediction equations*

Prediction performances of extant equations using the complete dataset, and the all-forage, high-forage, and low-forage subsets are presented in Table 3. In the complete subset, the equation from van Lingen et al. (2019), including corrections for dietary forage content and BW, was ranked first with a RMSE of 29.4%, good precision (MB = 0.02%), and reasonable accuracy (SB = 4.46%) (Table 3 and Fig. 1). This model was originally developed from a comprehensive global dataset. All remaining extant equations evaluated using the complete dataset had  $RSR > 1.00$  (Table S1 and Fig. S1, Supplementary Material). In the all-forage subset, the equation from Ramin and Huhtanen (2013) was the best-performing (Table 3 and Fig. 2). The second-ranked equation was that of van Lingen et al. (2019), but it had low precision (MB = 12.18%). The linear equation by Yan et al. (2009) was ranked third but had low accuracy (SB = 20.0%). The equation from Ribeiro et al. (2020) was ranked fourth, but it presented a large MB of 31.7% (Table 3 and Fig. S2). All remaining extant equations were not suitable to predict  $CH_4$  emission considering the all-forage subset including the IPCC (1997, 2006) Tier 2 equations (Table S2 and Fig. S2). The equation from Ellis et al. (2009) was the most accurate among extant equations for the high-forage subset (Table 3 and



Fig. 3). The equation from Benaouda et al. (2020) that included both DMI and NDF, was ranked second by RSR, but it had a large MB of 25.7%. The equation by van Lingen et al. (2019) also had  $RSR < 1.00$  but also had larger systematic biases ( $MB = 12.54\%$  and  $SB = 12.63\%$ ) (Table 3 and Fig. S3). All remaining extant equations evaluated using the high-forage subset had  $RSR > 1.00$  (Table S3 and Fig. S3). No extant equations were suitable for predicting  $CH_4$  emission in the low-forage subset with  $RSR > 1.00$  (Table S4 and Fig. S4).

Table 3

### 3.2.2. Comparison against extant prediction equations

Compared to the best extant equations, the most accurate  $CH_4$  emission models derived in the present study had a comparable accuracy with the all-forage subset, were outperformed with the high-forage, and outperformed with the low-forage subset. The equation from van Lingen et al. (2019) performed better than our ADG simple model (Eq. 1) for the complete dataset. For all-forage diets, the developed model 7 including both BW and FL had similar overall performance compared with those from Ramin and Huhtanen (2013) and van Lingen et al. (2019). The equation by Ellis et al. (2009) was the first-ranked extant equation and it outperformed our ADG model (Eq. 8) in the high-forage subset. Considering the low-forage subset, all extant equations were outperformed by models developed in the current analysis (Eqs. 9-11). Tier 2 equations from IPCC (1997, 2006) appeared unsuitable for predicting  $CH_4$  emission from beef cattle in the LAC region (Tables S1-S4). They had  $RSR > 1.00$  and were outperformed by the models developed in the present study for all subsets, highlighting an opportunity to improve  $CH_4$  inventories from LAC countries. Considering the fact that feed intake is barely obtainable in LAC beef ranches, the ADG simple or multiple regression models (Eqs. 1, 8, and 10) are good alternatives to predict  $CH_4$  emission from beef cattle under different dietary regimes with respect to forage inclusion level.

### 3.3. Methane yield

Table 4 shows CH<sub>4</sub> yield equations and model performance metrics. Increasing model complexity did not result in more accurate models for the complete dataset and high-forage subset. The FL simple models (Eqs. 12 and 14) were best predicting CH<sub>4</sub> yield. There were no simple regression models with RSR < 1.00 for the all-forage and low-forage subsets. The multiple regression model including both FL and dietary EE (Eq. 13) was the most accurate for the all-forage forage subset, whereas for the low-forage subset Eq. 15, which included dietary contents of GE, CP, and EE along with FL, showed the best predictive performance. From the extant equations for prediction of CH<sub>4</sub> yield, that of van Lingen et al. (2019) had a RSR < 1.00 for only the complete dataset and the low-forage subset with a high and low RMSPE of 34.4 and 21.9%, respectively (Table 5 and Fig. S5).

#### *Tables 4 and 5*

Similar to the results for CH<sub>4</sub> emission, the accuracy of newly-developed CH<sub>4</sub> yield models increased with model complexity for the all-forage and low-forage subsets, but not for the complete dataset and the high-forage subset. Most CH<sub>4</sub> modeling analyses have been focused on CH<sub>4</sub> emission rather than CH<sub>4</sub> yield (Congio et al., 2022a, 2022b). Our results indicate however that CH<sub>4</sub> yield on beef cattle fed typical LAC diets can also be predicted reasonably well using FI as predictor (Eqs. 12 and 14), combining it with dietary EE (Eq. 13), or with dietary GE, CP, and EE (Eq. 15). The newly-developed models in the present study outperformed the extant equation of van Lingen et al. (2019) for the all-forage, high-forage, and low-forage subsets.

#### *3.4. Research implications*

Research on CH<sub>4</sub> emissions from ruminants is relatively novel in the LAC region as reviewed by Congio et al. (2021). The dataset used in the present study reflected diets typically used for feeding beef cattle in LAC production systems but adding further research could improve accuracy of the current models. A more complete dietary nutrient

characterization in future research would eliminate the necessity to adopt table values from the literature to fill missing variables in dataset (Congio et al., 2022a).

Recently, meta-analysis studies also reported more accurate CH<sub>4</sub> prediction models when using digestibility variables as predictors (Benaouda et al., 2019, 2020; Ribeiro et al., 2020). We could not explore this kind of model in our study due to the absence of such data. Moreover, this poses a limitation to the applicability of such models in assisting CH<sub>4</sub> inventories in LAC countries is limited because such observations are very scarce in livestock operations. For these reasons the newly-derived models in the present study may be more useful and feasible options to follow.

Ideally, the comparison of developed models and extant equations in this kind of meta-analysis should be based on equal number of observations. However, in the present study, the use of ADG as predictor variable, that was included in only 75% of observations, as well as the evaluation of some equations, which presented overlapping of studies with our dataset, did not allow this ideal comparison scenario. As previously mentioned, these overlapped studies were removed from the evaluation of those extant equations in order to ensure independent evaluation. Moreover, we imagined that would be valuable to explore models including ADG because it is readily available on-farm. In cases of unequal amount of observations to compare model performances, the RSR is the most recommended statistic metric because it considers the data variability. This same approach was adopted by 'Global Network' studies by Benaouda et al. (2019) and van Lingen et al. (2019) in an attempt to maximize the number of observations to evaluate each model.

#### **4. Conclusions**

The current study is based on the largest dataset of individual animal data including CH<sub>4</sub> measurements for beef cattle from the LAC region, and thus represents the greatest endeavour to date to establish accurate models for predicting CH<sub>4</sub> emissions for beef cattle in

this region. Feed intake and ADG were the main predictors of CH<sub>4</sub> emission, whereas this was FL for CH<sub>4</sub> yield. Our most accurate CH<sub>4</sub> emission models outperformed IPCC Tier 2 equations (1997, 2006), whereas the CH<sub>4</sub> yield models outperformed extant equations, for all subsets. Simple and multiple regression models including ADG were accurate and probably a more feasible option to predict CH<sub>4</sub> emission when feed intake data are not available. Different individual extant equations were evaluated to be good options as well to predict CH<sub>4</sub> emission from beef cattle for the complete dataset, and the all-forage and high-forage subsets, but not for low-forage subset. The current developed models can replace IPCC Tier 2 equations thus allowing LAC countries to estimate CH<sub>4</sub> more accurately in their inventories.

### **Ethical approval**

Not applicable.

### **Funding**

The 'Latin America Methane Project' (LAMP) was funded by the New Zealand Government through AgResearch/New Zealand Agricultural Greenhouse Gas Research Centre (NZAGRC; grant S7-SOW21-Feed/Methane). The LAMP was a flagship in the 'Feed and Nutrition Network' of the 'Livestock Research Group' within the 'Global Research Alliance for Agricultural Greenhouse Gases'. The funder had no role in study design, data collection, decision to publish, or preparation of the manuscript. The funding sources that allowed the collaborators to carry out their projects are described in the Supplementary Material.

### **References**

Arndt, C., Hristov, A.N., Price, W.J., McClelland, S.C., Pelaez, A.M., Cueva, S.F., Oh, J., Dijkstra, J., Bannink, A., Bayat, A.R., Crompton, L.A., Eug`ene, M.A., Enahoro, D., Kebreab, E., Kreuzer, M., McGee, M., Martin, C., Newbold, C.J., Reynolds, C.K., Schwarm, A., Shingfield, K.J., Veneman, J.B., Y´añez-Ruiz, D.R., Yu, Z., 2022. Full

- adoption of the most effective strategies to mitigate methane emissions by ruminants can help meet the 1.5°C target by 2030 but not 2050. *Proc. Natl. Acad. Sci. U. S. A.* 119 (20), e2111294119. <https://doi.org/10.1073/pnas.2111294119>
- Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67, 1–51. <https://dx.doi.org/10.18637/jss.v067.i01>
- Benaouda, M., González-Ronquillo, M., Appuhamy, J., Kebreab, E., Molina, L.T., Herrera-Camacho, J., Ku-Vera, J.C., Ángeles-Hernández, J.C., Castelán-Ortega, O., 2020. Development of mathematical models to predict enteric methane emission by cattle in Latin America. *Livest. Sci.* 241, 104177. <https://doi.org/10.1016/j.livsci.2020.104177>
- Benaouda, M., Martin, C., Li, X., Kebreab, E., Hristov, A.N., Yu, Z., Yáñez-Ruiz, D.R., Reynolds, C.K., Crompton, L.A., Dijkstra, J., Bannink, A., Schwarm, A., Kreuzer, M., McGee, M., Lund, P., Hellwing, A.L.F., Veisbjerg, M.R., Moate, P.J., Bayat, A.R., Shingfield, K.J., Peiren, N., Eugène, M., 2019. Evaluation of the performance of existing mathematical models predicting enteric methane emissions from ruminants: animal categories and dietary mitigation strategies. *Anim. Feed Sci. Technol.*, 114207. <https://doi.org/10.1016/J.ANIFEEDSCI.2019.114207>
- Bibby, J., Toutenburg, T., 1977. *Prediction and Improved Estimation in Linear Models*. John Wiley & Sons, Chichester.
- Blaxter, K.L., Wainman, F.W., Dewey, P.J.S., Davidson, J., Denerley, H., Gunn, J.B., 1971. The effects of nitrogenous fertilizer on the nutritive value of artificially dried grass. *J. Agric. Sci.* 76, 307–319. <https://doi.org/10.1017/S0021859600025685>
- Charmley, E., Williams, S.R.O., Moate, P.J., Hegarty, R.S., Herd, R.M., Oddy, V.H., Reyenga, P., Staunton, K.M., Anderson, A., Hannah, M.C., 2016. A universal equation to predict methane production of forage-fed cattle in Australia. *Anim. Prod. Sci.* 56, 169–180. <https://doi.org/10.1071/AN1536>

- Congio, G.F.S., Bannink, A., Mayorga, O.L., Rodrigues, J.P.P., Bougouin, A., Kebreab, E., Silva, R.R., Maurício, R.M., da Silva, S.C., Oliveira, P.P.A., Muñoz, C., Pereira, L.G.R., Gómez, C., Ariza-Nieto, C., Ribeiro-Filho, H.M.N., Castelán-Ortega, O.A., Rosero-Noguera, J.R., Tieri, M.P., Rodrigues, P.H.M., Marcondes, M.I., Astigarraga, L., Abarca, S., Hristov, A.N., 2022a. Prediction of enteric methane production and yield in dairy cattle using a Latin America and Caribbean database. *Sci. Total Environ.* 825, 153982. <https://doi.org/10.1016/j.scitotenv.2022.153982>
- Congio, G.F.S., Bannink, A., Mayorga, O.L., Rodrigues, J.P.P., Bougouin, A., Kebreab, E., Carvalho, P.C.F., Abdalla, A.L., Monteiro, A.L.G., Kinyera, J.C., Gere, J.I., Gómez, C., Hristov, A.N., 2022b. Prediction of enteric methane production and yield in sheep using a Latin America and Caribbean database. *Livest. Sci.* 264, 105036. <https://doi.org/10.1016/j.livsci.2022.105036>
- Congio, G.F.S., Bannink, A., Mayorga Mogollón, O.L., Latin America Methane Project Collaborators, Hristov, A.N., 2021. Enteric methane mitigation strategies for ruminant livestock systems in the Latin America and Caribbean region: A meta-analysis. *J. Clean. Prod.* 312, 127693. <https://doi.org/10.1016/j.jclepro.2021.127693>
- Demment, M.W., Van Soest, P.J., 1985. A nutritional explanation for body-size patterns of ruminant and non-ruminant herbivores. *Am. Nat.* 125, 641–672. <https://doi.org/10.1086/284369>
- Ellis, J.L., Kebreab, E., Odongo, N.E., Beauchemin, K.A., McGinn, S., Nkrumah, J.D., Moore, S.S., Christopherson, R., Murdoch, G.K., McBride, B.W., Okine, E.K., France, J., 2009. Modeling methane production from beef cattle using linear and nonlinear approaches. *J. Anim. Sci.* 87, 1334–1345. <https://doi.org/10.2527/jas.2007-0725>
- Ellis, J.L., Kebreab, E., Odongo, N.E., McBride, B.W., Okine, E.K., France, J., 2007. Prediction of methane production from dairy and beef cattle. *J. Dairy Sci.* 90, 3456–3467.

- <https://doi.org/10.3168/jds.2006-675>
- Escobar-Bahamondes, P., Oba, M., Beauchemin, K.A., 2017. Universally applicable methane prediction equations for beef cattle fed high- or low-forage diets. *Can. J. Anim. Sci.* 97, 83–94. <https://doi.org/10.1139/cjas-2016-0042>
- FAOSTAT, 2020. ‘FAO statistical database.’ (Food and Agricultural Organization of the United Nations: Rome). <https://www.fao.org/faostat/en/#data> (accessed on 20 Sep 2021)
- Hristov, A.N., Kebreab, E., Niu, M., Oh, J., Bannink, A., Bayat, A.R., Boland, T., Brito, A.F., Casper, D., Crompton, L.A., Dijkstra, J., Eugène, M.A., Gransworthy, Ph.C., Haque, M.N., Hellwing, A.L.F., Huhtanen, P., Kreuzer, M., Kuhla, B., Lund, P., Madsen, J., Martin, C., Moate, P.J., Muetzel, S., Muñoz, C., Peiren, N., Powell, J.M., Reynolds, C.K., Schwarm, A., Shingfield, K.J., Storlien, T.M., Weisbjerg, M.R., Yáñez-Ruiz, D.R., Yu, Z., 2018. *Symposium review: Uncertainties in enteric methane inventories, measurement techniques, and prediction models.* *J. Dairy Sci.* 101, 6655–6674. <https://doi.org/10.3168/jds.2017-13526>
- Hristov, A.N., Oh, J., Firkins, J., Dijkstra, J., Kebreab, E., Waghorn, G., Makker, M.P.S., Adesogan, A.T., Yang, W., Lee, C., Gerber, P.J., Henderson, B., Tricarico, J.M., 2013. SPECIAL TOPICS - Mitigation of methane and nitrous oxide emissions from animal operations: I. A review of enteric methane mitigation options. *J. Anim. Sci.* 91, 5045–5069. <https://doi.org/10.2527/jas.2013-6583>
- Hristov, A.N., Price, W.J., Shafii, B., 2005. A meta-analysis on the relationship between intake of nutrients and body weight with milk volume and milk protein yield in dairy cows. *J. Dairy Sci.* 88, 2860–2869. [https://doi.org/10.3168/jds.S0022-0302\(05\)72967-2](https://doi.org/10.3168/jds.S0022-0302(05)72967-2)
- IPCC, 1997. Revised 1996 IPCC guidelines for national greenhouse gas inventories. Bracknell, UK: Intergovernmental Panel on Climate Change, IPCC/OECD/IEA. <https://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html> (accessed on 20 Sep 2021)


- IPCC, 2006. 2006 IPCC guidelines for national greenhouse gas inventories. IGES, Kanagawa, Japan: Intergovernmental Panel on Climate Change. <https://www.ipcc-nggip.iges.or.jp/public/2006gl/> (accessed on 20 Sep 2021)
- James, G., Witten, D., Hastie, T., Tibshirani, R., 2014. An Introduction to Statistical Learning: With Applications in R. Springer New York, US.
- Lin, L.I., 1989. A concordance correlation coefficient to evaluate reproducibility. *Biometrics* 45, 255–268. <https://doi.org/10.2307/2532051>
- Moraes, L.E., Strathe, A.B., Fadel, J.G., Casper, D.P., Kebreab, E., 2014. Prediction of enteric methane emissions from cattle. *Glob. Change Biol.* 20, 2140–2148. <https://doi.org/10.1111/gcb.12471>
- Moriyasi, D.N., Arnold, J.G., Van Liew, M.W., Bingham, R.L., Harmel, R.D., Veith, T.L., 2007. Model evaluation guidelines for systematic quantification of accuracy in watershed simulations. *T. ASABE* 50, 885–900. <https://doi.org/10.13031/2013.23153>
- Niu, M., Kebreab, E., Hristov, A.N., Oen, J., Arndt, C., Bannink, A., Bayat, A.R., Brito, A.F., Boland, T., Casper, D., Crompton, L.A., Dijkstra, J., Eugène, M.A., Garnsworthy, P.C., Haque, M.N., Hellwing, A.F., Huhtanen, P., Kreuzer, M., Kuhla, B., Lund, P., Madsen, J., Martin, C., McClelland, S.C., McGee, M., Moate, P.J., Muetzel, S., Muñoz, C., O’Kiely, P., Peiren, N., Reynolds, C.K., Schwarm, A., Shingfield, K.J., Storlien, T.M., Weisbjerg, M.R., Yáñez-Ruiz, D.R., Yu, Z., 2018. Prediction of enteric methane production, yield, and intensity in dairy cattle using an intercontinental database. *Glob. Change Biol.* 24, 3368–3389. <https://doi.org/10.1111/gcb.14094>
- Patra, A.K., 2017. Prediction of enteric methane emission from cattle using linear and non-linear statistical models in tropical production systems. *Mitig. Adapt. Strateg. Glob. Change* 22, 629–650. <https://doi.org/10.1007/s11027-015-9691-7>
- Patra, A.K., Lalhriatpuii, M., 2016. Development of statistical models for prediction of




- enteric methane emission from goats using nutrient composition and intake variables. *Agric. Ecosyst. Environ.* 215, 89–99. <https://doi.org/10.1016/j.agee.2015.09.018>
- R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/> (accessed on 20 Sep 2021)
- Ramin, M., Huhtanen, P., 2013. Development of equations for predicting methane emissions from ruminants. *J. Dairy Sci.* 96, 2476–2493. <https://dx.doi.org/10.3168/jds.2012-6095>
- Ribeiro, R.S., Rodrigues, J.P.P., Mauricio, R.M., Borges, A.L.C.C., Berchielli, T.T., Valadares-Filho, S.C., Machado, F.S., Campos, M.M., Tomich, T.R., Pereira, L.G.R., 2020. Predicting enteric methane production from cattle in the tropics. *Animal* 14, s438–s452. <https://doi.org/10.1017/S1751731120001743>
- Saunio, M., Stavert, A.R., Poulter, B., Fouquet, P., Canadell, J.G., Jackson, R.B., Raymond, P.A., Dlugokencky, E.J., Howeling, S., Patra, P.K., Ciais, P., Arora, V.K., Bastviken, D., Bergamaschi, P., Blake, D.R., Brailsford, G., Bruhwiler, L., Carlson, K.M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P.M., Covey, K., Curry, C.L., Etiope, G., Frankenberg, C., Geuney, N., Hegglin, M.I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K.M., Joos, F., Kleinen, T., Krummel, P.B., Langenfelds, R.L., Laruelle, G.G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O’Doherty, S., Parker, R.J., Peng, C., Peng, S., Peters, G.P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W.J., Rosentreter, J.A., Segers, A., Simpson, I.J., Shi, H., Smith, S.J., Steele, L.P., Thornton, B.F., Tian, H., Tohjima, Y., Tubiello, F.N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T.S., van Weele, M., van der Werf, G.R., Weiss, R.F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q.,

- Zhuang, Q., 2020. The Global Methane Budget 2000–2017. *Earth Syst. Sci. Data* 12, 1561–1623. <https://doi.org/10.5194/essd-12-1561-2020>
- UNEP and CAC, 2021. *Global Methane Assessment: Benefits and Costs of Mitigating Methane Emissions*. Nairobi: United Nations Environment Programme and Clean Air Coalition. <https://www.unep.org/resources/report/global-methane-assessment-benefits-and-costs-mitigating-methane-emissions> (accessed on 20 Sep 2021)
- van Lingen, H.J., Niu, M., Kebreab, E., Valadares-Filho, S.C., Rooke, J.A., Duthie, C-A, Schwarm, A., Kreuzer, M., Hynd, P.I., Caetano, M., Eugène, M., Martin, C., McGee, M., O’Kiely, P., Hünerberg, M., McAllister, T.A., Berchiesni, T.T., Messina, J.D., Peiren, N., Chaves, A.V., Charmley, E., Cole, N.A., Hales, K.E., Lee, S-S, Berndt, A., Reynolds, C.K., Crompton, L.A., Bayat, A-R, Yáñez-Ruiz, D.P., Yu, Z., Bannink, A., Dijkstra, J., Casper, D.P., Hristov, A.N., 2019. Prediction of enteric methane production, yield and intensity of beef cattle using an international database. *Agric. Ecosyst. Environ.* 283, 106575. <https://doi.org/10.1016/j.agee.2019.106575>
- Weiss, W.P., Tebbe, A.W., 2019. Estimating digestible energy values of feeds and diets and integrating those values into net energy systems. *TAS* 3(3), 953–961. <https://doi.org/10.1093/tas/tyy119>
- Yan, T., Porter, M.G., Mayne, C.S., 2009. Prediction of methane emission from beef cattle using data measured in indirect open-circuit respiration calorimeters. *Animal* 3, 1455–1462. <https://doi.org/10.1017/S175173110900473X>
- Zuur, A.F., Ieno, E.N., Elphick, C.S., 2010. A protocol for data exploration to avoid common statistical problems. *Methods Ecol. Evol.* 1, 3–14. <https://doi.org/10.1111/j.2041-210X.2009.00001.x>
- Zwillinger, D., Kokoska, S., 2000. *CRC Standard Probability and Statistics Tables and Formulae*. CRC Press, Boca Raton, US.

## Figure Captions




**Fig. 1.** Relationship between observed and predicted methane emission ( $\text{g d}^{-1}$ ) using the complete dataset. The black solid line is the identity line ( $y = x$ ), and the blue solid line is the fitted regression line for the relationship between observed and predicted values. For the interpretation of the references, the reader is referred to Table 2 (developed models), and Tables 3 and S1 (extant equations).




**Fig. 2.** Relationship between observed and predicted methane emission ( $\text{g d}^{-1}$ ) using the all-

forage (dietary forage content = 100%) subset. The black solid line is the identity line ( $y = x$ ), and the blue solid line is the fitted regression line for the relationship between observed and predicted values. For the interpretation of the references, the reader is referred to Tables 2 (developed models) and 3 (extant equations).



**Fig. 3.** Relationship between observed and predicted methane emission ( $\text{g d}^{-1}$ ) using the high-forage ( $94\% \geq$  dietary forage content  $\geq 54\%$ ) subset. The black solid line is the identity line ( $y = x$ ), and the blue solid line is the fitted regression line for the relationship between observed and predicted values. For the interpretation of the references, the reader is referred to Tables 2 (developed models) and 3 (extant equations).



**Fig. 4.** Relationship between observed and predicted methane emission ( $\text{g d}^{-1}$ ) using the low-forage ( $50\% \geq$  dietary forage content) subset. The black solid line is the identity line ( $y = x$ ), and the blue solid line is the fitted regression line for the relationship between observed and predicted values. For the interpretation of the references, the reader is referred to Table 2 (developed models).

**Table 1.** Summary statistics for the complete dataset, and the all-forage [dietary forage content (DFC) = 100%], high-forage (94%  $\geq$  DFC  $\geq$  54%), and low-forage (50%  $\geq$  DFC) subsets.

	Complete					All-forage					High-forage					Low-forage					
Item <sup>a</sup>	<i>n</i> <sup>b</sup>	Me	Min	Ma	SD <sup>b</sup>	<i>n</i>	Me	Min	Ma	SD	<i>n</i>	Me	Min	Ma	SD	<i>n</i>	Me	Min	Ma	SD	
		an	<sup>b</sup>	x <sup>b</sup>			an		x			an		x			an		x		
Animal variables																					
DMI (kg d <sup>-1</sup> )	110	7.40	1.3	14.	2.4	32	6.57	1.7	14.	2.3	51	7.49	1.3	14.	2.7	25	8.25	4.4	13.	1.3	
	0		7	6	4	5		7	6	5	6		7	3	4	9		4	1	6	
GEI (MJ d <sup>-1</sup> )	110	130	24.	263	43.	32	113	24.	23	38.	51	131	25.	26	49.	25	148	78.	22	24.	
	0		8		2	5		8	8	6	6		1	3	0	9		6	9	3	
BW (kg)	110	345	106	657	103	32	300	122	53	92.	51	351	106	65	99.	25	389	110	64	101	
	0					5			4	3	6			7	8	9			3		
FL (DMI as %	110	2.20	0.7	4.4	0.6	32	2.27	0.9	4.1	0.6	51	2.14	0.7	4.1	0.5	25	2.23	1.1	4.4	0.5	
BW)	0		51	6	00	5		27	5	97	6		51	4	60	9		1	6	39	
ADG (kg d <sup>-1</sup> )	825	1.01	0.0	2.7	0.4	25	1.01	0.3	2.0	0.3	36	0.91	0.0	2.3	0.4	20	1.17	0.3	2.7	0.3	
			92	9	31	4		57	0	60	8		5	92	1	75	3		33	9	79

Diet composition (% DM)																				
NDF	110	49.9	17.	85.	14.	32	60.1	41.	85.	11.	51	50.1	28.	73.	12.	25	36.6	17.	63.	10.
	0		2	7	2	5		1	7	0	6		3	4	1	9		2	4	2
EE	110	2.97	0.6	6.9	1.1	32	2.82	0.6	4.0	1.0	51	2.90	0.7	6.9	1.2	25	3.31	1.2	6.4	1.1
	0		70	9	8	5		70	0	8	6		30	9	5	9		4	6	0
CP	110	14.3	4.0	25.	3.4	32	15.2	4.0	25.	5.0	51	13.7	7.0	21.	2.4	25	14.2	9.8	20.	2.1
	0		9	1	4	5		6	1	8	6		8	2	1	9		5	0	3
Ash	110	7.31	3.2	13.	2.3	32	9.17	5.1	13.	1.8	51	7.18	3.5	12.	1.9	25	5.24	3.2	11.	1.5
	0		5	2	3	5		9	2	6	6		0	8	6	9		5	0	2
Forage	110	72.6	17.	100	23.	32	100	100	10	0	51	72.1	53.	94.	10.	25	39.1	17.	50.	10.
	0		3		7	5			0		6		9	4	8	9		3	0	0
GE (MJ kg <sup>-1</sup> DM)	110	17.6	14.	21.	0.9	32	17.3	14.	19.	0.6	51	17.6	14.	20.	1.0	25	18.0	16.	21.	0.8
	0		0	3	48	5		0	9	55	6		5	0	7	9		5	3	57
Methane emissions																				
CH <sub>4</sub> (g d <sup>-1</sup> )	110	153	38.	364	48.	32	171	57.	34	59.	51	139	38.	36	44.	25	156	71.	29	31.

	0		2		7	5		3	7	1	6		2	4	4	9		4	8	6
CH <sub>4</sub> per DMI (g kg <sup>-1</sup> )	110	22.1	8.7	49.	7.8	32	27.3	13.	48.	8.0	51	20.2	8.7	49.	7.4	25	19.3	9.9	37.	4.5
	0		8	5	0	5		0	1	4	6		8	5	1	9		7	0	0
Y <sub>m</sub> (% GEI)	110	7.01	2.6	15.	2.5	32	8.79	4.2	15.	2.6	51	6.42	2.5	15.	2.3	25	5.97	3.1	12.	1.3
	0		6	0	0	5		0	0	0	6		6	0	1	9		4	0	2

<sup>a</sup> DMI = dry matter intake; GEI = gross energy intake; BW = body weight; FL = feeding level; NDF = dietary neutral-detergent fiber content; EE = dietary ether extract content; CP = dietary crude protein content; GE = dietary gross energy; ADG = average daily gain; Y<sub>m</sub> = methane conversion factor.

<sup>b</sup> n = number of observations; Min = minimum; Max = maximum; SD = standard deviation.

**Table 2.** Methane emission (g d<sup>-1</sup>) prediction equations and model evaluation metrics for the complete dataset, and for the all-forage [dietary forage content (DFC) = 100%], high-forage (34% ≥ DFC ≥ 54%), and low-forage (50% ≥ DFC) subsets.

Eq.	Prediction equation <sup>a</sup>		Model performance <sup>b</sup>				
		n <sup>b</sup>	RSR	RMSPE, %	MB, %	SB, %	CCC
(1)	114 (7.02) + 33.0 (4.17) × ADG	825	0.99	32.3	1.57	1.74	0.11
	All-forage						

(2)	$20.0 (12.3) + 19.6 (1.32) \times \text{DMI}$	325	0.93	32.2	15.98	4.87	0.49
(3)	$65.2 (23.7) + 19.2 (1.38) \times \text{DMI} - 21.2 (8.37) \times \text{EE}$	325	0.92	32.0	7.28	7.33	0.48
(4)	$19.6 (12.3) + 1.15 (0.077) \times \text{GEI}$	325	0.92	31.8	15.86	4.05	0.50
(5)	$65.2 (23.6) + 1.12 (0.080) \times \text{GEI} - 21.2 (8.36) \times \text{EE}$	325	0.91	31.4	6.65	6.33	0.49
(6)	$- 83.9 (30.7) + 0.675 (0.093) \times \text{GEI} + 0.358 (0.036) \times \text{BW} + 6.53 (2.29) \times$ ash	254	0.88	28.6	17.86	2.18	0.54
(7)	$- 71.0 (20.9) + 0.570 (0.033) \times \text{BW} + 27.9 (4.26) \times \text{FL}$	254	0.85	27.6	14.09	0.82	0.54
High-forage							
(8)	$80.8 (9.81) + 63.6 (4.66) \times \text{ADG}$	368	0.99	31.8	0.17	11.67	0.35
Low-forage							
(9)	$85.1 (28.3) + 0.655 (0.070) \times \text{GEI} - 1.73 (1.76) \times \text{CP}$	259	0.99	20.1	0.29	6.01	0.22
(10)	$207 (33.8) + 28.7 (5.59) \times \text{ADG} - 6.21 (2.27) \times \text{CP}$	203	0.96	19.9	0.11	2.39	0.24
(11)	$10.3 (68.8) + 0.739 (0.084) \times \text{GEI} - 17.4 (5.99) \times \text{FL} + 11.3 (3.73) \times \text{GE} -$ $7.18 (1.97) \times \text{CP} - 9.37 (2.33) \times \text{EE}$	203	0.91	18.9	0.13	1.79	0.37

<sup>a</sup> ADG = average daily gain ( $\text{kg d}^{-1}$ ); DMI = dry matter intake ( $\text{kg d}^{-1}$ ); EE = dietary ether extract content (% DM); GEI = gross energy intake



(MJ d<sup>-1</sup>); BW = body weight (kg); ash = dietary ash content (% DM); FL = feeding level (DMI as % BW); CP = dietary crude protein content (% DM); GE = dietary gross energy (MJ kg<sup>-1</sup> DM).

<sup>b</sup>  $n$  = number of observations used to fit equations and for model evaluation; RSR = RMSPE-observations standard deviation ratio; RMSPE = root mean square prediction error (% of observed CH<sub>4</sub> production means); MB = mean bias (% of MSPE); SB = slope bias (% of MSPE); CCC = concordance correlation coefficient.

**Table 3.** Performance evaluation of extant equations to predict enteric CH<sub>4</sub> emission (g d<sup>-1</sup>) using the complete dataset, and the all-forage [dietary forage content (DFC) = 100%], and high-forage (94% ≥ DFC ≥ 54%) subsets.

Ran k	Reference	Equation <sup>a</sup>	$n^b$	RSR <sup>b</sup>	RMSPE, % <sup>b</sup>	MB, % <sup>b</sup>	SB, % <sup>b</sup>	CCC <sup>b</sup>
Complete								
1	van Lingen et al. (2019)	$- 6.41 + 11.3 \times \text{DMI} + 0.557 \times \text{For} + 0.0996 \times \text{BW}$	975	0.88	29.4	0.02	4.46	0.47
All-forage								
1	Ramin and Huhtanen (2013)	$(62 + 25 \times \text{DMI}) \times 0.714$	325	0.83	28.7	3.51	2.03	0.55
2	van Lingen et al. (2019)	$- 6.41 + 11.3 \times \text{DMI} + 0.557 \times \text{For} + 0.0996 \times \text{BW}$	325	0.84	29.0	12.18	0.57	0.49

3	Yan et al. (2009)	$(14.7 + 35.1 \times \text{DMI}) \times 0.714$	325	0.90	31.3	0.70	20.04	0.59
4	Ribeiro et al. (2020)	$(0.734 + 0.041 \times \text{GEI} + 0.009 \times \text{BW} - 0.04 \times \text{EE}) \div 0.05565$	289	0.95	31.7	31.75	0.30	0.49
High-forage								
1	Ellis et al. (2009)	$(2.29 + 0.67 \times \text{DMI}) \div 0.05565$	516	0.92	29.4	3.59	7.48	0.46
2	Benaouda et al. (2020)	$17.0 \times \text{DMI} + 0.03 \times \text{NDF}$	301	0.96	32.9	25.69	9.97	0.56
3	van Lingen et al. (2019)	$-6.41 + 11.3 \times \text{DMI} + 0.557 \times \text{For} + 0.0906 \times \text{BW}$	484	0.98	31.9	12.54	12.63	0.49

<sup>a</sup> DMI = dry matter intake ( $\text{kg d}^{-1}$ ); BW = body weight (kg); GEI = gross energy intake ( $\text{MJ d}^{-1}$ ); EE = dietary ether extract content (% DM); NDF = dietary neutral-detergent fiber content (% DM); For = dietary forage content (% DM).

<sup>b</sup>  $n$  = number of observations used for model evaluation; RSR = RMSPE-observations standard deviation ratio; RMSPE = root mean square prediction error (% observed  $\text{CH}_4$  production means); MB = mean bias (% MSPE); SB = slope bias (% MSPE); CCC = concordance correlation coefficient.

**Table 4.** Methane yield ( $\text{g kg}^{-1}$  DMI) prediction equations and model evaluation metrics for the complete dataset, and the all-forage [dietary forage content (DFC) = 100%], high-forage ( $94\% \geq \text{DFC} \geq 54\%$ ), and low-forage ( $50\% \geq \text{DFC}$ ) subsets.

Eq.	Prediction equation <sup>a</sup>	Model performance <sup>b</sup>
-----	----------------------------------	--------------------------------

Complete		$n^b$	RSR	RMSPE, %	MB, %	SB, %	CCC
(12)	$30.7 (1.04) - 4.35 (0.262) \times \text{FL}$	1100	0.97	34.2	1.65	0.21	0.16
All-forage							
(13)	$29.8 (2.64) - 6.31 (0.565) \times \text{FL} + 4.03 (1.20) \times \text{EE}$	325	0.88	26.0	3.14	0.15	0.40
High-forage							
(14)	$27.4 (1.22) - 3.28 (0.324) \times \text{FL}$	516	0.92	33.5	0.08	5.54	0.21
Low-forage							
(15)	$1.19 (10.2) - 3.20 (0.676) \times \text{FL} + 2.27 (0.547) \times \text{GE} - 0.853 (0.269) \times \text{CP} - 1.16 (0.292) \times \text{EE}$	203	0.90	19.9	1.24	0.58	0.37

<sup>a</sup> FL = feeding level (DMI as % BW); EE = dietary ether extract content (% DM); GE = dietary gross energy (MJ kg<sup>-1</sup> DM); CP = dietary crude protein content (% DM).

<sup>b</sup>  $n$  = number of observations used to fit equations and for model evaluation; RSR = RMSPE-observations standard deviation ratio; RMSPE = root mean square prediction error (% observed CH<sub>4</sub> production means); MB = mean bias (% MSPE); SB = slope bias (% MSPE); CCC = concordance correlation coefficient.

**Table 5.** Performance evaluation of van Lingen et al. (2019) equation<sup>a</sup> to predict enteric CH<sub>4</sub> yield (g kg<sup>-1</sup> DMI) using complete dataset, and the

all-forage [dietary forage content (DFC) = 100%], high-forage ( $94\% \geq \text{DFC} \geq 54\%$ ), and low-forage ( $50\% \geq \text{DFC}$ ) subsets.

Dataset or Subset	$n^b$	RSR <sup>b</sup>	RMSPE, % <sup>b</sup>	MB, % <sup>b</sup>	SB, % <sup>b</sup>	CCC <sup>b</sup>
Complete	975	0.96	34.4	1.14	5.49	0.11
All-forage	325	1.13	33.4	22.51	2.28	0.01
High-forage	484	1.00	37.7	2.72	0.84	0.03
Low-forage	166	0.99	21.9	2.01	0.09	0.03

<sup>a</sup>  $\text{CH}_4$  yield ( $\text{g kg}^{-1}$  DMI) =  $17.3 + 0.0565 \times \text{For}$ ; where For = dietary forage content (% DM).

<sup>b</sup>  $n$  = number of observations used for model evaluation; RSR = RMSE/observations standard deviation ratio; RMSPE = root mean square prediction error (% observed  $\text{CH}_4$  production means); MB = mean bias (% MSPE); SB = slope bias (% MSPE); CCC = concordance correlation coefficient.

**CRedit authorship contribution statement**


**Guilherme F.S. Congio:** Investigation, Methodology, Data curation, Formal analysis, Writing – original draft, Visualization, Writing – review & editing; **André Bannink:** Conceptualization, Methodology, Validation, Writing – review & editing, Supervision, Funding acquisition, Project administration; **Olga L. Mayorga:** Supervision, Project administration; **João P.P. Rodrigues, Adeline Bougouin, Ermias Kebreab:** Methodology, Formal analysis, Software, Visualization; **Paulo C.F. Carvalho, Telma T. Berchielli, Maria E.Z. Mercadante, Sebastião C. Valadares-Filho, Ana L.C.C. Borges, Alexandre Berndt, Paulo H.M. Rodrigues, Juan C. Ku-Vera, Isabel C. Molina-Botero, Jacobo Arango, Ricardo A. Reis, Sandra L. Posada-Ochoa, Thierry P. Tomich, Octavio A. Castelán-Ortega, Marcos I. Marcondes, Carlos Gómez, Henrique M.N. Ribeiro-Filho, José I. Gere, Claudia Ariza-Nieto, Luis A. Giraldo, Horacio Gonda, María E. Cerón-Cucchi, Olegario Hernández, Patricia Ricci:** Investigation, Data curation, Writing – review & editing; **Alexander N. Hristov:** Conceptualization, Methodology, Validation, Writing – review & editing, Supervision, Project administration, Funding acquisition.

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Graphical abstract



## HIGHLIGHTS

- ✓ Intake and ADG were the key predictors of beef cattle CH<sub>4</sub> production
- ✓ Separate models according to dietary forage content improved predictive ability
- ✓ Developed models were more accurate than IPCC Tier 2 equations for all subsets
- ✓ Developed models can allow LAC countries to improve the accuracy of their GHG inventories

Journal Pre-proof