

RESEARCH ARTICLE

Intestinal parasitic infections in a community from Pampa del Indio, Chaco (Argentina) and their association with socioeconomic and environmental factors

Carlos Matias Scavuzzo^{1,2,3*}, Cintia Delgado¹, Marcia Goy⁴, Favio Crudo¹, Ximena Porcasi^{2,3}, María Victoria Periago^{1,3}

1 Fundación Mundo Sano, Buenos Aires, Argentina, **2** Instituto de Altos Estudios Espaciales Mario Gulich, Universidad Nacional de Córdoba, Comisión Nacional de Actividades Espaciales, Córdoba, Argentina, **3** Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina, **4** Hospital Dr. Dante Tardelli, Pampa del Indio, Chaco, Argentina

* matiasscavuzzo@fcm.unc.edu.ar



OPEN ACCESS

Citation: Scavuzzo CM, Delgado C, Goy M, Crudo F, Porcasi X, Periago MV (2023) Intestinal parasitic infections in a community from Pampa del Indio, Chaco (Argentina) and their association with socioeconomic and environmental factors. PLoS ONE 18(6): e0285371. <https://doi.org/10.1371/journal.pone.0285371>

Editor: Clement Ameh Yaro, University of Uyo, NIGERIA

Received: July 14, 2022

Accepted: April 21, 2023

Published: June 29, 2023

Copyright: © 2023 Scavuzzo et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data and methods generated or analyzed during this study are included in this published article.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Neglected tropical diseases are a group of 20 disabling diseases, which, in particular, are the most common chronic infections in the most vulnerable people. This study aimed to characterize the infection by intestinal parasites (IPs) in dwellings from a peri-urban neighborhood in Pampa del Indio, Chaco (Argentina), and its association with socioeconomic and environmental variables. Single stool samples were collected from all individuals older than 1 year through household visits and processed using coprological sedimentation and flotation techniques. Standardized questionnaires were used at the household level to collect socio-economic information. Environmental variables were obtained from the Planetscope image, Landsat 8 images and remote sensors, while land-use layers were obtained through the use of a maximum likelihood algorithm. Stool samples were provided by 314 individuals. The prevalence of IPs found was 30.6% (n = 96), with a predominance of *Giardia lamblia* (12.7%, n = 40) and *Hymenolepis nana* (7.6%, n = 24). The only soil-transmitted helminth found was *Strongyloides stercoralis* with a 2.5% prevalence (n = 8). Individuals of adult age (> 18 years) were 0.65 times less likely to present parasitic infections with respect to children and adolescents. The only environmental variable that was closely associated with the presence of IPs, was the Normalized Difference Water Index (NDWI), a measure of humidity; being higher around houses with positive individuals. Most of the IPs found in this study were of water-borne transmission and those transmitted directly from person-to-person, therefore fecal contamination is present. We believe that the low prevalence of STH in this area, which requires a passage through the soil, is related to the environmental characteristics, which are unsuitable for the development/permanence of the infective stages of these parasites. The geospatial data and tools used herein proved to be useful for the study of the relationship between the different factors that influence the presence of IPs in a community, from an eco-health approach.

Introduction

Neglected tropical diseases (NTDs) are a group of 20 disabling diseases recognized by the World Health Organization (WHO), which are the most common chronic infections in the most vulnerable people [1–4]. These include soil-transmitted helminths (STH), such as *Ascaris lumbricoides*, *Trichuris trichiura*, and hookworm, which are intestinal parasites (IPs). Other IPs are the protozoans, like *Giardia intestinalis*, *Entamoeba histolytica*, and *Cryptosporidium* spp. [5]), which can also limit the health and nutritional status of their hosts, leading to iron deficiency anemia, growth and cognitive retardation [6–11]. A separate but equally important case to consider is the STH *Strongyloides stercoralis*, given its worldwide prevalence and burden [12]. Although this species was not originally contemplated within the control strategy designed by the WHO for STH, due to specific requirements for diagnosis and treatment; it was included in 2021 [13].

Evidence shows that NTDs, poverty, and certain combinations of ecological, social, political, and economic determinants are strongly correlated, and these factors explain the emergence of hotspots worldwide, especially in developing countries and vulnerable communities [14–16]. Although IPs as a group are not included in the list of NTDs, only STHs are, other IPs have also been found to be associated with similar determinants. Research conducted among urban, peri-urban, and rural populations in Argentina have observed different prevalence rates of IPs, ranging from 0.5 to 88.9%, according to socioeconomic level, sanitary and environmental conditions, and water supply [17–32], with a predominance of STHs in northeast and northwest provinces of the country. Nonetheless, despite having STH prevalence greater than 20% in some areas of the country, mass drug administration (MDA) programs were implemented for a very short period of time (2005–2007) and then discontinued [33].

An in-depth analysis of the complex systems involved in infectious diseases, as well as their causes and consequences, requires more integrative paradigms, such as the ecosystem approach to human health (eco-health or one-health), which incorporates ecological, biological, and social factors as well as their possible interactions [34]. One tool that is useful for this type of approach is the concept of landscape epidemiology, which is widely used in the area of geomatics and remote sensing (RS) to refer to environmental conditions such as land cover, land use, and composition, climatic and geographic characteristics, among others [35–38]. A great variety of environmental conditions can be readily obtained from RS (satellite images and products), making them useful in various fields, such as the preparation of risk maps to guide health effectors and prioritize resources for different infectious diseases [39–43].

Cross-sectoral collaboration, including education, nutrition, and agriculture, has strengthened the control of certain infectious diseases [44]. Working to overcome the impact of many infectious diseases, especially those that affect children, represents a largely untapped development opportunity to alleviate poverty for many populations and thus have a direct impact on achieving international collaborative agendas such as the Sustainable Development Goals (SDGs) or the new roadmap for NTDs [44, 45]. The province of Chaco in Argentina is part of the Gran Chaco Ecoregion, which is considered a hotspot for NTDs, especially Chagas Disease (ChD) [14, 46]. The city of Pampa del Indio has been extensively studied with respect to ChD [47], with a focus on vector surveillance and control, but there is a lack of published studies on IPs and deworming campaigns have not been implemented in this area.

Given the relationship between the environment, humans, certain social determinants and the transmission of IPs, this study aims to identify the IP infections present in individuals from a neighborhood of Pampa del Indio, Chaco, Argentina; and determine if there are any associations between the presence of IPs and different socioeconomic and environmental variables. For this purpose, we have collected socioeconomic data through a household

questionnaire, parasite presence through the analysis of fecal samples; and land-use and environmental indexes derived from remote sensed data that potentially play a role in the presence of IPs as determined in previous studies [24, 43, 47–51].

Materials and methods

Study area

Fieldwork was conducted in a peri-urban neighborhood of Pampa del Indio (Lat: -26.0473; Long: -59.9416), Chaco province, northeastern Argentina, located in the transition between the humid and dry Chaco region. The municipality was inhabited by approximately 22,000 people in 2013. Official records from the 2001 and 2010 decennial census indicated that the population of Pampa del Indio municipality increased markedly from 11,558 to about 18,000 people, respectively (annual population growth rate, 4.9%) [52, 53].

The climate is currently continental, mostly warm, with more rainfall in the summer. The average annual temperature is 22.8 °C (average minimum and maximum, 16.9 and 29.3 °C). Annual rainfall has historically been 954 mm. The landscape is flat and consists mainly of a mosaic of crop patches mixed with the native dry forest that has undergone varying degrees of degradation, and occasional water bodies and swamps [54–56]. Houses included in this study are located on the periphery of Pampa del Indio, in the peri-urban neighborhood of Parque Industrial, an area surrounded by agricultural fields mixed with patches of native forest subject to varying degrees of degradation (Fig 1). Based on this characterization, this neighborhood was selected as a first step to determine the presence of IPs in the area since there was no previous data on these infections in the population.

Study design

The study was designed as a cross-sectional study which was conducted between June 2016 and January 2017. Parque Industrial is composed of 22 blocks and houses were randomly selected assuring at least four houses from each block were included, one from each of the four sides of each block. Each house was georeferenced and characteristics of each household, as well as demographic data from each of the inhabitants, was collected through the use of a standardized questionnaire. The questionnaire was divided into different sections and covered aspects related to education, occupation, household characteristics (including source of drinking water, type of toilet, as well as construction materials) and the presence/ownership of domestic animals.

If a household declined to participate, the next household on that same side of the block was visited. Sterile and leak-proof wide-cap containers without any fixative were distributed to all the members of the families which showed interest in participating as evidenced through the signed consent/assent. During the initial visit, instructions were given on the ideal way to collect stool samples, by defecating on a clean surface (bag or paper), without contact with water, urine or dirt and then with a wooden spatula to place a good amount of sample into the container. Participants were told the team would return on the following day to retrieve the samples and that they were to be kept in a shaded area of the house. A single stool sample was collected for each individual and houses were visited at least on three occasions in order to give individuals time to produce the sample. The samples were transported in a cool box to the Clinical Analysis Laboratory of the public hospital “Dr. Dante Tardelli” of Pampa del Indio, where all the samples were analyzed macroscopically for the presence of parasitic forms and then processed by three different concentration methods including sedimentation (Telemann), salt flotation (Willis), and sugar flotation (Sheather), as previously described [57]. Infected individuals were clinically evaluated by the project physician and treatment was




Fig 1. Study area in Pampa del Indio, Chaco (Argentina). The neighborhood included in this study was a peri-urban area called Parque Industrial. Map data © 2020 Google, base map obtained through QuickMapServices QGIS plugin—QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>. Contains information from OpenStreetMap and OpenStreetMap Foundation, which is made available under the Open Database License.

<https://doi.org/10.1371/journal.pone.0285371.g001>

provided by the project following national guidelines from the National Ministry of Health [58] which indicate treatment with mebendazole or metronidazole depending on the IP.

Ethical considerations

This study was approved by the Institutional Review Board (IRB) of the National Center of Medical Genetics (Centro Nacional de Genética Médica), Administration of Laboratories and Institutes of Health (Administración Nacional de Laboratorios e Institutos de Salud—ANLIS) “Dr. Carlos G. Malbrán” of the National Ministry of Health of Argentina (protocol approved unanimously 27 May 2016). Written informed consent was obtained from all participating adults and from a parent or guardian of every child under 16 years of age. Moreover, written assent was obtained from children between the ages of 6 and 15 years of age, inclusive.

Criteria for inclusion in the study included being older than 1 year and having signed the appropriate informed consents and assents. Exclusion criteria included working or living outside the area of study for more than a week at a time or having behavioral, cognitive or psychiatric issues that may affect the ability to understand and adhere to the study protocol.

Deworming is not routinely performed in this area, therefore this was not considered as criteria for exclusion.

Environmental characteristics

The land-use map of the study area was obtained using the Planetscope image (with a spatial resolution of 3.5 m; radiometrically and geometrically calibrated) [59]. Using a maximum likelihood algorithm, a land-use layer regarding 4 classes based on updated literature [60] and the relationship with parasitic transmission cycles was obtained. These classes (C) were: C1) bushy or thicker vegetation, C2) low and sparse vegetation, C3) bare soil, and C4) urban construction. For the image classification, training (20 pixels for each class) and validation points were obtained from Airbus images. Finally, a black and white image (in a dichotomous/binary manner) was generated for each class separately and the statistics of the neighborhoods of each house were extracted.

Additionally, several spectral indexes to characterize the environment surrounding each house were generated using Landsat 8 image collections from the Google Earth Engine platform [61, 62]. These environmental variables were calculated as the annual averages of year 2016; and are commonly used in spatial epidemiology and disease mapping [88]: Normalized Vegetation Index (NDVI) as a proxy of vegetation cover, Normalized Difference Water Index (NDWI) as a proxy of wetness, Normalized Snow Differential Index (NDSI) as a proxy for soil characteristics, and Normalized Burnt Area Index (NBRT) as a proxy for soil temperature, were included.

To extract the indices from the vicinity of each house, a buffer of 50 m around each house was used through zone statistics tools provided by Qgis software; spatial and environmental analysis was performed with open source Qgis software version 3.10 [63]. Descriptive and exploratory statistical analyses were performed with Stata software version 15.1 [64]. Moreover, in order to evaluate whether there were significant differences among means (quantitative) and proportions (categorical) between sociodemographic characteristics and the presence of intestinal parasites, a t-test was used.

Spatial analysis

To visualize the spatial dimension of infection by IPs, the number of cases per household was represented through heat maps. Later, a spatial cluster analysis was performed with Satscan software [65] using a Bernoulli distribution model to explore the distribution of households with infected and non-infected individuals. If spatial clusters were detected, the environmental indexes obtained through RS for the households within the significant cluster were compared with those of the households outside the cluster, using independent mean tests.

To determine the association of the different variables and the presence or absence of IPs, logistic regression was used. Through initial bivariate logistic regression models, the relationship between different variables were tested in order to discard covariates and proceed with the multivariate logistic regression. Finally, a negative binomial model with robust deviancy, using the quasi maximum likelihood method was used to model the risk of infection of IPs.

Results

A total of 127 dwellings were surveyed and 525 containers were distributed throughout the study period. Participation was 59.8%, given that 314 individuals from 107 of the households provided stool samples for analysis. IPs were found in 96 individuals (30.6%), many of them polyparasitized with more than one species ($n = 35$, 11.1%), not all of them pathogenic

Table 1. Descriptive prevalence of intestinal parasites in inhabitants from Parque Industrial, Pampa del Indio, Chaco (Argentina), 2016–2017.

Parasitological description (N = 314 individuals)	
	No. (%)
Positive	96 (30.6)
Negative	218 (69.4)
Mono-infections	61 (19.5)
<i>Endolimax nana</i> ¹	10 (3.2)
<i>Entamoeba coli</i> ¹	13 (4.1)
<i>Blastocystis</i> spp. ¹	4 (1.3)
<i>Chilomastix mesnili</i> ¹	3 (1.0)
<i>Giardia lamblia</i>	17 (5.4)
<i>Enterobius vermicularis</i>	3 (1.0)
<i>Hymenolepis nana</i>	8 (2.5)
<i>Strongyloides stercoralis</i>	3 (1.0)
Double infections	28 (8.9)
<i>E. coli</i> ¹ / <i>E. nana</i> ¹	3 (1.0)
<i>E. nana</i> ¹ / <i>C. mesnili</i> ¹	1 (0.3)
<i>G. lamblia</i> / <i>Blastocystis</i> spp. ¹	1 (0.3)
<i>G. lamblia</i> / <i>E. coli</i> ¹	3 (1.0)
<i>G. lamblia</i> / <i>E. nana</i> ¹	5 (1.7)
<i>G. lamblia</i> / <i>Hymenolepis nana</i>	7 (2.2)
<i>H. nana</i> / <i>C. mesnili</i> ¹	2 (0.6)
<i>H. nana</i> / <i>E. coli</i> ¹	1 (0.3)
<i>H. nana</i> / <i>E. nana</i> ¹	1 (0.3)
<i>S. stercoralis</i> / <i>C. mesnili</i> ¹	1 (0.3)
<i>S. stercoralis</i> / <i>E. nana</i> ¹	1 (0.3)
<i>S. stercoralis</i> / <i>H. nana</i>	2 (0.6)
Triple-infections	7 (2.2)
<i>G. lamblia</i> / <i>E. coli</i> ¹ / <i>E. nana</i> ¹	3 (1.0)
<i>G. lamblia</i> / <i>H. nana</i> / <i>E. coli</i> ¹	1 (0.3)
<i>G. lamblia</i> / <i>H. nana</i> / <i>E. nana</i> ¹	1 (0.3)
<i>E. vermicularis</i> / <i>H. nana</i> / <i>G. lamblia</i>	1 (0.3)
<i>S. stercoralis</i> / <i>G. lamblia</i> ¹ / <i>E. nana</i> ¹	1 (0.3)

No.: Number.

¹Non-pathogenic species for which the patient is clinically evaluated to determine if treatment is needed, depending on medical judgment. We have included *Blastocystis* spp. as a non-pathogenic species given there is still controversy and subtyping of this protist was not an aim of the current study [47].

<https://doi.org/10.1371/journal.pone.0285371.t001>

(Table 1). The only STH species found was *S. stercoralis*; this parasite was detected in 8 individuals (2.5%).

Data collected through the standardized questionnaires showed that only 4% of the participants were employed in agriculture, and only 7% had a vegetable garden, while the water for irrigation came from the public water supply. Almost all of the households (95%), had access to gas, either natural or bottled, although the smallest proportion was supplemented with firewood (6.5%), while all households had access to the electrical network. Also, 100% reported that the source of water for drinking, cooking, and handwashing was from the public drinking water network and none of them treated the water at home before drinking it (either by boiling

or adding drops of bleach). With respect to excreta disposal, 100% had a latrine and 85% were improved with septic tank drainage. With respect to animals, 74% had at least 2 dogs and 42% had never dewormed them. It was noted that 76% of the households do not have cats. In the same sense, 98% did not raise animals and no one had animal pens.

Table 2 shows the description of the sample according to the presence or absence of IPs. Variables were analyzed both at the individual and household level, given that family income and characteristics of the household are common to all the inhabitants of each house. At the individual level, there was a higher proportion of infected children and adolescents (67.7%), 1 to 17 years of age, in comparison to adults (32.3%), older than 18 years. There was also a higher percentage of infected individuals with incomplete primary education (63.5%), although these differences were not statistically significant. From the 314 participants, 52 individuals reported having an intestinal parasitic infection on previous occasions, of which 71.2% ($n = 37$) were treated. With respect to the characteristics at the household level, no statistically significant differences were found between the proportions of houses with any positive cases and negative houses without infection.

The analysis of the study area with respect to the environmental variables (Fig 2) shows that the most frequent land cover class was C2 (low and sparse vegetation), followed by C3 (bare soil) and some isolated patches of C1 (bushy or thicker vegetation). This analysis also shows that most of the households were located adjacent to or on bare soil (C3) or areas with urban construction (C4). Table 3 shows the association between the environmental variables around the households with and without individuals infected with IPs. NDWI was the environmental variable more closely associated with infected individuals ($p = 0.05$); showing lower humidity around their homes.

For the spatial analysis, a total of 69 households were included given that not all households were geolocalized, this comprised a total of 302 inhabitants. This analysis showed a spatial pattern of clustering in a north-south direction, with a higher proportion of positive individuals in the northern part of the neighborhood. The output of the spatial cluster analysis presented one significant cluster covering an area of 0.16 km, including 17 households with 73 individuals. In this cluster, according to the quasi maximum likelihood method, there should be 23 individuals with IPs, but the number of observed cases was actually 11, thus determining a ratio of 0.48 and establishing a relative risk of 0.41 with a spatial prevalence of 15.1% (Fig 3). The spatial cluster with the lowest significant presence of cases (in light green) is observed in Fig 3, containing a greater proportion of houses with fewer relative risk. These houses within the low risk cluster for IPs had a significantly higher NDVI ($p < 0.01$) and NDWI ($p < 0.01$) than those houses outside the cluster; this is probably due to the large adjacent area to the left of the cluster presenting thicker vegetation. With respect to the land use classes, the houses within this lower risk cluster fell under class C2 (low and sparse vegetation ($p < 0.01$)).

The analysis of the variables and the output of the logistic regression (Table 4), shows the crude odds ratios (ORs) that were statistically significant or close to significance in the bivariate models and their 95% confidence intervals (CI). Both income from a retirement fund or pension, as well as age were significantly associated with the absence of IPs, supporting the presence of these variables in a subsequent multivariate logistic model. As observed, those individuals older than 18 years had 0.6 times less chance (protective factor) of presenting IPs in comparison to children and adolescents. Similarly, a person receiving a retirement fund or pension had 0.44 times less chance of being infected with IPs. These two variables, age and pension fund, showed a statistically significant adjusted interaction (OR, $p = 0.03$), showing to be a protective factor, since the risk of infection with IPs is decreased by 0.24 (95% CI: 0.06–0.87). The environmental variables NDWI and NDVI were also included in the multivariate regression given their near significance ($p = 0.05$).

Table 2. Sociodemographic characteristics of the study population, according to presence or absence of parasites, Parque Industrial neighborhood, Pampa del Indio, Chaco, 2016–2017.

	Complete sample	Infected	Non-infected	p-value for the difference in proportions (T-test) between individuals with and without infection
Individual samples	n = 314	n = 96	n = 218	
Age (years), [n (%)]				
Children and adolescents (1–17)	191 (60.8)	65 (67.7)	126 (57.8)	0.09
Adults (> = 18)	123 (39.2)	31 (32.3)	92 (42.2)	0.09
Gender [n (%)]				
Female	180 (57.3)	53 (55.2)	127 (58.3)	0.61
Male	134 (42.7)	43 (44.8)	91 (41.7)	0.61
Education [n (%)]				
Incomplete elementary school	209 (66.5)	61 (63.5)	148 (67.9)	0.54
Finished elementary school	77 (24.5)	26 (27.1)	51 (23.4)	0.30
Finished middle school	14 (4.5)	6 (6.3)	8 (3.7)	0.32
No data	14 (4.5)	3 (3.1)	11 (5.0)	
Parasitic antecedents [n (%)]				
Yes	52 (16.6)	17 (17.7)	35 (16.1)	0.96
No	191 (60.8)	63 (65.6)	128 (58.7)	0.96
No data	71 (22.6)	16 (16.7)	55 (25.2)	
Previous treatment [n (%)]				
Yes	37 (71.2)	10 (58.8)	27 (77.1)	0.54
No	5 (9.6)	2 (11.8)	3 (8.6)	0.54
No data	10 (19.2)	5 (29.4)	5 (14.3)	
Households sampled	N = 107	N = 49	N = 43	
Household income [n (%)]				
Day laborer				
Yes	56 (55.4)	24 (53.3)	24 (58.5)	0.71
No	45 (44.6)	21 (46.6)	17 (41.5)	0.72
Social plan beneficiaries				
Yes	68 (63.5)	31 (63.2)	26 (60.4)	0.82
No	33 (30.8)	14 (28.6)	15 (34.8)	0.72
No data	6 (5.6)	4 (8.1)	2 (4.6)	
Receive retirement fund or pension				
Yes	16 (14.9)	4 (8.1)	9 (20.9)	0.57
No	85 (79.4)	41 (83.7)	32 (74.4)	0.32
No data	6 (5.61)	4 (8.2)	2 (4.6)	
Roof [n (%)]				

(Continued)

Table 2. (Continued)

	Complete sample	Infected	Non-infected	p-value for the difference in proportions (T-test) between individuals with and without infection
Metal sheets	99 (92.5)	44 (89.8)	40 (93.2)	0.57
Adobe and wood	8 (7.4)	5 (10.1)	3 (6.9)	0.87
Floor [n (%)]				
Concrete	71 (66.3)	33 (67.5)	29 (67.4)	0.99
Concrete and adobe	21 (19.6)	9 (18.4)	6 (13.9)	0.81
Ceramic	9 (8.4)	3 (6.1)	6 (13.9)	0.72
No data	6 (5.6)	4 (8.2)	2 (4.6)	
Wall [n (%)]				
Adobe and wood	21 (19.8)	9 (18.3)	6 (14.3)	0.83
Bricks	79 (74.5)	36 (73.5)	34 (80.9)	0.46
No data	6 (5.7)	4 (8.2)	2 (4.8)	

<https://doi.org/10.1371/journal.pone.0285371.t002>




Fig 2. Environmental information derived from RS data: The land-use classification and Landsat environmental indices. Map data © 2020 Google, base map obtained through QuickMapServices QGIS plugin—QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>. Contains information from OpenStreetMap and OpenStreetMap Foundation, which is made available under the Open Database License.

<https://doi.org/10.1371/journal.pone.0285371.g002>

Table 3. Summary of the environmental characteristics in the immediate vicinity of the households with and without individuals infected with intestinal parasites from Parque Industrial, Pampa del Indio, Chaco.

	Environmental characteristics in the immediate vicinity of households with infected individuals	Environmental characteristics in the immediate vicinity of households without infected individuals	p-value for the difference in means
Mean of NDVI [Me.(SD)] [CI 95%]	0.388 (0.005) [0.377–0.400]	0.400 (0.003) [0.393–0.408]	0.08
Mean of NDWI [Me.(SD)] [CI 95%]	0.034 (0.004) [0.026–0.043]	0.044 (0.002) [0.038–0.049]	0.05
Mean of NDSI [Me.(SD)] [CI 95%]	-0.374 (0.002) [-0.379- -0.368]	-0.372 (0.001) [-0.375- -0.369]	0.69
Mean of NBRT [Me.(SD)] [CI 95%]	0.957 (0.0006) [0.956–0.958]	0.958 (0.0003) [0.957–0.958]	0.20
Mean of C1: urban areas [Me.(SD)] [CI 95%]	0.340 (0.012) [0.316–0.364]	0.322 (0.007) [0.310–0.341]	0.30
Mean of C2: bare soil [Me.(SD)] [CI 95%]	0.295 (0.008) [0.278–0.312]	0.308 (0.006) [0.295–0.321]	0.24
Mean of C3: vigorous vegetation [Me.(SD)] [CI 95%]	0.054 (0.006) [0.041–0.066]	0.053 (0.004) [0.045–0.062]	0.96
Mean of C4: less dense and sparse vegetation [Me.(SD)] [CI 95%]	0.310 (0.008) [0.293–0.327]	0.312 (0.005) [0.301–0.323]	0.85

Me.: Mean; SD: Standard Deviation; CI: Confidence Interval; NDVI: Normalized Difference Vegetation Index; NDWI: Normalized Difference Water Index; NDSI: Normalized Difference Snow Differential Index; NBRT: Normalized Burnt Area Index. Land cover classes: C1 = bushy or thicker vegetation, C2 = low and sparse vegetation, C3 = bare soil and C4 = urban construction.

<https://doi.org/10.1371/journal.pone.0285371.t003>

The negative binomial regression model, using the number of infected individuals per household as the independent variable, presented very good performance metrics (prob >Chi² = 0.000; Pseudo R² MacFadden = 0.239; R² = 0.56; Prob >LR = 0.02) and included 9 of the 16 variables collected in the study as predictor variables (Table 5). The model predicts a mean of 1.4 infected per household (SD +/-0.5, max: 8 persons), similar to the real values. A normal and zero-centered distribution of residuals was obtained. This result can be understood as a complementary validation of the model's performance to the analyzed dataset.

Discussion

In this study, a public health problem, the presence of IPs, was addressed using RS as a novel tool in a neighborhood of Pampa del Indio, Chaco (Argentina); evidencing the useful approaches these geospatial tools can provide to landscape epidemiology for health [39–43]. There is a lack of data on the prevalence of IPs in Chaco province, except for two small studies conducted in Resistencia, the capital city, more than 18 years ago [66, 67], and another study conducted in rural areas, adjacent to Pampa del Indio, in 2018 [21]. These previous studies found IPs, both protozoan and helminths species, while STH were found only in the studies from Resistencia. Therefore, this is the first report of the presence of IPs in the city of Pampa del Indio.

The total prevalence of IPs found in this study was 30.6% (n = 96), with 35 individuals presenting multiple infections (11.1%). Two of the most prevalent parasites were *G. lamblia* (12.7%, n = 40) and *H. nana* (7.6%, n = 24). Overall, the prevalence of STH was low, with a total of 8 participants infected with *S. stercoralis* (2.5%), this species was also detected in the studies from Resistencia with similar prevalences, although in those studies, they also detected hookworm, *A. lumbricoides* and, to a lesser extent, *T. trichiura* [66, 67]. Given that a specific method for *S. stercoralis*, such as agar plate or Baermann, was not used, it might have been




Fig 3. Purely spatial pattern analysis performed with SaTScan™ software, version 5.1.3. Map data © 2020 Google, base map obtained through QuickMapServices QGIS plugin—QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>. Contains information from OpenStreetMap and OpenStreetMap Foundation, which is made available under the Open Database License.

<https://doi.org/10.1371/journal.pone.0285371.g003>

underdiagnosed. Moreover, since this STH can be maintained in an individual for many years due to autoinfection and lack of treatment, these individuals might have gotten infected elsewhere.

The low prevalence of STH in this region of Chaco, as evidenced by the current study and a study conducted in adjacent rural areas [21], might be related to the environmental variables of the area, which, as observed in a previous study from Santiago de Estero (Argentina), are not

Table 4. Risk of intestinal parasites: Bivariate logistic models at the individual level in a neighborhood from Pampa del Indio, Chaco (Argentina) during 2016–2017.

	OR	P-value	CI (95%)
Household income from retirement fund or pension	0.44	0.04	0.18–0.99
Age (years): Adults (18 or more)	0.65	0.03	0.39–0.95
NDWI: Normalized Difference Water Index	0.002	0.05	0.002–1.02
NDVI: Normalized Difference Vegetation Index	0.02	0.05	0.0002–1.08

OR: Odds Ratio; CI: Confidence Interval.

<https://doi.org/10.1371/journal.pone.0285371.t004>

Table 5. Components and influence of the negative binomial risk model (number of infected individuals per household) in a neighborhood of Pampa del Indio, Chaco (Argentina) during 2016–2017.

Predictor variables	Modification in risk prediction of people infected per household
Parasitological variables	
Previous infection	decreases
Previous parasitic treatment	decreases
Socioeconomic variables	
Economic income (day laborer)	decreases
Building characteristics (ceramic floor)	decreases
Building characteristics (dirt floor)	increases
Environmental variables	
NBRT (proxy of temperature)	decreases
Class 1 (urban area)	decreases
Class 2 (bare soil)	decreases
Class 3 (dense vegetation)	decreases
Class 4 (low and sparse vegetation)	increases

NBRT: Normalized Burnt Area Index

<https://doi.org/10.1371/journal.pone.0285371.t005>

conducive to the maintenance of the infective stages of these parasites in the soil [57]. Despite the presence of improved water and sanitation in the neighborhood studied, the individuals had a 30.6% prevalence of other IPs, evidencing the presence of fecal contamination, yet the only STH detected was *S. stercoralis*. Regarding the land use at the immediate vicinity of individual households, a higher proportion of people infected with IPs (except STH) live in an area with sparse vegetation and with households located adjacent to or on bare soil or urban area. The average Normalized Difference Vegetation Index (NDVI) and Normalized Difference Water Index (NDWI) in the vicinity of those dwellings was 0.39, and 0.03 respectively; both values are lower than vicinities without IP infections. This could explain the low prevalence of STH which are more sensitive to the external environment. Previous studies have shown that the characteristics of the area we describe in this study are not conducive to the development of STH with respect to precipitation, soil, temperature, and vegetation cover [43, 68–77]. The environmental characteristics of the study area and the presence of improved latrines in 85% of the households seem to act as protective factors with respect to STH infection.

The use of a detailed resolution of 0.5 m for the land use product, proved to be useful for the risk analysis presented herein (Table 5), since it allowed visualization of the spatial co-occurrence of four land covers (urban, bare soil, as well as both sparse and vigorous vegetation) in a very small peri-urban area. The analysis showed that all four classes were significantly associated with the number of infected individuals per household, allowing the study of the environment-parasite relationship at a very fine scale.

Although the prevalence of STH was low, the presence of other IPs is evident, especially those which are waterborne, like *Giardia* spp., and those that can be transmitted directly from person to person, i.e. *H. nana*. Moreover, the prevalence of infection between children and adults was significantly different, being higher in children; individuals of adult age had 0.65 times less chance of presenting an intestinal parasite in comparison to children and adolescents. This agrees with previous reports [51, 57, 78, 79] and is linked to behavioral differences between children and adults, since they are more likely to play in the soil and have lower hygienic standards [80–83]. Additionally, the socioeconomic characteristics of the dwellings and the physical environment (economic income and floor type) were also associated with the number of infected individuals in this neighborhood from Pampa del Indio which is composed

mostly of material houses in a peri-urban area, with access to a water network, an electrical network and latrines.

Access to clean water and sanitation has been previously associated with IP infection [84–90]. In this study area, even though all the participants declared obtaining drinking water from the public water network, there were many individuals infected with amoebae and/or *Giardia* spp., evidencing either a sub-optimal quality of the water for drinking and cleaning of fruits and vegetables, due to either contamination from the water treatment plant to the households, contamination during storage of water in the household, or to improper functioning of the water treatment plant. The infection with *Giardia* spp. could also be due to zoonotic genotypes, as previously found in an area from Puerto Iguazú (Misiones, Argentina) [79] and given that many participants had domestic animals in their households. Nonetheless, this aspect escaped the aim of the current study.

The geospatial data and tools used in this study proved to be useful for the study of the relationship between the different factors that influence the presence of IPs in a community, from an eco-health or one-health approach. Moreover, most of the geospatial data and tools used are freely available and obtained by RS; this is very relevant in the field of epidemiology when it comes to balancing resources spent on data collection in the field. The risk analysis model used in this study shows that the dimension that most influences the prediction of the model are the environmental variables, specifically those coming from the classification carried out to validate the criteria established to differentiate the classes. Precisely these environmental characteristics (soil, vegetation, and humidity) have been previously associated with the presence of STH in Argentina at the national scale [43].

A limitation of this study, although the sample size was representative of the neighborhood Parque Industrial of Pampa del Indio, is that it may not be extrapolated to the entire city; therefore, the results were worked at a very small scale, both at the household level and spatially. As mentioned above, another limitation specific for the presence of *S. stercoralis* is that a coprological technique specific for this parasite was not used and therefore the prevalence might be underestimated.

Conclusion

Most of the IPs found in this study were of water-borne transmission and those transmitted directly from person-to-person which is usually associated with quality of the water used for drinking and food manipulation, hygiene and overcrowding. There was a low prevalence of STH in this area, despite a moderate prevalence of other IPs, with only a few cases of *S. stercoralis*. We believe that this is related to the environmental characteristics of the area, which are unsuitable for the development/permanence of the infective stages of STHs in the soil, given that deworming is not routinely performed in this area and fecal contamination is occurring.

The spatial analysis of the IPs found showed a marked north to south distribution of IPs with a predominant low-risk cluster in the south. Given that sometimes it's difficult to obtain large scale epidemiological data, it is relevant to generate models with local and geospatial data that can learn from free and open access data sources to estimate an accurate probability of occurrence, providing a better understanding of risk and prioritizing situations of greater vulnerability for evidence-based, strategic, and effective implementation of public policies in general, and health policies in particular.

Acknowledgments

We would like to thank Fundación Mundo Sano (Pampa del Indio site) for its collaboration and support in this study. We are most grateful to the community leaders for facilitating

participation and contact with the community. We would also like to thank Marcelo Abril for his institutional support, as Executive Director of Fundación Mundo Sano.

Author Contributions

Conceptualization: Carlos Matias Scavuzzo, María Victoria Periago.

Data curation: Carlos Matias Scavuzzo, María Victoria Periago.

Formal analysis: Carlos Matias Scavuzzo, Ximena Porcasi, María Victoria Periago.

Funding acquisition: Carlos Matias Scavuzzo, María Victoria Periago.

Investigation: Carlos Matias Scavuzzo, Cintia Delgado, Marcia Goy, Favio Crudo, María Victoria Periago.

Methodology: Carlos Matias Scavuzzo, Ximena Porcasi, María Victoria Periago.

Project administration: Carlos Matias Scavuzzo, María Victoria Periago.

Resources: Carlos Matias Scavuzzo, Marcia Goy, Favio Crudo, María Victoria Periago.

Software: Carlos Matias Scavuzzo, Ximena Porcasi, María Victoria Periago.

Supervision: Carlos Matias Scavuzzo, Ximena Porcasi, María Victoria Periago.

Validation: Carlos Matias Scavuzzo, Ximena Porcasi, María Victoria Periago.

Visualization: Carlos Matias Scavuzzo, Ximena Porcasi, María Victoria Periago.

Writing – original draft: Carlos Matias Scavuzzo, María Victoria Periago.

Writing – review & editing: Carlos Matias Scavuzzo, Cintia Delgado, María Victoria Periago.

References

1. Molyneux DH. "Neglected" diseases but unrecognised successes challenges and opportunities for infectious disease control. *Lancet*. 2004; 364(9431):380–3. Available from: [http://dx.doi.org/10.1016/S0140-6736\(04\)16728-7](http://dx.doi.org/10.1016/S0140-6736(04)16728-7) PMID: 15276399
2. Molyneux DH, Hotez PJ, Fenwick A. "Rapid-impact interventions": how a policy of integrated control for Africa's neglected tropical diseases could benefit the poor. *PLoS Medicine*. 2005; 2(11):e336. Available from: <http://dx.doi.org/10.1371/journal.pmed.0020336> PMID: 16212468
3. Hotez P, Ottesen E, Fenwick A, Molyneux D. The neglected tropical diseases: the ancient afflictions of stigma and poverty and the prospects for their control and elimination. *Advances in Experimental Medicine and Biology*. 2006; 582:23–33. Available from: http://dx.doi.org/10.1007/0-387-33026-7_3 PMID: 16802616
4. Hotez PJ, Molyneux DH, Fenwick A, Ottesen E, Ehrlich Sachs S, Sachs JD. Incorporating a rapid-impact package for neglected tropical diseases with programs for HIV/AIDS, tuberculosis, and malaria. *PLoS Medicine* [Internet]. 2006; 3(5):e102. Available from: <http://dx.doi.org/10.1371/journal.pmed.0030102> PMID: 16435908
5. Haque R. Human intestinal parasites. *Journal of Health Population and Nutrition*. 2007; 25(4):387–91. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2754014/> PMID: 18402180
6. Pawlowski Z. S., Schad G. A., Stott G. J., & World Health Organization. (1991). Hookworm infection and anaemia: approaches to prevention and control. World Health Organization. <https://apps.who.int/iris/handle/10665/40857>
7. Hall A, Hewitt G, Tuffrey V, de Silva N. A review and meta-analysis of the impact of intestinal worms on child growth and nutrition. *Maternal and Child Nutrition*. 2008; 4(s1):118–236. Available from: <http://dx.doi.org/10.1111/j.1740-8709.2007.00127.x> PMID: 18289159
8. Crompton DWT, Nesheim MC. Nutritional impact of intestinal helminthiasis during the human life cycle. *Annual Review of Nutrition*. 2002; 22(1):35–59. Available from: <http://dx.doi.org/10.1146/annurev.nutr.22.120501.134539> PMID: 12055337
9. Joseph SA, Casapia M, Montresor A, Rahme E, Ward BJ, Marquis GS, et al. The effect of deworming on growth in one-year-old children living in a soil-transmitted helminth-endemic area of Peru: A

- randomized controlled trial. PLoS Neglected Tropical Diseases. 2015; 9(10):e0004020. Available from: <http://dx.doi.org/10.1371/journal.pntd.0004020> PMID: 26426270
10. Liu C, Lu L, Zhang L, Luo R, Sylvia S, Medina A, et al. Effect of deworming on indices of health, cognition, and education among schoolchildren in rural China: A cluster-randomized controlled trial. American Journal of Tropical Medicine and Hygiene. 2017;16–0354. Available from: <http://dx.doi.org/10.4269/ajtmh.16-0354> PMID: 28093533
 11. Zavala GA, Rosado JL, Doak CM, Caamaño M del C, Campos-Ponce M, Ronquillo D, et al. Energy and food intake are associated with specific intestinal parasitic infections in children of rural Mexico. Parasitology International. 2017; 66(6):831–6. Available from: <http://dx.doi.org/10.1016/j.parint.2017.07.005> PMID: 28743469
 12. Krolewiecki AJ, Lammie P, Jacobson J, Gabrielli A-F, Levecke B, Socias E, et al. A public health response against *Strongyloides stercoralis*: Time to look at soil-transmitted helminthiasis in full. PLoS Neglected Tropical Diseases. 2013; 7(5):e2165. Available from: <http://dx.doi.org/10.1371/journal.pntd.0002165> PMID: 23675541
 13. WHO World Health Organization. Diagnostic methods for the control of strongyloidiasis: virtual meeting, 29 September 2020. <https://apps.who.int/iris/bitstream/handle/10665/340265/9789240016538-eng.pdf?sequence=1>
 14. Hotez PJ. Ten global “hotspots” for the neglected tropical diseases. PLoS Neglected Tropical Diseases. 2014; 8(5):e2496. Available from: <http://dx.doi.org/10.1371/journal.pntd.0002496> PMID: 24873825
 15. Fernández MDP, Gaspe MS, Gürtler RE. Inequalities in the social determinants of health and Chagas disease transmission risk in indigenous and creole households in the Argentine Chaco. Parasites and Vectors. 2019; 12(1):184. Available from: <http://dx.doi.org/10.1186/s13071-019-3444-5> PMID: 31029147
 16. Hotez PJ, Fenwick A, Molyneux D. The new COVID-19 poor and the neglected tropical diseases resurgence. Infectious Diseases of Poverty. 2021; 10(1):10. Available from: <http://dx.doi.org/10.1186/s40249-020-00784-2> PMID: 33509283
 17. Zonta ML, Navone GT, Oyhenart EE. Parasitosis intestinales en niños de edad preescolar y escolar: situación actual en poblaciones urbanas, periurbanas y rurales en Brandsen, Buenos Aires, Argentina. Parasitología latinoamericana. 2007; 62(1–2). Available from: <http://dx.doi.org/10.4067/s0717-77122007000100009>
 18. Gamboa MI, Navone GT, Orden AB, Torres MF, Castro LE, Oyhenart EE. Socio-environmental conditions, intestinal parasitic infections and nutritional status in children from a suburban neighborhood of La Plata, Argentina. Acta Tropica. 2011; 118(3):184–9. Available from: <http://dx.doi.org/10.1016/j.actatropica.2009.06.015>
 19. Dib J, Oquilla J, Lazarte SG, Gonzalez SN. Parasitic prevalence in a suburban school of Famailá, Tucumán, Argentina. International Scholarly of Research Network Microbiology. 2012; 2012:1–4. Available from: <http://dx.doi.org/10.5402/2012/560376>
 20. Gamboa María I, Giambelluca Luis A, Navone Graciela T. Distribución espacial de las parasitosis intestinales en la ciudad de La Plata, Argentina. Medicina (B. Aires). 2014 Oct [citado 2022 Sep 02]; 74 (5): 363–370. Disponible en: http://www.scielo.org.ar/scielo.php?script=sci_arttext&pid=S0025-76802014000500003&lng=es.
 21. Richards LR, Delgado C, Goy M, Liang S, Periago MV. Prevalence of intestinal parasites and related risk factors in rural localities from Pampa del Indio, Chaco, Argentina. UF Journal of Undergraduate Research. 2019; 21(1). Available from: <http://dx.doi.org/10.32473/ufjur.v21i1.107939>
 22. Rivero MR, De Angelo C, Nuñez P, Salas M, Liang S. Intestinal parasitism and nutritional status among indigenous children from the Argentinian Atlantic Forest: Determinants of enteroparasites infections in minority populations. Acta Tropica. 2018; 187:248–56. Available from: <http://dx.doi.org/10.1016/j.actatropica.2018.08.015> PMID: 30125528
 23. Rivero MR, De Angelo C, Nuñez P, Salas M, Motta CE, Chiaretta A, et al. Environmental and socio-demographic individual, family and neighborhood factors associated with children intestinal parasitoses at Iguazú, in the subtropical northern border of Argentina. PLoS Neglected Tropical Diseases. 2017; 11(11):e0006098. Available from: <http://dx.doi.org/10.1371/journal.pntd.0006098>
 24. Echazú A, Juárez M, Vargas PA, Cajal SP, Cimino RO, Heredia V, et al. Albendazole and ivermectin for the control of soil-transmitted helminths in an area with high prevalence of *Strongyloides stercoralis* and hookworm in northwestern Argentina: A community-based pragmatic study. PLoS Neglected Tropical Diseases. 2017; 11(10):e0006003. Available from: <http://dx.doi.org/10.1371/journal.pntd.0006003>
 25. Cociancic P, Zonta ML, Navone GT. A cross-sectional study of intestinal parasitoses in dogs and children of the periurban area of La Plata (Buenos Aires, Argentina): Zoonotic importance and implications in public health. Zoonoses Public Health. 2018; 65(1):e44–53. Available from: <http://dx.doi.org/10.1111/zph.12408> PMID: 28984036

26. Cimino RO, Jeun R, Juarez M, Cajal PS, Vargas P, Echazú A, et al. Identification of human intestinal parasites affecting an asymptomatic peri-urban Argentinian population using multi-parallel quantitative real-time polymerase chain reaction. *Parasites and Vectors*. 2015; 8(1):380. Available from: <http://dx.doi.org/10.1186/s13071-015-0994-z> PMID: 26183074
27. Echazú A, Bonanno D, Juarez M, Cajal SP, Heredia V, Caropresi S, et al. Effect of poor access to water and sanitation as risk factors for soil-transmitted helminth infection: Selectiveness by the infective route. *PLoS Neglected Tropical Diseases*. 2015; 9(9):e0004111. Available from: <http://dx.doi.org/10.1371/journal.pntd.0004111> PMID: 26421865
28. Barda B, Cajal P, Villagran E, Cimino R, Juarez M, Krolewiecki A, et al. Mini-FLOTAC, Kato-Katz and McMaster: three methods, one goal; highlights from north Argentina. *Parasites and Vectors*. 2014; 7(1):271. Available from: <http://dx.doi.org/10.1186/1756-3305-7-271> PMID: 24929554
29. Zonta ML, Oyhenart EE, Navone GT. Socio-environmental variables associated with malnutrition and intestinal parasitoses in the child population of Misiones, Argentina: Environment, Malnutrition and Enteroparasitoses. *American Journal of Human Biology*. 2014; 26(5):609–16. Available from: <http://dx.doi.org/10.1002/ajhb.22570>
30. Oyhenart EE, Garraza M, Bergel ML, Torres MF, Castro LE, Luis MA, et al. Caracterización del estado nutricional, enteroparasitosis y condiciones socio-ambientales de la población infanto-juvenil del partido de la plata. *Revista argentina de antropología biológica*. <http://www.scielo.org.ar/pdf/raab/v15n1/v15n1a05.pdf>
31. Molina N, Pezzani B, Ciarmela M, Orden A, Rosa D, Apezteguía M, et al. Intestinal parasites and genotypes of *Giardia intestinalis* in school children from Berisso, Argentina. *Journal of Infection in Developing Countries*. 2011; 5(07):527–34. Available from: <https://jdc.org/index.php/journal/article/view/1660>
32. Krolewiecki AJ, Ramanathan R, Fink V, McAuliffe I, Cajal SP, Won K, et al. Improved diagnosis of *Strongyloides stercoralis* using recombinant antigen-based serologies in a community-wide study in northern Argentina. *Clinical and Vaccine Immunology*. 2010; 17(10):1624–30. Available from: <https://pubmed.ncbi.nlm.nih.gov/20739501/>
33. Altchegh J, Fernández G, Guarnera EA, Gutiérrez N, Pizzi H, Taranto N. Geohelminthiasis en la República Argentina. Buenos Aires: Área de Difusión del PROAPS-REMEDIAAR; 2007. http://186.33.221.24/medicamentos/files/modulo_geohelminth_baja.pdf
34. Gebreyes WA, Dupouy-Camet J, Newport MJ, Oliveira CJB, Schlesinger LS, Saif YM, et al. The global one health paradigm: Challenges and opportunities for tackling infectious diseases at the human, animal, and environment interface in low-resource settings. *PLoS Neglected Tropical Diseases*. 2014; 8(11):e3257. Available from: <http://dx.doi.org/10.1371/journal.pntd.0003257> PMID: 25393303
35. Ostfeld RS, Glass GE, Keesing F. Spatial epidemiology: an emerging (or re-emerging) discipline. *Trends in Ecology and Evolution*. 2005; 20(6):328–36. Available from: <https://doi.org/10.1016/j.tree.2005.03.009> PMID: 16701389
36. Elliott P, Cuzick J, English D, Stern R. *Geographical and environmental epidemiology*. Oxford University Press; 1996.
37. Lawson AB, Biggeri A, Bohning D, Lesaffre E, Viel J-F, Bertollini R, editors. *Disease mapping and risk assessment for public health*. 1st ed. Chichester, England: John Wiley & Sons; 2000.
38. Thomas DC. *Statistical methods in environmental epidemiology*. London, England: Oxford University Press; 2009.
39. Polop F, Provencal C, Scavuzzo M, Lamfri M, Calderón G, Polop J. On the relationship between the environmental history and the epidemiological situation of Argentine hemorrhagic fever. *Ecological Research*. 2008; 23(1):217–25. Available from: <http://dx.doi.org/10.1007/s11284-007-0371-2>
40. Porcasi X, Rotela CH, Introini MV, Frutos N, Lanfri S, Peralta G, et al. An operative dengue risk stratification system in Argentina based on geospatial technology. *Geospatial Health*. 2012; 6(3):31. Available from: <http://dx.doi.org/10.4081/gh.2012.120> PMID: 23032281
41. Estallo EL, Benitez EM, Lanfri MA, Scavuzzo CM, Almiron WR. MODIS Environmental Data to Assess Chikungunya, Dengue, and Zika Diseases Through *Aedes (Stegomia) aegypti* Oviposition Activity Estimation. *IEEE Journal of Selected Topics in Applied Earth Observation and Remote Sensing*. 2016; 9(12):5461–6. Available from: <http://dx.doi.org/10.1109/jstars.2016.2604577>
42. Rotela C, Lopez L, Frías Céspedes M, Barbas G, Lighezzolo A, Porcasi X, et al. Analytical report of the 2016 dengue outbreak in Córdoba city, Argentina. *Geospatial Health*. 2017; <http://dx.doi.org/10.4081/gh.2017.564>
43. Alvarez Di Fino EM, Rubio J, Abril MC, Porcasi X, Periago MV. Risk map development for soil-transmitted helminth infections in Argentina. *PLoS Neglected Tropical Diseases*. 2020; 14(2):e0008000. Available from: <http://dx.doi.org/10.1371/journal.pntd.0008000> PMID: 32040473
44. Waage J, Banerji R, Campbell O, Chirwa E, Collender G, Dieltiens V, et al. The Millennium Development Goals: a cross-sectoral analysis and principles for goal setting after 2015. *Lancet*. 2010; 376(9745):991–1023. Available from: [http://dx.doi.org/10.1016/s0140-6736\(10\)61196-8](http://dx.doi.org/10.1016/s0140-6736(10)61196-8)

45. WHO World Health Organization. (2020). Ending the neglect to attain the sustainable development goals: a road map for neglected tropical diseases 2021–2030: overview. <https://apps.who.int/iris/bitstream/handle/10665/338565/9789240026582-rus.pdf>
46. S Gaspe M, Provecho YM, Cardinal MV, del Pilar Fernández M, Gürtler RE (2015) Ecological and Sociodemographic Determinants of House Infestation by *Triatoma infestans* in Indigenous Communities of the Argentine Chaco. *PLoS Neglected Tropical Diseases*. 9(3): e0003614. <https://doi.org/10.1371/journal.pntd.0003614> PMID: 25785439
47. Chammartin F, Guimarães LH, Scholte RG, Bavia ME, Utzinger J, Vounatsou P. Spatio-temporal distribution of soil-transmitted helminth infections in Brazil. *Parasites and Vectors*. 2014; 7(1):440. Available from: <http://dx.doi.org/10.1186/1756-3305-7-440> PMID: 25230810
48. Anegagrie M, Lanfri S, Aramendia AA, Scavuzzo CM, Herrador Z, Benito A, et al. Environmental characteristics around the household and their association with hookworm infection in rural communities from Bahir Dar, Amhara Region, Ethiopia. *PLoS Neglected Tropical Diseases*. 2021; 15(6):e0009466. Available from: <http://dx.doi.org/10.1371/journal.pntd.0009466> PMID: 34157019
49. Campbell SJ, Nery SV, Wardell R, D'Este CA, Gray DJ, McCarthy JS, et al. Water, Sanitation and Hygiene (WASH) and environmental risk factors for soil-transmitted helminth intensity of infection in Timor-Leste, using real time PCR. *PLoS Neglected Tropical Diseases*. 2017; 11(3):e0005393. Available from: <http://dx.doi.org/10.1371/journal.pntd.0005393> PMID: 28346536
50. Chaiyos J, Suwannatrai K, Thinkhamrop K, Pratumchart K, Sereewong C, Tesana S, et al. MaxEnt modeling of soil-transmitted helminth infection distributions in Thailand. *Parasitology Research*. 2018; 117(11):3507–17. Available from: <http://dx.doi.org/10.1007/s00436-018-6048-7> PMID: 30120589
51. Cardinal MV, Enriquez GF, Macchiaverna NP, Argibay HD, Fernández MDP, Alvedro A, et al. Long-term impact of a ten-year intervention program on human and canine *Trypanosoma cruzi* infection in the Argentine Chaco. *PLoS Neglected Tropical Diseases*. 2021; 15(5):e0009389. Available from: <http://dx.doi.org/10.1371/journal.pntd.0009389> PMID: 33979344
52. INDEC. Censo nacional de población. Hogares y Viviendas. Buenos Aires, Argentina; 2001.
53. INDEC. Censo nacional de población. Hogares y Viviendas. Buenos Aires, Argentina; 2010.
54. Gurevitz JM, Ceballos LA, Gaspe MS, Alvarado-Otegui JA, Enriquez GF, Kitron U, et al. Factors affecting infestation by *Triatoma infestans* in a rural area of the humid Chaco in Argentina: A multi-model inference approach. *PLoS Neglected Tropical Diseases*. 2011; 5(10):e1349. Available from: <http://dx.doi.org/10.1371/journal.pntd.0001349>
55. Cardinal MV, Ceballos LA, Gaspe MS, Orozco MM, Alvarado-Otegui JA, Gürtler RE, et al. Heterogeneities in the Ecoepidemiology of *Trypanosoma cruzi* Infection in Rural Communities of the Argentinean Chaco. *American Journal of Tropical Medicine and Hygiene*. 2014; 90(6):1063–73. Available from: <http://dx.doi.org/10.4269/ajtmh.13-0251>
56. Naumann M. Atlas del Gran Chaco Sudamericano. Buenos Aires, Argentina; 2006: Sociedad Alemana de Cooperación Técnica (GTZ). ErreGé & Asoc.
57. Periago MV, García R, Astudillo OG, Cabrera M, Abril MC. Prevalence of intestinal parasites and the absence of soil-transmitted helminths in Añatuya, Santiago del Estero, Argentina. *Parasites and Vectors*. 2018 [cited 2022 Aug 24]; 11(1):638. Available from: <https://doi.org/10.1186/s13071-018-3232-7>
58. Molla E, Mamo H. Soil-transmitted helminth infections, anemia and undernutrition among schoolchildren in Yirgacheffe, South Ethiopia. *BMC Research Notes*. 2018; 11(1). Available from: <http://dx.doi.org/10.1186/s13104-018-3679-9> PMID: 30103797
59. Ministerio de Salud de la Nación. Atención Primaria de la Salud. Boletín PROAPS-REMIAR. 2001; 2 (14). Available from: <http://medicamentos.msal.gov.ar/files/boletin14.pdf>
60. Pbc Planet Labs. PlanetScoupe images. Planet Imagery Product Specifications. 2022. <https://www.planet.com/products/>
61. De Roeck E, Van Coillie F, De Wulf R, Soenen K, Charlier J, Vercruyse J, et al. Fine-scale mapping of vector habitats using very high resolution satellite imagery: a liver fluke case-study. *Geospatial Health*. 2014; 8(3):S671–83. Available from: <https://www.geospatialhealth.net/index.php/gh/article/view/296> PMID: 25599638
62. Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*.
63. QGIS.org. 2022. QGIS Geographic Information System. QGIS Association. <http://www.qgis.org>
64. StataCorp. 2017. Stata Statistical Software: Release 15. College Station, TX: StataCorp LLC.
65. Kulldorff M, Heffernan R, Hartman J, Assunção R, Mostashari F. A space–time permutation scan statistic for disease outbreak detection. *PLoS Medicine*. 2005; 2(3):e59. Available from: <http://dx.doi.org/10.1371/journal.pmed.0020059> PMID: 15719066

66. Matzkin R, Galván M, Miranda O, Merino D, Balbachán S. Parasitosis entéricas en una población escolar periurbana de Resistencia, Chaco. *Chaco Comunicaciones Científicas y Tecnológicas*. 2000; 40:197–200.
67. Ledesma AE, Fernández G. Enteroparasitosis: Factores predisponentes en población infantil de la Ciudad de Resistencia, Chaco. *UNNE Comunicaciones Científicas y Tecnológicas*. 2004.
68. Scavuzzo CM, Scavuzzo JM, Campero MN, Anegagrie M, Aramendia AA, Benito A, et al. Feature importance: Opening a soil-transmitted helminth machine learning model via SHAP. *Infectious Diseases Modelling*. 2022; 7(1):262–76. Available from: <http://dx.doi.org/10.1016/j.idm.2022.01.004> PMID: 35224316
69. Raso G. Assessment, mapping and prediction of the spatial distribution of parasitic infections in western Côte d'Ivoire and implications for integrated control. University of Basel; 2004.
70. Brooker S, Clements ACA, Bundy DAP. Global epidemiology, ecology and control of soil-transmitted helminth infections. In: *Advances in Parasitology*. Elsevier; 2006. p. 221–61.
71. Matthys B, Bobieva M, Karimova G, Mengliboeva Z, Jean-Richard V, Hoimnazarova M, et al. Prevalence and risk factors of helminths and intestinal protozoa infections among children from primary schools in western Tajikistan. *Parasites and Vectors*. 2011; 4(1). Available from: <http://dx.doi.org/10.1186/1756-3305-4-195> PMID: 21981979
72. Ovutor O, Helen I, Awi-waadu G. Assessment of physico-chemical parameters of soils in fallowing farmlands and pit toilet environments as it affects the abundance of geohelminths in emohua local government area, rivers state, Nigeria. *Annual Research and Review in Biology*. 2017; 14(3):1–10. Available from: <http://dx.doi.org/10.9734/arrb/2017/31546>
73. Manz KM, Clowes P, Kroidl I, Kowuor DO, Geldmacher C, Ntinginya NE, et al. *Trichuris trichiura* infection and its relation to environmental factors in Mbeya region, Tanzania: A cross-sectional, population-based study. *PLoS One*. 2017; 12(4):e0175137. Available from: <http://dx.doi.org/10.1371/journal.pone.0175137>
74. Wardell R, Clements ACA, Lal A, Summers D, Llewellyn S, Campbell SJ, et al. An environmental assessment and risk map of *Ascaris lumbricoides* and *Necator americanus* distributions in Manufahi District, Timor-Leste. *PLoS Neglected Tropical Diseases*. 2017; 11(5):e0005565. Available from: <http://dx.doi.org/10.1371/journal.pntd.0005565>
75. Mota KCP, Grama DF, Fava MNM, Úngari LP, Faria ESM, Cury MC. Distribution and risk factors of Ascarididae and other geohelminths in the soil of Uberlândia, Minas Gerais, Brazil. *Revista del Instituto de Medicina Tropical Sao Paulo*. 2018; 60(0). Available from: <http://dx.doi.org/10.1590/s1678-9946201860017> PMID: 29694601
76. Sedionoto B, Wasessombat S, Punsawad C, Anamart W. Environmental factors and prevalence of hookworm infection and strongyloidiasis in rural East Kalimantan, Indonesia. *E3S Web Conference, International Conference on Energy, Environment, Epidemiology and Information System*. 2019;125:04001.
77. Gamboa MI, Basualdo JA, Kozubsky L, Costas E, Cueto Rua E, Lahitte HB. Prevalence of intestinal parasitosis within three population groups in La Plata, Argentina. *Euro Journal of Epidemiology*. 1998; 14(1):55–61. Available from: <http://dx.doi.org/10.1023/a:1007479815249> PMID: 9517874
78. Visser S, Giatti LL, de Carvalho RAC, Guerreiro JCH. Study of the association between socio-environmental factors and the prevalence of intestinal parasitosis in the suburbs of the city of Manaus in the state of Amazonas, Brazil. *Ciencia y Sociedad Coletiva*. 2011; 16(8):3481–92. Available from: <https://www.proquest.com/openview/a6e4aa2eb6e4640a0a8508983ae72a56/1?pq-origsite=gscholar&cbl=2034998>
79. Candela E, Goizueta C, Periago MV, Muñoz-Antoli C. Prevalence of intestinal parasites and molecular characterization of *Giardia intestinalis*, *Blastocystis spp.* and *Entamoeba histolytica* in the village of Fortín Mbororé (Puerto Iguazú, Misiones, Argentina). *Parasites and Vectors*. 2021; 14(1). Available from: <http://dx.doi.org/10.1186/s13071-021-04968-z>
80. Tadege B, Mekonnen Z, Dana D, Tiruneh A, Sharew B, Dereje E, et al. Assessment of the nail contamination with soil-transmitted helminths in schoolchildren in Jimma Town, Ethiopia. *PLoS One*. 2022; 17(6):e0268792. Available from: <http://dx.doi.org/10.1371/journal.pone.0268792> PMID: 35767573
81. Eyasu A, Molla M, Kefale B, Sisay W, Andargie Y, Kebede F, et al. Prevalence and associated risk factors of endoparasites among under-five children in Debre Tabor comprehensive specialized hospital, Debre Tabor, Northwest Ethiopia: A cross-sectional study. *Journal of Parasitology Research*. 2022; 2022:6917355. Available from: <https://www.hindawi.com/journals/jpr/2022/6917355/> PMID: 35535044
82. Hajare ST, Chekol Y, Chauhan NM. Assessment of prevalence of *Giardia lamblia* infection and its associated factors among government elementary school children from Sidama zone, SNNPR, Ethiopia. *PLoS One*. 2022; 17(3):e0264812. Available from: <http://dx.doi.org/10.1371/journal.pone.0264812>

83. Gizaw Z, Yalaw AW, Bitew BD, Lee J, Bisesi M. Fecal indicator bacteria along multiple environmental exposure pathways (water, food, and soil) and intestinal parasites among children in the rural northwest Ethiopia. *BMC Gastroenterol* [Internet]. 2022; 22(1):84. Available from: <http://dx.doi.org/10.1186/s12876-022-02174-4> PMID: 35220951
84. Freeman MC, Chard AN, Nikolay B, Garn JV, Okoyo C, Kihara J, et al. Associations between school- and household-level water, sanitation and hygiene conditions and soil-transmitted helminth infection among Kenyan school children. *Parasites and Vectors*. 2015; 8(1). Available from: <http://dx.doi.org/10.1186/s13071-015-1024-x> PMID: 26248869
85. Bartram J, Cairncross S. Hygiene, sanitation, and water: Forgotten foundations of health. *PLoS Medicine*. 2010; 7(11):e1000367. Available from: <http://dx.doi.org/10.1371/journal.pmed.1000367> PMID: 21085694
86. Ngure FM, Reid BM, Humphrey JH, Mbuya MN, Pelto G, Stoltzfus RJ. Water, sanitation, and hygiene (WASH), environmental enteropathy, nutrition, and early child development: making the links. *Annals of the New York Academy of Science*. 2014; 1308(1):118–28. Available from: <http://dx.doi.org/10.1111/nyas.12330> PMID: 24571214
87. Strunz EC, Addiss DG, Stocks ME, Ogden S, Utzinger J, Freeman MC. Water, sanitation, hygiene, and soil-transmitted helminth infection: A systematic review and meta-analysis. *PLoS Medicine*. 2014; 11(3):e1001620. Available from: <http://dx.doi.org/10.1371/journal.pmed.1001620> PMID: 24667810
88. Chin YT, Lim YAL, Chong CW, Teh CSJ, Yap IKS, Lee SC, et al. Prevalence and risk factors of intestinal parasitism among two indigenous sub-ethnic groups in Peninsular Malaysia. *Infectious Diseases of Poverty*. 2016; 5(1). Available from: <http://dx.doi.org/10.1186/s40249-016-0168-z> PMID: 27430215
89. Tekalign E, Bajiro M, Ayana M, Tiruneh A, Belay T. Prevalence and intensity of soil-transmitted helminth infection among rural community of southwest Ethiopia: A community-based study. *Biomed Research International*. 2019; 2019:1–7. Available from: <http://dx.doi.org/10.1155/2019/3687873>
90. Vilchez Barreto PM, Santivañez S, García HH, Lescano AG, O'Neal SE, Moyano L-M, et al. Prevalence, age profile, and associated risk factors for *Hymenolepis nana* infection in a large population-based study in northern Peru. *American Journal of Tropical Medicine and Hygiene*. 2017; 97(2):583–6. Available from: <http://dx.doi.org/10.4269/ajtmh.16-0939> PMID: 28829724