

# Primer relevamiento de hongos aislados de suelos e insectos plagas asociados a cultivos orgánicos de cáñamo industrial (*Cannabis sativa* L.) en Paraguay

First survey of fungi isolated from soils and pest insects associated with organic plantations of industrial hemp (*Cannabis sativa* L.) in Paraguay

Claudia Elena González<sup>1\*</sup> , David Santiago Gayoso<sup>1</sup> , Alan Salinas<sup>2</sup> , Gustavo A. Bich<sup>3</sup> , María Lorena Castrillo<sup>3</sup> 



<sup>1</sup>Universidad Nacional de Itapúa. Facultad de Ciencias y Tecnología. Encarnación, Paraguay.

<sup>2</sup>Evona S.A. Ciudad del Este, Alto Paraná, Paraguay.

<sup>3</sup>Universidad Nacional de Misiones. Facultad de Ciencias Exactas Químicas y Naturales. Instituto de Biotecnología Misiones. Misiones, Argentina.

DOI: 10.57201/IEUNA2313332

Sección: Comunicación Corta

\*Autor correspondiente:  
claudiagonzalez@cyt.uni.edu.py

Editor de área:

Andrea Alejandra Arrúa  
Alvarenga, Universidad Nacional  
de Asunción.

Editor invitado:

Guillermo Enciso, Centro de  
Desarrollo e Innovación  
Tecnológica (CEDIT)

Recibido:


16 de febrero de 2023

Aceptado:

19 de junio de 2023

Recibido en versión modificada:

22 de junio de 2023

Este es un artículo  
publicado en acceso  
abierto bajo una licencia  
Creative Commons "CC  
BY 4.0". 

Declaración de conflicto: Los  
autores declaran no tener conflicto  
de intereses.

e-ISSN 2709-0817

Como citar: González, C. E. ,  
Gayoso, D. S., Salinas, A., Bich,  
G. A. & Castrillo, M. L. (2023).  
Primer relevamiento de hongos  
aislados de suelos e insectos  
plagas asociados a cultivos  
orgánicos de cáñamo industrial  
(*Cannabis sativa* L.) en  
Paraguay. Rev. Invest. y Est. -  
UNA, 14(1), 105-111.

**Abstract.** The cultivation of *Cannabis sativa* has gained importance in recent years in Paraguay and other Latin American countries where legal regulations allow its cultivation for medicinal use, as fiber, and also as food. Since several companies apply the organic cultivation model to guarantee quality and certify their product, it is necessary to study biological control alternatives that guarantee sustainable agriculture. In this study, microorganisms present in the soil of *Cannabis sativa* cultivation and those that infected pests found on the plant were isolated and identified. The serial dilution method was used to isolate microorganisms from the soil and direct seeding on specific culture media to isolate them from the bodies of pest insects. The isolated genera from the soil were *Rhizopus*, *Aspergillus*, *Absidia*, *Alternaria*, *Mucor*, *Penicillium* and *Fusarium*; the isolated genera from pest insects were *Cladosporium*, *Alternaria*, *Aspergillus*, and *Fusarium*. This study is the first report on fungi in soils and pest insects in industrial hemp cultivation in Paraguay.

**Keywords:** agroecosystem, management, organic, Paraguay.

**Resumen.** El cultivo de *Cannabis sativa* ha tomado una gran importancia en los últimos años en Paraguay y en otros países de América Latina donde las reglamentaciones legales permiten su cultivo para usos medicinales, como fibra y también como alimento. Debido a que varias empresas aplican el modelo de cultivo orgánico para garantizar la calidad y además para llegar a certificar su producto es necesario estudiar alternativas de control biológico que garanticen una agricultura sostenible. En este estudio se aislaron e identificaron los microorganismos presentes en el suelo de cultivo de *Cannabis sativa* y también aquellos que infectaron a los insectos plagas encontrados sobre la planta. Se utilizó el método de dilución seriada para aislar microorganismos del suelo y para aislarlos de los cuerpos de insectos plagas la siembra directa en medios de cultivos específicos. Los géneros aislados del suelo fueron *Rhizopus*, *Aspergillus*, *Absidia*, *Alternaria*, *Mucor*, *Penicillium* y *Fusarium*; los géneros aislados de insectos plaga fueron *Cladosporium*, *Alternaria*, *Aspergillus* y *Fusarium*. Este es la primera descripción sobre la presencia de hongos en suelos e insectos plagas en cultivo de cáñamo industrial en Paraguay

**Palabras clave:** agroecosistema, manejo, orgánico, Paraguay.

## Introduction


In recent years, non-psychoactive industrial hemp (*Cannabis sativa* L.) has experienced significant growth due to its usefulness in the textile, medicinal, and food industries (Rupasinghe et al., 2020). The cultivation of hemp in Paraguay has increased eight times, ranging from 600 ha in 2019 to 5000 ha in 2022 (Prohibition Partner, 2022). The reasons include its decriminalization and high demand in domestic and international markets (Reuters, 2023). The rapid expansion of organic crops has increased focus on pest management research. Nevertheless, limited information on the subject presents significant challenges to producers in effectively managing diseases caused by fungal and insect pests, which emerge as a result of high crop density and specific management practices implemented in the field (Sunoj Valiaparambil et al., 2023; Zheljzkov et al., 2023). Soil is considered one of the main sources of inoculum of phytopathogens (Johns et al., 2021). The phytopathogen *Fusarium oxysporum* has been found in soils of *C. sativa* crops in North America, Israel, and Italy, raising concerns among hemp growers worldwide due to the significant commercial losses. This pathogen can infect the plant from its seedling stage and spread through infected mother plants, generating varying degrees of aggressiveness depending on the haplotypes (Jerushalmi et al., 2022). Previous studies in *C. sativa* have revealed the presence of 12 haplotypes of *Fusarium cannabis*, of which five have been reported to cause significant economic losses due to plant diseases and mycotoxin residues (Gwinn et al., 2022). Hemp crops are also vulnerable to pest, such as the red spider mite (*Tetranychus urticae*), which can significantly reduce production and commercial product quality. To control this pest growers, use mites from the genera *Neoseiulus californicus* and *Phytoseiulus persimilis* as biological control agents. In addition, hemp farmers explored the potential of entomopathogenic fungi, whose effect as bioinsecticides is well-known in other crops (Jimenez Lara, 2022). In 2019 in Ecuador, *Tetranychus urticae*, caused 60- 80% losses in strawberry crops, which motivated the evaluation of the entomopathogenic activity of *Bacillus subtilis* in vitro (Mendoza-Léon et al., 2019). While the effects of fungi such as *Beauveria bassiana* and *Metarhizium anisopliae* as biocontrol agents are well-known, bacteria have been little studied. The results obtained were very encouraging and suggest that this technique could be used as a biocontrol alternative for other crops. (Pacheco Hernández et al., 2019). In Paraguay, during research looking to obtain biological control agents from the compost used to grow *Cannabis* seedlings, several genera of filamentous fungi were isolated and molecularly identified as *Aspergillus* sp., *Penicillium* sp., *Cladosporium* sp., *Geomyces* sp., *Plectosphaerella* sp., and *Rasamsonia* sp. These genera are common in crop soils, although *Cladosporium* sp. should be considered a potential threat in *C. sativa* as it is a pathogen in other crops (Álvarez Beauchamp, 2022). This research aimed to isolate and identify microscopic fungi associated with the soils and insects of organic non-psychoactive industrial hemp crops on a farm in the Eastern Region of Paraguay to characterize the diversity.

## Materials and methods

### Sampling site

The sampling places were selected according to farmers that develop organic crops and previous investigations agreements. Samples were collected at nine georeferenced points selected randomly in the certified organic soils belonging to EVONA S.A. (Figure 1). The production fields were located in Hernandarias city, 7.32

km from route PY07 Km 28 (25°18'7"S 54°37'54"W). Georeferencing was performed using the Android operating system application GPS Fields Area Measure PRO.



**Figure 1.** Sampling site of hemp plantation. A) Georeferenced sampling points, Hernandarias, Paraguay. B) Types of soils. C) Detail of parcel of sampling. D) Selected hem plantation.

### Soil samples

According to data published by the Departamento de Planificación Ambiental del Ministerio de Agricultura (Paraguay), the soil type belongs to the Oxisol classification, soils derived from basalt that show an undulating profile in the form of hills followed by flat areas, the content of mica, apatites, feldspars and ferromagnesian minerals must be below 10% since they are weatherable.

Subsequently, soil samples weighing 200 grams each, 1800 grams for all, were extracted for pH determination and microbiological analysis. The soil samples came from places where hemp was grown between April and August 2022, in plots where hemp cultivation is anticipated in the future, and from residual substrate. The sample preparation followed the methodology proposed by Jerke et al., (2019). The soil samples were collected in magnetic-closure bags for transport to the Ciencia y Tecnología laboratory at the Universidad Nacional de Itapúa.

Solutions were prepared in sterile distilled water from nine soil samples to measure pH. Twenty grams of each sample was weighed and added to a beaker containing 50 ml of distilled water. The mixture was stirred with a glass rod for 2 minutes and left to settle for 15 minutes. The pH of the supernatant was measured using a QUIMIS Q400AS 240V-10W benchtop pH meter. To isolate fungi from the soil samples prepared for hemp

cultivation, serial dilutions of  $10^{-1}$ ,  $10^{-2}$ , and  $10^{-3}$  were prepared in sterile distilled water. They were then seeded onto nutrient agar plates containing chloramphenicol antibiotic (150 mg/l). The plates were labeled and incubated at 25°C for 5 to 7 days until colony growth was observed (Gonzalez et al., 2013; Humber, 1997).


### Insect samples

All dead insects found on the plants or laying on the soil were collected. The insect samples were transported sterile jars to the Ciencia y Tecnología laboratory (Universidad Nacional Itapúa). They were processed using the wet chamber technique and the direct puncture technique of the soft tissues, which were then cultured on potato dextrose agar (PDA) agar plates. The insects were identified in the field through morphological features.

Fungal colonies were observed both macroscopically and microscopically. For microscopic observations, a compound microscope (Zeiss Primo Stard Trinocular®) with an incorporated digital camera was used to capture images at magnifications of 400X and 1000X. The taxonomic guide of Dr. Humber (1997) was used for the microscopic identification of the fungal genus. The fungal colonies were observed in the Faculty of Science and Technology laboratory at UNI.

## Results and Discussion

A total of nine samples of oxisol soils and 12 samples of insects were collected from the hemp plantations (Figure 2). The main insects genera found were *Diabrotica*, *Empoasca*, *Naupactus* and *Autographa* (personal communication with Ing. Alan Salinas, EVONA S.A.).



**Figure 2.** Examples of organisms-inhabiting in hemp plantations sampled in Paraguay. A and B fungal colonies isolated from soil samples. C and D insects collected from hem plantations.

Of the 42 fungal colonies growing in the Petri dishes with PDA from soil samples, 30 were confirmed through microscopic examination to represent one of the seven genera: a *Rhizopus*, *Aspergillus*, *Absidia*, *Alternaria*, *Mucor*, *Penicillium*, and *Fusarium*. In all the oxisol soils from the hemp plantation the genera *Fusarium*, *Aspergillus*, *Penicillium* and *Cladosporium* were found. The absence of entomopathogenic fungi of the species *B. bassiana* and *M. anisopliae* may be attributed to the acidic conditions of the soil and the high temperatures encountered at the various sampling sites. From the collected insects *Cladosporium*, *Alternaria*, *Aspergillus*, and *Fusarium* were isolated. In our study, all the samples were collected during the continuous dry season prevailing in our region, with a remarkably absence of regular rainfalls, which may alter the occurrence of many groups of

microorganisms from samples of soil. It has been suggested that the presence of *B. bassiana* to be very low during periods of scarce rainfall and intense drought, increasing during rainy months with higher humidity and temperatures of 20-30°C, conditions favorable for spore dispersal and germination. However, the genus *M. anisopliae* was found to invade tissues of dead insects when temperatures reached 40°C. These data highlight the importance of climatic conditions in the development of some pathogens and emphasize the need to consider them in future studies of pathogenicity related to hemp cultivation (García-García et al., 2011). Another important aspect to consider is that the pH in the analyzed soils was rather acid, varying between 5.2 and 5.71. The field frequency of entomopathogenic fungi was low, which could be related to crop management, environmental conditions, availability of susceptible hosts, pathogen virulence, and host crop characteristics, among others (Borges, 2007; Vänninen, 1996). Climatic factors such as moderate temperatures, abundant rainfalls, and high humidity are correlated with the occurrence of entomopathogenic fungi (Clifton et al., 2015; Jaronski, 2010; Yaginuma, 2007). Similarly, Silva Guerra et al., (2009) reported the effect of moderated temperatures and high percentages of humidity favours the persistence of conidia in soil, where *M. anisopliae* was found to be more resistant than *M. acridum*. The presence of *Penicillium*, *Alternaria*, *Aspergillus*, and *Fusarium* genera in soil and pest insect tissues would merit further study of their pathogenicity under controlled laboratory conditions to deepen knowledge of their potential use as biocontrol agents of hemp crop pests.

## Conclusion

This work has made it possible to know the fungal diversity present in soils and pest insects associated with organic crops of industrial hemp (*Cannabis sativa*, L.) for the first time in Hernandarias, Alto Paraná, Paraguay. All of the isolated and identified fungi of the genera *Alternaria*, *Rhizopus*, *Absidia*, *Mucor*, *Penicillium*, *Aspergillus*, *Fusarium*, and *Cladosporium* have been stored in the InBioMis culture collection, Misiones Argentina, and the Faculty of Science and Technology of the National University of Itapúa for future molecular identification assays.

The present work allowed us to establish links between the National University of Itapúa, the Institute of Biotechnology of Misiones, Argentina, and the company Evona S.A. so that we can continue researching and interacting in the future for the benefit of organic production of industrial hemp.

## Contribución de los autores

**Conceptualization:** C.E.G., G.A.B., M.L.C. **Experiment design:** C.E.G., D.S.G, G.A.B., M.L.C. **Experiment execution:** C.E.G., D.S.G., A.S. **Experiment verification:** C.E.G., D.S.G **Data analysis/interpretation:** C.E.G., D.S.G, G.A.B., M.L.C. **Statistical analysis:** **Manuscript preparation:** C.E.G., G.A.B. **Manuscript editing and revision:** C.E.G., G.A.B., M.L.C. **Approval of the final manuscript version:** C.E.G., G.A.B., M.L.C.

## Referencias Bibliográficas

- Álvarez Beauchamp, M. H. (2022). Adaptabilidad y desempeño del cañamo industrial (*Cannabis sativa* L.) y cuantificación de los principales cannabinoides.
- Borges, L. (2007). *Eficiência de Beauveria bassiana (bals.) vuill. (Deuteromycota) para o controle de Hedypathes betulinus (Klug) (Coleoptera: Cerambycidae) em erva-mate, Ilex paraguariensis St. Hil. (Aquifoliaceae)* [Tesis Doctoral]. Universidad Federal de Paraná.
- Clifton, E. H., Jaronski, S. T., Hodgson, E. W., & Gassmann, A. J. (2015). Abundance of Soil-Borne Entomopathogenic Fungi in Organic and Conventional Fields in the Midwestern USA with an Emphasis on the Effect of Herbicides and Fungicides on Fungal Persistence. *PLOS ONE*, *10*(7), e0133613. <https://doi.org/10.1371/journal.pone.0133613>
- García-García, M. A., Cappello García, S., Leshner Gordillo, J. M., & Molina Martínez, R. F. (2011). Aislamiento y caracterización morfológica de los hongos entomopatógenos *Beauveria bassiana* y *Metarhizium anisopliae*. *Horizonte Sanitario*, *10*(2), 21–28. <http://www.redalyc.org/articulo.oa?id=457845138002>
- Gonzalez, E., Martinez, C., Castrillo, M., Fonseca, M., Zapata, P., & Villalba, L. (2013). Optimización del protocolo para el aislamiento de hongos de pudrición blanca.
- Gwinn, K. D., Hansen, Z., Kelly, H., & Ownley, B. H. (2022). Diseases of *Cannabis sativa* Caused by Diverse *Fusarium* Species. *Frontiers in Agronomy*, *3*(January), 1–17. [doi.org/10.3389/fagro.2021.796062](https://doi.org/10.3389/fagro.2021.796062)
- Humber, R. (1997). Fungi: Preservation of cultures. In L. A. Lacey (Ed.), *Manual of Techniques in Invertebrate Pathology* (Second Ed., pp. 317–328).
- Jaronski, S. T. (2010). Role of fungal ecology in the inundative use of entomopathogenic fungi. *BioControl*, *55*, 159–185.
- Jerke, G., Horiński, M. A., Castrillo, M. L., & Chade, M. E. (2019). *Guía de prácticas de laboratorio 1º cuatrimestre / 2da. Ed. Cátedra de Microbiología e Inmunología*.
- Jerushalmi, S., Maymon, M., O'Donnell, K., & Freeman, S. (2022). Members of the *Fusarium oxysporum* Complex Causing Wilt Symptoms in Medical Cannabis in Israel, Italy, and North America Comprise a Polyphyletic Assemblage. *106*(10), 2656–2662. <https://doi.org/10.1094/PDIS-01-22-0155-RE>
- Jimenez Lara, A. D. (2022). Alternativas de manejo para la araña roja (*Tetranychus urticae*), en el cultivo de cañamo (*Cannabis sativa*). <http://dspace.utb.edu.ec/handle/49000/11331>
- Johns, L. E., Goldman, G. H., Ries, L. N. A., & Brown, N. A. (2021). Nutrient sensing and acquisition in fungi: mechanisms promoting pathogenesis in plant and human hosts. *Fungal Biology Reviews*, *36*, 1–14. <https://doi.org/10.1016/j.fbr.2021.01.002>
- Mendoza-Léon, D., Dobronski-Arcos, J., Vásquez-Freytez, C., Frutos-Pinto, V., & Paredes-Carreño, S. (2019). Control de *Tetranychus urticae* Koch (Acari: Tetranychidae) con *Bacillus subtilis* en hojas de fresa (*Fragaria vesca*). *Agronomía Costarricense*, *43*(1), 125–133.
- Pacheco Hernández, Ma. de L., Reséndiz Martínez, J. F., & Arriola Padilla, V. J. (2019). Organismos entomopatógenos como control biológico en los sectores agropecuario y forestal de México: una revisión Entomopathogenic organisms for pest control in the mexican agriculture, livestock and forest sectors: a review Introducción El uso indiscriminado. *Revista Mexicana de Ciencias Forestales*, *10*(56), 1–32.
- Prohibition Partner. (2022). *Paraguay Reigns Over the Latin American Hemp Market*. <https://prohibitionpartners.com/2022/09/28/paraguay-reigns-over-the-latin-american-hemp-market/>
- Reuters. (2023). *Paraguay marijuana festival sets high hopes for legalization*. <https://www.nasdaq.com/articles/paraguay-marijuana-festival-sets-high-hopes-for-legalization>
- Rupasinghe, H. P. V., Zheljaskov, V. D., Davis, A., Kumar, S. K., & Murray, B. (2020). Industrial Hemp (*Cannabis sativa* subsp. *sativa*) as an Emerging Source for Value-Added Functional Food Ingredients and Nutraceuticals. *Molecules*, *25*(Figure 1), 1–24.
- Silva Guerra, D. M., Duarte Pires, A. P., & Alves Lima, E. A. de L. (2009). Persistence of *Metarhizium anisopliae* spp. in soil under different conditions of temperature and humidity. *Revista Caatinga*, *22*(2).
- Sunoj Valiamparambil Sebastian, J., Dong, X., Trostle, C., Pham, H., Joshi, M. V., Jessup, R. W., Burow, M. D., & Provin, T. L. (2023). Hemp Agronomy: Current Advances, Questions, Challenges, and Opportunities. *Agronomy*, *13*(2), 1–26. <https://doi.org/10.3390/agronomy13020475>
- Vänninen, I. (1996). Distribution and occurrence of four entomopathogenic fungi in Finland: effect of geographical location, habitat type and soil type. *Mycological Research*, *100*(1), 93–101. [https://doi.org/10.1016/S0953-7562\(96\)80106-7](https://doi.org/10.1016/S0953-7562(96)80106-7)
- Yaginuma, K. (2007). Seasonal Occurrence of Entomopathogenic Fungi in Apple Orchard Not Sprayed with Insecticides. *Japanese Journal of Applied Entomology and Zoology*, *51*(3), 213–220. <https://doi.org/10.1303/jjaez.2007.213>
- Zheljaskov, V. D., Sikora, V., Noller, J., Latkovi, D., Ocamb, C. M., & Koren, A. (2023). *Industrial Hemp (Cannabis sativa L.) Agronomy and Utilization: A Review*.