

Journal of Dairy Research

<http://journals.cambridge.org/DAR>

Additional services for *Journal of Dairy Research*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Histological examination of non-lactating bovine udders inoculated with *Lactobacillus perolens* CRL 1724

Ignacio D Frola, Matías S Pellegrino, Gabriel Magnano, José A Giraudó, María C Espeche, María EF Nader-Macias and Cristina I Bogni

Journal of Dairy Research / *FirstView* Article / November 2012, pp 1 - 8

DOI: 10.1017/S0022029912000581, Published online: 09 November 2012

Link to this article: http://journals.cambridge.org/abstract_S0022029912000581

How to cite this article:

Ignacio D Frola, Matías S Pellegrino, Gabriel Magnano, José A Giraudó, María C Espeche, María EF Nader-Macias and Cristina I Bogni Histological examination of non-lactating bovine udders inoculated with *Lactobacillus perolens* CRL 1724. *Journal of Dairy Research*, Available on CJO 2012 doi:10.1017/S0022029912000581

Request Permissions : [Click here](#)

Histological examination of non-lactating bovine udders inoculated with *Lactobacillus perolens* CRL 1724

Ignacio D Frola¹, Matías S Pellegrino¹, Gabriel Magnano², José A Giraudo², María C Espeche³,
María EF Nader-Macias³ and Cristina I Bogni^{1*}

¹ Department of Microbiology and Immunology, Faculty of Cs. Ex. Fco-Qcas y Naturales, University of Río Cuarto, Ruta 8 Km 601, X5804ZAB Río Cuarto, Córdoba, Argentina

² Department of Animal Pathology, Faculty of Agronomy and Veterinary, University of Río Cuarto, Ruta 8 Km 601, X5804ZAB Río Cuarto, Córdoba, Argentina

³ Department of Preventive Microbiology, CERELA-CONICET (Centro de Referencia para Lactobacilos-Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina), Chacabuco 145, 4000 San Miguel de Tucumán, Argentina

Received 16 December 2011; accepted for publication 02 July 2012

The effect of intramammary inoculation of *Lactobacillus perolens* CRL 1724 on bovine udders at drying off was evaluated through histological examination of the canal and cistern tissues. The persistence of the strain in the udder 7 d post inoculation was also determined. *Lb. perolens* CRL 1724 was recovered from all mammary quarters and no clinical signs or teat damage were observed after inoculation of 10^6 cfu/ml. The udders showed a normal structural aspect and there were no modifications of the milk appearance. *Lb. perolens* CRL 1724 cells were evidenced on the surface of the epithelial cells of the cistern without causing any morphological modifications or cell alterations. *Lb. perolens* CRL 1724 produces a mild inflammatory reaction, characterized by recruitment of neutrophils to the epithelial zone and a slight hyperaemia into blood vessels. This preliminary study provides important information for further studies directed towards the inclusion of *Lb. perolens* CRL 1724 in the design of probiotic products for preventing bovine mastitis in non-lactating dairy cows.

Keywords: Bovine mastitis, *Lactobacillus perolens* CRL 1724, adherence, non-lactating dairy cows.

Bovine mastitis, one of the most persistent, prevalent and expensive diseases of dairy cows (Viguier et al. 2009; Derong & Zhao, 2010) is defined as 'inflammation of the bovine udder' (Fetrow, 2000). This disease causes considerable distress to the animal, decreased milk production and major economic losses on dairy farms worldwide (Crispie et al. 2008). Dairy cows are highly susceptible to mastitis, mainly during the dry period (Ryan et al. 1999) and antibiotics are currently administered at the drying period to help to eliminate subclinical cases and to prevent the establishment of new intramammary infections (Twomey et al. 2000).

The use of antibiotics for prophylactic treatment is being subjected to considerable debate all over the world because of its perceived connection with the emergence of antibiotic resistance in bacteria, particularly with the increased prevalence of organisms such as methicillin-resistant

Staphylococcus aureus, which are prevalent in nosocomial infections in humans (David & Daum, 2010). Moreover the use of antibiotics to control mastitis generates the appearance of residues in the milk of treated cows (Dalton, 2006). Such concerns have prompted the World Health Organization (WHO) and United Nations Organization for Food and Agriculture Organization (FAO), in the 34th session of the Codex Alimentarius Commission, to increase measures aimed at reducing antimicrobial resistance generated for antibiotic therapies in bovines (González-Gracia, 2011). Some researchers (Browning et al. 1990; Crispie et al. 2004a; Klostermann et al. 2008) have recommended that dry cow antibiotic therapy should not be used as a routine prophylactic measure but rather restricted only to the treatment of infected cows.

During recent years there has been increased interest in developing alternative approaches to the prevention of intramammary infections, particularly in dry cows (Crispie et al. 2004b, 2005; Dallard et al. 2010). In this sense, there is a wide variety of proposals that include the application of vaccines (Giraudo et al. 1997; Pellegrino et al. 2010; Pereira

*For correspondence; e-mail: cbogni@exa.unrc.edu.ar

et al. 2011) and some immunomodulators (Larsen et al. 2004; Dallard et al. 2010). Other measures include the use of 'organic' products, such as the use of probiotic microorganisms (Klostermann et al. 2008) or some of the metabolic or bioactive compounds they are able to produce (Crispie et al. 2005).

For the identification of the beneficial strains that can be used as probiotics, there are many criteria and laboratory assays, many of them recommended by international organizations (FAO & WHO, 2002; International Scientific Association for Probiotics and Prebiotics (ISAPP, 2009). One of the characteristics to be applied for the selection of probiotic strains is the capability of adhesion to epithelial cells (Otero & Nader-Macías, 2007; Both et al. 2010). Adhesion is one of the strategies that promote the further colonization of the bacteria. Also the adhesion of the lactobacilli to the epithelium is the first step in the formation of a barrier or a biofilm to prevent undesirable microbial colonization (Lebeer et al. 2007). Adherence of *Lactobacillus* to epithelial surfaces has been studied in vaginal epithelial cells because of their competition with pathogens for the receptor sites for adhesion (Osset et al. 2001; Nader-Macías et al. 2007).

In earlier work, *Lb. perolens* CRL 1724 was isolated from bovine milk and characterized as a potential probiotic strain (Espeche et al. 2009; Frola et al. 2011). Frola et al. (2011) showed that in vitro *Lb. perolens* CRL 1724 was able to inhibit and co-aggregate with microorganisms considered as major bovine mastitis causing pathogens. Moreover *Lb. perolens* CRL 1724 showed a high efficacy in vitro to adhere to bovine teat canal epithelial cells and was recovered from all the mammary quarters in vivo. No clinical signs or teat damage were observed in lactating cows after the inoculation of 10^6 cfu/ml.

The aim of the present study was to evaluate whether intramammary inoculation of *Lactobacillus perolens* CRL 1724 in dry-period cows produced some type of structural modifications in the gland, using histological examination. The persistence of the strain in the udders 7 d post inoculation was also determined.

Materials and methods

Bacterial strain and culture conditions

Lb. perolens CRL 1724 (Centro de Referencia para Lactobacilos Culture Collection) used in this study was isolated and characterized as potentially probiotic from milk of healthy Holstein cows and characterized according to Espeche et al. (2009) and Frola et al. (2011).

Lb. perolens CRL 1724, resistant to streptomycin, was grown in Man, Rogosa and Sharpe broth (MRS, Britania) at 37 °C for 18 h, and stored in milk yeast extract (MYE) (10 g low fat milk, 0.5 g yeast extract and 1 g glucose per 100 ml) with 12% glycerol at -20 °C. Before performing the experimental assays, bacteria were subcultured three times, every 12–14 h at 37 °C in MRS broth. The bacterial

inoculum was prepared as follows: a culture of the strain (10^9 cfu/ml) incubated for 18 h at 37 °C in MRS broth was centrifuged and the bacterial pellet was washed twice with saline solution (0.8% NaCl). Cells were suspended in 5 ml of saline solution to obtain a concentration of 10^9 cfu/ml. The concentrated preparation was serially diluted in saline solution to 10^6 cfu/ml. This inoculum was fractionated and stored at 4 °C until inoculation was performed (a period no longer than 2 h) (Frola et al. 2011).

Intramammary inoculation of Lb. perolens CRL 1724 at drying off

The study was carried out on an experimental dairy farm in Córdoba, Argentina. The experimental farm operated under conventional management. Five clinically healthy non-lactating Holstein cows were used for the assays. One of the cows from the experimental dairy was used to evaluate the effect of *Lb. perolens* CRL 1724 after intramammary inoculation and the other four cows were used for histological examinations; one of them belonged to the experimental farm and the other three cows were selected from a commercial dairy farm. The animals were of parity 1 and 4 and were in late lactation. Cows were milked twice daily and produced an average of 14 kg milk/d before interruption of lactation. The animals were selected based on previous bacteriological studies and somatic cell counts (SCC). All the quarters used in this work were free of major mastitis-causing pathogens (MCPs) and with SCC in individual quarters <200 000 cells/ml. The animals were inoculated after evening milking. The cleaning of the udders before inoculation, and inoculation procedure were performed following the methodology described by Frola et al. (2011). The unit of study was the mammary quarter. After intramammary inoculation, four of the animals (of parity 1 and showing reproductive problems) were removed from the herd and sent to slaughter at the end of the trial.

One cow was first used to evaluate whether intramammary inoculation of *Lb. perolens* CRL 1724 produced some type of effect on the udder, by evaluation of the clinical signs, milk appearance, SCC and recovery of viable lactobacilli in the milk. Taking in account previous results (Frola et al. 2011) three quarters were infused once on day 0 (D0) with 1 ml containing 10^6 cfu of *Lb. perolens* CRL 1724. One quarter was used as control. To minimize animal handling and to follow animal welfare best practices, no infusion was inoculated in the control quarter.

Three quarters of each one of the remaining four cows were infused with 1 ml of 10^6 cfu of *Lb. perolens* CRL 1724 for histological examination. One quarter was used as control and samples were taken 2 d after intramammary inoculation.

Sampling and bacterial recovery

Before inoculation, foremilk samples were collected from each quarter according to the National Mastitis Council

procedure (National Mastitis Council, 2004) immediately before milking. Milk or dry secretion samples were transported refrigerated (a period no longer than 2 h) to the laboratory and immediately 10 µl was plated onto blood-agar (TSA with 5% of sheep blood) and incubated at 37 °C for 24 h. Bacteria were characterized by standard biochemical tests (Bergey & Holt, 1994). SCC was performed with a Somacount 300 (Bentley) according to the revised protocol of the 148A method C, fluoro-opto-electronic (International Dairy Federation Laboratory, 1995). In all cows, milk or dry secretion samples were obtained 2 d before infusion (D – 2), immediately prior to infusion (D0) and until 2 d (D2) post infusion. One cow, used to evaluate whether the intramammary inoculation of *Lb. perolens* CRL 1724 produces some type of effect on the udder, was sampled daily until day 7 (D7).

Serial dilutions of samples in saline solution were streaked on MRS agar plates in duplicate and incubated at 37 °C for 24–48 h under microaerophilic conditions (5% CO₂, 95% air) for *Lactobacillus* isolation. Isolated colonies were identified as *Lb. perolens* CRL 1724 by phenotypic tests (Gram stain, morphology, catalase activity, nitrate reduction, indole production) and by determination of streptomycin resistance.

Clinical observations and animal care

Clinical signs were monitored throughout the experiment by a veterinarian, every 8 h during the first 24 h and daily until the end of the assay. General attitude and appetite of the cows were observed. The udders were palpated for soreness, swelling, hardness and heat, and the appearance of milk and dry secretion was assessed visually for clots and changes in colour or composition. Animals were cared for in accordance with The International Guiding Principles for Biomedical Research Involving Animals (1985).

Histological examination of the dry udder

Four inoculated cows were sacrificed and examined histologically before sending them to the slaughterhouse. Two representative samples of canal and cistern of bovine mammary gland were removed and fixed in 10% buffered formalin and 2.5% glutaraldehyde (in 0.1 mol/l phosphate buffer, pH 7.4) during 24 h at room temperature for histological examination and transmission electronic microscopy (TEM).

For histological examination, fragments fixed were embedded in paraffin and cut in 5-µm sections. Sections were stained with haematoxylin and eosin (H-E) according to Grignaschi et al. (1983), toluidine blue and Gram, placed on a glass slide and covered with cover-slip according to a method standardized in our laboratory. Samples were examined under high resolution optical microscopy (HROM) and pictures were taken with a Powershot G6, 7.1 megapixels (Canon INC, Japón) digital camera and the software AxioVision Release 4.6.3 (Carl Zeiss, Alemania).

For TEM examination, fixed tissue fragments were fixed overnight in 1% osmium tetroxide and afterwards treated with an aqueous solution of 2% uranyl acetate for 40 min. After fixation, tissues were gradually dehydrated in a series of alcohol solutions of increasing strength, passed through acetone and embedded in Spurr resin. Ultrathin sections were stained with uranyl acetate and lead citrate and examined with Siemens Elmiskop 101 transmission electron microscope.

Results

Effect of intramammary inoculation of *Lb. perolens* CRL 1724

To evaluate the in-vivo performance of *Lb. perolens* CRL 1724, the tolerance of udder to the inoculation of 10⁶ cfu/ml of lactobacilli was first determined in one non-lactating cow. The concentration used was tested previously in lactating cows (Frola et al. 2011). The inoculum was well tolerated by the animal: no clinical signs or teat damage were observed and the udders presented a normal aspect. There were no changes in the appearance of milk secretions. SCC in milk samples was 4.5 × 10⁶ cells/ml after 24 h of intramammary inoculation and decreased to a normal value (1 × 10⁵ cells/ml) until the end of the assay (7 d post inoculation). *Lb. perolens* CRL 1724 was recovered until the end of the assay and from all the inoculated quarters. The highest bacterial recovery value (25 cfu/ml) was obtained 48 h after intramammary inoculations. All the quarters inoculated were negative for mastitis causing pathogen isolation during the whole trial period.

Histological examination

Intramammary inoculation of *Lb. perolens* CRL 1724 evaluated by epithelial tissue sections from the teat canal of the udders (analysed by histological preparations) showed, in all cows, the presence of a stratified squamous epithelium (flat cells) with an important keratin layer (Fig. 1a, c). No difference was found between inoculated and non-inoculated quarters and both quarters presented scarce numbers of blood cells in blood vessels (Fig. 1b, d) in connective tissue. No bacteria were observed after H-E and Gram stains (data not shown).

On the other hand, the epithelial tissue sections obtained from mammary gland cistern, in all samples analysed, were characterized by the presence of an epithelial lining composed of one or two layers of cuboidal or columnar cells without keratin layer. In all cows, mammary gland cistern showed differences between the inoculated and non-inoculated quarter (Fig. 2). The presence of some neutrophils was observed in the connective and epithelial tissue. In this sense, a higher number of neutrophils were observed near the epithelial zone (Fig. 2b, c). Another characteristic detected was a blood vessels hyperaemia which, together with neutrophils, denoted a mild inflammatory reaction




Fig. 1. High resolution optical microscopy showing haematoxylin and eosin stained from representative epithelial tissue sample of bovine mammary gland canal after intramammary inoculation with *Lactobacillus perolens* CRL 1724. (a) and (b) non-inoculated with *Lb. perolens* CRL 1724. Presence of keratin superficial layer (K) and connective tissue with presence of empty blood vessel (black arrow) (20 \times and 40, respectively). (c) and (d) epithelial tissue samples and connective tissue from inoculated quarter without change compared with controls (20 \times and 40, respectively). The bar in each panel is equivalent to 10 μm .

(data not shown). Bacteria adhered to the surface of the cistern epithelial tissue were observed in some quarters inoculated with *Lb. perolens* CRL 1724 (Fig. 2d). This result was confirmed for Gram stain (data not shown).

There were no ultrastructural modifications in the epithelial tissue cells of the canal and cistern of all mammary gland analysed after *Lb. perolens* CRL 1724 inoculation (Figs. 3 & 4, respectively). TEM and RHOM examinations did not show ultrastructural changes, necrosis or apoptosis of epithelial cells, which might be evidenced by the shrinkage of the nucleus and chromatin condensation (pyknosis), outbreak of the nucleus (cariorexix) or the appearance of small cytoplasm vacuoles due to water accumulation. No changes were observed in the nucleus, nucleolus, nuclear membrane and cytoplasmic membrane of the all samples assayed.

Discussion

Bovine mastitis currently produces a serious health and economic problem in dairy farms around the world and antibiotics and disinfectants are routinely used to fight this disease. The generation of non-antibiotic formulations for the treatment and prevention of mastitis in cows has the potential to reduce the veterinary dependence on antibiotics

in the control of this persistent and costly disease (Ryan et al. 1999). In this sense, the use of probiotic bacteria has been widely studied as a novel approach to prevent infections in animals, especially in the gastrointestinal and vaginal tract (Otero & Nader-Macías, 2007; Walsh et al. 2008).

Previous reports showed the isolation and characterization of *Lb. perolens* CRL 1724 to be a potential probiotic strain (Espeche et al. 2009; Frola et al. 2011). *Lb. perolens* CRL 1724 was able to inhibit and co-aggregate in vitro with microorganisms considered as major bovine mastitis causing pathogens. Moreover *Lb. perolens* CRL 1724 showed a high efficacy to adhere to bovine teat canal epithelial cells in vitro. Intramammary inoculation of 10⁶ cfu/ml of the strain in lactating cows showed no udder clinical signs or teat damage and bacteria could be recovered from milk until 15 d after challenge.

In this work the inoculation of 10⁶ cfu/ml of *Lb. perolens* CRL 1724 was also well tolerated by the udders at the beginning of the dry period. No clinical signs or teat damage were observed in the inoculated quarters and the udders presented a normal aspect. There were no changes in the appearance of milk secretions. These results differ from those obtained by Beecher et al. (2009) who reported that all the lactating cows inoculated with *Lactococcus lactis* at 10⁸ cfu/ml experienced swollen udder quarters. Similar results were observed in lactating cows by Frola et al.




Fig. 2. High resolution optical microscope photographs showing haematoxylin and eosin stained from representative epithelial tissue sample of bovine mammary gland cistern after intramammary inoculation with *Lactobacillus perolens* CRL 1724. (a) non-inoculated (100×). (b), (c) and (d) epithelial tissue of inoculated quarter. (b) and (c) presence of neutrophils near to the epithelial zone (black arrows) (100×). (d) bacterial adhered to the surface of epithelial tissue (greys arrows) (100×). The bar in each panel is equivalent to 10 μm.

(2011), where a short-term significant increase in SCC was observed 1 d post inoculation as a normal reaction of the udder, returning to normal values at the end of the trial. In this sense, Crispie et al. (2008) and Klostermann et al. (2008) reported increased values of SCC in the first 2 d after the inoculation of 10^9 cfu/ml of *Lc. lactis* which decreased on days 5 and 7 post inoculation, respectively.

In the present study, *Lb. perolens* CRL 1724 was recovered from the udders until the end of the assay. This could indicate that the strain persisted in the udders. Beecher et al. (2009) observed the recovery of *Lc. lactis* for a period of 72 h post inoculation.

Different authors studied the effect of lactic acid bacteria in the control of mastitis utilizing, for example, a commercial probiotic for oral use in calves (Greene et al. 1991) or substances such as bacteriocins produced by lactic acid bacteria isolated from different niches (Crispie et al. 2005; Cao et al. 2007). In contrast, the host specificity and ecological niche specificity has been shown by different research groups (Ocaña et al. 1999; Otero et al. 2006; Zoetendal et al. 2006). In this work, the effect of intramammary inoculation of a lactobacilli strain isolated from bovine milk on the epithelial tissue of the udders was determined. The preliminary results obtained in this work show that *Lb. perolens* CRL 1724 did not cause damage in the epithelium cells.

Differences between epithelial tissue sections of the cistern of the udder from inoculated and non-inoculated quarters were observed. Neutrophils were observed close to the epithelium and together with hyperaemia of blood vessels denoted a mild inflammatory reaction. In this connection Crispie et al. (2008) showed that administration of the lactococcal culture into the mammary glands of uninfected animals elicited an immunomodulatory effect, with substantial recruitment of polymorphonuclear lymphocytes to the infused quarters and concluded that the mechanism by which the live culture can provide host protection against mastitis infection may be associated with its ability to elicit a rapid immune response. In addition, Beecher et al. (2009) obtained a massive immune response, with an increase in pro-inflammatory genes as IL-1b, IL-8 and CXCR1, after inoculation of a strain of *Lc. lactis* in lactating cows and argued that this result may be one of the immunomodulatory mechanisms by which the bacterium confers its therapeutic effect. On the other hand, according to Sordillo & Streicher (2002) the mammary gland is protected by a variety of defence mechanisms and one of them is innate immunity. This mechanism is mediated by the physical barrier of the teat end, macrophages, neutrophils, natural killer (NK) cells, and certain soluble factors, and if these function adequately, most pathogens are readily eliminated within a short




Fig. 3. Transmission electron micrographs showing representative sample of teat canal epithelial cells after intramammary inoculation with *Lactobacillus perolens* CRL 1724 (a) non-inoculated, nucleus (n), nucleolus (white arrows) and nuclear membrane (black arrows) of epithelial cells (4000 \times). (b) epithelial cells of inoculated quarter with nucleus (n), nucleolus (white arrows) and membrane nuclear (black arrows) without modifications (6000 \times). (K): Keratin.




Fig. 4. High resolution optical microscope photographs showing toluidine blue stained of representative sample of tissue cells of bovine mammary gland cistern after intramammary inoculation with *Lactobacillus perolens* CRL 1724. (a) non-inoculated, mammary cells (black arrows) and blood vessel with scarce presence of blood cells (white arrow) (100 \times). (b) mammary tissue cells of inoculated quarter without morphological modifications in nucleus, nucleolus or membrane nuclear (black arrows); full blood vessel (white arrow) (100 \times). The bar in each panel is equivalent to 10 μm .

period of time and before the specific immune system is activated.

Adhesion to epithelial cells is an important step in both pathogenic infection and probiotic colonization of different

mucosal surfaces such as gastrointestinal, urogenital and respiratory tracts. Adhesion of probiotic microorganisms to the mucosa and the antagonism against pathogens by interference mechanisms have been related to many of the

health benefits attributed to probiotics. In consequence, the ability to adhere to epithelial cells is considered an important criterion for in-vitro selection of probiotics (Morelli, 2000).

Several studies have suggested that *Lactobacillus* adherence is mediated by proteins associated to the external protein S-layer (Henriksson et al. 1991; Frece et al. 2005) while others have suggested a role for lipoteichoic acid and carbohydrate (Chan et al. 1985). Zárate & Nader-Macias (2006) argue that the adhesion to epithelial cells by pathogens is considered an important prerequisite for the onset of infections in the different tracts and that it is the first step in the colonization of the vaginal surface by probiotic microorganisms. Chan et al. (1985) and Reid et al. (1987) affirm that this 'anti-infective' mechanism may involve the blockage of pathogen adherence by both steric hindrance and competition for receptors in the urogenital tract. Bernet et al. (1994) reported that adherent lactic acid bacterial strains may hinder the cell association and invasion by bacterial pathogens and explained such inhibitory effects of lactobacilli by a mechanism of non-specific steric hindrance on the receptors for pathogens. Other reports also showed that lactic acid bacterial strains with adhesion ability may hinder the contact between the epithelial cells and the pathogenic bacteria (Hudault et al. 1997; Coconnier et al. 2000). In consequence, when selecting lactobacilli for probiotic purposes, the adherent strains are preferred in order to form a film on the epithelial tract as a biological barrier against colonization of pathogenic bacteria.

In this work, bacteria on the epithelial tissue surface of inoculated cistern were observed, but no bacteria were observed in the epithelial tissue sections obtained from teat canal. This absence may be due to the keratin layer that coats the canal tissue and would prevent bacterial adherence to the epithelial surfaces as reported by Sandholm & Korhonen (1995). No inoculated samples of both cistern and canal were free of bacteria.

Lb. perolens CRL 1724 (10^6 cfu/ml) did not cause structural or morphological modifications, lesions or necrosis in the epithelial tissue cells of canal and cistern of bovine mammary gland. These observations suggest that intramammary inoculation of *Lb. perolens* CRL 1724 does not produce any damage to the epithelial cells covering the bovine mammary gland tissue studied. This condition, together with the mild inflammation observed at cistern tissue is a promissory result for inclusion of *Lb. perolens* CRL 1724 in a probiotic formulation.

Conclusions

In conclusion, our preliminary results show that *Lb. perolens* CRL 1724 could adhere in vivo to mammary gland cistern of non-lactating cows and generate a mild inflammatory reaction, characterized by the presence of neutrophils in the area close to the epithelia, without causing structural or ultrastructural modifications, lesions or necrosis in tissue cells of bovine mammary gland teat canal and cistern. This

study provides important information for further studies directed towards including *Lb. perolens* CRL 1724 in the design of a probiotic product for preventing bovine mastitis in non-lactating dairy cows.

This work was supported by SECYT-UNRC, MINCYT Córdoba PID280. These are the results obtained from the project 'Design of a probiotic product for bovine mastitis prevention' signed between CONICET and UNRC Res. 2907. Ignacio Daniel Froila is recipients of a fellowship from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

References

- Beecher C, Daly M, Berry DP, Klostermann K, Flynn J & Meaney W 2009 Administration of a live culture of *Lactococcus lactis* DPC 3147 into the bovine mammary gland stimulates the local host immune response, particularly IL-1 and IL-8 gene expression. *Journal of Dairy Research* **76** 340–348
- Bergey DH, Holt JG 1994 *Bergey's Manual of Determinative Bacteriology*. Baltimore, MD, USA: Lippincott Williams & Wilkins
- Bernet MF, Brassart D, Neeser JR & Servin AL 1994 *Lactobacillus acidophilus* LA1 binds to cultured human intestinal cell lines and inhibits cell attachment and cell invasion by enterovirulent bacteria. *Gut* **35** 483–489
- Both E, Gyorgy E, Kibédi-Szabó CZ, Tamás E, Ábrahám B, Miklóssy I & Lányi S 2010 Acid and bile tolerance, adhesion to epithelial cells of probiotic microorganisms. *U.P.B. Science Bulletin Series B* **72** 37–44
- Browning JW, Mein GA, Barton M, Nicholls TJ & Brightling P 1990 Effects of antibiotic therapy at drying off on mastitis in the dry period and in early lactation. *Australian Veterinary Journal* **67** 440–442
- Cao LT, Wu JQ, Xie F, Hu SH & Mo Y 2007 Efficacy of nisin in treatment of clinical mastitis in lactating dairy cows. *Journal of Dairy Science* **90** 3980–3985
- Chan RCY, Reid G, Irvin RT, Bruce AW & Costerton JW 1985 Competitive exclusion of uropathogens from human uroepithelial cells by *Lactobacillus* whole cells and cell wall fragments. *Infection and Immunity* **47** 84–89
- Coconnier MH, Lievin V, Lorrot M & Servin AL 2000 Antagonistic activity of *Lactobacillus acidophilus* LB against intracellular *Salmonella enterica* serovar *Typhimurium* infecting human enterocyte-like Caco-2/TC-7 cells. *Applied and Environmental Microbiology* **66** 1152–1157
- Crispie F, Flynn J, Ross RP, Hill C & Meaney WJ 2004a Update on the development of a novel dry cow therapy using a bismuth-based intramammary teat seal in combination with the bacteriocin lactacin 3147. *Irish Veterinary Journal* **57** 652–656
- Crispie F, Flynn J, Ross PR, Hill C & Meaney WJ 2004b Dry cow therapy with a non-antibiotic intramammary teat seal – a review. *Irish Veterinary Journal* **57** 412–418
- Crispie F, Twomey D, Flynn J, Hill C, Ross P & Meaney W 2005 The lantibiotic lactacin 3147 produced in a milk-based medium improves the efficacy of a bismuth-based teat seal in cattle deliberately infected with *Staphylococcus aureus*. *Journal of Dairy Research* **72** 159–167
- Crispie F, Alonso-Gómez M, Collette O'Loughlin, Klostermann K, Flynn J, Arkins S, Meaney W, Ross RP & Hill C 2008 Intramammary infusion of a live culture for treatment of bovine mastitis: effect of live lactococci on the mammary immune response. *Journal of Dairy Research* **75** 374–384
- Dallard BE, Ortega HH, Iguzquiza IA, Salvetti NR, Quaino OA & Calvino LF 2010 The effect of a single intramammary infusion of a biological response modifier in cows at drying off. *Veterinary Research Communications* **34** 519–532
- Dalton JC 2006 Antibiotic residue prevention in milk and dairy beef. *Western Dairy News* **6** 79
- David MZ & Daum RS 2010 Community-associated methicillin-resistant *Staphylococcus aureus*: Epidemiology and clinical consequences of an emerging epidemic. *Clinical Microbiology Reviews* **23** 616–687

- Derong SYZ & Zhao HY** 2010 Prevalence of bacterial infection responsible for bovine mastitis. *African Journal of Microbiology Research* **4** 1110–1116
- Espeche MC, Otero MC, Sesma F & Nader-Macias MEF** 2009 Screening of surface properties and antagonistic substances production by lactic acid bacterial isolated from the mammary gland of healthy and mastitis cows. *Veterinary Microbiology* **135** 346–357
- FAO, WHO** 2002 *Guidelines for the Evaluation of Probiotics in Food*. Geneva, Switzerland: Food and Agriculture Organization of the United Nations and World Health Organization Working Group
- Fetrow J** 2000 Mastitis: an economic consideration. In *Proceedings of the 29th Annual Meeting of the National Mastitis Council*, Atlanta: Ga. National Mastitis Council. pp. 3–47. Madison, Wisconsin
- Frece J, Kos B, Svetec IK, Zgaga Z, Mrsa V & Suskovic J** 2005 Importance of S-layer proteins in probiotic activity of *Lactobacillus acidophilus* M92. *Journal of Applied Microbiology* **98** 285–292
- Frola ID, Pellegrino MS, Espeche MC, Giraud JA, Nader-Macias MEF & Bogni CI** 2011 Effects of intramammary inoculation of *Lactobacillus perolens* CRL1724 in lactating cows udders. *Journal of Dairy Research* **78** 1–9
- Giraud JA, Calzolari A, Rampone H, Rampone A, Giraud AT & Bogni C** 1997 Field trials of a vaccine against bovine mastitis. 1. Evaluation in heifers. *Journal of Dairy Science* **80** 845–853
- González-Gracia R** 2011 La OMS y la FAO quieren mayor vigilancia de los antibióticos para el ganado. Un problema de seguridad alimentaria mundial. <http://albeitar.portalveterinaria.com/noticia/10289/ACTUALIDAD/oms-fao-quieren-mayor-vigilancia-antibioticos-ganado.html> (Accessed 9 December 2011)
- Greene WA, Gano AM, Smith KL, Hogan JS & Todhunter A** 1991 Comparison of probiotic and antibiotic intramammary therapy of cattle with elevated somatic cell counts. *Journal of Dairy Science* **74** 2976–2981
- Grignaschi V, Diaz N, Alonso M, Lardo M & Licero G** 1983 Citomorfología y Citoquímica Hemáticas. Edit. Britania. Argentina
- Henriksson A, Szewzyk R & Conway PL** 1991 Characteristics of the adhesive determinants of *Lactobacillus fermentum* 104. *Applied and Environmental Microbiology* **57** 499–502
- Hudault S, Lievin V, Bernet-Camard MF & Servin AL** 1997 Antagonistic activity exerted *in vitro* and *in vivo* by *Lactobacillus casei* (strain GC) against *Salmonella typhimurium* C5 infection. *Applied and Environmental Microbiology* **63** 513–518
- International Dairy Federation Laboratory** 1995 *Milk and Milk Products: Detection of Salmonella*. IDF Standard 93B:1005. Brussels, Belgium
- International Guiding Principles for Biomedical Research Involving Animals** 1985 http://www.cioms.ch/publications/guidelines/1985_texts_of_guidelines.htm. (Accessed 2 September 2011)
- International Scientific Association for Probiotics and Prebiotics (ISAPP)** 2009 Probiotics: a consumer guide for making smart choices. http://www.isapp.net/docs/Consumer_Guidelines-probiotic.pdf. (Accessed 8 June 2009)
- Klostermann K, Crispie F, Flynn J, Ross RP, Hill C & Meaney W** 2008 Intramammary infusion of a live culture of *Lactococcus lactis* for treatment of bovine mastitis: comparison with antibiotic treatment in field trials. *Journal of Dairy Research* **75** 365–373
- Larsen MW, Moser C, Hoiby N, Song Z & Kharazmi A** 2004 Ginseng modulates the immune response by induction of interleukin-12 production. *Acta Pathologica Microbiologica et Immunologica* **112** 369–373
- Lebeer S, Verhoeven TLA, Perea-Vélez M, Vanderleyden J & De Keersmaecker SCJ** 2007 Impact of environmental and genetic factors on biofilm formation by the probiotic strain *Lactobacillus rhamnosus* GG. *Applied and Environmental Microbiology* **73** 6768–6775
- Morelli L** 2000 In vitro selection of probiotic lactobacilli: a critical appraisal. *Current Issues in Intestinal Microbiology* **1** 59–67
- National Mastitis Council** 2004 *Microbiological procedures for the diagnosis of bovine udder infection and determination of milk quality*. Fourth Edition. Inc., Arlington, VA, USA, pp. 1–47.
- Nader-Macias MEF, Ocaña VS, Juárez Tomás MS & Silva de Ruiz C** 2007 In *Fundamentos Biológicos, Procesos y Biotecnología de Bacterias Lácticas* (Ed. G Pérez Martínez). España: CSIC
- Ocaña VS, Bru E, Ruiz Holgado A & Nader-Macias ME** 1999 Surface characteristics of lactobacilli isolated from human vagina. *Journal of General and Applied Microbiology* **45** 203–212
- Osset J, García E, Bartolomé RM & Andreu A** 2001 Role of *Lactobacillus* as protector against vaginal candidiasis. *Medicina Clínica* **117** 285–288
- Otero MC, Morelli L & Nader-Macias ME** 2006 Probiotic properties of bovine vaginal lactic acid bacteria to prevent metritis. *Letters in Applied Microbiology* **43** 91–97
- Otero MC & Nader-Macias ME** 2007 *Lactobacillus* adhesion to epithelial cells from bovine vagina. In *Communicating Current Research and Educational Topics and Trends in Applied Microbiology*. pp. 749–757. (Ed. A Méndez-Vilas). Badajoz, Spain: Formatex
- Pellegrino M, Giraud J, Raspanti C, Odierno L & Bogni C** 2010 Efficacy of immunization against bovine mastitis using a *Staphylococcus aureus* avirulent mutant vaccine. *Vaccine* **28** 4523–4528
- Pereira U, Oliveira D, Mesquita L, Costa G & Pereira L** 2011 Efficacy of *Staphylococcus aureus* vaccines for bovine mastitis: A systematic review. *Veterinary Microbiology* **148** 117–124
- Reid G, Cook R & Bruce A** 1987 Examination of strains of *Lactobacilli* for properties that may influence bacterial interference in the urinary tract. *Journal of Urology* **138** 330–335
- Ryan MP, Flynn J, Hill C, Ross RP & Meaney WJ** 1999 The natural food grade inhibitor lactacin 3147 can prevent mastitis in non-lactating dairy cows. *Journal of Dairy Science* **82** 2625–2631
- Sandholm M & Korhonen H** 1995 Infection of the udder-Udder inflammation. In *The Bovine Udder and Mastitis*. pp. 37–48. (Ed. M Sandholm). Helsinki: University of Helsinki Press
- Sordillo LM & Streicher KL** 2002 Mammary gland immunity and mastitis susceptibility. *Journal of Mammary Gland Biology and Neoplasia* **7** 135–146
- Twomey DP, Wheelock AI, Flynn J, Meaney WJ, Hill C & Ross RP** 2000 Protection against *Staphylococcus aureus* mastitis in dairy cows using a bismuth-based teat seal containing the bacteriocin, Lactacin 3147. *Journal of Dairy Science* **83** 1981–1988
- Viguié C, Arora S & Gilmartin N** 2009 Mastitis detection: current trends and future perspectives. *Trends Biotechnology* **27** 486–493
- Walsh MC, Gardiner GE, Hart OM, Lawlor PG, Daly M & Lynch B** 2008 Predominance of a bacteriocin-producing *Lactobacillus salivarius* component of a five-strain probiotic in the porcine ileum and effects on host immune phenotype. *FEMS Microbiology Ecology* **64** 317–327
- Zárate G & Nader-Macias ME** 2006 Influence of probiotic vaginal lactobacilli on in vitro adhesion of urogenital pathogens to vaginal epithelial cells. *Letters in Applied Microbiology* **43**(2) 174–180
- Zoetendal EG, Vaughan EE & de Vos WM** 2006 A microbial world within us. *Molecular Microbiology* **59** 1639–1650