SYNCHRONISING THE TIMING OF CARBON CYCLE, VOLCANISM AND PACING OF THE EARTH'S ORBIT DURING THE EARLY CRETACEOUS

Mathieu Martinez^{1*} | Beatriz Aguirre-Urreta² | Guillaume Dera³ | Marina Lescano² | Julieta Omarini⁴ | Maisa Tunik⁴ | Luis O'Dogherty⁵ | Roque Aguado⁶ | Miguel Company⁷ | Stéphane Bodin⁸

- 1 Université de Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000 Rennes, France; e-mail: *mathieu.martinez@univ-rennes1.fr
- 2| Instituto de Estudios Andinos Don Pablo Groeber, CONICET & Universidad de Buenos Aires, Ciudad Universitaria, Pabellón 2, 1428 Buenos Aires, Argentina
- 3 GET, Université Paul Sabatier, CNRS UMR 5563, IRD, Toulouse, France
- 4| Instituto de Investigación en Paleobiología y Geología, CONICET & Universidad de Río Negro, Sede Alto Valle, 8332 General Roca, Río Negro, Argentina
- 5| Departamento Ciencias de la Tierra, Universidad de Cádiz, CASEM, 11510 Puerto Real, Spain
- 6| Departamento de Geología y CEACTEMA, Universidad de Jaén, Campus Científico-Tecnológico de Linares, 23700 Linares, Spain
- 7| Departamento de Estratigrafía y Paleontología, Universidad de Granada,
 18002 Granada, Spain
- 8| Department of Geoscience, Aarhus University, Høegh-Guldbergs Gade 2, 8000 Aarhus C, Denmark

The Weissert Event (mid-Valanginian), the Faraoni Event (latest Hauterivian), the mid-Barremian and Taxy events in the Barremian are referred to as Episodes of Environmental Change (EECs) (Föllmi 2012). These episodes are associated with accelerated hydrolysis cycles, increased levels of continental weathering and temperature-triggered fertilization of water masses. Other events were recognized in the Berriasian to Barremian, suggesting that these events punctuated the Early Cretaceous (Kujau et al. 2013; Charbonnier et al. 2016; Martinez 2018; Martinez et al. 2020). Uncertainties in the geological time scales, however, preclude the full understanding of the onset, unfolding and termination of EECs. Here, we analyze the amplitude modulation of precession cycles recorded in two Hauterivian sedimentary series in France and Spain to provide a comprehensive and accurate time scale of the Valanginian–Barremian interval based on the stable

405-kyr eccentricity cycle. The new time scale proposed here significantly differs from the Geologic Time Scale 2020 (GTS 2020) (Gradstein et al. 2021). According to our astrochonological framework, the Weissert Event started at 134.56±0.18 Ma, in perfect synchronicity with the peak of volcanic activity of the Paraná-Etendeka Large Igneous Province. The above-mentioned EECs are within a pacing of 2.40 myr of the detrital supply and carbon isotope variations recorded in bulk rock and belemnite rostra. Hence, long eccentricity cycles were key parameters in the regulation of climate and carbon cycles during the Early Cretaceous through changes in the detrital and nutrient supply, oceanic fertilization, organic carbon storage and global sea level. We also demonstrate that the humid peak related to the Weissert Event is driven by the pacing of the long orbital cycles, in spite of the emplacement of the Paraná-Etendeka province. Nevertheless, in comparison to other EECs of the Valanginian-Barremian, the Weissert Event appears to be a singularly long event with profound impact on climate and marine ecosystems. We posit that this is a consequence of the concomitant effect of the emplacement of the Paraná-Etendeka province and the long orbital cycles.

REFERENCES

Charbonnier, G., Duchamp-Alphonse, S., Adatte, T., Föllmi, K. B., Spangenberg, J. E., Gardin, S., Galbrun, B. and Colin, C. 2016. Eccentricity paced monsoon-like system along the north-western Tethyan margin during the Valanginian (Early Cretaceous): new insights from detrital and nutrient fluxes into the Vocontian Basin (SE France). Palaeogeography, Palaeoclimatology, Palaeoecology, 443, 145-155.

Föllmi, K.B. 2012. Early Cretaceous life, climate and anoxia. Cretaceous Research, 35, 230-257.

Gradstein, F.M., Ogg, J.G., Schmitz, M.D. and Ogg, G.M. 2020. Geologic time scale 2020, 1357 pp. Elsevier BV; Amsterdam, The Netherlands.

Kujau, A., Heimhofer, U., Hochuli, P.A., Pauly, S., Morales, C., Adatte, T., Föllmi, K., Ploch, I. and Mutterlose, J. 2013. Reconstructing Valanginian (Early Cretaceous) mid-latitude vegetation and climate dynamics based on spore–pollen assemblages. Review of Palaeobotany and Palynology, 197, 50-69.

Martinez, M. 2018. Mechanisms of preservation of the eccentricity and longer-term Milankovitch cycles in detrital supply and carbonate production in hemipelagic marl-limestone alternations. In: Montenari, M. (Ed.), Stratigraphy & Timescales, vol. 3, 189-218. Elsevier Academic Press; London, United Kingdom.

Martinez, M., Aguado, R., Company, M., Sandoval, J. and O'Dogherty, L. 2020. Integrated astrochronology of the Barremian Stage (Early Cretaceous) and its biostratigraphic subdivisions. Global and Planetary Change, 195, 103368.