



MYO1C, MYO6 AND MYO18A ARE NECESSARY FOR CHLAMYDIA TRACHOMATIS DEVELOPMENT

Cuervo Bustamante ME¹, Del Balzo D^{1,2}, Damiani MT^{1,2}, Capmany A¹.

¹ Instituto de Instituto de Bioquímica y Biotecnología, Facultad de Ciencias Médicas, UNCUYO. ² Laboratorio de Bioquímica e Inmunidad, IMBECU-CONICET. E-mail: anahicapmany@hotmail.com

Chlamydia trachomatis (CT) is an obligate intracellular bacterium and the most frequent bacterial agent of sexually transmitted infections. The latest research has placed CT as a risk factor of cellular transformation, which could lead to cervical or ovarian cancer development. This bacterium induces several alterations in the host cell such as inhibition of apoptosis and cytokinesis, a decrease in molecules involved in cell adhesion, and loss of front–rear polarity in migrating cells. The cytoskeleton and the associated proteins are the main factors that ensure polarized trafficking and a correct cell division. Particularly, myosins stand out, not only because of their role in actin cytoskeleton arrangement but also because of their implication in vesicular transport. By confocal microscopy, we observed that MYO1C is recruited to the chlamydial inclusion at 24 h post-infection. We recently published that MYO1C stabilizes actin at the Golgi apparatus facilitating the arrival of incoming transport carriers at this organelle. Strikingly, CT establishes a close relationship with the Golgi apparatus, receiving from this organelle a continuous supply of vesicles loaded with essential nutrients. Thus, CT could recruit this myosin as a strategy to ensure the arrival of post-Golgi vesicles. Interestingly, the knockdown of MYO1C impairs the CT development, assessed by flow cytometry and confocal microscopy. Moreover, this function could be shared with MYO6 and MYO18A that are also necessary for the normal chlamydial development and function of the Golgi apparatus. Our results suggest that MYO1C, MYO6, and MYO18A are manipulated by CT to ensure its development.

CB-P11

CHLAMYDIA TRACHOMATIS CAUSES THE INTRACELLULAR REDISTRIBUTION OF MHC-I AND IMPAIRS ANTIGEN PRESENTATION IN DENDRITIC CELLS

Del Balzo D¹; Capmany A¹; Cebrián I², Damiani MT¹.

¹Laboratorio de Bioquímica e Inmunidad. Instituto de Bioquímica y Biotecnología. FCM-UNCuyo. IMBECU-CONICET. ²IHEM-CONICET, FCEN- UNCUYO. Mendoza. E-mail: ddelbalzo@mendoza-conicet.gob.ar

Chlamydia trachomatis (CT) is the most frequent bacterial cause of sexually transmitted infections worldwide. This highly adapted intracellular bacterium has evolved multiple strategies to hide inside cells. However, little is known about the molecular mechanisms underlying CT evasion of the immune response. Dendritic cells (DCs) are the most efficient antigen-presenting cells of the immune system and an essential link between innate and adaptive immunity. Therefore, DCs could play a key role in CT's clearance. In this study, we analyzed in CT-infected DCs the process of cross-presentation, in which exogenous antigens are associated with MHC-I molecules to activate CD8⁺ T lymphocytes. By confocal microscopy and flow cytometry-based approaches, we observed, after chlamydial infection, a decrease in MHC-I molecules exposed at the plasma membrane while they are redistributed intracellularly. However, the total amount of MHC-I molecules did not change after infection, as assessed by western blot analysis. Finally, we found that CT-infected DCs were less efficient than non-infected ones to cross-present the model antigen Ovalbumin, as measured colorimetrically by the activation of the antigen-specific CD8⁺ T cell hybrid called B3Z. Altogether, these findings indicate that CT infection impairs antigen-cross presentation in DCs through the disturbance of MHC-I transport.

CB-P12

NOVEL CAFFEINE ANALOGS AS POTENTIAL LEADERS ON THE CHOLINERGIC SYSTEM

Fabiani C¹, Biscussi B², Munafó JP¹, Salvador G¹, Corradi J¹, Murray AP², Antollini SS¹.

¹INIBIBB, Dpto. Biol. Bioq. Farmacia; ²INQUISUR, Dpto. Qca. Univ. Nac. del Sur – CONICET, Bahía Blanca, Argentina. E-mail: cfabiani@inibibb-conicet.gob.ar

Cholinergic deficit is regarded as an important factor responsible for Alzheimer's disease symptoms. Two molecular targets for the treatment of this disease are acetylcholinesterase (AChE) and nicotinic receptor (nAChR). We previously demonstrated that caffeine has a dual effect on muscle and $\alpha 7$ nAChRs, behaving as a weak agonist at low concentrations and as a negative modulator at high concentrations. Furthermore, it is well-known that caffeine also acts as an inhibitor of AChE. The aim of this work was to synthesize more potent caffeine analogs with a dual effect on the cholinergic system by inhibiting AChE and potentiating nAChRs. With this objective, a theophylline fragment, resembling the caffeine chemical structure, was connected with a pyrrole fragment, which is present in the nicotinic chemical structure, through homologation from 3 to 6 carbon atoms (C_n). We first tested the capacity of the different compounds to inhibit the AChE. We found that whether theophylline alone inhibited the enzyme, pyrrolidine did not. With respect to C_n , they all can inhibit the AChE at concentrations of 100, 200, and 400 μ M, having C6 the strongest effect. We then explored if theophylline, pyrrolidine, and C_n influence the nAChR conformational state. To this end, we used the AChR conformational-sensitive fluorescence probe crystal violet (CrV) and AChR-rich membranes from *Torpedo californica*. We found that whether pyrrolidine induced changes in the K_D values of CrV taking the nAChR to a state close to the desensitized one at concentrations of 200 and 400 μ M, theophylline did not show a significant change in the K_D value. The combined analogs also produced changes in the K_D values of CrV. This effect was dependent on the length of homologation, being C5 and C6 the most potent analogs with effect at concentrations lower than 50 nM. To understand the molecular mechanism underlying the conformational changes of the nAChR, we expressed adult muscle nAChR in HEK293 cells and performed single-channel recordings with different C_n concentrations. We found that C5 activated muscle nAChR at very low concentrations (from 0.01 pM). At the highest tested concentration (30 μ M), we observed a decrease in the mean open duration, which suggests that C5 also acts as an open channel blocker. As a partial conclusion, we can say that we have synthesized more potent caffeine analogs through the combination of caffeine and nicotinic structures. The effect of theophylline on AChE, the effect of pyrrole on AChR, and the effect of C_n on