

A conceptual framework for nomenclatural stability and validity of medically important fungi: a proposed global consensus guideline for fungal name changes supported by ABP, ASM, CLSI, ECMM, ESCMID-EFISG, EUCAST-AFST, FDLC, IDSA, ISHAM, MMSA, and MSGERC

Sybren de Hoog,^{1,2,3,4,5,6} Thomas J. Walsh,^{6,7,8,9,10,11,12,13,14,15} Sarah A. Ahmed,^{1,2,6} Ana Alastruey-Izquierdo,^{6,16,17} Barbara D. Alexander,^{10,18} Maiken Cavling Arendrup,^{19,20} Esther Babady,^{10,21} Feng-Yan Bai,^{22,23,24,25,26,27} Joan-Miquel Balada-Llasat,^{10,28} Andrew Borman,²⁹ Anuradha Chowdhary,^{17,30} Andrew Clark,^{10,31} Robert C. Colgrove,^{32,33} Oliver A. Cornely,^{12,17,34,35} Tanis C. Dingle,^{10,13,36} Philippe J. Dufresne,^{10,13,37} Jeff Fuller,^{10,38} Jean-Pierre Gangneux,^{12,39} Connie Gibas,⁴⁰ Heather Glasgow,^{10,41} Yvonne Gräser,⁴² Jacques Guillot,^{26,43} Andreas H. Groll,^{17,44} Gerhard Haase,⁴⁵ Kimberly Hanson,^{10,46} Amanda Harrington,^{10,47} David L. Hawksworth,^{48,49,50,51,52,53} Randall T. Hayden,^{10,13,41} Martin Hoenigl,^{11,12,54,55,56} Vit Hubka,⁵⁷ Kristie Johnson,^{10,58} Julianne V. Kus,^{10,59,60} Ruoyu Li,^{3,5,15,17,20,24} Jacques F. Meis,^{1,15,34,35} Michaela Lackner,^{6,61} Fanny Lanternier,⁶² Sixto M. Leal Jr.,^{10,11,13,63} Francesca Lee,^{10,31} Shawn R. Lockhart,^{10,64} Paul Luethy,^{10,58} Isabella Martin,^{10,65} Kyung J. Kwon-Chung,⁶⁶ Wieland Meyer,^{9,67} M. Hong Nguyen,^{10,11,14,68} Luis Ostrosky-Zeichner,^{11,69} Elizabeth Palavecino,^{10,70} Preeti Pancholi,^{10,28} Peter G. Pappas,^{11,63} Gary W. Procop,^{10,13,71,72} Scott A. Redhead,^{9,73} Daniel D. Rhoads,^{74,75,76} Stefan Riedel,^{10,77} Bryan Stevens,^{10,68} Kaede Ota Sullivan,^{10,78} Paschalis Vergidis,^{10,79} Emmanuel Roilides,^{6,12,17,80} Amir Seyedmousavi,^{10,17,26,81} Lili Tao,^{10,82} Vania A. Vicente,⁴ Roxana G. Vitale,^{83,84} Qi-Ming Wang,⁸⁵ Nancy L. Wengenack,^{10,79} Lars Westblade,^{10,86} Nathan Wiederhold,^{10,11,13,14,40} Lewis White,⁸⁷ Christina M. Wojewoda,⁸⁸ Sean X. Zhang^{6,10,89}

AUTHOR AFFILIATIONS See affiliation list on p. 16.

ABSTRACT The rapid pace of name changes of medically important fungi is creating challenges for clinical laboratories and clinicians involved in patient care. We describe two sources of name change which have different drivers, at the species versus the genus level. Some suggestions are made here to reduce the number of name changes. We urge taxonomists to provide diagnostic markers of taxonomic novelties. Given the instability of phylogenetic trees due to variable taxon sampling, we advocate to maintain genera at the largest possible size. Reporting of identified species in complexes or series should where possible comprise both the name of the overarching species and that of the molecular sibling, often cryptic species. Because the use of different names for the same species will be unavoidable for many years to come, an open access online database of the names of all medically important fungi, with proper nomenclatural designation and synonymy, is essential. We further recommend that while taxonomic discovery continues, the adaptation of new name changes by clinical laboratories and clinicians be reviewed routinely by a standing committee for validation and stability over time, with reference to an open access database, wherein reasons for changes are listed in a transparent way.

KEYWORDS nomenclature, taxonomy, fungi

Development of the modern naming system in medical mycology

Advanced and novel diagnostic and research methods, particularly nucleic-acids sequencing, have revolutionized microbial taxonomy. Since rearrangements in the Tree of Life are closely linked to the names of the newly recognized entities, name

Editor Romney M. Humphries, Vanderbilt University Medical Center, Nashville, Tennessee, USA

Address correspondence to Thomas J. Walsh, thomaswalshmd@gmail.com, Sybren de Hoog, sybren.dehoog@radboudumc.nl, or Sean X. Zhang, szhang28@jhmi.edu.

The authors declare no conflict of interest.

Published 26 October 2023

Copyright © 2023 de Hoog et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

changes are inevitable. This happens everywhere in microbial taxonomy and is the result of scientific progress. In the kingdom *Fungi*, taxa that were previously delineated by morphological features or physiological profiles are being recharacterized by sequence data underlying their phylogenetic positions. However, unlike most prokaryotic microbes, fungi exhibit a wealth of distinguishing phenotypic characteristics which have served as the basis for accruing clinical data over the centuries. Many fungi have a complicated life cycle, with one or more asexual and sexual forms of sporulation that have very different appearances and are produced under different growth conditions. In the past, it was often not known that these different forms, or morphs, belonged to the same species. This problem was mitigated by the development of a naming system with two separate categories: one name for the sexual morph (teleomorph), and one (or more) for each asexual morph (anamorph). As sexuality is mostly expressed in the environment, and asexual morphs are often preponderant *in vitro* and in the animal host, the connection between the two or more life forms remained problematic despite increasing availability of DNA data, awaiting experimental establishment. Taxonomic categories above the species level were based on the sexual morph and prioritized, while the separate system for asexual ones was generally acknowledged to be artificial—although some authors also published names of families and even higher ranks based only on asexual morphs. Although imperfect, abandoning the knowledge that has been accrued during 270 years of mycological research under this “old” nomenclature system is more painful than in other areas of research, particularly as the old nomenclature is critically linked to patient management.

Challenges and limitations of molecular taxonomy for medically important fungi

With DNA sequencing and other non-microscopic methods, able to firmly establish species, the system with dual names became increasingly obsolete. After some 20 years of debate, and strongly divided opinions voiced by a Special Committee on alternate names (1), followed by a symposium “One Fungus One Name” held in Amsterdam, in 2011, many of the mycological community came to a consensus to abandon the classical separate naming of different morphs of the same species with the “Amsterdam Declaration” (2). Subsequently, this new direction was formally proposed on the debate “floor” by Redhead (3), disputed, and, finally, adopted by the international community at the International Botanical Congress held in Melbourne in 2011. The change was effective from 2011 and was retro-active (4). This enabled a closer representation of the natural system than phenotypic characterization. However, since many names used in medical and veterinary mycology were in the asexual-based artificial system, the need arose for a fundamentally different approach in systematization. Formal procedures were introduced to keep the number of changes within limits through protected lists of names developed by working groups operating under the auspices of the International Commission on the Taxonomy of Fungi. The ensuing name changes of the combined transition toward molecular phylogeny and priority of the oldest name irrespective of sexual states are dramatic, leaving few names unaffected among the medical fungi. Comparing the second and fourth editions of the *Atlas of Clinical Fungi*, of the names accepted in 2020 (5), only 30% were still used in the same sense as in 2000. This demonstrates an overwhelming impact of molecular taxonomy on medical mycology, larger than any other discipline within clinical microbiology practice.

Numerous authors have expressed their concerns about the pace of name change in mycology and provided lists of recommended names primarily focused on the species level (6–10), while others recommended to accept and adopt generic name changes (11–13). The present paper aims to mitigate some of these problematic shifts with proposals, which we think are practicable and minimally disruptive for the advancement of science. Certainly, naming of medically important fungi needs a thorough renovation. In many historical handbooks and guidelines, grand subdivisions are based on obsolete morphological criteria, such as “*Dematiaceae*,” a family designation for fungi with often

large conidia and conidiophores carrying melanin in the cell wall, or “*Coelomyces*,” a class rank name for those forming asexual cup- or flask-like spore-bearing structures anywhere in their asexual life cycle.

Molecular phylogeny has enabled major advances in providing logical coherence between taxonomic entities (14) and has led to better understanding of the origin, relationships, and properties of fungi—including whether they are likely to be pathogens of concern or mycotoxin producers. Phylogeny sheds light upon such potentially shared ecological strategies between species. Whenever obvious benefits of the new naming system exist, name changes are easily accepted by stakeholders in the medical community. An example of one such easy adoption is the change of the intracellular pathogen *Talaromyces marneffe* separated from the strictly saprobic *Penicillium* species (15). The natural classification of dermatophytes (16) also met limited resistance. Another example is the segregation of morphologically similar, but phylogenetically extremely remotely related (i.e., ascomycetous versus basidiomycetous) fungi, now placed in separate genera [*Geotrichum/Trichosporon* (17); *Sporothrix/Quambalaria* (18)]. A more recent example can be found in the replacement of the genus name *Phialemoniopsis/Phialemonium* by *Thyridium* which turned out to be the name given earlier to the sexual morph (19).

However, reestablishing the correct systematic position of tens of thousands fungal names cannot be done by a simple declaration, and in some groups of fungi, molecular data is not easy to obtain. The above-mentioned, well-accepted examples share their application of a holistic, biological approach to taxonomy, demonstrating that the newly separated groups are fundamentally different in life cycle, habitat choice and clinically relevant parameters, with molecular phylogeny as a supporting feature facilitating definite identifications. This demonstrates an optimal approach to reach meaningful taxonomic name changes in mycology. In contrast, resistance against change invariably entails cases where the change involves a relatively homogeneous group without clear character difference, which is divided exclusively based on phylogeny. An example is the rearrangement of the ecologically similar (20) genera *Curvularia* and *Bipolaris* where the multi-character phenotypic separation did not match with molecular barcoding (21). Conversely, species of *Chaetomium* were assigned to several novel genera (22). The latter study made these rearrangements with a data set containing strains from indoor habitats alone, implying that criteria other than phylogeny were insignificant; such cases should be reconsidered using a wider range of named species and ecologies.

We advocate that name changes of medically important fungi should be meaningful and carefully applied to reduce the potential for confusion by those responsible for patient care. Name changes are effective and will reach wide application only when based on fundamental differences that have shaped evolution and have clinical relevance. Two levels of diversity, i.e., that of the species and of the genus, have entirely different drivers (7, 8) which are discussed separately, i.e., the subdivision of a species, and the rearrangement and splitting of genera.

Species diversity: The fragmenting epithet

The borderline between species is not clear-cut, and there are over 30 different species concepts used across biology (23). The classical biological concept is not directly applicable to microbes. Genetic composition and viability of progeny may vary anywhere between 0% and 100% due to all sorts of constraints. Sexually hyperactive strains are able to mate beyond established species borders (24, 25), and this is recognized as hybridization in some plant pathogenic fungi where hybrid nomenclature has been introduced. Conversely, progeny may lack genetic recombination (26–28) or may produce sterile hybrids unable to mate (29, 30). Crosses appearing to be sexual may be uniparental, underlining that fungi are able to propagate asexually over extended periods. Many fungi have alternative sexual strategies, which do not require an opposite mating partner (31). Ideally, a species is genetically focused by genetic interaction, but

most fungi multiply parts of their life cycle clonally to increase successful genotypes. Asexual reproduction may be dominant, and sexuality often remains cryptic.

The near-absence of sexuality leads to fragmentation of the species into numerous molecular siblings. In taxonomic practice in fungi, the siblings are recognized by mutations in sections of barcoding loci, such as the rDNA ITS regions, *CAM*, *rPB1*, *rPB2*, *TEF1*, or *TUB2* (*BenA*) (32). Genealogical concordance enables *in silico* detection of sexual recombination, providing an operational criterion to verify the species borderline. In taxonomic practice of large data sets, however, most authors use concatenated barcoding sequences to provide molecular distances that are judged sufficient for novel species description. For example, depending on the genes studied, numerous siblings were observed within the genera *Cladosporium* (33) distinguishing 54 siblings in the *C. cladosporioides* complex and *Fusarium* (34) distinguishing 74 siblings in the *F. fujikuroi* complex. In plant pathogens, clonal expansion may occur after horizontal gene transfer of pathogenicity islands (35) leading to lineages, which are sometimes known as special forms. These emerging genotypes may be sampled more frequently as a result and have a higher chance of being detected, giving the false impression of obligate host adaptation. However, the responsible horizontally transferred accessory chromosomes can be dispensable (36), and the species can infect another susceptible host, as in the case of the banana-fruit infecting pathogen *Fusarium musae* that has been encountered in human infection and suggested as a possible health concern (37–39).

Decreased sexuality leads to higher clonal diversity (40, 41). An example from clinical fungi is given by dermatophytes, which lose sexuality upon adaptation to the human host, and whose mating types evolve differentially (42, 43). Conversely, numerous genotypes without ecological or clinical differentiation may emerge within a single species (44). Thus, phylogenetic distance in a particular gene region alone is not always a sufficient criterion for novel species description. Distinction of clones based only on molecular difference answers epidemiological questions and is essential to reveal sources of contamination and routes of infection. Speciation, however, is a slower process, with slight differences in ecological preferences of drivers of future separation that ultimately leads to loss of recombination ability. Many barcoding markers are not transcribed and do not result in different evolutionarily relevant properties; they just are a proxy of relevant differences elsewhere in the genome. Addition of more genes should increase the stability of the tree (45).

Species are an amalgamation of genetically deviating lineages, bordered by increasing inability to recombine and produce viable progeny. In contrast to the phenotypic approach, molecular data enable recognition of each lineage. Rather than species, the concept of “species complex” has become a trend. A complex combines molecular siblings without known gene flow under a single umbrella. In most fungal groups, the potential ability of hybridization and recombination between siblings has not been tested. Multi-locus analysis of non-transcribed markers provides potential diversity which is close to infinite (46). Novel names for the individual lineages may conceal the close affinity to the classical species of the aggregate.

Suggestions for the stability of species names in medical mycology

Given the above-described fragmentation of species into named molecular siblings and to promote stability of species names, taxonomy should be separated from epidemiology. Taxonomy and epidemiology both involve microbial diversity and overlap because species are distinguished from intraspecific lineages genetic separation which is difficult to establish. Rules of botanical nomenclature such as on typification need to be evaluated retrospectively and numerous debatable cases will appear (47). In addition, medical mycology covers only a very small fragment of global biodiversity studies, and endeavoring to impose rules from a medical perspective to the much larger study areas in agriculture, ecology, and industry and involving the entire fungal kingdom is unproductive. Rather, we would suggest adoption of the following recommendations:

- For closely related entities, official nomenclatural categories such as subspecies, variety, and form are available. Individual clones and genotypes below the concordant species level are preferably numbered rather than named (48) as long as they do not fulfill a number of criteria that are in use to define species (outlined above). This enhances the link to the overarching species name and facilitates the connection with existing literature.
- For the description of closely related species, data besides phylogenetic distance, such as the absence of recombination by genealogical concordance, and presence of phenotypic, ecological, clinical, or evolutionarily relevant parameters should be included.
- The species complex, also known as a species aggregate or series, is a practical recommendation rather than a solid scientific conclusion and, thus, may be subject to improved circumscription, covering more siblings defined with the ex-type strain as reference.
- To establish the classical species as the portal toward all existing literature, it is recommended that the overarching species complex name is always mentioned, followed by the sibling's or cryptic species name.
- As species names are linked to (epi)type material as the ultimate reference, this should be accessible to investigation, and hence, deposition of a living culture in a reference collection should become mandatory in mycology for those fungi that can be cultured; if the type is a drawing or in inaccessible fungarium or deteriorated herbarium material and, therefore, unrecognizable molecularly, epi-typification or deposition of a reference culture is strongly encouraged.
- Mycologists introducing changes near the species level should be required to provide criteria with which the species at hand can be recognized, i.e., a diagnosis, as is already recommended in the *Code*.
- Name changes generally acquire wide recognition only after the underlying taxonomy has been confirmed in peer-reviewed papers by separate groups of researchers using trees with different taxon sampling. Name stability and validation should be held to the highest possible standard for medically important fungi where clinical decisions based upon that information directly influence patient management.
- As taxonomic science progresses, a committee composed of clinical microbiologists, physicians, medical mycologists, and taxonomists would review proposed name changes for medical relevance, validity, and stability for adoption in clinical laboratories and patient care.

Genus diversity: the phylogenetic framework

At the genus level, the problems are entirely different. While species are subdivided because diverse entities are distinguishable by modern methodology, at the genus level, the central question is the position in the phylogenetic Tree of Life. To this aim, markers with a lower mutation rate are used, particularly those of the ribosomal repeat. Numerous methods have been developed to optimally reflect the course of evolution (49). Different gene trees are expected to be concordant albeit with different resolution. Reclassification of a fungus leads to a new combination of a genus name where the original species epithet is maintained. Consequently, generic rearrangements have become a major source of naming instability.

In contrast to the species, there are no operational molecular criteria to define and circumscribe a genus. All taxonomic entities above the species level can be introduced at will. Indeed, different traditions in the respective research areas have led to genera enormously deviating in size and level of intrageneric diversity. Construction of phylogenetic trees has become the nearly exclusive approach in establishing the position and size of genera. However, as phylogenetic trees are fundamentally relative, being based on mutual comparison of its members, generally using limited numbers of barcoding gene regions which are for diagnostics rather than for taxonomy, they

suffer from inherent instability during the early years of molecular exploration. Although methods of tree reconstruction are highly sophisticated, the underlying taxon sampling effect causes variation with every selection of objects in the tree. Sampling may be relatively complete in well-known groups with a long history of research, but a balanced overview of extant diversity concerns selected groups with practical significance rather than complete genera.

In common taxonomic practice, the main generic parameter is phylogenetic distance. In this approach, genera are statistically supported aggregates of similar species as recognizable clades by conserved, single or concatenated markers. Without analysis of concordance of genes, molecular phylogeny alone is one-dimensional, similar to the obsolete approach of using only microscopic morphology. Different levels of diversity have largely been determined by the taxonomic history of the genus at hand. Many newly created genera are clades that contain just a few species. Rearrangement of such clades requires creation of other genera; small genera comprising just a few species, thus, have a large risk of further fragmentation. As an example, *Scopulariopsis* and *Microascus* were previously distinguished as being asexual or sexual, respectively; presently, the names are used for two groups that are molecularly distinct, with a small genus *Pithoascus* between the clades that contain the type species of the two genera (50). Phenotypic descriptions of *Scopulariopsis* and *Microascus* in the new concepts are nearly identical. In such a case, synonymy of *Pithoascus* with the oldest generic name, *Microascus*, would have required just a few name changes of rare fungi and may be a more pragmatic approach. Another example is the afore-mentioned separation of *Curvularia* and *Bipolaris*, where phylogeny did not match with classical phenotypes. Prioritizing phylogeny, Manamgoda et al. (21) reshuffled species of both genera, but a more parsimonious solution would have been the recognition that the bipartition apparently does not exist, giving priority to the single genus name *Curvularia*.

In contrast, the large genus *Aspergillus* with its characteristic conidiophores was recognized as a group since its establishment in 1809 and now comprises numerous smaller clades. The genus is monophyletic (51), but distances between ultimate members of the genus clade are much larger than usual. In addition, fungi without the characteristic aspergillus-like conidiophores may appear to contain similar genotypes, leading to merging of such genera (e.g., *Phialosimplex*, *Polypaecilum*) with *Aspergillus* (52). Broader generic circumscriptions cover smaller ones; genera then do not fall apart but tend to become larger. In general, it may be nomenclaturally advantageous to maintain broad generic concepts for taxa like *Aspergillus*, *Chaetomium*, and *Fusarium* in their classical sense, as long as they are monophyletic. Smaller genera will appear synonymous, but this usually involves a lower number of name changes.

In the yeasts, classical genera are mostly not monophyletic. The clinically relevant genera that have been classified in *Candida* on the basis of physiological criteria, phylogenetically were found to belong to eight families (5), and therefore, the genus in the traditional sense is untenable. Some genera could coincide with the current level of family ranks, e.g., *Debaryomycetaceae* containing the single genus *Candida* which might keep future instability within limits.

Genera will be meaningfully distinguished when the clades are consistently supported by high bootstrap values. This was successful with the dermatophytes mentioned above (16), but statistical support of the backbone of trees is often lacking or minimal. For example, the family *Herpotrichiellaceae* (black yeasts and allies) in *Chaetothyriales* has been analyzed repeatedly using different genes and data sets but achieving poor statistical support below the family level (53). The authors decided to leave the genera defined by morphology despite the known disagreement with phylogeny. This provides temporary nomenclatural stability, but such genera remain highly polyphyletic, and treating unrelated species under the same generic name, as done, e.g., by Thitla et al. (38) for *Exophiala*, cannot be recommended in the long run.

There may be an optimal size of genera. Extensive divisions resulting in unique genera for many medically important species risks loss of phylogenetic coherence and

ease of recognition. On the other hand, it is difficult to develop diagnostic criteria for genera that are too broad.

Suggestions for stability of the genus concept in medical mycology

- Genera are preferably based on phenotypic characteristics supplemental to molecular distance, reflecting main ecological and medical significance, or evolutionary or other behavioral trends, such that their separation is likely to be widely understood and accepted.
- Names of large-sized genera such as *Aspergillus* tend to be more stable than small genera containing just a few species, based upon phylogenetic distance alone; provided that the genus is monophyletic, it is recommended to maintain the size of genera if there is no compelling reason to break them up.
- Small and phenotypically similar genera in the same clade can be combined if this reduces the number of necessary name changes.
- Authors introducing name changes at the generic level are requested to provide or select criteria with which the genus at hand can be recognized.
- Name changes are preferably adopted in routine practice only after the underlying taxonomy has been confirmed in several papers by independent authors; an evaluating Committee under the auspices of ISHAM is preferable.
- Remember that there are procedures under the *Code* to avoid the necessity to replace a well-known genus name by a less known one, using a conservation process, while also a change of the original type species is possible.

DISCUSSION

Taxonomy is not a galactic spaceship operating in a scientific vacuum. On the contrary: thousands of diagnostic laboratories worldwide in all areas of microbiology apply the outcome of this research on a daily basis for patient care. There are few areas of fundamental science with comparable practical implications. Nomenclatural changes should, therefore, be scientifically sound and based on convincing evidence (54, 55), taking into account phenotypic, biological, and ecological features as well as clinically relevant characteristics of pathogenicity and specifically antifungal resistance patterns. In the absence of established scientific criteria for delimiting genera, the default should be that proposed reclassification should benefit the clinical user. An ongoing discussion relevant to medical mycology concerns the bipartition of *Fusarium* in *Fusarium s.str.* and *Neocosmospora*. The latter is an interesting showcase of the consequences of generic fragmentation. Although the phylogenetic distance between *Neocosmospora* and *Fusarium* may justify their separation, it necessitates the maintenance of 21 additional fusarium-like genera, which are phylogenetically more distant from *Fusarium s. str.* than *Neocosmospora*. The separation of *Neocosmospora* is not necessary for the medical community to be adopted immediately; this can wait until taxonomists stabilize the classification over years of investigation and consensus. A similar appeal was made by Redhead (56), who in proposing a reclassification of coprinus-like species, stated that we do not need to adopt these names immediately; instead, we may be conservative initially. Following molecular testing by different laboratories, the pathogens named from asexual cultures, *Hormographiella aspergillata* and *H. verticillata*, were subsequently synonymized with the mushroom species *Coprinopsis cinereus* and *Coprinellus domesticus*, respectively.

At the species level, we are able to show massive amounts of difference below the level of genetic exchange, which remains the ultimate criterion for conspecificity (organisms belonging to the same species). Although this borderline is vague, subject to many exceptions, and often difficult to establish in research areas outside microbiology, it is uncommon to attribute species status to interbreeding entities. In several groups, more entities can be distinguished than is taxonomically meaningful. In dermatophytes, Tang et al. (48) made a recommendation to formally name only those siblings that are relevant to clinical practice. For example, Bian et al. (57) synonymized several of

the described entities in the *Aspergillus niger* clade and Sklenář et al. (58) in the *A. versicolores* clade. The requirement to distinguish large numbers of molecular siblings without clinical relevance would have a negative impact on clinical practice and patient care when the siblings are only recognizable by experimental, non-microscopic methods such as multi-locus sequencing and have no broader significance.

How should a species name be reported? For routine clinical practice, it has been claimed that identification at species complex level is usually sufficient (6, 9) although this may depend on the fungal group at hand or the clinical questions (59). Sequencing or MALDI-ToF MS (Matrix Assisted Laser Desorption/Ionization coupled to time-of-flight mass spectrometry) may directly identify the sibling name, without being aware that this is a member of a particular species complex. As MALDI-ToF MS results may be dependent on the spectral database used for comparison, it is recommended to mention this database with version number along with the identification in the lab report for retrospective analysis. Since nearly all major pathogenic species comprise subspecific entities, both the species complex and the sibling should be identified in clinical laboratory reports, with the siblings reported either by name or lineage number as applicable. Reporting would, therefore, appear as follows: *Trichophyton interdigitale* (member of *T. mentagrophytes* species complex), or, in case of an unnamed lineage: *Trichophyton mentagrophytes* species complex (molecular sibling ITS XIV). For the numerous species that are not part of a complex, the single recommended name is sufficient. Given the fact that the size of a genus lacks clear parameters, in practice for some fungi, several names may remain in use concomitantly until phylogenetic disputes can be resolved, as highlighted by the example of *Fusarium/Neocosmospora* (60–63).

At times, this situation may become very confusing. For example, *Mucor elegans* is synonymous with *Actinomucor elegans* and *Rhizopus elegans*, but *Apophysomyces elegans* is a completely different species. One solution to reduce naming confusion is to provide an easily usable and complete tool where current and prior names can be found and mapped to which species complex they belong. A list of currently recommended names for medical fungi is available open access at www.atlasclinicalfungi.org/, subheader nomenclature, also providing synonyms and affiliation to species complexes. The database has a comfortable search function as well as a printable version of the list recommended names. Use of the database prevents the necessity to mention old and new names in clinical reports. A committee of clinical microbiologists, physicians, and medical mycologists has been formed under the auspices of the International Society for Human and Animal Mycology (ISHAM), Working Groups on Nomenclature and Fungal Diagnostics, where names will be discussed, reviewed, and adjudicated for medical relevance, validity, and stability, consulting specialists on the topic. Recommendations will be publicized for transparency. We strongly recommend deposition of representative strains in the established fungal culture collections for future investigation.

The stability of names remains a much-debated issue, and there is no simple solution. Names will inevitably change. However, the changes should have some benefit. There are ways to mitigate some of the detrimental effects, but recommendations tend to have only temporary effect. As a recommendation to taxonomists, we suggest that rules in the *Code* (64, 65) could be used more than they are now to promote stability. For example, changing types of genera is possible, and names can be protected so that they cannot be replaced by older but later discovered synonyms. Careful consideration and restraint are required at the taxonomic research side, but the clinical user also has to accept the reality that frequently more than one name may persist in the literature for the same fungus for quite some time to come. As long as there is no consensus, some laboratories mention old and new names in their reports. Kidd et al. (12) suggested a 5-y transition period.

Nomenclatural databases, i.e., *Index Fungorum* (www.indexfungorum.org) and *Mycobank* (www.mycobank.org) apply latest names of new fungi and adjustments but refrain from providing recommendations of the usefulness of the changes. Therefore, a list of recommended names (Table 1; www.atlasclinicalfungi.org/nomenclature) has been proposed with the following considerations: (i) the table covers a list of medically

TABLE 1 Common and medically important fungi^a

Classical name most commonly used in clinical laboratories ^b	Alternative name appeared in literature (anamorph, teleomorph, synonym, synanamorph, or obsolete name)	Recommended name to be reported for clinical use ^c
<i>Acremonium egyptiacum</i>	<i>Acremonium sclerotigenum</i>	<i>Acremonium egyptiacum</i>
<i>Acremonium kiliense</i>	<i>Cephalosporium kiliense</i> , <i>Sarocladium kiliense</i>	<i>Sarocladium kiliense</i>
<i>Acremonium recifei</i>	<i>Xenoacremonium recifei</i>	<i>Acremonium recifei</i>
<i>Acremonium strictum</i>	<i>Cephalosporium acremonium</i> , <i>Sarocladium strictum</i>	<i>Sarocladium strictum</i>
<i>Acrophialophora fusispora</i>	<i>Paecilomyces fusisporus</i>	<i>Acrophialophora fusispora</i>
<i>Acrophialophora levis</i>		<i>Acrophialophora levis</i>
<i>Actinomyces elegans</i>	<i>Mucor elegans</i> , <i>Rhizopus elegans</i>	<i>Actinomyces elegans</i>
<i>Alternaria alternata</i>		<i>Alternaria alternata</i>
<i>Alternaria infectoria</i>		<i>Alternaria infectoria</i>
<i>Alternaria tenuissima</i>	<i>Helminthosporium tenuissimum</i>	<i>Alternaria tenuissima</i>
<i>Aphanoascus keratinophilus</i>	<i>Chryso sporium keratinophilum</i>	<i>Aphanoascus keratinophilus</i>
<i>Apophysomyces elegans</i>		<i>Apophysomyces elegans</i>
<i>Apophysomyces trapeziformis</i>		<i>Apophysomyces trapeziformis</i>
<i>Apophysomyces variabilis</i>		<i>Apophysomyces variabilis</i>
<i>Arthrimum arundinis</i>	<i>Apiospora arundinis</i>	<i>Arthrimum arundinis</i>
<i>Arthrographis kalrae</i>	<i>Oididendron kalrae</i>	<i>Arthrographis kalrae</i>
<i>Aspergillus calidoustus</i>		<i>Aspergillus calidoustus</i>
<i>Aspergillus flavus</i>	<i>Aspergillus oryzae</i>	<i>Aspergillus flavus</i>
<i>Aspergillus fischeri</i>	<i>Aspergillus fischerianus</i> , <i>Neosartorya fischeri</i>	<i>Aspergillus fischeri</i> [member of <i>A. fumigatus</i> series (species complex)]
<i>Aspergillus fumigatus</i>		<i>Aspergillus fumigatus</i>
<i>Aspergillus glaucus</i>	<i>Eurotium herbariorum</i>	<i>Aspergillus glaucus</i>
<i>Aspergillus lentulus</i>		<i>Aspergillus lentulus</i>
<i>Aspergillus nidulans</i>	<i>Emericella nidulans</i>	<i>Aspergillus nidulans</i>
<i>Aspergillus niger</i>		<i>Aspergillus niger</i>
<i>Aspergillus sclerotiorum</i>		<i>Aspergillus sclerotiorum</i>
<i>Aspergillus sydowii</i>		<i>Aspergillus sydowii</i>
<i>Aspergillus terreus</i>		<i>Aspergillus terreus</i>
<i>Aspergillus thermomutatus</i>	<i>Aspergillus fischeri</i> var. <i>thermomutatus</i> , <i>Neosartorya pseudofischeri</i>	<i>Aspergillus thermomutatus</i>
<i>Aspergillus tubingensis</i>		<i>Aspergillus tubingensis</i> [member of <i>A. niger</i> series (species complex)]
<i>Aspergillus udagawae</i>	<i>Neosartorya udagawae</i>	<i>Aspergillus udagawae</i>
<i>Aspergillus unguis</i>	<i>Emericella unguis</i>	<i>Aspergillus unguis</i>
<i>Aspergillus ustus</i>		<i>Aspergillus ustus</i>
<i>Aspergillus versicolor</i>		<i>Aspergillus versicolor</i>
<i>Aureobasidium melanogenum</i>	<i>Aureobasidium pullulans</i> var. <i>melanogenum</i>	<i>Aureobasidium melanogenum</i>
<i>Aureobasidium pullulans</i>		<i>Aureobasidium pullulans</i>
<i>Basidiobolus ranarum</i>		<i>Basidiobolus ranarum</i>
<i>Beauveria bassiana</i>		<i>Beauveria bassiana</i>
<i>Bipolaris australiensis</i>	<i>Curvularia australiensis</i>	<i>Bipolaris australiensis</i>
<i>Bipolaris hawaiiensis</i>	<i>Cochliobolus hawaiiensis</i> , <i>Curvularia hawaiiensis</i>	<i>Bipolaris hawaiiensis</i>
<i>Bipolaris spicifera</i>	<i>Drechslera spicifera</i> , <i>Cochliobolus spicifer</i> , <i>Curvularia spicifera</i>	<i>Bipolaris spicifera</i>
<i>Blastomyces dermatitidis</i>	<i>Ajellomyces dermatitidis</i>	<i>Blastomyces dermatitidis</i>
<i>Blastomyces gilchristii</i>		<i>Blastomyces gilchristii</i> (member of <i>B. dermatitidis</i> complex)
<i>Candida albicans</i>		<i>Candida albicans</i>
<i>Candida auris</i>		<i>Candida auris</i>
<i>Candida bracarensis</i>	<i>Nakaseomyces bracarensis</i>	<i>Candida bracarensis</i> or <i>Nakaseomyces bracarensis</i> (member of <i>C. glabrata</i> complex) ^d
<i>Candida dubliniensis</i>		<i>Candida dubliniensis</i>

(Continued on next page)

TABLE 1 Common and medically important fungi^a (Continued)

Classical name most commonly used in clinical laboratories ^b	Alternative name appeared in literature (anamorph, teleomorph, synonym, synanamorph, or obsolete name)	Recommended name to be reported for clinical use ^c
<i>Candida duobushaemulonis</i>	<i>Candida duobushaemulonii</i> , <i>Candida duobushaemuli</i>	<i>Candida duobushaemuli</i> (member of <i>C. haemuli</i> complex)
<i>Candida fabianii</i>	<i>Hansenula fabianii</i> , <i>Pichia fabianii</i> , <i>Lindnera fabianii</i> , <i>Cyberlindnera fabianii</i>	<i>Candida fabianii</i> or <i>Cyberlindnera fabianii</i> ^d
<i>Candida famata</i>	<i>Debaryomyces hansenii</i>	<i>Candida famata</i> or <i>Debaryomyces hansenii</i> ^d
<i>Candida fermentati</i>	<i>Torula fermentati</i> , <i>Pichia caribbica</i> , <i>Meyerozyma caribbica</i>	<i>Candida fermentati</i> (member of <i>C. guilliermondii</i> complex)
<i>Candida glabrata</i>	<i>Torulopsis glabrata</i> , <i>Nakaseomyces glabratus</i>	<i>Candida glabrata</i> or <i>Nakaseomyces glabratus</i> ^d
<i>Candida guilliermondii</i>	<i>Pichia guilliermondii</i> , <i>Meyerozyma guilliermondii</i>	<i>Candida guilliermondii</i> or <i>Meyerozyma guilliermondii</i> ^d
<i>Candida guilliermondii</i> var. <i>membranifaciens</i>	<i>Pichia ohmeri</i> , <i>Yamadazyma ohmeri</i> , <i>Kodamaea ohmeri</i>	<i>Kodamaea ohmeri</i> ^d
<i>Candida haemulonis</i>	<i>Torulopsis haemulonis</i> , <i>Candida haemulonii</i> , <i>Candida haemuli</i>	<i>Candida haemuli</i> (member of <i>C. haemuli</i> complex)
<i>Candida inconspicua</i>	<i>Torulopsis inconspicua</i> , <i>Pichia cactophila</i>	<i>Candida inconspicua</i>
<i>Candida kefyr</i>	<i>Candida pseudotropicalis</i> , <i>Kluyveromyces marxianus</i>	<i>Candida kefyr</i> or <i>Kluyveromyces marxianus</i> ^d
<i>Candida krusei</i>	<i>Issatchenkia orientalis</i> , <i>Pichia kudriavzevii</i>	<i>Candida krusei</i> or <i>Pichia kudriavzevii</i> ^d
<i>Candida lambica</i>	<i>Pichia fermentans</i>	<i>Candida lambica</i> or <i>Pichia fermentans</i> ^d
<i>Candida lipolytica</i>	<i>Yarrowia lipolytica</i>	<i>Candida lipolytica</i> or <i>Yarrowia lipolytica</i> ^d
<i>Candida lusitanae</i>	<i>Clavispora lusitanae</i>	<i>Candida lusitanae</i> or <i>Clavispora lusitanae</i> ^d
<i>Candida metapsilosis</i>		<i>Candida metapsilosis</i> (member of <i>C. parapsilosis</i> complex)
<i>Candida nivariensis</i>	<i>Nakaseomyces nivariensis</i>	<i>Candida nivariensis</i> or <i>Nakaseomyces nivariensis</i> (member of <i>C. glabrata</i> complex) ^d
<i>Candida norvegensis</i>	<i>Pichia norvegensis</i>	<i>Candida norvegensis</i> or <i>Pichia norvegensis</i> ^d
<i>Candida orthopsilosis</i>		<i>Candida orthopsilosis</i> (member of <i>C. parapsilosis</i> complex)
<i>Candida parapsilosis</i>		<i>Candida parapsilosis</i>
<i>Candida pelliculosa</i>	<i>Hansenula anomala</i> , <i>Pichia anomala</i> , <i>Wickerhamomyces anomalus</i>	<i>Candida pelliculosa</i> or <i>Hansenula anomala</i> ^d
<i>Candida rugosa</i>	<i>Diutina rugosa</i>	<i>Candida rugosa</i> or <i>Diutina rugosa</i> ^d
<i>Candida tropicalis</i>		<i>Candida tropicalis</i>
<i>Candida utilis</i>	<i>Torulopsis utilis</i> , <i>Pichia jadinii</i> , <i>Lindnera jadinii</i> , <i>Cyberlindnera jadinii</i>	<i>Candida utilis</i> or <i>Cyberlindnera jadinii</i> ^d
<i>Chaetomium atrobrunneum</i>	<i>Amesia atrobrunnea</i>	<i>Chaetomium atrobrunneum</i>
<i>Chaetomium brasiliense</i>	<i>Ovatospora brasiliensis</i>	<i>Chaetomium brasiliense</i>
<i>Chaetomium globosum</i>		<i>Chaetomium globosum</i>
<i>Chaetomium perlucidum</i>	<i>Parachaetomium perlucidum</i>	<i>Chaetomium perlucidum</i>
<i>Chaetomium strumarium</i>	<i>Achaetomium strumarium</i>	<i>Chaetomium strumarium</i>
<i>Chrysonilia sitophila</i>	<i>Monila sitophila</i> , <i>Neurospora sitophila</i>	<i>Neurospora sitophila</i>
<i>Chrysosporium queenslandicum</i>	<i>Uncinocarpus queenslandicus</i> , <i>Brunneospora queenslandica</i>	<i>Brunneospora queenslandica</i>
<i>Chrysosporium zonatum</i>	<i>Uncinocarpus orissae</i>	<i>Chrysosporium zonatum</i>
<i>Cladophialophora bantiana</i>	<i>Xylohypha bantiana</i> , <i>Cladosporium bantianum</i> , <i>Cladosporium trichoides</i>	<i>Cladophialophora bantiana</i>
<i>Cladophialophora boppi</i>	<i>Taeniolella boppii</i>	<i>Cladophialophora boppii</i>
<i>Cladophialophora carrionii</i>	<i>Cladosporium carrionii</i>	<i>Cladophialophora carrionii</i>
<i>Cladosporium cladosporioides</i>		<i>Cladosporium cladosporioides</i>
<i>Cladosporium herbarum</i>		<i>Cladosporium herbarum</i>
<i>Cladosporium sphaerospermum</i>		<i>Cladosporium sphaerospermum</i>
<i>Coccidioides immitis</i>		<i>Coccidioides immitis</i>
<i>Coccidioides posadasii</i>		<i>Coccidioides posadasii</i>
<i>Cokeromyces recurvatus</i>		<i>Cokeromyces recurvatus</i>
<i>Collectotrichum coccodes</i>		<i>Collectotrichum coccodes</i>

(Continued on next page)

TABLE 1 Common and medically important fungi^a (Continued)

Classical name most commonly used in clinical laboratories ^b	Alternative name appeared in literature (anamorph, teleomorph, synonym, synanamorph, or obsolete name)	Recommended name to be reported for clinical use ^c
<i>Colletotrichum gloeosporioides</i>		<i>Colletotrichum gloeosporioides</i>
<i>Conidiobolus coronatus</i>		<i>Conidiobolus coronatus</i>
<i>Cryptococcus adeliensis</i>	<i>Naganishia adeliensis</i>	<i>Naganishia adeliensis</i>
<i>Cryptococcus albidus</i>	<i>Cryptococcus albidus</i> var. <i>albidus</i> , <i>C. albidus</i> var. <i>diffluens</i> , <i>Cryptococcus diffluens</i> , <i>Cryptococcus genitalis</i> , <i>Naganishia albida</i>	<i>Naganishia albida</i>
<i>Cryptococcus gattii</i>		<i>Cryptococcus gattii</i>
<i>Cryptococcus laurentii</i>	<i>Torula laurentii</i> , <i>Papiliotrema laurentii</i>	<i>Papiliotrema laurentii</i>
<i>Cryptococcus liquefaciens</i>	<i>Torulopsis liquefaciens</i> , <i>Naganishia liquefaciens</i>	<i>Naganishia liquefaciens</i>
<i>Cryptococcus neoformans</i>		<i>Cryptococcus neoformans</i>
<i>Cryptococcus uniguttulatus</i>	<i>Filobasidium uniguttulatum</i>	<i>Filobasidium uniguttulatum</i>
<i>Cunninghamella bertholletiae</i>		<i>Cunninghamella bertholletiae</i>
<i>Curvularia geniculata</i>	<i>Cochliobolus geniculatus</i>	<i>Curvularia geniculata</i>
<i>Curvularia lunata</i>	<i>Cochliobolus lunatus</i>	<i>Curvularia lunata</i>
<i>Curvularia pallescens</i>	<i>Cochliobolus pallescens</i>	<i>Curvularia pallescens</i>
<i>Cylindrocarpon cyanescens</i>	<i>Fusarium cyanescens</i> , <i>Neocosmospora cyanescens</i>	<i>Cylindrocarpon cyanescens</i>
<i>Cylindrocarpon destructans</i>	<i>Ilyonectria destructans</i>	<i>Cylindrocarpon destructans</i>
<i>Drechslera biseptata</i>	<i>Pyrenophora biseptata</i>	<i>Drechslera biseptata</i>
<i>Emmonsia crescens</i>	<i>Emmonsia parva</i> var. <i>crescens</i> , <i>Ajellomyces crescens</i> , <i>Emergomyces crescens</i>	<i>Emmonsia crescens</i>
<i>Emmonsia parva</i>	<i>Chrysosporium parvum</i> , <i>Blastomyces parvus</i>	<i>Blastomyces parvus</i>
<i>Emmonsia pasteuriana</i>	<i>Emergomyces pasteurianus</i>	<i>Emergomyces pasteurianus</i>
<i>Engyodontium album</i>	<i>Tritirachium album</i> , <i>Beauveria alba</i> , <i>Parengyodontium album</i>	<i>Parengyodontium album</i>
<i>Epicoccum nigrum</i>	<i>Epicoccum purpurascens</i> , <i>Phoma epicoccina</i>	<i>Epicoccum nigrum</i>
<i>Epidermophyton floccosum</i>	<i>Acrothecium floccosum</i> , <i>Trichothecium floccosum</i>	<i>Epidermophyton floccosum</i>
<i>Exophiala dermatitidis</i>	<i>Wangiella dermatitidis</i>	<i>Exophiala dermatitidis</i>
<i>Exophiala jeanselmei</i>		<i>Exophiala jeanselmei</i>
<i>Exophiala lecanii-corni</i>	<i>Exophiala jeanselmei</i> var. <i>lecanii-corni</i>	<i>Exophiala lecanii-corni</i>
<i>Exophiala oligosperma</i>		<i>Exophiala oligosperma</i>
<i>Exophiala spinifera</i>	<i>Phialophora spinifera</i> , <i>Rhinochloidiella spinifera</i>	<i>Exophiala spinifera</i>
<i>Exserohilum rostratum</i>	<i>Helminthosporium rostratum</i> , <i>Exserohilum longirostratum</i> , <i>Exserohilum macginnisii</i>	<i>Exserohilum rostratum</i>
<i>Fonsecaea monophora</i>		<i>Fonsecaea monophora</i>
<i>Fonsecaea pedrosoi</i>	<i>Fonsecaea compacta</i>	<i>Fonsecaea pedrosoi</i>
<i>Fusarium chlamydosporum</i>		<i>Fusarium chlamydosporum</i>
<i>Fusarium dimerum</i>	<i>Bisifusarium dimerum</i>	<i>Fusarium dimerum</i>
<i>Fusarium falciforme</i>	<i>Acremonium falciforme</i> , <i>Neocosmospora falciforme</i>	<i>Fusarium falciforme</i> (member of <i>F. solani</i> complex)
<i>Fusarium incarnatum</i>	<i>Fusarium semitectum</i>	<i>Fusarium incarnatum</i> (member of <i>F. incarnatum-equi-seti</i> complex)
<i>Fusarium keratoplasticum</i>	<i>Neocosmospora keratoplastica</i>	<i>Fusarium keratoplasticum</i> (member of <i>F. solani</i> complex)
<i>Fusarium lichenicola</i>	<i>Cylindrocarpon lichenicola</i> , <i>Neocosmospora lichenicola</i>	<i>Fusarium lichenicola</i> (member of <i>F. solani</i> complex)
<i>Fusarium neocosmosporiellum</i>	<i>Neocosmospora vasinfecta</i>	<i>Fusarium neocosmosporiellum</i> (member of <i>F. solani</i> complex)
<i>Fusarium oxysporum</i>		<i>Fusarium oxysporum</i>
<i>Fusarium petrophilum</i>	<i>Neocosmospora petroliphila</i>	<i>Fusarium petrophilum</i> (member of <i>F. solani</i> complex)
<i>Fusarium proliferatum</i>		<i>Fusarium proliferatum</i> (member of <i>F. fujikuroi</i> complex)
<i>Fusarium solani</i>	<i>Neocosmospora solani</i>	<i>Fusarium solani</i>
<i>Fusarium verticillioides</i>	<i>Fusarium moniliforme</i> , <i>Gibberella moniliformis</i>	<i>Fusarium verticillioides</i> (member of <i>F. fujikuroi</i> complex)

(Continued on next page)

TABLE 1 Common and medically important fungi^a (Continued)

Classical name most commonly used in clinical laboratories ^b	Alternative name appeared in literature (anamorph, teleomorph, synonym, synanamorph, or obsolete name)	Recommended name to be reported for clinical use ^c
<i>Geomyces destructans</i>	<i>Pseudogymnoascus destructans</i>	<i>Pseudogymnoascus destructans</i>
<i>Geomyces pannorum</i>	<i>Chrysosporium pannorum</i> , <i>Pseudogymnoascus pannorum</i>	<i>Geomyces pannorum</i>
<i>Geotrichum candidum</i>	<i>Galactomyces candidus</i> , <i>Dipodascus geotrichum</i>	<i>Geotrichum candidum</i>
<i>Geotrichum capitatum</i>	<i>Dipodascus capitatus</i> , <i>Blastoschizomyces capitatus</i> , <i>Saprochaetea capitata</i> , <i>Magnusiomyces capitatus</i>	<i>Magnusiomyces capitatus</i>
<i>Geotrichum clavatum</i>	<i>Saprochaete clavate</i> , <i>Magnusiomyces clavatus</i>	<i>Magnusiomyces clavatus</i>
<i>Gymnascella hyalinospora</i>	<i>Narasimhella hyalinospora</i> , <i>Gymnoascus hyalinosporus</i>	<i>Gymnascella hyalinospora</i>
<i>Histoplasma capsulatum</i>	<i>Ajellomyces capsulatus</i>	<i>Histoplasma capsulatum</i>
<i>Hormographiella aspergillata</i>	<i>Coprinus cinereus</i> , <i>Coprinopsis cinerea</i>	<i>Coprinopsis cinerea</i>
<i>Hormographiella verticillata</i>	<i>Coprinus domesticus</i> , <i>Coprinellus domesticus</i>	<i>Coprinellus domesticus</i>
<i>Hortaea werneckii</i>	<i>Exophiala werneckii</i> , <i>Phaeoannellomyces werneckii</i>	<i>Hortaea werneckii</i>
<i>Lasiodiplodia theobromae</i>	<i>Botryosphaeria rhodina</i>	<i>Lasiodiplodia theobromae</i>
<i>Lecythophora hoffmannii</i>	<i>Coniochaeta hoffmannii</i>	<i>Lecythophora hoffmannii</i>
<i>Lecythophora mutabilis</i>	<i>Coniochaeta mutabilis</i>	<i>Lecythophora mutabilis</i>
<i>Lichtheimia corymbifera</i>	<i>Absidia corymbifera</i>	<i>Lichtheimia corymbifera</i>
<i>Lodderomyces elongisporus</i>	<i>Saccharomyces elongisporus</i>	<i>Lodderomyces elongisporus</i> ^d
<i>Lomentospora prolificans</i>	<i>Scedosporium prolificans</i> , <i>Scedosporium inflatum</i>	<i>Lomentospora prolificans</i>
<i>Madurella grisea</i>	<i>Trematosphaeria grisea</i>	<i>Trematosphaeria grisea</i>
<i>Madurella mycetomatis</i>	<i>Madurella mycetomi</i>	<i>Madurella mycetomatis</i>
<i>Malassezia furfur</i>		<i>Malassezia furfur</i>
<i>Malassezia globosa</i>		<i>Malassezia globosa</i>
<i>Malassezia pachydermatis</i>		<i>Malassezia pachydermatis</i>
<i>Malassezia restricta</i>		<i>Malassezia restricta</i>
<i>Malassezia slooffiae</i>		<i>Malassezia slooffiae</i>
<i>Malassezia sympodialis</i>		<i>Malassezia sympodialis</i>
<i>Malbranchea pulchella</i>		<i>Malbranchea pulchella</i>
<i>Microascus cinereus</i>	<i>Scopulariopsis cinereus</i>	<i>Microascus cinereus</i>
<i>Microascus cirrosus</i>		<i>Microascus cirrosus</i>
<i>Microsporium audouinii</i>		<i>Microsporium audouinii</i>
<i>Microsporium canis</i>	<i>Microsporium canis</i> var. <i>distortum</i> , <i>Arthroderma otae</i> , <i>Microsporium equinum</i>	<i>Microsporium canis</i>
<i>Microsporium cookei</i>	<i>Paraphyton cookei</i>	<i>Paraphyton cookei</i>
<i>Microsporium ferrugineum</i>		<i>Microsporium ferrugineum</i>
<i>Microsporium gallinae</i>	<i>Lophophyton gallinae</i>	<i>Lophophyton gallinae</i>
<i>Microsporium gypseum</i>	<i>Nannizzia gypsea</i>	<i>Nannizzia gypsea</i>
<i>Microsporium nanum</i>	<i>Arthroderma obtusum</i> , <i>Nannizzia nana</i>	<i>Nannizzia nana</i>
<i>Microsporium persicolor</i>	<i>Arthroderma persicolor</i> , <i>Nannizzia persicolor</i>	<i>Nannizzia persicolor</i>
<i>Mortierella wolfii</i>	<i>Actinomortierella wolfii</i>	<i>Mortierella wolfii</i>
<i>Mucor circinelloides</i>		<i>Mucor circinelloides</i>
<i>Mucor indicus</i>		<i>Mucor indicus</i>
<i>Mucor irregularis</i>	<i>Rhizomucor variabilis</i>	<i>Mucor irregularis</i>
<i>Mucor ramosissimus</i>		<i>Mucor ramosissimus</i> (member of <i>M. circinelloides</i> complex)
<i>Mucor velutinosus</i>		<i>Mucor velutinosus</i> (member of <i>M. circinelloides</i> complex)
<i>Myceliophthora thermophila</i>	<i>Chrysosporium thermophilum</i> , <i>Thermothelomyces thermophila</i>	<i>Myceliophthora thermophila</i>
<i>Nigrospora sphaerica</i>		<i>Nigrospora sphaerica</i>
<i>Nodulisporium griseobrunneum</i>	<i>Hypoxyton griseobrunneum</i>	<i>Hypoxyton griseobrunneum</i>
<i>Ochroconis gallopava</i>	<i>Dactylaria gallopava</i> , <i>Dactylaria constricta</i> var. <i>gallopava</i> , <i>Verruconis gallopava</i>	<i>Verruconis gallopava</i>

(Continued on next page)

TABLE 1 Common and medically important fungi^a (Continued)

Classical name most commonly used in clinical laboratories ^b	Alternative name appeared in literature (anamorph, teleomorph, synonym, synanamorph, or obsolete name)	Recommended name to be reported for clinical use ^c
<i>Ochroconis mirabilis</i>	<i>Ochroconis musae</i> , <i>Scolecobasidium musae</i> , <i>Scolecobasidium mirabilis</i>	<i>Scolecobasidium mirabilis</i>
<i>Onychocola canadensis</i>	<i>Arachnomyces nodosetosus</i>	<i>Arachnomyces nodosetosus</i>
<i>Paecilomyces formosus</i>		<i>Paecilomyces formosus</i>
<i>Paecilomyces lilacinus</i>	<i>Purpureocillium lilacinum</i>	<i>Purpureocillium lilacinum</i>
<i>Paecilomyces variotii</i>	<i>Paecilomyces spectabilis</i> , <i>Byssoschlamys spectabilis</i>	<i>Paecilomyces variotii</i>
<i>Paracoccidioides brasiliensis</i>		<i>Paracoccidioides brasiliensis</i>
<i>Paracoccidioides lutzii</i>		<i>Paracoccidioides lutzii</i> (member of <i>P. brasiliensis</i> complex)
<i>Penicillium chrysogenum</i>		<i>Penicillium chrysogenum</i>
<i>Penicillium citrinum</i>		<i>Penicillium citrinum</i>
<i>Penicillium marneffei</i>	<i>Talaromyces marneffei</i>	<i>Talaromyces marneffei</i>
<i>Penicillium purpureogenum</i>	<i>Talaromyces purpureogenus</i>	<i>Talaromyces purpureogenus</i>
<i>Phaeoacremonium parasiticum</i>	<i>Phialophora parasitica</i> , <i>Togninia parasitica</i>	<i>Phaeoacremonium parasiticum</i>
<i>Phialemonium curvatum</i>	<i>Phialemonium dimorphosporum</i> , <i>Thyridium curvatum</i> , <i>Phialemoniopsis curvata</i>	<i>Thyridium curvatum</i>
<i>Phialemonium atrogriseum</i>	<i>Acremonium atrogriseum</i>	<i>Phialemonium atrogriseum</i>
<i>Phialemonium obovatum</i>		<i>Phialemonium obovatum</i>
<i>Phialophora americana</i>	<i>Capronia semiimmersa</i> , <i>Cadophora americana</i>	<i>Phialophora americana</i>
<i>Phialophora europaea</i>	<i>Cyphellophora europaea</i>	<i>Cyphellophora europaea</i>
<i>Phialophora richardsiae</i>	<i>Pleurostomophora richardsiae</i> , <i>Pleurostoma richardsiae</i>	<i>Pleurostoma richardsiae</i>
<i>Phialophora verrucosa</i>		<i>Phialophora verrucosa</i>
<i>Phoma cruris-hominis</i>		<i>Phoma cruris-hominis</i>
<i>Phoma herbarum</i>	<i>Phoma muscivora</i>	<i>Phoma herbarum</i>
<i>Piedraia hortae</i>		<i>Piedraia hortae</i>
<i>Pithomyces chartarum</i>	<i>Pseudopithomyces chartarum</i>	<i>Pithomyces chartarum</i>
<i>Prototheca wickerhamii</i>		<i>Prototheca wickerhamii</i> ^d
<i>Pseudozyma aphidis</i>	<i>Moesziomyces aphidis</i>	<i>Moesziomyces aphidis</i>
<i>Pyrenochaeta romeroi</i>	<i>Medicopsis romeroi</i>	<i>Medicopsis romeroi</i>
<i>Pythium insidiosum</i>		<i>Pythium insidiosum</i> ^e
<i>Ramichloridium schulzeri</i>	<i>Myrmecridium schulzeri</i>	<i>Myrmecridium schulzeri</i>
<i>Rasamsonia aegroticola</i>		<i>Rasamsonia aegroticola</i>
<i>Rasamsonia argillacea</i>	<i>Penicillium argillaceum</i> , <i>Geosmithia argillacea</i>	<i>Rasamsonia argillacea</i> (member of <i>R. argillacea</i> complex)
<i>Rhinocladiella aquaspersa</i>		<i>Rhinocladiella aquaspersa</i>
<i>Rhinocladiella mackenziei</i>	<i>Ramichloridium mackenziei</i>	<i>Rhinocladiella mackenziei</i>
<i>Rhinocladiella similis</i>		<i>Rhinocladiella similis</i>
<i>Rhizomucor miehei</i>		<i>Rhizomucor miehei</i>
<i>Rhizomucor pusillus</i>		<i>Rhizomucor pusillus</i>
<i>Rhizopus arrhizus</i>	<i>Rhizopus oryzae</i>	<i>Rhizopus arrhizus</i>
<i>Rhizopus azygosporus</i>		<i>Rhizopus azygosporus</i>
<i>Rhizopus microsporus</i>	<i>Rhizopus rhizopodiformis</i>	<i>Rhizopus microsporus</i>
<i>Rhizopus schipperae</i>		<i>Rhizopus schipperae</i>
<i>Rhizopus stolonifer</i>		<i>Rhizopus stolonifer</i>
<i>Rhodotorula glutinis</i>	<i>Torulopsis glutinis</i>	<i>Rhodotorula glutinis</i>
<i>Rhodotorula minuta</i>	<i>Torula minuta</i> , <i>Cystobasidium minutum</i>	<i>Rhodotorula minuta</i>
<i>Rhodotorula mucilaginosa</i>	<i>Rhodotorula rubra</i> , <i>Torula mucilaginosa</i>	<i>Rhodotorula mucilaginosa</i>
<i>Saccharomyces cerevisiae</i>	<i>Saccharomyces boulardii</i>	<i>Saccharomyces cerevisiae</i>
<i>Saksenaia vasiformis</i>		<i>Saksenaia vasiformis</i>
<i>Scedosporium apiospermum</i>	<i>Pseudallescheria apiosperma</i>	<i>Scedosporium apiospermum</i> (member of <i>S. apiospermum</i> complex)

(Continued on next page)

TABLE 1 Common and medically important fungi^a (Continued)

Classical name most commonly used in clinical laboratories ^b	Alternative name appeared in literature (anamorph, teleomorph, synonym, synanamorph, or obsolete name)	Recommended name to be reported for clinical use ^c
<i>Scedosporium aurantiacum</i>		<i>Scedosporium aurantiacum</i>
<i>Scedosporium boydii</i>	<i>Pseudallescheria boydii</i>	<i>Scedosporium boydii</i> (member of <i>S. apiospermum</i> complex)
<i>Schizophyllum commune</i>		<i>Schizophyllum commune</i>
<i>Schizophyllum radiatum</i>		<i>Schizophyllum radiatum</i>
<i>Scopulariopsis brevicaulis</i>	<i>Microascus brevicaulis</i>	<i>Scopulariopsis brevicaulis</i>
<i>Scopulariopsis brumptii</i>	<i>Microascus paisii</i>	<i>Scopulariopsis brumptii</i>
<i>Scytalidium dimidiatum</i>	<i>Hendersonula toruloidea</i> , <i>Nattrassia mangiferae</i> , <i>Neoscytalidium dimidiatum</i>	<i>Neoscytalidium dimidiatum</i>
<i>Sporobolomyces salmonicolor</i>	<i>Sporidiobolus salmonicolor</i>	<i>Sporobolomyces salmonicolor</i>
<i>Sporothrix brasiliensis</i>		<i>Sporothrix brasiliensis</i>
<i>Sporothrix cyanescens</i>	<i>Fugomyces cyanescens</i> , <i>Cerinosterus cyanescens</i> , <i>Quambalaria cyanescens</i>	<i>Quambalaria cyanescens</i>
<i>Sporothrix globosa</i>		<i>Sporothrix globosa</i>
<i>Sporothrix luriei</i>	<i>Sporothrix schenckii</i> var. <i>luriei</i>	<i>Sporothrix luriei</i>
<i>Sporothrix mexicana</i>		<i>Sporothrix mexicana</i>
<i>Sporothrix schenckii</i>		<i>Sporothrix schenckii</i>
<i>Sporotrichum pruinosum</i>	<i>Chrysosporium pruinosum</i> , <i>Phanerochaete chrysosporium</i>	<i>Phanerochaete chrysosporium</i>
<i>Syncephalastrum racemosum</i>		<i>Syncephalastrum racemosum</i>
<i>Trichoderma harzianum</i>		<i>Trichoderma harzianum</i>
<i>Trichoderma longibrachiatum</i>		<i>Trichoderma longibrachiatum</i>
<i>Trichophyton benhamiae</i>	<i>Arthroderma benhamiae</i>	<i>Trichophyton benhamiae</i>
<i>Trichophyton concentricum</i>		<i>Trichophyton concentricum</i> (member of <i>T. benhamiae</i> complex)
<i>Trichophyton equinum</i>		<i>Trichophyton equinum</i> (member of <i>T. tonsurans</i> complex)
<i>Trichophyton indotineae</i>	<i>Trichophyton mentagrophytes</i> ITS Type VIII	<i>Trichophyton indotineae</i>
<i>Trichophyton interdigitale</i>	<i>Trichophyton krajenii</i>	<i>Trichophyton interdigitale</i> (member of <i>T. mentagrophytes</i> complex)
<i>Trichophyton mentagrophytes</i>		<i>Trichophyton mentagrophytes</i>
<i>Trichophyton rubrum</i>		<i>Trichophyton rubrum</i>
<i>Trichophyton schoenleinii</i>		<i>Trichophyton schoenleinii</i>
<i>Trichophyton soudanense</i>		<i>Trichophyton soudanense</i> (member of <i>T. rubrum</i> complex)
<i>Trichophyton tonsurans</i>		<i>Trichophyton tonsurans</i>
<i>Trichophyton verrucosum</i>		<i>Trichophyton verrucosum</i>
<i>Trichophyton violaceum</i>	<i>Trichophyton yaoundei</i>	<i>Trichophyton violaceum</i> (member of <i>T. rubrum</i> complex)
<i>Trichosporon asahii</i>		<i>Trichosporon asahii</i>
<i>Trichosporon dermatis</i>	<i>Cutaneotrichosporon dermatis</i>	<i>Trichosporon dermatis</i> or <i>Cutaneotrichosporon dermatis</i> ^d
<i>Trichosporon inkin</i>		<i>Trichosporon inkin</i>
<i>Trichosporon loubieri</i>	<i>Apiotrichum loubieri</i>	<i>Apiotrichum oubieri</i>
<i>Trichosporon mucoides</i>	<i>Cutaneotrichosporon mucoides</i>	<i>Trichosporon mucoides</i> or <i>Cutaneotrichosporon mucoides</i> ^d
<i>Trichosporon mycotoxinivorans</i>	<i>Apiotrichum mycotoxinivorans</i>	<i>Apiotrichum mycotoxinivorans</i>
<i>Tritirachium oryzae</i>	<i>Beauveria oryzae</i> , <i>Tritirachium roseum</i>	<i>Tritirachium oryzae</i>
<i>Ulocladium botrytis</i>	<i>Alternaria botrytis</i>	<i>Alternaria botrytis</i>
<i>Ulocladium chartarum</i>	<i>Alternaria chartarum</i>	<i>Alternaria chartarum</i>
<i>Veronaea botryosa</i>		<i>Veronaea botryosa</i>

(Continued on next page)

TABLE 1 Common and medically important fungi^a (Continued)

Classical name most commonly used in clinical laboratories ^b	Alternative name appeared in literature (anamorph, teleomorph, synonym, synanamorph, or obsolete name)	Recommended name to be reported for clinical use ^c
<i>Verticillium dahliae</i>		<i>Verticillium dahliae</i>

^aWe recommend Table 1 for the following reasons: (i) the table covers a list of medically important fungi that are considered as most commonly encountered in the clinical labs and listed in seven Medical Mycology Textbooks widely used in the clinical laboratories; (ii) the current recommendation on how to report these fungi in the table is based on the rationale illustrated in the manuscript and consensus agreement among authors after an extensive review; (3) as clinical laboratories are struggling with how to report these fungi to clinicians due to nomenclature variation in the literature, the table may serve as a current reference to guide clinical laboratories on how to report these fungi; (4) the names listed in the table are not fixed names but rather representing nomenclature stability; they will be reviewed and updated periodically by an international committee representative of clinical microbiologists, physicians, medical mycologists, and taxonomists. Note: a complete overview of pathogenic and opportunistic species with descriptions and references can be found in Atlas of Clinical Fungi, 4th ed. 2020.

^bBased on a list of textbooks and reference materials used in clinical labs: (i) Larone's Medically Important Fungi 6th Ed by Thomas J Walsh, Randall T. Hayden, Davise H. Larone; (ii) Manual of Clinical Microbiology 12th Ed; (iii) CAP (College of American Pathologists) Master list of Fungi; (iv) CAP Color Atlas of Mycology by Gordon L. Love, Julie A. Ribes; (v) Guide to Clinically Significant Fungi by Deanna A. Sutton, Annette W. Fothergill, Michael G. Rinaldi; (vi) Doctor Fungus online reference (<https://drfungus.org/knowledge-base-category/fungi-descriptions/>); (vii) Identifying fungi: a clinical laboratory handbook 2nd Ed by Guy St-Germain, Richard Summerbell; (viii) Identification of pathogenic fungi 2nd Ed by Colin K. Campbell, Elizabeth M. Johnson, David W. Warnock; (ix) Descriptions of medical fungi 4th Ed by Sarah Kidd, Catriona Halliday, David Ellis. Furthermore, only the ones that have been reported causing human infection are included.

^cThe recommended names to be reported for clinical use are based on current treatment guidelines and the Atlas of Clinical Fungi (<https://www.clinicalfungi.org>). The fungi listed in the table represent a list of the common and medically important ones from the database of the ATLAS (>6,000 clinical fungi). Species level identification on some of them may not be readily achievable based on morphological or phenotypic features and thus will have to rely on additional tools, e.g., DNA sequencing identification or MALDI-TOF MS to obtain reliable species identification.

^dThese names will be further reviewed by an international committee representative of clinical microbiologists, physicians, medical mycologists, and taxonomists.

^eNot a fungus.

important fungi that are considered most commonly encountered in clinical laboratories based on their inclusion in seven textbooks of medical mycology and two reference resources (CAP master list and Doctor Fungus online reference); (ii) the current recommendations on how to report these fungi in the table is based on the rationale illustrated in the present manuscript and consensus agreement among authors after an extensive consultation; (iii) as clinical laboratories are struggling with how to report these fungi to patient-care providers due to nomenclature variation in the literature, the table may serve as a current reference to guide clinical laboratories on how to report these fungi; (iv) the names listed in the table are not fixed names but rather represent nomenclature stability. We recommend that this list, including our author's consensus reporting recommendations, could be used to guide initial efforts by an upcoming international committee representative of clinical microbiologists, physicians, medical mycologists, and taxonomists. This committee will function as a governing body to curate future fungal name changes for clinical use. We also recommend special focus on reporting some names (e.g., some *Candida* species) of which the present authors were unable to reach consensus using the methods discussed in this document.

Incorporating name changes that are validated and stable is essential for clinical laboratories and patient care. Other well-established groups which review relevant scientific advances for application in clinical laboratories include the Clinical and Laboratory Standards Institute (CLSI) and the European Committee on Antimicrobial Susceptibility Testing (EUCAST). These organizations provide additional levels of standardization and validation of diagnostic changes before introduction into clinical laboratories. Clinical practice will benefit from clear guidance from the international community on the adaption to nomenclatural changes of medically important fungi. Future association with the database of SNOMED CT and LONIC codes might be considered, thereby facilitating international digital data exchange and standardization of patient reports.

We recognize it is important to disseminate nomenclature changes to clinical community timely and accurately. Education required not only for the clinician but also for the taxonomist that has an insufficient eye for the user. Published lists are an alternative but are forgotten within a few years. We, therefore, think that the most parsimonious and sustainable solution is a readily accessible database, which is easy to find and easy to handle. Only a single website should be remembered. The contents of the database will be supervised by a committee of ISHAM (International Society for Human and Animal Mycology), composed of researchers, as well as of users, and decisions are made in consultancy with taxonomic specialists. Furthermore, the

committee will partner with the societies that support this document and use their proper media channels to deliver any recommended nomenclature changes to the clinicians and clinical laboratories widely without any delay.

ACKNOWLEDGMENTS

We thank the following people for their contributions of reviewing this document. Members of the Laboratory Practice Subcommittee of the American Society of Microbiology (ASM): Audrey N. Schuetz (Mayo Clinic, Rochester, MN, U.S.A.) and Laura Filkins (University of Texas Southwestern Medical Center, TX, U.S.A.). Members of EUCAST-AFST Subcommittee: Sevtap Arikan-Akdagli (Turkey), Lena Rós Ásmundsdóttir (Iceland), Francesco Barchiesi (Italy), Jørgen Vildershøj Bjørnholt (Norway), Jochem Buil (The Netherlands), Mariana Castanheira (U.S.A., USCAST representative), Erja Chryssanthou (Sweden), Eric Dannaoui (France), Nathalie Friberg (Finland), Christian Giske (Sweden), Jesus Guinea (Spain), Oliver Kurzai (Germany), Cornelia Lass-Floerl (Austria), Katrien Lagrou (Belgium), Pavlina Lyskova (Czech Republic), Mihai Mares (Romania), Tadeja Matos (Slovenia), Joseph Meletiadis (Greece), Caroline Moore (U.K.), Konrad Muehlethaler (Switzerland), Georgia Vriani (Greece), and Johannes Wagener (Ireland). Members of bioMerieux: Nancy Moss, Valerie Collin. Members of Bruker Scientific LLC: Markus Kostrzewa, Gongyi Shi. Yu Quan is acknowledged for editorial assistance.

AUTHOR AFFILIATIONS

¹Radboudumc-CWZ Centre of Expertise for Mycology, Nijmegen, the Netherlands

²Foundation Atlas of Clinical Fungi, Hilversum, the Netherlands

³Department of Dermatology and Venereology, Peking University First Hospital, Beijing, China

⁴Department of Basic Pathology, Federal University of Paraná, Curitiba, Brazil

⁵Research Center for Medical Mycology, Peking University, Beijing, China

⁶International Society for Human and Animal Mycology (ISHAM), Working Group Nomenclature

⁷Center for Innovative Therapeutics and Diagnostics, Richmond, Virginia, USA

⁸University of Maryland School of Medicine, Baltimore, Maryland, USA

⁹Nomenclature Committee for Fungi, International Mycological Association (IMA)

¹⁰Fungal Diagnostics Laboratory Consortium (FDLC)

¹¹Mycoses Study Group, Education and Research Consortium (MSG-ERC)

¹²European Confederation of Medical Mycology (ECMM)

¹³Clinical and Laboratory Standards Institute (CLSI)

¹⁴Medical Mycological Society of the Americas (MMSA)

¹⁵ISHAM Working Group on Diagnostics

¹⁶Mycology Reference Laboratory, Spanish National Centre for Microbiology, Madrid, Spain

¹⁷Fungal Infection Study Group, European Society of Clinical Microbiology and Infectious Diseases (EFISG/ESCMID), Basel, Switzerland

¹⁸Departments of Medicine and Pathology, Duke University, Durham, North Carolina, USA

¹⁹Department of Microbiology and Infection Control, Statens Serum Institut, Copenhagen, Denmark; Department of Clinical Microbiology, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark

²⁰Antifungal Susceptibility Testing Subcommittee of European Committee of Antimicrobial Susceptibility Testing (EUCAST-AFST)

²¹Clinical Microbiology Service, Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, USA

²²Mycology Committee of Chinese Society for Microbiology

²³Institute of Microbiology, State Key Laboratory of Mycology, Chinese Academy of Sciences, Beijing, China

²⁴Medical Mycology Society of Chinese Medicine and Education Association

- ²⁵Asia Pacific Society for Medical Mycology
- ²⁶ISHAM Working Group Veterinary Mycology and One Health
- ²⁷Mycological Society of China (MSC)
- ²⁸Clinical Microbiology at The Ohio State University Wexner Medical Center, Columbus, Ohio, USA
- ²⁹National Mycology Reference Laboratory, Public Health England, Bristol, United Kingdom
- ³⁰Department of Microbiology, National Reference Laboratory for Antimicrobial Resistance in Fungal Pathogens, Medical Mycology Unit, Vallabhbhai Patel Chest Institute, University of Delhi, Delhi, India
- ³¹University of Texas Southwestern Medical Center, Dallas, Texas, USA
- ³²Division of Infectious Diseases, Mount Auburn Hospital, Cambridge, Massachusetts, USA
- ³³Infectious Diseases Society of America (ISDA)
- ³⁴University of Cologne, Faculty of Medicine, Institute of Translational Research, Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases, Cologne, Germany
- ³⁵Department I of Internal Medicine, University of Cologne, Excellence Center for Medical Mycology, Cologne, Germany
- ³⁶Alberta Precision Laboratories, Public Health Laboratory, Calgary, Alberta, Canada
- ³⁷Mycology Department, Laboratoire de Santé Publique du Québec, Institut National de Santé Publique du Québec (INSPQ), Sainte-Anne-de-Bellevue, Québec, Canada
- ³⁸Department of Pathology and Laboratory Medicine, London Health Sciences Center, London, Ontario, Canada
- ³⁹Department of Mycology, Centre Hospitalier Universitaire de Rennes, Rennes, France
- ⁴⁰University of Texas Health Science Center, San Antonio, Texas, USA
- ⁴¹Clinical and Molecular Microbiology, Department of Pathology, St. Jude Children's Research Hospital, Memphis, Tennessee, USA
- ⁴²Department of Parasitology (Charité), Institute of Microbiology and Hygiene, Humboldt University, Berlin, Germany
- ⁴³Oniris, École Nationale Vétérinaire, Agroalimentaire et de l'Alimentation Nantes-Atlantique, Nantes, France
- ⁴⁴Infectious Disease Research Program, Department of Pediatric Hematology and Oncology and Center for Bone Marrow Transplantation, University Children's Hospital, Münster, Germany
- ⁴⁵Laboratory Diagnostic Center, RWTH Aachen University Hospital, Aachen, Germany
- ⁴⁶University of Utah School of Medicine, Salt Lake City, Utah, USA
- ⁴⁷Loyola University Health System, Loyola University Chicago, Maywood, Illinois, USA
- ⁴⁸Royal Botanic Gardens, Kew, Richmond, Surrey, United Kingdom
- ⁴⁹Natural History Museum, London, United Kingdom
- ⁵⁰University of Southampton, Southampton, United Kingdom
- ⁵¹Jilin Agricultural University, Changchun, China
- ⁵²General Committee for Nomenclature, International Botanical Congress (IBC)
- ⁵³Advisory Board of International Commission on the Taxonomy of Fungi (ICTF)
- ⁵⁴Division of Infectious Diseases, Medical University of Graz, Graz, Austria
- ⁵⁵Translational Medical Mycology Research Unit, ECMM Excellence Center for Medical Mycology, Medical University of Graz, Graz, Austria
- ⁵⁶European Hematology Association, Specialized Working Group for Infections in Hematology, The Hague, the Netherlands
- ⁵⁷Department of Botany, Charles University, Prague, Czechia
- ⁵⁸Clinical Microbiology Laboratory, UMMC Laboratories of Pathology, University of Maryland School of Medicine, Baltimore, Maryland, USA
- ⁵⁹Public Health Ontario, Toronto, Ontario, Canada
- ⁶⁰Department of Laboratory Medicine and Pathobiology, Canada and University of Toronto, Toronto, Ontario, Canada

- ⁶¹Institute of Hygiene and Medical Microbiology, Medical University of Innsbruck, Innsbruck, Austria
- ⁶²Institut Pasteur, Paris, France
- ⁶³University of Alabama at Birmingham, Birmingham, Alabama, USA
- ⁶⁴Centers for Disease Control and Prevention, Atlanta, Georgia, USA
- ⁶⁵Dartmouth Health, Lebanon, New Hampshire, USA
- ⁶⁶National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA
- ⁶⁷Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands
- ⁶⁸University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA
- ⁶⁹University of Texas Health Science Center at Houston, Houston, Texas, USA
- ⁷⁰Clinical Microbiology Laboratory, Wake Forest Baptist Medical Center, Winston-Salem, North Carolina, USA
- ⁷¹The American Board of Pathology, Tampa, Florida, USA
- ⁷²American Board of Pathology (ABP)
- ⁷³National Mycological Herbarium, Ottawa Research and Development Centre, Science and Technology Branch, Agriculture & Agri-Food Canada, Ottawa, Ontario, Canada
- ⁷⁴Cleveland Clinic Lerner College of Medicine, Case Western Reserve University, Cleveland, Ohio, USA
- ⁷⁵Department of Laboratory Medicine, Cleveland Clinic, Cleveland, Ohio, USA
- ⁷⁶Infection Biology Program, Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio, USA
- ⁷⁷Beth Israel Deaconess Medical Center, Boston, Massachusetts, USA
- ⁷⁸Lewis Katz School of Medicine, Temple University, Philadelphia, Pennsylvania, USA
- ⁷⁹Mayo Clinic, Rochester, Minnesota, USA
- ⁸⁰Hippokraton Hospital, Thessaloniki, Greece
- ⁸¹Microbiology Service, Department of Laboratory Medicine, Clinical Center, National Institutes of Health, Bethesda, Maryland, USA
- ⁸²Vanderbilt University Medical Center, Nashville, Tennessee, USA
- ⁸³Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), Buenos Aires, Argentina
- ⁸⁴Unidad de Parasitología, Sector Micología, Hospital J.M. Ramos Mejía, Buenos Aires, Argentina
- ⁸⁵Engineering Laboratory of Microbial Breeding and Preservation of Hebei Province, School of Life Sciences, Institute of Life Sciences and Green Development, Hebei University, Baoding, China
- ⁸⁶Department of Pathology and Laboratory Medicine, Weill Cornell Medical College, New York, USA
- ⁸⁷Public Health Wales Microbiology, Cardiff, United Kingdom
- ⁸⁸Department of Pathology and Laboratory Medicine, University of Vermont Medical Center, Burlington, Vermont, USA
- ⁸⁹Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

AUTHOR ORCID*s*

- Sybre de Hoog  <http://orcid.org/0000-0002-5344-257X>
- Thomas J. Walsh  <http://orcid.org/0000-0002-4142-8711>
- Sarah A. Ahmed  <http://orcid.org/0000-0003-0155-8113>
- Ana Alastruey-Izquierdo  <http://orcid.org/0000-0001-8651-4405>
- Maiken Cavling Arendrup  <http://orcid.org/0000-0002-4747-0144>
- Esther Babady  <http://orcid.org/0000-0003-4399-1249>
- Andrew Borman  <http://orcid.org/0000-0003-0585-5721>
- Andreas H. Groll  <http://orcid.org/0000-0003-1188-393X>
- Kimberly Hanson  <http://orcid.org/0000-0001-9790-277X>
- Randall T. Hayden  <http://orcid.org/0000-0002-3073-4115>

Martin Hoenigl [id http://orcid.org/0000-0002-1653-2824](http://orcid.org/0000-0002-1653-2824)
Julianne V. Kus [id http://orcid.org/0000-0001-6033-7244](http://orcid.org/0000-0001-6033-7244)
Jacques F. Meis [id http://orcid.org/0000-0003-3253-6080](http://orcid.org/0000-0003-3253-6080)
Sixto M. Leal Jr. [id http://orcid.org/0000-0001-6939-9572](http://orcid.org/0000-0001-6939-9572)
Shawn R. Lockhart [id http://orcid.org/0000-0002-4383-5994](http://orcid.org/0000-0002-4383-5994)
Paul Luethy [id http://orcid.org/0000-0001-8636-0258](http://orcid.org/0000-0001-8636-0258)
Kyung J. Kwon-Chung [id http://orcid.org/0000-0001-6671-0994](http://orcid.org/0000-0001-6671-0994)
M. Hong Nguyen [id http://orcid.org/0000-0003-2065-3239](http://orcid.org/0000-0003-2065-3239)
Preeti Pancholi [id http://orcid.org/0000-0001-9053-0264](http://orcid.org/0000-0001-9053-0264)
Gary W. Procop [id http://orcid.org/0000-0002-3164-2968](http://orcid.org/0000-0002-3164-2968)
Daniel D. Rhoads [id http://orcid.org/0000-0002-7636-5191](http://orcid.org/0000-0002-7636-5191)
Stefan Riedel [id http://orcid.org/0000-0002-0573-4957](http://orcid.org/0000-0002-0573-4957)
Paschalis Vergidis [id http://orcid.org/0000-0002-6581-3956](http://orcid.org/0000-0002-6581-3956)
Roxana G. Vitale [id http://orcid.org/0000-0001-8425-0856](http://orcid.org/0000-0001-8425-0856)
Nancy L. Wengenack [id http://orcid.org/0000-0003-2853-8219](http://orcid.org/0000-0003-2853-8219)
Lewis White [id http://orcid.org/0000-0003-3056-4205](http://orcid.org/0000-0003-3056-4205)
Christina M. Wojewoda [id http://orcid.org/0000-0002-7761-4612](http://orcid.org/0000-0002-7761-4612)
Sean X. Zhang [id http://orcid.org/0000-0003-1166-2563](http://orcid.org/0000-0003-1166-2563)

AUTHOR CONTRIBUTIONS

Sybre de Hoog, Conceptualization, Formal analysis, Methodology, Supervision, Validation, Writing – original draft, Writing – review and editing | Thomas J. Walsh, Writing – review and editing | Sarah A. Ahmed, Writing – review and editing | Ana Alastruey-Izquierdo, Writing – review and editing | Barbara D. Alexander, Writing – review and editing | Maiken Cavling Arendrup, Writing – review and editing | Esther Babady, Writing – review and editing | Feng-Yan Bai, Writing – review and editing | Joan-Miquel Balada-Llasat, Writing – review and editing | Andrew Borman, Writing – review and editing | Anuradha Chowdhary, Writing – review and editing | Andrew Clark, Writing – review and editing | Robert C. Colgrove, Writing – review and editing | Oliver A. Cornely, Writing – review and editing | Tanis C. Dingle, Writing – review and editing | Philippe J. Dufresne, Writing – review and editing | Jeff Fuller, Writing – review and editing | Jean-Pierre Gangneux, Writing – review and editing | Connie Gibas, Writing – review and editing | Heather Glasgow, Writing – review and editing | Yvonne Gräser, Writing – review and editing | Jacques Guillot, Writing – review and editing | Andreas H. Groll, Writing – review and editing | Gerhard Haase, Writing – review and editing | Kimberly Hanson, Writing – review and editing | Amanda Harrington, Writing – review and editing | David L. Hawksworth, Writing – review and editing | Randall T. Hayden, Writing – review and editing | Martin Hoenigl, Writing – review and editing | Vit Hubka, Writing – review and editing | Kristie Johnson, Writing – review and editing | Julianne V. Kus, Writing – review and editing | Ruoyu Li, Writing – review and editing | Jacques F. Meis, Writing – review and editing | Michaela Lackner, Writing – review and editing | Francesca Lee, Writing – review and editing | Shawn R. Lockhart, Writing – review and editing | Paul Luethy, Writing – review and editing | Isabella Martin, Writing – review and editing | Kyung J. Kwon-Chung, Writing – review and editing | Wieland Meyer, Writing – review and editing | M. Hong Nguyen, Writing – review and editing | Luis Ostrosky-Zeichner, Writing – review and editing | Elizabeth Palavecino, Writing – review and editing | Preeti Pancholi, Writing – review and editing | Peter G. Pappas, Writing – review and editing | Gary W. Procop, Writing – review and editing | Scott A. Redhead, Writing – review and editing | Daniel D. Rhoads, Writing – review and editing | Stefan Riedel, Writing – review and editing | Bryan Stevens, Writing – review and editing | Kaede Ota Sullivan, Writing – review and editing | Paschalis Vergidis, Writing – review and editing | Emmanuel Roilides, Writing – review and editing | Amir Seyedmousavi, Writing – review and editing | Lili Tao, Writing – review and editing | Vania A. Vicente, Writing – review and editing | Roxana G. Vitale, Writing – review and editing | Qi-Ming Wang, Writing – review and editing | Nancy L. Wengenack,

Writing – review and editing | Lars Westblade, Writing – review and editing | Nathan Wiederhold, Writing – review and editing | Christina M. Wojewoda, Writing – review and editing.

REFERENCES

- Redhead SA. 2010. Report on the special committee on the nomenclature of fungi with a pleomorphic life cycle. *TAXON* 59:1863–1866. <https://doi.org/10.1002/tax.596017>
- Hawksworth DL, Crous PW, Redhead SA, Reynolds DR, Samson RA, Seifert KA, Taylor JW, Wingfield MJ, Abaci O, Aime C, et al. 2011. The Amsterdam declaration on fungal nomenclature. *IMA Fungus* 2:105–112. <https://doi.org/10.5598/imafungus.2011.02.01.14>
- Flann C, Turland NJ, Monro AM. 2014. Report on botanical nomenclature—Melbourne 2011. XVIII international botanical congress, Melbourne: nomenclature section, 18–22 July 2011. *PhytoKeys*:1–289. <https://doi.org/10.3897/phytokeys.41.8398>
- McNeill J, Barrie F, Buck W, Demoulin V, We G, Hawksworth D, Herendeen P, Knapp S, Marhold K, Prado J. 2012. International code of nomenclature for algae, fungi and plants (Melbourne code). Vol. 154. Koeltz scientific books, Königstein.
- Guarro J, Gené J, Ahmed SA, Al-Hatmi AMS, Figueras MJ, Vitale RG. 2020. Atlas of clinical fungi. 4th ed, p 1599. Foundation Atlas of Clinical Fungi, Hilversum.
- Chen M, Zeng J, De Hoog GS, Stielow B, Gerrits Van Den Ende AHG, Liao W, Lackner M. 2016. The ‘species complex’ issue in clinically relevant fungi: a case study in *Scedosporium apiospermum*. *Fungal Biol* 120:137–146. <https://doi.org/10.1016/j.funbio.2015.09.003>
- de Hoog GS, Chaturvedi V, Denning DW, Dyer PS, Frisvad JC, Geiser D, Gräser Y, Guarro J, Haase G, Kwon-Chung K-J, Meis JF, Meyer W, Pitt JI, Samson RA, Taylor JW, Tintelnot K, Vitale RG, Walsh TJ, Lackner M, ISHAM Working Group on Nomenclature of Medical Fungi. 2015. Name changes in medically important fungi and their implications for clinical practice. *J Clin Microbiol* 53:1056–1062. <https://doi.org/10.1128/JCM.02016-14>
- de Hoog GS, Haase G, Chaturvedi V, Walsh TJ, Meyer W, Lackner M. 2013. Taxonomy of medically important fungi in the molecular era. *Lancet Infect Dis* 13:385–386. [https://doi.org/10.1016/S1473-3099\(13\)70058-6](https://doi.org/10.1016/S1473-3099(13)70058-6)
- Kwon-Chung KJ, Bennett JE, Wickes BL, Meyer W, Cuomo CA, Wollenburg KR, Bicanic TA, Castañeda E, Chang YC, Chen J, et al. 2017. The case for adopting the “species complex” nomenclature for the etiologic agents of cryptococcosis. *mSphere* 2:e00357-16. <https://doi.org/10.1128/mSphere.00357-16>
- Yurkov A, Alves A, Bai F-Y, Boundy-Mills K, Buzzini P, Čadež N, Cardinali G, Casaregola S, Chaturvedi V, Collin V, et al. 2021. Nomenclatural issues concerning cultured yeasts and other fungi: why it is important to avoid unneeded name changes. *IMA Fungus* 12:18. <https://doi.org/10.1186/s43008-021-00067-x>
- Borman AM, Johnson EM. 2021. Name changes for fungi of medical importance, 2018 to 2019. *J Clin Microbiol* 59:e01811-20. <https://doi.org/10.1128/JCM.01811-20>
- Kidd SE, Abdolrasouli A, Hagen F. 2023. Fungal nomenclature: managing change is the name of the game. Oxford University Press, USA.
- Kidd SE, Halliday CL, McMullan B, Chen S-A, Elvy J. 2021. New names for fungi of medical importance: can we have our cake and eat it too. *J Clin Microbiol* 59:e02730-20. <https://doi.org/10.1128/JCM.02730-20>
- Schoch CL, Sung G-H, López-Giráldez F, Townsend JP, Miadlikowska J, Hofstetter V, Robbertse B, Matheny PB, Kauff F, Wang Z, Gueidan C, et al. 2009. The ascomycota tree of life: a phylum-wide phylogeny clarifies the origin and evolution of fundamental reproductive and ecological traits. *Syst Biol* 58:224–239. <https://doi.org/10.1093/sysbio/syp020>
- Samson RA, Yilmaz N, Houbraken J, Spierenburg H, Seifert KA, Peterson SW, Varga J, Frisvad JC. 2011. Phylogeny and nomenclature of the genus *Talaromyces* and taxa accommodated in *Penicillium* subgenus *Biverticillium*. *Stud Mycol* 70:159–183. <https://doi.org/10.3114/sim.2011.70.04>
- de Hoog GS, Dukik K, Monod M, Packeu A, Stubbe D, Hendrickx M, Kupsch C, Stielow JB, Freeke J, Göker M, Rezaei-Matehkolaei A, Mirhendi H, Gräser Y. 2017. Toward a novel multilocus phylogenetic taxonomy for the dermatophytes. *Mycopathologia* 182:5–31. <https://doi.org/10.1007/s11046-016-0073-9>
- Weijman AC. 1979. Carbohydrate composition and taxonomy of *Geotrichum*, *Trichosporon* and allied genera. *Antonie Van Leeuwenhoek* 45:119–127. <https://doi.org/10.1007/BF00400785>
- de Beer ZW, Begerow D, Bauer R, Pegg GS, Crous PW, Wingfield MJ. 2006. Phylogeny of the Quambalariaceae fam. nov., including important eucalyptus pathogens in South Africa and Australia. *Stud Mycol* 55:289–298. <https://doi.org/10.3114/sim.55.1.289>
- Sugita R, Tanaka K. 2022. *Thyridium* revised: synonymisation of *Phialemoniopsis* under *Thyridium* and establishment of a new order, thyridiales. *MycKeys* 86:147–176. <https://doi.org/10.3897/mycokeys.86.78989>
- Connally A, Smith D, Marek S, Wu Y, Walker N. 2022. Phylogenetic evaluation of *Bipolaris* and *Curvularia* species collected from turfgrasses. *Int Turfgrass Soc Res J* 14:916–930. <https://doi.org/10.1002/its2.16>
- Manamgoda DS, Cai L, McKenzie EHC, Crous PW, Madrid H, Chukeatirote E, Shivas RG, Tan YP, Hyde KD. 2012. A phylogenetic and taxonomic re-evaluation of the *Bipolaris-Cochliobolus-Curvularia* complex. *Fungal Div* 56:131–144. <https://doi.org/10.1007/s13225-012-0189-2>
- Wang XW, Houbraken J, Groenewald JZ, Meijer M, Andersen B, Nielsen KF, Crous PW, Samson RA. 2016. Diversity and taxonomy of *Chaetomium* and chaetomium-like fungi from indoor environments. *Stud Mycol* 84:145–224. <https://doi.org/10.1016/j.smyco.2016.11.005>
- Zachos FE. 2016. Species concepts in biology. Vol. 801. Springer. <https://doi.org/10.1007/978-3-319-44966-1>
- Anzawa K, Kawasaki M, Mochizuki T, Ishizaki H. 2010. Successful mating of trichophyton rubrum with *Arthroderma simii*. *Med Mycol* 48:629–634. <https://doi.org/10.3109/13693780903437884>
- Hubka V, Barrs V, Dudová Z, Sklenář F, Kubátová A, Matsuzawa T, Yaguchi T, Horie Y, Nováková A, Frisvad JC, Talbot JJ, Kolařík M. 2018. Unravelling species boundaries in the *Aspergillus viridinutans* complex (section *Fumigati*): opportunistic human and animal pathogens capable of interspecific hybridization. *Persoonia* 41:142–174. <https://doi.org/10.3767/persoonia.2018.41.08>
- Hartmann FE, Duhamel M, Carpentier F, Hood ME, Foulongne-Oriol M, Silar P, Malagnac F, Grognet P, Giraud T. 2021. Recombination suppression and evolutionary strata around mating - type loci in fungi: documenting patterns and understanding evolutionary and mechanistic causes. *New Phytol* 229:2470–2491. <https://doi.org/10.1111/nph.17039>
- Yadav V, Sun S, Heitman J. 2021. Uniparental nuclear inheritance following bisexual mating in fungi. *Elife* 10:e66234. <https://doi.org/10.7554/eLife.66234>
- Zhao L, de Hoog S, Hagen F, Kang Y, Al-Hatmi AMS. 2019. Species borderlines in *Fusarium* exemplified by *F. circinatum*/*F. subglutinans*. *Fungal Genet Biol* 132:103262. <https://doi.org/10.1016/j.fgb.2019.103262>
- Samarasinghe H, Xu J. 2018. Hybrids and hybridization in the *Cryptococcus neoformans* and *Cryptococcus gattii* species complexes. *Infect Genet Evol* 66:245–255. <https://doi.org/10.1016/j.meegid.2018.10.011>
- Samarasinghe H, You M, Jenkinson TS, Xu J, James TY. 2020. Hybridization facilitates adaptive evolution in two major fungal pathogens. *Genes* 11:101. <https://doi.org/10.3390/genes11010101>
- Wilson AM, Wilken PM, van der Nest MA, Steenkamp ET, Wingfield MJ, Wingfield BD. 2015. Homothallism: an umbrella term for describing diverse sexual behaviours. *IMA Fungus* 6:207–214. <https://doi.org/10.5598/imafungus.2015.06.01.13>
- Stielow JB, Lévesque CA, Seifert KA, Meyer W, Iriny L, Smits D, Renfurm R, Verkley GJM, Groenewald M, Chaduli D, et al. 2015. One fungus, which genes? Development and assessment of universal primers for potential secondary fungal DNA barcodes. *Persoonia* 35:242–263. <https://doi.org/10.3767/003158515X689135>
- Sandoval-Denis M, Sutton DA, Martin-Vicente A, Cano-Lira JF, Wiederhold N, Guarro J, Gené J. 2015. *Cladosporium* species recovered

- from clinical samples in the United States. *J Clin Microbiol* 53:2990–3000. <https://doi.org/10.1128/JCM.01482-15>
34. Wang MM, Crous PW, Sandoval-Denis M, Han SL, Liu F, Liang JM, Duan WJ, Cai L. 2022. *Fusarium* and allied genera from China: species diversity and distribution. *Persoonia* 48:1–53. <https://doi.org/10.3767/persoonia.2022.48.01>
 35. Hudson O, Waliullah S, Fulton JC, Ji P, Dufault NS, Keinath A, Ali ME. 2021. Marker development for differentiation of *Fusarium oxysporum* f. sp. niveum race 3 from races 1 and 2. *Int J Molec Sci* 22:822. <https://doi.org/10.3390/ijms22020822>
 36. Vlaardingerbroek I, Beerens B, Schmidt SM, Cornelissen BJC, Rep M. 2016. Dispensable chromosomes in *Fusarium oxysporum* f. sp. *lycopersici*. *Mol Plant Pathol* 17:1455–1466. <https://doi.org/10.1111/mpp.12440>
 37. Esposito MC, Prigitano A, Tortorano AM. 2016. *Fusarium musae* as cause of superficial and deep-seated human infections. *J Mycol Med* 26:403–405. <https://doi.org/10.1016/j.mycmed.2016.02.021>
 38. Thitla T, Kumla J, Khuna S, Lumyong S, Suwannarach N. 2022. Species diversity, distribution, and phylogeny of *Exophiala* with the addition of four new species from Thailand. *J Fungi* 8:766. <https://doi.org/10.3390/jof8080766>
 39. Verbeke V, Bourgeois T, Lodewyck T, Van Praet J, Lagrou K, Reynders M, Nulens E. 2020. Successful outcome of disseminated *Fusarium* musae fungemia with skin localization treated with liposomal amphotericin B and voriconazole in a patient with acute myeloid leukemia. *Mycopathologia* 185:1085–1089. <https://doi.org/10.1007/s11046-020-00499-w>
 40. Gräser Y, De Hoog S, Summerbell RC. 2006. Dermatophytes: recognizing species of clonal fungi. *Med Mycol* 44:199–209. <https://doi.org/10.1080/13693780600606810>
 41. Taylor JW, Jacobson DJ, Kroken S, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. 2000. Phylogenetic species recognition and species concepts in fungi. *Fungal Genet Biol* 31:21–32. <https://doi.org/10.1006/fgbi.2000.1228>
 42. Kandemir H, Dukik K, Hagen F, Ilkit M, de Hoog G. 2019. A Polyphasic approach for the differentiation *Trichophyton tonsurans* and *T. Equinum*. *Mycopathologia* 185:113–122.
 43. Kosanke S, Hamann L, Kupsch C, Moreno Garcia S, Chopra A, Gräser Y. 2018. Unequal distribution of the mating type (MAT) locus idiomorphs in dermatophyte species. *Fungal Genet Biol* 118:45–53. <https://doi.org/10.1016/j.fgb.2018.07.003>
 44. Taghipour S, Pchelin IM, Zarei Mahmoudabadi A, Ansari S, Katirae F, Rafiei A, Shokohi T, Abastabar M, Taraskina AE, Kermani F, Diba K, Nouripour-Sisakht S, Najafzadeh MJ, Pakshir K, Zomorodian K, Ahmadi K, Rezaei-Matehkolaei A. 2019. *Trichophyton mentagrophytes* and T interdigitale genotypes are associated with particular geographic areas and clinical manifestations. *Mycoses* 62:1084–1091. <https://doi.org/10.1111/myc.12993>
 45. Kuramae EE, Robert V, Echavari-Erasun C, Boekhout T. 2007. Cophenetic correlation analysis as a strategy to select phylogenetically informative proteins: an example from the fungal kingdom. *BMC Evol Biol* 7:134. <https://doi.org/10.1186/1471-2148-7-134>
 46. van Dam P, de Sain M, Ter Horst A, van der Gragt M, Rep M. 2018. Use of comparative genomics-based markers for discrimination of host specificity in *Fusarium oxysporum*. *Appl Environ Microbiol* 84:e01868-17. <https://doi.org/10.1128/AEM.01868-17>
 47. Vilela R, de Hoog S, Bensch K, Bagagli E, Mendoza L. 2023. A taxonomic review of the genus *Paracoccidioides*, with focus on the uncultivable species. *PLOS Negl Trop Dis* 17:e0011220. <https://doi.org/10.1371/journal.pntd.0011220>
 48. Tang C, Ahmed SA, Deng S, Zhang L, Zoll J, Al-Hatmi AMS, Meis JF, Thakur R, Kang Y, de Hoog GS. 2022. Detection of emerging genotypes in *Trichophyton mentagrophytes* species complex: a proposal for handling biodiversity in dermatophytes. *Front Microbiol* 13:960190. <https://doi.org/10.3389/fmicb.2022.960190>
 49. James TY, Stajich JE, Hittinger CT, Rokas A. 2020. Toward a fully resolved fungal tree of life. *Annu Rev Microbiol* 74:291–313. <https://doi.org/10.1146/annurev-micro-022020-051835>
 50. Sandoval-Denis M, Gené J, Sutton DA, Cano-Lira JF, de Hoog GS, Decock CA, Wiederhold NP, Guarro J. 2016. Redefining *Microascus*, *Scopulariopsis* and allied genera. *Persoonia* 36:1–36. <https://doi.org/10.3767/003158516X688027>
 51. Kocsubé S, Perrone G, Magistà D, Houbraeken J, Varga J, Szigeti G, Hubka V, Hong S-B, Frisvad JC, Samson RA. 2016. *Aspergillus* is monophyletic: evidence from multiple gene phylogenies and extrolites profiles. *Stud Mycol* 85:199–213. <https://doi.org/10.1016/j.simyco.2016.11.006>
 52. Houbraeken J, Kocsubé S, Visagie CM, Yilmaz N, Wang X-C, Meijer M, Kraak B, Hubka V, Bensch K, Samson RA, Frisvad JC. 2020. Classification of *Aspergillus*, *Penicillium*, *Talaromyces* and related genera (*Eurotiales*): an overview of families, genera, subgenera, sections, series and species. *Stud Mycol* 95:5–169. <https://doi.org/10.1016/j.simyco.2020.05.002>
 53. Quan Y, Muggia L, Moreno LF, Wang M, Al-Hatmi AMS, da Silva Menezes N, Shi D, Deng S, Ahmed S, Hyde KD, Vicente VA, Kang Y, Stielow JB, de Hoog S. 2020. A re-evaluation of the Chaetothyriales using criteria of comparative biology. *Fungal Div* 103:47–85. <https://doi.org/10.1007/s13225-020-00452-8>
 54. Lücking R, Aime MC, Robbertse B, Miller AN, Ariyawansa HA, Aoki T, Cardinali G, Crous PW, Druzhinina IS, Geiser DM, et al. 2020. Unambiguous identification of fungi: where do we stand and how accurate and precise is fungal DNA barcoding? *IMA Fungus* 11:14. <https://doi.org/10.1186/s43008-020-00033-z>
 55. Lücking R, Aime MC, Robbertse B, Miller AN, Aoki T, Ariyawansa HA, Cardinali G, Crous PW, Druzhinina IS, Geiser DM, et al. 2021. Fungal taxonomy and sequence-based nomenclature. *Nat Microbiol* 6:971. <https://doi.org/10.1038/s41564-021-00921-z>
 56. Redhead SA. 2001. Bully for *Coprinus* – a story of manure, minutiae and molecules. *Field Mycol* 2:118–126. [https://doi.org/10.1016/S1468-1641\(10\)60532-4](https://doi.org/10.1016/S1468-1641(10)60532-4)
 57. Bian C, Kusuya Y, Sklenář F, D'hooge E, Yaguchi T, Ban S, Visagie C, Houbraeken J, Takahashi H, Hubka V. 2022. Reducing the number of accepted species in *Aspergillus* series *Nigri*. *Stud Mycol* 102:95–132. <https://doi.org/10.3114/sim.2022.102.03>
 58. Sklenář F, Glässnerová K, Jurjević Ž, Houbraeken J, Samson RA, Visagie CM, Yilmaz N, Gené J, Cano J, Chen AJ, Nováková A, Yaguchi T, Kolařík M, Hubka V. 2022. Taxonomy of *Aspergillus* series *Versicolores*: species reduction and lessons learned about intraspecific variability. *Stud Mycol* 102:53–93. <https://doi.org/10.3114/sim.2022.102.02>
 59. Hagen F, Lumbsch HT, Arsic Arsenijevic V, Badali H, Bertout S, Billmyre RB, Bragulat MR, Cabañes FJ, Carbia M, Chakrabarti A, et al. 2017. Importance of resolving fungal nomenclature: the case of multiple pathogenic species in the *Cryptococcus* genus. *mSphere* 2:e00238-17. <https://doi.org/10.1128/mSphere.00238-17>
 60. Crous PW, Lombard L, Sandoval-Denis M, Seifert KA, Schroers H-J, Chaverri P, Gené J, Guarro J, Hirooka Y, Bensch K, Kema GHJ, et al. 2021. *Fusarium*: more than a node or a foot-shaped basal cell. *Stud Mycol* 98:100116. <https://doi.org/10.1016/j.simyco.2021.100116>
 61. Geiser DM, Al-Hatmi AMS, Aoki T, Arie T, Balmas V, Barnes I, Bergstrom GC, Bhattacharyya MK, Blomquist CL, Bowden RL, et al. 2021. Phylogenetic analysis of a 55.1-kb 19-gene dataset resolves a monophyletic *Fusarium* that includes the *Fusarium solani* species complex. *Phytopathology* 111:1064–1079. <https://doi.org/10.1094/PHYTO-08-20-0330-LE>
 62. Geiser DM, Aoki T, Bacon CW, Baker SE, Bhattacharyya MK, Brandt ME, Brown DW, Burgess LW, Chulze S, Coleman JJ, et al. 2013. One fungus, one name: defining the genus *Fusarium* in a scientifically robust way that preserves longstanding use. *Phytopathology* 103:400–408. <https://doi.org/10.1094/PHYTO-07-12-0150-LE>
 63. O'Donnell K, Al-Hatmi AMS, Aoki T, Brankovics B, Cano-Lira JF, Coleman JJ, de Hoog GS, Di Pietro A, Frandsen RJN, Geiser DM, et al. 2020. No to *Neocosmospora*: phylogenomic and practical reasons for continued inclusion of the *Fusarium solani* species complex in the genus *Fusarium*. *mSphere* 5:e00810-20. <https://doi.org/10.1128/mSphere.00810-20>
 64. May TW, Redhead SA, Bensch K, Hawksworth DL, Lendemer J, Lombard L, Turland NJ. 2019. Chapter F of the *International Code of Nomenclature for algae, fungi, and plants* as approved by the 11th International mycological Congress, San Juan, Puerto Rico, July 2018. *IMA Fungus* 10:21. <https://doi.org/10.1186/s43008-019-0019-1>
 65. Turland N, Wiersma J, Barrie F, Greuter W, Hawksworth D, Herendeen P, Knapp S, Kusber W-H, Li D-Z, Marhold K, May T, McNeill J, Monro A, Prado J, Price M, Smith G. 2018. Adopted by the Nineteenth International Botanical Congress. *International code of nomenclature for algae, fungi, and plants* (shenzhen code). Koeltz Botanical Books, Shenzhen, China. <https://doi.org/10.12705/Code.2018>