

Tomato mutants sensitive to abiotic stress display different abscisic acid content and metabolism during germination

Andrea Andrade, Oscar Masciarelli, Sergio Alemano, Virginia Luna and Guillermina Abdala*

Laboratorio de Fisiología Vegetal, Departamento de Ciencias Naturales, Facultad de Ciencias Exactas, Físico-Químicas y Naturales, Universidad Nacional de Río Cuarto, 5800 Río Cuarto, Argentina

Abstract

We report the determination of abscisic acid (ABA) and its metabolites, phaseic acid (PA), dihydrophaseic acid (DPA) and ABA glucose ester (ABA-GE), in non-dormant dry and imbibed seeds of tomato (*Solanum lycopersicum* Mill.) cv. Moneymaker (wild type), and its *tss1*, *tss2* and *tos1* mutants. High ABA in dry seeds may originate from ABA accumulation in the sheath tissue, which was in contact with an ABA-containing medium, the endocarpus. The highest germination percentages at 72 h, observed in *tss1* and *tss2*, coincided with minimal ABA content. Wild-type and mutant seeds showed different ABA and catabolic patterns, and these were correlated with their sensitivity to abiotic stress. Whereas dry seeds showed a high basal ABA, imbibed seeds showed higher ABA metabolite content, particularly DPA. The dramatic decrease of ABA following seed imbibition suggests an activation of ABA catabolism during the early stages of the germination process. The observed variation of ABA metabolites among dry and imbibed seeds of *Solanum lycopersicum* cv. Moneymaker and its *tss1*, *tss2* and *tos1* mutants shows that ABA metabolism is differentially regulated in these genotypes.

Keywords: abiotic stress, abscisic acid, abscisic acid metabolites, germination, *Solanum lycopersicum*, tomato mutants

Introduction

Several metabolites of abscisic acid (ABA) have been isolated from various plant materials. ABA catabolism proceeds through two major pathways: the

'oxidative pathway' involving oxidation at different positions, and the 'conjugation pathway'. The pathway used depends on the plant species, developmental stage or tissue type. The oxidative pathway, considered the more common in plant catabolism (Nambara and Marion-Poll, 2005), is initiated by hydroxylation at C-8' to produce 8'-hydroxy-ABA (8'-HOABA), which rearranges to phaseic acid (PA) (Cutler and Krochko, 1999; Todoroki *et al.*, 2000). PA is subsequently reduced to dihydrophaseic acid (DPA) and/or its analogue, *epi*-dihydrophaseic acid (*epi*-DPA) (Zeevaert *et al.*, 1991).

Recently, Zhou *et al.* (2004) described a new oxidative pathway in which hydroxylation of ABA occurred at the 9'-methyl group of ABA, as well as at the 7'- and 8'-methyl groups. The new metabolite isolated from the plant extract was identified as the closed form of 9'-hydroxy-ABA and was named *neophaseic acid* (neoPA). Levels of neoPA were high in immature tomato seeds, but disappeared at later stages of seed development, suggesting further metabolism (Zaharia *et al.*, 2005).

ABA and hydroxyl-ABA can be conjugated as glucose esters (ABA-GE) or glycosides. ABA-GE, a major ABA-conjugated metabolite, is formed by an ABA glucosyltransferase (Xu *et al.*, 2002). The abundance of ABA-GE is low in comparison with ABA, and it may be part of a reversible inactivation reaction used to regenerate free ABA (Sauter *et al.*, 2002). The fate of ABA-GE is still unknown (Zaharia *et al.*, 2005), although it has been hypothesized to contribute to ABA homeostasis in plant cells (Chiwocha *et al.*, 2005).

The hormonal activity reported for some ABA metabolites (Zhou *et al.*, 2004) has raised the intriguing possibility that these metabolites may mediate hormonal effects hitherto associated only with ABA *per se*. Thus, further elucidation of the physiological roles of ABA requires quantitative analysis of not only ABA, but also of its metabolites (Setha *et al.*, 2005).

*Correspondence

Fax: 0054-358-4676230

Email: gabdala@exa.unrc.edu.ar

In angiosperms, the ABA that accumulates during mid-maturation of seeds is synthesized in both zygotic and maternal tissues; thus, the latter is most likely transported from the mother plant. The site of ABA synthesis specifies its physiological action. Maternal ABA promotes reserve accumulation, and embryonic ABA induces seed dormancy and desiccation tolerance (Lefebvre *et al.*, 2006). Many mutations affecting plant responses to a particular hormone have been identified through genetic analysis, including several mutants with altered sensitivity to ABA (Leung and Giraudat, 1998; Moller and Chua, 1999). A number of salt (NaCl)-hypersensitive mutants have been isolated in *Arabidopsis thaliana* (Zhu *et al.*, 1998) and *Solanum lycopersicum* (Borsani *et al.*, 2001, 2002).

In *Solanum lycopersicum*, these mutants have lost certain salt-tolerance mechanisms, and are therefore more sensitive to salt stress than the wild-type cv. MoneyMaker. The *tss1* (*tomato salt sensitive 1*) mutant is hypersensitive to growth inhibition by Na⁺ but not to osmotic stress induced by mannitol, whereas *tss2* (*tomato salt sensitive 2*) is hypersensitive to both ionic and osmotic stresses (Borsani *et al.*, 2001). Another mutant, *tos1* (*tomato osmotic sensitive*), has a single recessive nuclear mutation responsible for its hypersensitivity to general osmotic stress (Borsani *et al.*, 2002). These three mutants are all similar to the wild type in their growth characteristics, with the distinctive phenotypes becoming evident only under stress conditions. Borsani *et al.* (2002) observed that growth inhibition by applied ABA was greater in *tss2* (hypersensitive to ABA) than in the wild type, *tss1*, or *tos1*. They proposed that the decreased sensitivity of *tos1* seedlings and increased sensitivity of *tss2* seedlings to ABA could result from either abnormal ABA metabolism or altered ABA signal transduction, and that *tss2* and *tos1* are not ABA-deficient mutants.

ABA clearly plays a key role in seed germination, but the contribution of ABA catabolism to the regulation of endogenous ABA content in this process is not clear. Likewise, in the above tomato mutants, modifications in ABA metabolism during seed imbibition, and the possible role of ABA metabolism in germination and sensitivity to osmotic and ionic stress, remain to be studied. Thus, we have investigated: (1) the correlation between endogenous ABA content and germination response of non-dormant seeds of cv. MoneyMaker and the hypersensitive tomato mutants *tss1*, *tss2*, and *tos1*; (2) ABA, PA, DPA, and ABA-GE accumulation during the imbibition process; and (3) possible relationships between ABA content and ABA metabolic patterns with previously reported differential sensitivity of the mutants.

Materials and methods

Plant materials

Seeds of tomato (*Solanum lycopersicum* Mill.) cv. MoneyMaker (wild type), and its homozygous *tss1*, *tss2*, and *tos1* mutants, generously provided by Dr Miguel A. Botella (Department of Molecular Biology and Biochemistry, University of Málaga, Spain), were used for the assays. Two alleles of the *tss1* mutant were reported by Borsani *et al.* (2001); in our experiments the allele *tss1-1* was used. Variability due to different growth and seed storage conditions was avoided by growing wild-type and mutant plants side by side. Seeds were extracted from mature fruits, fermented in the juice for 1 d, rinsed, dried at room temperature and stored at 4°C for 6 months. For each genotype, three biological replicates of equivalent age were processed.

Imbibition and germination assays

A simple random design consisting of ten imbibition times was used. For each experiment, 20 seeds were placed in a 7-ml flask with 300 µl distilled water (three replicate flasks per experiment). The number of germinated seeds was recorded daily at 12, 24, 48, 72, 96, 120, 144, 168, 192 and 216 h, and expressed as a percentage of the total (%G). Germination was defined as the presence of a clearly visible radicle protrusion (Ni and Bradford, 1993). Experiments were performed under controlled conditions in a Conviron G 30 germination chamber (Winnipeg, Manitoba, Canada) at 25 ± 1°C, relative humidity 80%, in the dark.

Results were analysed using the program Statgraphics Plus, version 3 (Manugistics, Rockville, Maryland, USA). Germination data as percentage of total number of seeds were arcsine transformed. A one-way ANOVA procedure was used to compare the effects of different imbibition times in distilled water. Normality was verified by the Shapiro–Wilk test. Homogeneity of variance was verified by the Bartlett test. *P* values ≤ 0.05 were considered statistically significant.

Endogenous concentrations of ABA, PA, DPA and ABA-GE were determined in dry and imbibed seeds at 12, 24, 48 and 72 h. The total volume of imbibition water was tested for the presence of free ABA at each imbibition time.

Abscisic acid extraction procedure

ABA was extracted and pre-purified by a modified protocol of Luna *et al.* (1993). The equivalent of 200 mg dry weight of dry or imbibed seeds was ground in a mortar with liquid nitrogen and 20 ml imidazole buffer (pH 7) plus 2,6-di-tert-butyl-p-cresol as

antioxidant. Fifty nanograms of [$^2\text{H}_6$]ABA (OChemIm Ltd, Olomouc, Czech Republic) were added as internal standard, and the sample was incubated overnight at 4°C to allow extraction and standard equilibration. After centrifugation and isopropanol evaporation, the aqueous fraction was loaded on to a conditioned amino anion exchange minicolumn [BAKERBOND speTM Amino (NH₂), Mallinckrodt Baker, Inc., Phillipsburg, New Jersey, USA] and washed sequentially with 6 ml each of hexane, ethyl acetate and acetonitrile. These fractions were discarded, and then ABA was eluted with methanol:acetic acid (95:5, v/v) and evaporated to dryness.

HPLC procedure for abscisic acid purification

Dried extracts were dissolved in 100 μl of elution solvent [methanol:water:acetic acid (70:30:0.1 v/v)], and separated on a preparative high-performance liquid chromatography (HPLC) system (KNK-500, Konic Instruments, Barcelona, Spain) equipped with an RP C₁₈ column (μ -Bondapack, 300 \times 3.9 mm internal diameter, 5 μm particle size; Waters Associates, Milford, Massachusetts, USA) coupled to a spectrometry system (UV-Vis) with diode array detector. Samples were subsequently eluted at a flow rate of 1 ml min⁻¹ with an isocratic mixture of methanol:water:acetic acid (70:30:0.1 v/v). Fractions corresponding to ABA (retention time: 2.77–5.00 min), as determined spectrophotometrically at 262 nm on a previous HPLC run, were pooled and dried.

Phaseic acid, dihydrophaseic acid and ABA-glucose ester extraction and purification

PA, DPA and ABA-GE were extracted and pre-purified as described by Zhou *et al.* (2003) with modifications. The equivalent of 200 mg dry weight of dry or imbibed seeds was ground in a mortar with liquid nitrogen and extracted with 3 ml of acetone:water:acetic acid (80:19:1 v/v). One hundred nanograms of [$^2\text{H}_3$]PA, [$^2\text{H}_3$]DPA and [$^2\text{H}_3$]ABA-GE (NRC-Plant Biotechnology Institute, Saskatoon, Canada) were added as internal standards. Extracts were transferred to 50 ml tubes, centrifuged at 5000 rpm for 15 min, and the supernatant was collected and evaporated at 35°C under vacuum in a SpeedVac. Dried extracts were dissolved in 100 μl methanol:acetic acid (99:1 v/v), then mixed with 900 μl of 1% acetic acid. The samples were filtered through a syringe filter tip and purified with 3 ml BondElut-C18 cartridges (Varian, Palo Alto, California, USA) on a vacuum manifold at a flow-rate less than 1 ml min⁻¹. The cartridges were conditioned with 1.5 ml methanol and equilibrated with 1.5 ml methanol:water:acetic acid (10:89:1 v/v). Samples (each ~1.5 ml volume) were loaded on to the cartridges

and washed with 1.5 ml methanol:water:acetic acid (10:89:1 v/v). ABA metabolites were eluted by 1.5 ml methanol:water:acetic acid (80:19:1 v/v) and collected in a 2-ml flat-bottom Eppendorf tube. The eluate was dried under vacuum at 35°C during centrifugation.

GC-MS-SIM

Samples containing ABA or ABA metabolites were dissolved in 100 μl of derivatization solution *N,O*-bis(trimethylsilyl)trifluoroacetamide (BSTFA) and then converted to methylester-trimethylsilylether (MeTMSi) derivatives. The samples were placed in an oven with increasing temperature from 70°C to 90°C for 30 min, and 1 μl was injected split-splitless in a gas chromatography-mass spectrometry with selected ion monitoring (GC-MS-SIM) system (Hewlett Packard 5890 Series II GC, with capillary direct interface to a 5972 Mass Selective Detector) equipped with a 25 m Chrompack CPSil 19 capillary column (internal diameter 0.25 mm; film thickness 0.22 μm). Carrier gas: He at flow rate of 1 ml min⁻¹; GC injector temperature: 280°C; oven temperature: initially maintained at 100°C for 1 min, then increased from 100 to 195°C at a rate of 15°C min⁻¹, then at a rate of 4°C min⁻¹ up to 280°C. For ABA quantification, ions 196 (deutero) and 190 (proto) were monitored at 9–10 min; for PA, ions 299 and 296 were monitored at 9–10 min; for DPA, ions 297 and 294 were monitored at 10–11 min; for ABA-GE, ions 268 and 263 were monitored at 11–12 min. Quantification of ABA-GE was performed as derivatized conjugate. For unequivocal determination, back-up ions (labelled ABA 134, labelled PA 279, labelled DPA 281, labelled ABA-GE 430) were monitored for peak confirmation. The amounts of endogenous ABA, PA, DPA and ABA-GE were calculated by comparison of the peak areas of the ions for the [^2H] isotope internal standard versus its proto counterpart, and the analysis was done in triplicate.

Data were expressed as means \pm SE of three independent determinations of three samples extracted from experimental treatments (imbibition time) and analysed using the Statgraphics Plus program. ABA and metabolite content in dry and imbibed seeds, as well as effects of different imbibition times on endogenous ABA and metabolite content, were compared by one-way ANOVA and Tukey's HSD test. *P* values \leq 0.05 were considered to be statistically significant.

Results

Endogenous content of ABA and its metabolites in dry seeds

Endogenous contents of ABA and its metabolites in dry seeds of cv. MoneyMaker (wild type), and its

tss1, *tss2*, and *tos1* mutants, are summarized in Fig. 1. ABA content was highest in wild-type seeds followed by *tss1* and *tos1*, and lowest in *tss2* [$732 \text{ pmol (g DW)}^{-1}$]. Differences in ABA content were significant between wild type and *tss2* and *tos1*, and among all mutants. PA, DPA and ABA-GE were detected in wild-type and mutant seeds. ABA metabolites were much lower than ABA in all genotypes. Mutants *tss2* and *tos1* showed similar distribution patterns, with PA being the least abundant metabolite. In the wild type, the total metabolites [PA + DPA + ABA-GE = $1.149 \text{ pmol (g DW)}^{-1}$] content was 83-fold lower than that of ABA (Fig. 1A). In *tss1* and *tos1*, metabolite contents were, respectively, 84-fold and 37-fold lower than that of ABA (Fig. 1B and D). In contrast, the difference between ABA and total metabolite level [$650 \text{ pmol (g DW)}^{-1}$] in *tss2* was only slight (Fig. 1C).

DPA was consistently the most abundant ABA metabolite, and was higher in the wild type than in the mutants. Among the mutants, DPA level was highest in *tss2*. PA was a minor component in all genotypes, and conjugated ABA were twofold higher in the wild type than in the mutants.

Germination of wild-type and mutant seeds

The time course of germination of tomato wild-type and mutant seeds is shown in Fig. 2. None of the genotypes showed germination before 24 h of imbibition. In *tss1*, germination percentage was 33% at 48 h and 41% at 72 h of imbibition. *Tss2* showed 5% germination at 48 h and 62% germination at 72 h. The slowest germination was found for the wild type and *tos1* mutant: 23% and 25%, respectively, after 72 h of imbibition. Differences in germination percentage at 72 h were significant between the wild type and *tss2*, and between *tss2* and *tos1*. At 216 h, germination for the wild type and *tss2* mutant was 100%, and for the *tss1* and *tos1* mutants it was around 70%.

Endogenous levels of ABA in imbibed seeds and imbibition water

We studied free ABA content during the imbibition process, and possible secretion of ABA from seeds into the imbibition water. During the imbibition time course, wild-type seeds showed a significant ABA




Figure 1. Abscisic acid (ABA) and ABA metabolite content [phaseic acid (PA), dihydrophaseic acid (DPA) and ABA glucose ester (ABA-GE)] of dry tomato seeds. (A) *Solanum lycopersicum* Mill. cv. Moneymaker (wild type); (B) *tss1* mutant; (C) *tss2* mutant; (D) *tos1* mutant. Data are means of three replicates with SEs ($P \leq 0.05$).




Figure 2. Germination time course of wild-type, *tss1*, *tss2* and *tos1* tomato seeds. Data are means of three replicates with SEs. Values with the same letter at 72 h are not significantly different ($P \leq 0.05$). Data points ≥ 96 h are from single observations, due to lack of seed material.

increase from 12 to 48 h, a maximum [$749 \text{ pmol (g DW)}^{-1}$] at 48 h, and decrease to $468 \text{ pmol (g DW)}^{-1}$ at 72 h. Secretion of ABA into the water was low and sustained during the imbibition process (Fig. 3A).

During *tss1* imbibition, seed ABA increased sharply at 24 h, reached a maximum [$754 \text{ pmol (g DW)}^{-1}$] at 48 h, then decreased sharply at 72 h. There were no significant changes in ABA secretion into the imbibition water (Fig. 3B).

ABA content in *tss2* imbibed seeds was considerably lower than that in wild type and *tss1*, and no significant variations were observed during the course of imbibition. Interestingly, ABA was secreted at much higher levels than found in the imbibed seeds (Fig. 3C).

In *tos1* imbibed seeds, ABA content was highest [$186 \text{ pmol (g DW)}^{-1}$] at 12 h, and was slightly lower at 24 h and thereafter. *Tos1* seeds also secreted high amounts of ABA into the imbibition water. ABA secreted by *tos1* seeds reached a peak of $496 \text{ pmol (g DW)}^{-1}$ at 24 h, 200% higher than the level of free ABA measured in the seeds at the same time. Similar values were found at 48 and 72 h (Fig. 3D).

Thus, levels of ABA secreted by *tos1* and *tss2* into imbibition water were high, as compared to




Figure 3. Abscisic acid content of imbibed tomato seeds and their incubation water during the germination time course: (A) cv. MoneyMaker (wild type); (B) *tss1* mutant; (C) *tss2* mutant; (D) *tos1* mutant. Results are expressed in pmol (g DW)^{-1} for seeds and in pmol ml^{-1} for imbibition water. Data are means of three replicates with SEs ($P \leq 0.05$).

endogenous levels in the seeds at various times. In contrast, patterns of secreted and endogenous ABA level for *tss1* were similar to those of the wild type.

Endogenous levels of PA, DPA and ABA-GE in imbibed seeds

ABA metabolites from barley are low in liquid medium, making identification difficult (Jacobsen *et al.*, 2002). However, in tomato seeds, ABA metabolites in imbibed seeds were much higher than ABA, and their profiles differed among the genotypes.

In wild-type tomato seeds, DPA was the major metabolite, increasing significantly from 2006 up to 7370 pmol(g DW)⁻¹ during the time course of imbibition. PA was very low at 12 h and 48 h, and declined to near zero at 72 h. ABA-GE was not detectable at any imbibition time (Fig. 4A).

Imbibed *tss1* seeds had the lowest levels of ABA metabolites among the four genotypes. DPA was again the major metabolite, with a maximal value

[1326 pmol(g DW)⁻¹] at 48 h. PA reached its maximum [837 pmol(g DW)⁻¹] at 72 h. ABA-GE was detectable only at 48 and 72 h (Fig. 4B).

In *tss2*, DPA was again the major metabolite, and was unusual in showing its maximal level [8407 pmol(g DW)⁻¹] at 12 h, after which it gradually decreased. PA and ABA-GE were minor metabolites (Fig. 4C).

In *tos1* DPA was also the major metabolite, showing high levels at 12 and 48 h and low levels at 24 and 72 h, coincident with increased PA. The *tos1* mutant was the only genotype producing high levels of ABA-GE, which was notable at 24 h and maximal [7850 pmol(g DW)⁻¹] at 72 h (Fig. 4D).

Discussion

The different basal levels of ABA and metabolites may be correlated with differential germination capacities between the wild type and the several mutant




Figure 4. Abscisic acid (ABA) metabolite content [phaseic acid (PA), dihydrophaseic acid (DPA) and ABA glucose ester (ABA-GE)] of imbibed tomato seeds: (A) cv. Moneymaker (wild type); (B) *tss1* mutant; (C) *tss2* mutant; (D) *tos1* mutant. Data are means of three replicates with SEs ($P \leq 0.05$).

genotypes. ABA content in dry seeds was always higher than for imbibed seeds. For example, ABA in wild-type and *tss1* dry seeds were 67- and 50-fold higher, respectively, than after 48 h of imbibition. This suggests that ABA is synthesized or accumulated during seed formation. ABA present in mature seeds is generally considered to be a remnant of the ABA pools that are involved in the regulation of seed development (Hilhorst, 1995). Some studies indicated that the osmotic environment experienced by the seeds during development, more than ABA content, determines whether tomato seeds progress into events associated with germination or remain arrested at the completion of development (Berry and Bewley, 1992; Groot and Karssen, 1992). Jacobsen *et al.* (2002) have also failed to demonstrate a consistent relationship between ABA content of mature barley dry grains and germinability.

During imbibition, all tomato genotypes showed an abrupt decrease in ABA compared to dry seeds. Further, the ABA decline was accompanied by a corresponding increase in ABA metabolites, mainly DPA and ABA-GE, and there were qualitative and quantitative differences among the four genotypes. In contrast, for imbibed seeds of *Arabidopsis thaliana* mutant *etr1-2* (mutation that confers ethylene insensitivity and affects the ABA metabolic pathway), the decline in ABA was not accompanied by an increase in any of the ABA catabolites monitored (Chiwocha *et al.*, 2005).

The delayed germination of wild-type seeds may be due to not only high basal ABA content, but also newly synthesized ABA during imbibition, since ABA levels increased sharply up to 48 h, before declining at 72 h when visible germination was present. These findings are consistent with those of Grappin *et al.* (2000), who showed that some of the endogenous ABA detected in *Nicotiana plumbaginifolia* seeds resulted from *de novo* synthesis during early imbibition. In non-dormant and dormant *Arabidopsis* seeds, there was a decrease of ABA during early imbibition, although the rate of decrease was slower in the dormant seeds. Three days after sowing, the ABA content in dormant seeds increased and then remained high. Therefore, dormant and non-dormant seeds differ in their ability to neosynthesize ABA following imbibition (Ali-Rachedi *et al.*, 2004).

Dry and imbibed seeds of the *tss1* mutant had ABA contents similar to that of the wild type. Amounts of ABA and its metabolism may have a reduced influence on germination of this mutant, considering its faster germination at 48 h relative to the other genotypes. Valpuesta and Botella (2007) suggested that *TSS1* may encode for a Ca^{2+} sensor. A member of the *Arabidopsis calmodulin-like* (CML) gene family, CML24, was recently reported to act downstream from ABA perception, perhaps mediating cellular

responses to ABA-induced Ca^{2+} fluctuations, thereby delaying germination and seedling growth (Delk *et al.*, 2005). In the present study, *tss1* exhibited high endogenous ABA during imbibition, but germinated earlier than the wild type. Thus, the low sensitivity of *tss1* to endogenous ABA could be due to the mutation affecting a Ca^{2+} sensor similar to CML24, perhaps resulting in reduced intracellular signalling.

While most varieties of mature tomato seeds are generally non-dormant, the ABA-deficient *sitiens* (*sit^w*) tomato mutant seeds have been observed to germinate about a day earlier than wild-type seeds (Downie *et al.*, 1999). In this work, the highest germination percentages obtained with *tss2* seeds were associated with low ABA content in dry and imbibed seeds, suggesting that this mutant has an active ABA catabolism, and perhaps a low rate of ABA biosynthesis. The *tss2* mutant is hypersensitive to exogenous ABA, and the *TSS2* locus has been suggested to encode a protein that negatively regulates ABA signalling (Borsani *et al.*, 2001). A similar pattern of ABA biosynthesis was obtained with *tos1* imbibed seeds; supporting the suggestion of Borsani *et al.* (2002) that *tos1* is not an ABA-deficient mutant.

While Jacobsen *et al.* (2002) observed leakage of ABA and PA from barley seeds into the imbibition liquid, major decreases in ABA content in the seeds were not reflected by increased ABA or PA in the liquid. In our experiments, ABA diffused readily from *tos1* seeds into the liquid medium, and its level was twofold higher than that in *tss1* or *tss2*. The *tos1* mutant (characterized by sensitivity to osmotic stress) could have altered cellular membrane properties that result in an appreciable loss of ABA from the seed. That said, reduced ABA content in the seed of *tos1* had no effect on germination at 72 h, i.e. germination percentages for *tos1* were similar to those of the wild type, and lower than that of *tss1* or *tss2*. Borsani *et al.* (2002) proposed that the osmotic hypersensitivity of *tos1* is due to a defect at the intracellular level of the ABA-dependent signalling pathway, which is necessary for osmotic tolerance, and not to reduced seed ABA. In view of the high ABA leakage from *tos1* seeds into the medium, our data suggest that either extracellular ABA perception or signalling could also be altered in this mutant.

Millar *et al.* (2006) suggested that ABA-mediated physiological processes are correlated with fluctuating endogenous levels of the hormone, dynamically maintained by continual synthesis, transport and catabolism. The sharp decrease in ABA content following seed imbibition observed in the present study suggests activation of catabolic pathways, in agreement with previous reports (Jacobsen *et al.*, 2002; Kushiro *et al.*, 2004; Zhou *et al.*, 2004; Nambara and Marion-Poll, 2005). In barley and *Arabidopsis*, the

decrease of ABA content during seed imbibition is associated with increases in PA (Jacobsen *et al.*, 2002; Kushiro *et al.*, 2004) or PA metabolites.

Of interest here is the finding that rapid ABA catabolism during *Arabidopsis* seed imbibition is associated with increased mRNA levels of *CYP707A2*, a member of the *CYP707A* gene family that encodes ABA 8'-hydroxylases (Kushiro *et al.*, 2004). This increase in *AtCYP707A2* expression was also associated with the rapid decline of ABA in germinating seeds. More recently, Okamoto *et al.* (2006) reported that *CYP707A1* and *CYP707A2* are complementary and necessary for seed dormancy and germination control in *Arabidopsis*.

We identified PA and DPA in dry and imbibed wild-type and mutant seeds by GC-MS-SIM, confirming that 8'-hydroxylation of ABA is the main catabolic route. In imbibed wild-type seeds, all of the PA was metabolized to DPA, with DPA rising throughout the imbibition time course. Reduced ABA and higher DPA were observed in *tss2* (sensitive to ionic and osmotic stress) and in *tos1* (which is sensitive to osmotic stress), relative to *tss1*. These results suggest, for *tss2* and *tos1*, that the correlation between sensitivity to osmotic stress and endogenous ABA/DPA differs from that of *tss1*. In *tss1*, which is hypersensitive to ionic stress but not to general osmotic stress, the 8'-hydroxylation pathway was almost inactive, yielding the lowest PA and DPA values among the studied genotypes. It should be noted that the earlier germination of *tss1* was coincident with its low ABA catabolism, which is in agreement with the findings of Chiwocha *et al.* (2003), e.g. that 8'-hydroxylation and DPA formation may represent a minor pathway under conditions that promote germination.

DPA was the main ABA metabolite in apple seeds and sweet cherry fruit and seeds, with levels higher than those of ABA, suggesting that DPA, via feedback inhibition, may help regulate ABA in seeds (Setha *et al.*, 2005). Our observation that PA is a minor compound in imbibed seeds may be explained by its rapid reduction to DPA, as was also seen for germinating seeds of *Brassica napus* (Zhou *et al.*, 2003). In the present study, only *tos1* seeds showed major changes in ABA metabolism during imbibition. The decline in DPA levels at 24 and 72 h may have resulted from further metabolism to its 4'-glucoside, which was not monitored in this study. Such a reaction has been reported in tissues of several species (Walton and Li, 1995).

The wild type was the only genotype in which ABA catabolism by glucosylation was not detected. In contrast, ABA-GE was the predominant ABA catabolite in lettuce seeds during germination (Chiwocha *et al.*, 2003), and the high ABA-GE in germinating wild-type *Arabidopsis* seeds suggests that this is the

major reason for decreased ABA accumulation during germination (Chiwocha *et al.*, 2005). Finally, for germinating maize kernels, Wang *et al.* (2002) found that conjugation to ABA-GE was not a major pathway for endogenous ABA catabolism. Thus, ABA catabolic pathways vary considerably among different plant species, or even in different tissues of the same species. While ABA-GE has been considered to be biologically inert, and thus not constitute a stored form (Dietz *et al.*, 2000), Sauter *et al.* (2002) provided evidence that ABA-GE contributes to ABA homeostasis. Adding further support for a role of ABA-GE in homeostasis is the finding that the β -glucosidase homologue *AtBG1* generates ABA from ABA-GE (Lee *et al.*, 2006).

In our study increased ABA-GE was measured in *tos1* seeds from 24 to 72 h of imbibition. This indicates that the ABA glucosylation pathway is active in *tos1* and, together with an active 8'-hydroxylation pathway, likely accounts for the very low level of free ABA during germination. Interestingly, at 72 h *tos1* showed the same germination response as the wild type, even though their pattern of ABA metabolism was very different. An active glucosylation pathway in *tos1* seeds may be a direct response to the mutation for this osmotic stress-sensitive mutant.

In conclusion, wild-type and mutant seeds exhibited differing ABA contents and different ABA metabolism patterns that may be related to their differential sensitivity to abiotic stress. ABA metabolism differs in dry and imbibed seeds of wild-type tomato cv. MoneyMaker and the *tss1*, *tss2* and *tos1* mutants, and metabolism to DPA is the most important pathway for ABA catabolism.

Acknowledgements

We thank Dr Miguel A. Botella (University of Málaga, Spain) for kindly providing the tomato seeds used in this study. This work was supported by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), SECYT-UNRC and UNRC-ANPCYT grants awarded to G.A. and fellowships from CONICET to A.A.

References

- Ali-Rachedi, S., Bouinot, D., Wagner, M-H., Bonnet, M., Sotta, B., Grappin, P. and Jullien, M. (2004) Changes in endogenous abscisic acid levels during dormancy release and maintenance of mature seeds: studies with the Cape Verde Islands ecotype, the dormant model of *Arabidopsis thaliana*. *Planta* **219**, 479–488.
- Berry, T. and Bewley, D. (1992) A role for the surrounding fruit tissues in preventing the germination of tomato

- (*Lycopersicon esculentum*) seeds. *Plant Physiology* **100**, 951–957.
- Borsani, O., Cuartero, J., Fernández, J.A., Valpuesta, V. and Botella, M.A.** (2001) Identification of two loci in tomato reveals distinct mechanisms for salt tolerance. *Plant Cell* **13**, 873–888.
- Borsani, O., Cuartero, J., Fernández, J.A., Valpuesta, V. and Botella, M.A.** (2002) Tomato *tos1* mutation identifies a gene essential for osmotic tolerance and abscisic acid sensitivity. *Plant Journal* **32**, 905–914.
- Chiwocha, S., Abrams, S., Ambrose, S., Cutler, A., Loewen, A., Ross, A. and Kermodé, A.** (2003) A method for profiling classes of plant hormones and their metabolites using liquid chromatography–electrospray ionization tandem mass spectrometry: an analysis of hormone regulation of thermodormancy of lettuce (*Lactuca sativa* L.) seeds. *Plant Journal* **35**, 405–417.
- Chiwocha, S., Cutler, A., Abrams, S.R., Ambrose, S.J., Yang, J., Ross, A.R.S. and Kermodé, A.R.** (2005) The *etr1-2* mutation in *Arabidopsis thaliana* affects the abscisic acid, auxin, cytokinin and gibberellin metabolic pathways during maintenance of seed dormancy, moist-chilling and germination. *Plant Journal* **42**, 35–48.
- Cutler, A.J. and Krochko, J.E.** (1999) Formation and breakdown of ABA. *Trends in Plant Science* **4**, 472–478.
- Delk, N., Johnson, K., Chowdhury, N. and Braam, J.** (2005) *CLM24* regulated in expression by diverse stimuli, encodes a potential Ca^{2+} sensor that functions in responses to abscisic acid, day length, and ion stress. *Plant Physiology* **139**, 240–253.
- Dietz, K.-J., Sauter, A., Wichert, K., Messdaghi, D. and Hartung, W.** (2000) Extracellular β -glucosidase activity in barley involved in the hydrolysis of ABA glucose conjugate in leaves. *Journal of Experimental Botany* **51**, 937–944.
- Downie, B., Gurusinge, S. and Bradford, K.** (1999) Internal anatomy of individual tomato seeds: relationship to abscisic acid and germination physiology. *Seed Science Research* **9**, 117–128.
- Grappin, P., Bouinot, D., Sotta, B., Miginiac, E. and Jullien, M.** (2000) Control of seed dormancy in *Nicotiana glauca*: post-imbibition abscisic acid synthesis imposes dormancy maintenance. *Planta* **210**, 279–285.
- Groot, S.P.C. and Karszen, C.M.** (1992) Gibberellins regulate seed germination in tomato by endosperm weakening: a study with gibberellin-deficient mutants. *Planta* **171**, 525–531.
- Hilhorst, H.W.M.** (1995) A critical update on seed dormancy. I: primary dormancy. *Seed Science Research* **5**, 61–73.
- Jacobsen, J.V., Pearce, D.W., Poole, A.T., Pharis, R. and Mander, L.N.** (2002) Abscisic acid, phaseic acid and gibberellin contents associated with dormancy and germination in barley. *Physiologia Plantarum* **115**, 428–441.
- Kushiro, T., Okamoto, M., Nakabayashi, K., Yamagishi, K., Kitamura, S., Asami, T., Hirai, N., Koshiba, T., Kamiya, Y. and Nambara, E.** (2004) The *Arabidopsis* cytochrome P450 *CYP707A* encodes ABA 8'-hydroxylases: key enzymes in ABA catabolism. *EMBO Journal* **23**, 1647–1656.
- Lee, K.H., Piao, H.L., Kim, H.-Y., Choi, S.M., Jiang, F., Hartung, W., Hwang, I., Kwak, J.M., Lee, I.-J. and Hwang, I.** (2006) Activation of glucosidase via stress-induced polymerization rapidly increases active pools of abscisic acid. *Cell* **126**, 1109–1120.
- Lefebvre, V., North, H., Frey, A., Sotta, B., Seo, M., Okamoto, M., Nambara, E. and Marion-Poll, A.** (2006) Functional analysis of *Arabidopsis* *NCED6* and *NCED9* genes indicates that ABA synthesized in the endosperm is involved in the induction of seed dormancy. *Plant Journal* **45**, 309–319.
- Leung, J. and Giraudat, J.** (1998) Abscisic acid signal transduction. *Annual Review of Plant Physiology and Plant Molecular Biology* **49**, 199–222.
- Luna, M.V., Soriano, M.D., Bottini, R., Sheng, C. and Pharis, R.** (1993) Levels of endogenous gibberellins, abscisic acid, indol-3-acetic acid and naringenin during dormancy of peach flower buds. *Acta Horticulturae* **329**, 265–267.
- Millar, A., Jacobsen, J., Ross, J., Helliwell, C., Poole, A., Scofield, G., Reid, J. and Gubler, F.** (2006) Seed dormancy and ABA metabolism in *Arabidopsis* and barley: the role of ABA 8'-hydroxylase. *Plant Journal* **45**, 942–954.
- Moller, S.G. and Chua, N.H.** (1999) Interactions and intersections of plant signaling pathways. *Journal of Molecular Biology* **293**, 219–234.
- Nambara, E. and Marion-Poll, A.** (2005) Abscisic acid biosynthesis and catabolism. *Annual Review of Plant Physiology and Plant Molecular Biology* **56**, 165–185.
- Ni, B.-R. and Bradford, K.J.** (1993) Germination and dormancy of abscisic acid- and gibberellin-deficient mutant tomato (*Lycopersicon esculentum*) seeds. *Plant Physiology* **101**, 607–617.
- Okamoto, M., Kuwahara, A., Seo, M., Kushiro, T., Asami, T., Hirai, N., Kamiya, Y., Koshiba, T. and Nambara, E.** (2006) *CYP707A1* and *CYP707A2*, which encode abscisic acid 8'-hydroxylases, are indispensable for proper control of seed dormancy and germination in *Arabidopsis*. *Plant Physiology* **141**, 97–107.
- Sauter, A., Abrams, S.R. and Hartung, W.** (2002) Structural requirements of abscisic acid (ABA) and its impact on water flow during radial transport of ABA analogues through maize roots. *Plant Growth Regulation* **21**, 50–59.
- Setha, S., Kondo, S., Hirai, N. and Ohgashi, H.** (2005) Quantification of ABA and its metabolites in sweet cherries using deuterium-labeled internal standards. *Plant Growth Regulation* **45**, 183–188.
- Todoroki, Y., Hirai, N. and Ohgashi, H.** (2000) Analysis of isomerization process of 8'-hydroxyabscisic acid and its 3'-fluorinated analog in aqueous solutions. *Tetrahedron* **56**, 1649–1653.
- Valpuesta, V. and Botella, M.A.** (2007) Identificación de genes esenciales para la tolerancia a estrés hídrico y salino. Laboratorio de Bioquímica y Biotecnología Vegetal. Universidad de Málaga. España. Available at <http://www.bmbq.uma.es/lbbv/tomate.htm> (accessed April 2007.)
- Walton, D.C. and Li, Y.** (1995) Abscisic acid biosynthesis and metabolism. pp. 140–157 in Davies, P.J. (Ed.) *Plant hormones: physiology, biochemistry and molecular biology*. Norwell, MA, Kluwer.
- Wang, Z., Mambelli, S. and Setter, T.** (2002) Abscisic acid catabolism in maize kernels in response to water deficit

- at early endosperm development. *Annals of Botany* **90**, 623–630.
- Xu, Z.-J., Nakajima, M., Suzuki, Y. and Yamaguchi, I.** (2002) Cloning and characterization of the abscisic acid-specific glucosyltransferase gene from Adzuki bean seedlings. *Plant Physiology* **129**, 1285–1295.
- Zaharia, L.I., Walker-Simmons, M.K., Rodríguez, C.N. and Abrams, S.R.** (2005) Chemistry of abscisic acid, abscisic acid catabolites and analogs. *Journal of Plant Growth Regulation* **24**, 274–284.
- Zeevaart, J.A.D., Rock, C.D., Fantauzzo, F., Heath, T.G. and Gage, D.A.** (1991) Metabolism of abscisic acid and its physiological implications. pp. 39–52 in Davies, W.J.; Jones, H.G. (Eds) *Abscisic acid: physiology and biochemistry*. Oxford, BIOS Scientific.
- Zhou, R., Squires, T.M., Ambrose, S.J., Abrams, S.R., Ross, A.R.S. and Cutler, A.** (2003) Rapid extraction of abscisic acid and its metabolites for liquid chromatography–tandem mass spectrometry. *Journal of Chromatography A* **1010**, 75–85.
- Zhou, R., Cutler, A., Ambrose, S.J., Galka, M.M., Nelson, K.M., Squires, T.M., Loewen, M.K., Juadhav, A.S., Ross, A.R., Taylor, D.C. and Abrams, S.R.** (2004) A new abscisic acid catabolic pathway. *Plant Physiology* **134**, 361–369.
- Zhu, J.-K., Liu, J. and Xiong, L.** (1998) Genetic analysis of salt tolerance in *Arabidopsis thaliana*: evidence of a critical role for potassium nutrition. *Plant Cell* **10**, 1181–1192.

Received 3 April 2008

accepted after revision 19 September 2008

© 2008 Cambridge University Press