

Geometric approach to the Moore-Penrose inverse and the polar decomposition of perturbations by operator ideals

Eduardo Chiumiento and Pedro Massey

Abstract

We study the Moore-Penrose inverse of perturbations by a symmetrically-normed ideal of a closed range operator on a Hilbert space. We show that the notion of essential codimension of projections gives a characterization of subsets of such perturbations in which the Moore-Penrose inverse is continuous with respect to the metric induced by the operator ideal. These subsets are maximal satisfying the continuity property, and they carry the structure of real analytic Banach manifolds, which are acted upon transitively by the Banach-Lie group consisting of invertible operators associated with the ideal. This geometric construction allows us to prove that the Moore-Penrose inverse is indeed a real bianalytic map between infinite-dimensional manifolds. We use these results to study the polar decomposition of closed range operators from a similar geometric perspective. At this point we prove that operator monotone functions are real analytic in the norm of any symmetrically-normed ideal. Finally, we show that the maps defined by the operator modulus and the polar factor in the polar decomposition of closed range operators are real analytic fiber bundles.

2020 MSC: Primary 22E65, 58B10 - Secondary 47A55, 47B10

Keywords: Moore-Penrose inverse; polar decomposition; essential codimension; symmetrically-normed ideal; homogeneous space; real analytic map.

Contents

1	Introduction	1
2	Preliminaries	4
3	Moore-Penrose inverse	6
3.1	Continuity of the Moore-Penrose inverse	6
3.2	Geometric structure of maximal continuity sets	9
3.3	Real analyticity of the Moore-Penrose inverse	14
4	Polar decomposition	17
4.1	Real analyticity of operator monotone functional calculus	17
4.2	Geometric structure of positive perturbations	22
4.3	Operator modulus and polar factor as real analytic fiber bundles	24
5	Appendix	28

1 Introduction

Let \mathcal{H} be a separable complex infinite-dimensional Hilbert space, and let $\mathcal{B}(\mathcal{H})$ be the algebra of bounded linear operators on \mathcal{H} . We denote by $\mathcal{CR} \subset \mathcal{B}(\mathcal{H})$ the set of closed range operators. Let

\mathfrak{S} be a symmetrically-normed ideal on \mathcal{H} equipped with a norm $\|\cdot\|_{\mathfrak{S}}$. For a fixed $A \in \mathcal{CR}$, we consider the set of closed range operators that are perturbations of A by operators in the ideal \mathfrak{S} , namely

$$\mathcal{CR} \cap (A + \mathfrak{S}) = \{B \in \mathcal{CR} : B - A \in \mathfrak{S}\}.$$

A natural metric is defined by $d_{\mathfrak{S}}(B_1, B_2) = \|B_1 - B_2\|_{\mathfrak{S}}$, for $B_1, B_2 \in \mathcal{CR} \cap (A + \mathfrak{S})$. For $X \in \mathcal{B}(\mathcal{H})$, we write $X = V_X|X|$ for its (unique) polar decomposition, where V_X is a partial isometry with the same nullspace as X , usually known as the polar factor, and $|X| = (X^*X)^{1/2}$ is the operator modulus. Denote by \mathcal{CR}^+ the set of closed range positive operators and \mathcal{PI} the set of partial isometries on \mathcal{H} . In the present paper we introduce a geometric framework to study the continuity and real analyticity of the following maps:

$$\mu : \mathcal{CR} \cap (A + \mathfrak{S}) \rightarrow \mathcal{CR}, \mu(B) = B^\dagger \text{ (Moore-Penrose inverse);}$$

$$\alpha : \mathcal{CR} \cap (A + \mathfrak{S}) \rightarrow \mathcal{CR}^+, \alpha(B) = |B| \text{ (operator modulus);}$$

$$v : \mathcal{CR} \cap (A + \mathfrak{S}) \rightarrow \mathcal{PI}, v(B) = V_B \text{ (polar factor).}$$

Our geometric constructions to study these maps fit into the context of Banach manifolds related to operator theory; in particular, we deal with real analytic homogeneous spaces of Banach-Lie groups associated with operator ideals (see, e.g., [6, 10, 13, 12, 11, 26, 39, 40, 35]). The notion of essential codimension of a pair of projections [15], or in other words the Fredholm index of a pair of projections [1, 9], plays a crucial role throughout the present work. Remarkably, the essential codimension appears as a useful tool in a variety of problems in operator theory and geometry such as Kadison Pythagorean's Theorem [32], equivalence of quasi-free states [44], unitary equivalence of projections [9, 43], geodesics in the Grassmann manifold and Toeplitz kernels [2, 3] and restricted diagonalization [21, 37].

Previous related results. The Moore-Penrose inverse and the polar decomposition are ubiquitous in linear algebra, matrix analysis and operator theory. The continuity and differentiability of the Moore-Penrose inverse have been extensively discussed. The map defined by taking the Moore-Penrose inverse of complex matrices of size $d \geq 1$ is continuous at a matrix A when one restricts its domain to the set of all the matrices with rank equal to $\text{rank}(A)$. Notice that the product group $\mathcal{GL}(d) \times \mathcal{GL}(d)$ of invertible matrices of size $d \geq 1$ acts transitively on the set of matrices of (constant) rank equal to $\text{rank}(A)$ by $(G, K) \cdot A = GAK^{-1}$. Hence, the set of matrices of rank equal to $\text{rank}(A)$ turns out to be a connected set that admits a real analytic manifold structure. Furthermore, for any differentiable map of a real parameter taking values in the manifold of matrices with constant rank, the composition of this map with the Moore-Penrose inverse is also differentiable. These results have interesting consequences in the perturbation theory of matrices (see [28, 45, 47]).

In the infinite-dimensional case, Labrousse and Mbekhta [34] proved that the maps given by the Moore-Penrose inverse and the polar factor are continuous at $A \in \mathcal{CR} \subset \mathcal{B}(\mathcal{H})$ if and only if A is injective or surjective. Other works on the continuity of the Moore-Penrose inverse on Hilbert spaces deal with the reduced modulus minimum, or consider the projections onto the range or nullspace in place of the notion of rank [8, 18, 30]. More generally, we refer to [14, 33, 36] for extensions of this circle of ideas to Banach algebras. Recently, the stability of the Moore-Penrose invertibility under compact perturbations has been studied in [31].

The theory of Banach-Lie groups and their homogeneous spaces provide an interesting point of view for understanding several objects in operator theory. In this direction we mention three concrete motivations for our work. First, the differential geometry of generalized inverses investigated by Andruchow, Corach and Mbekhta [4]. Second, the study of several metrics on the set of closed range operators by Corach, Maestripietri and Mbekhta [24]. In particular, they introduced on \mathcal{CR} an action of the product group of invertible operators $\mathcal{GL}(\mathcal{H}) \times \mathcal{GL}(\mathcal{H})$, and they showed that for any $A \in \mathcal{CR}$ the orbits

$$\mathcal{O}(A) = \{GAK^{-1} : G, K \in \mathcal{GL}(\mathcal{H})\} \tag{1}$$

are topological homogeneous spaces. The metrics considered on \mathcal{CR} in this result are given by $d_R(B_1, B_2) = \|B_1 - B_2\| + \|P_{R(B_1)} - P_{R(B_2)}\|$, or $d_N(B_1, B_2) = \|B_1 - B_2\| + \|P_{N(B_1)} - P_{N(B_2)}\|$, for $B_1, B_2 \in \mathcal{CR}$, where the norm on each term is the operator norm. Here $P_{N(B_i)}$ and $P_{R(B_i)}$ denote the projections onto the nullspace and range of B_i , respectively. These metrics allow the construction of continuous local cross-sections for the action, defined in terms of the Moore-Penrose inverse. Third, the work on generalized inversion due to Beltiță, Goliński, Jakimowicz and Pelletier [12]. Recall that the invertible group of a Banach algebra is a manifold, which is actually an open set of the algebra, and the inversion map is complex analytic on this manifold by the holomorphic functional calculus. So the authors proposed to understand the Moore-Penrose inverse in Banach algebras as an inversion with some pathologies. This lead them to an application of the theory of Banach-Lie groupoids, with particular emphasis on the case of C^* -algebras.

The results of this paper. For a symmetrically-normed ideal \mathfrak{S} , we consider the Banach-Lie groups $\mathcal{G}\ell_{\mathfrak{S}} := \mathcal{G}\ell(\mathcal{H}) \cap (I + \mathfrak{S})$ and $\mathcal{U}_{\mathfrak{S}} := \mathcal{U}(\mathcal{H}) \cap (I + \mathfrak{S})$, where $\mathcal{G}\ell(\mathcal{H})$ and $\mathcal{U}(\mathcal{H})$ are the full invertible and unitary groups, respectively. We denote by $[P : Q]$ the essential codimension of two orthogonal projections P and Q . In this context, we consider perturbations of closed range operators \mathcal{CR} , positive closed range operators \mathcal{CR}^+ and partial isometries \mathcal{PI} by the symmetrically-normed ideal \mathfrak{S} . For fixed operators $A \in \mathcal{CR}$, $C \in \mathcal{CR}^+$ and $V \in \mathcal{PI}$, we will show that their set of perturbations can be decomposed as the following disjoint unions $\mathcal{CR} \cap (A + \mathfrak{S}) = \cup_{k \in \mathbb{J}_A} \mathcal{C}_k(A)$, $\mathcal{CR}^+ \cap (C + \mathfrak{S}) = \cup_{k \in \mathbb{J}_C} \mathcal{P}_k(C)$ and $\mathcal{PI} \cap (V + \mathfrak{S}) = \cup_{k \in \mathbb{J}_V} \mathcal{V}_k(V)$, where the sets on the unions are defined using the essential codimension as

$$\begin{aligned} \mathcal{C}_k(A) &:= \{B \in \mathcal{CR} \cap (A + \mathfrak{S}) : [P_{N(B)} : P_{N(A)}] = k\}; \\ \mathcal{P}_k(C) &:= \{D \in \mathcal{CR}^+ \cap (C + \mathfrak{S}) : [P_{N(D)} : P_{N(C)}] = k\}; \\ \mathcal{V}_k(V) &:= \{X \in \mathcal{PI} \cap (V + \mathfrak{S}) : [P_{N(X)} : P_{N(V)}] = k\}. \end{aligned}$$

The set of indices \mathbb{J}_A , \mathbb{J}_C and \mathbb{J}_V in the previous unions are always infinite subsets of \mathbb{Z} , and depend on the dimension of the nullspace, range and corange of the operators A , C and V .

The main results of this paper are the following:

- The sets $\mathcal{C}_k(A)$ and $\mathcal{P}_k(C)$ are Banach manifolds. Indeed, they admit the structure of real analytic homogeneous spaces, which are also submanifolds of natural affine spaces (Theorems 3.14 and 4.10).
- The map $\mu : \mathcal{C}_k(A) \rightarrow \mathcal{C}_k(A^\dagger)$, $\mu(B) = B^\dagger$, is a real bianalytic map between Banach manifolds (Theorem 3.18).
- The maps $\alpha : \mathcal{C}_k(A) \rightarrow \mathcal{P}_k(|A|)$, $\alpha(B) = |B|$, and $v : \mathcal{C}_k(A) \rightarrow \mathcal{V}_k(V_A)$, $v(B) = V_B$, are real analytic fiber bundles between Banach manifolds (Theorem 4.17).

Before proving these results, which hold for every $k \in \mathbb{J}_A$, we show that these maps are (well-defined and) continuous. Indeed, the choice of the sets of the form $\mathcal{C}_k(A)$ is not arbitrary, these are dense connected subsets of $\mathcal{CR} \cap (A + \mathfrak{S})$, which are maximal with respect to the continuity property of the Moore-Penrose inverse (Theorems 3.4 and 3.9). The sets $\mathcal{C}_k(A)$ can be roughly described as formed by those closed range operators that are perturbations of A by elements in \mathfrak{S} and have ‘the same rank with respect to A ’; these are *local* conditions induced by A and \mathfrak{S} (as opposed to the condition of merely having a fixed rank of fixed nullity). To the best of our knowledge, the above results on these three maps are also new in the context of finite-dimensional manifolds.

The Banach manifold structures of the sets $\mathcal{C}_k(A)$, $\mathcal{P}_k(C)$ and $\mathcal{V}_k(V)$ are not evident from their definitions. Each $\mathcal{C}_k(A)$ admits a transitive action of $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$ which consists of restricted versions of the orbits in (1). In particular, $\mathcal{C}_0(A)$ (the set containing A) has the following characterization:

$$\mathcal{C}_0(A) = \{GAK^{-1} : G, K \in \mathcal{G}\ell_{\mathfrak{S}}\}. \quad (2)$$

This fact is related to our previous work on restricted orbits of closed range operators [22]. Similar results for unitary orbits, associated to operator ideals, of normal operators have been recently obtained in [11]. In contrast to the larger orbits in (1), there is no need to introduce metrics such as d_R or d_N to construct continuous local cross sections for the map $\pi_0 : \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}} \rightarrow \mathcal{C}_0(A)$, $\pi_0(G, K) = GAK^{-1}$. Once these sections are constructed, we can further endow an orbit like (2) with the structure of real analytic homogeneous space that is also a submanifold of $A + \mathfrak{S}$. On the other hand, the Banach manifold structure of the sets $\mathcal{P}_k(C)$ is given in terms of congruence orbits of the group $\mathcal{G}\ell_{\mathfrak{S}}$ (Theorem 4.10). For instance when $k = 0$, we obtain

$$\mathcal{P}_0(C) = \{GCG^* : G \in \mathcal{G}\ell_{\mathfrak{S}}\}. \quad (3)$$

The motivation for considering the restricted orbits in (2) and (3) comes from previous work on partial isometries [19, 20], where the above defined sets $\mathcal{V}_k(V)$, $k \in \mathbb{J}_V$, were proved to be orbits of the product group $\mathcal{U}_{\mathfrak{S}} \times \mathcal{U}_{\mathfrak{S}}$. Finally, we observe that the real analyticity of the operator modulus depends on the real analyticity of the square root on the sets $\mathcal{P}_k(C)$. We present a more general statement for operator monotone functions in Corollary 4.12. This follows from Theorem 4.6, which in turn is based on earlier work of Ando and van Hemmen on perturbations by symmetrically-normed ideals [7].

The paper is organized as follows. Section 2 is devoted to notation and preliminary results. In Section 3 we introduce several geometric structures and prove the main results on the Moore-Penrose inverse. In Section 4 we establish the results on operator monotone functions, the operator modulus and the polar factor.

2 Preliminaries

Let \mathcal{H} be an infinite-dimensional (complex separable) Hilbert space, and let $\mathcal{B}(\mathcal{H})$ be the algebra of bounded operators on \mathcal{H} . Given $A \in \mathcal{B}(\mathcal{H})$ we write $N(A)$ and $R(A)$ for the nullspace and range of A , respectively. The orthogonal projection onto a closed subspace \mathcal{S} is denoted by $P_{\mathcal{S}}$.

Moore-Penrose inverse and polar decomposition. The set of all closed range operators on \mathcal{H} is given by

$$\mathcal{CR} = \{A \in \mathcal{B}(\mathcal{H}) : R(A) \text{ is a closed subspace}\}.$$

An operator $B \in \mathcal{B}(\mathcal{H})$ is said to be the *Moore-Penrose inverse* of $A \in \mathcal{B}(\mathcal{H})$ if it satisfies that $ABA = A$, $BAB = B$, $(AB)^* = AB$ and $(BA)^* = BA$. If the Moore-Penrose exists, then it is uniquely determined, and we denote it by $B = A^\dagger$. It is not difficult to check that $A \in \mathcal{B}(\mathcal{H})$ admits a Moore-Penrose inverse if and only if $A \in \mathcal{CR}$. Furthermore, $AA^\dagger = P_{R(A)}$ and $A^\dagger A = P_{N(A)^\perp}$, whenever $A \in \mathcal{CR}$. The following useful identity was proved by Wedin [47]:

$$A^\dagger - B^\dagger = -A^\dagger(A - B)B^\dagger + (A^*A)^\dagger(A^* - B^*)(I - BB^\dagger) + (I - A^\dagger A)(A^* - B^*)(BB^*)^\dagger. \quad (4)$$

For $A \in \mathcal{B}(\mathcal{H})$, $A \neq 0$, the reduced minimum modulus is given by $\gamma(A) = \min_{\lambda \in \sigma(|A|) \setminus \{0\}} \lambda$, where $\sigma(|A|)$ is the spectrum of $|A|$. Equivalently, $\gamma(A) = \inf\{\|Af\| : f \in N(A)^\perp, \|f\| = 1\}$. Recall that for $A \in \mathcal{B}(\mathcal{H})$, we have $A \in \mathcal{CR}$ if and only if $\gamma(A) > 0$. In such case, $\gamma(A) = \|A^\dagger\|^{-1}$, where $\|\cdot\|$ denotes the operator norm. Also, it holds $\gamma(A) = \gamma(|A|) = \gamma(|A^*|) = \gamma(A^*)$, which in particular gives that A^* , $|A|$ and $|A^*|$ have closed range if A has closed range.

An operator $X \in \mathcal{B}(\mathcal{H})$ is a partial isometry if $\|Xf\| = \|f\|$, for all $f \in N(X)^\perp$. This is equivalent to having that XX^* is an (orthogonal) projection, or X^*X is an (orthogonal) projection. We write

$$\mathcal{PI} = \{X \in \mathcal{B}(\mathcal{H}) : X \text{ is partial isometry}\}.$$

The *polar decomposition* of an operator $A \in \mathcal{B}(\mathcal{H})$ is the factorization $A = V_A|A|$, where $|A| = (A^*A)^{1/2}$ is the *operator modulus* and $V_A \in \mathcal{PI}$ is the unique partial isometry that further satisfies

the condition $N(V_A) = N(A)$. In the case where $A \in \mathcal{CR}$, we observe that $|A| \in \mathcal{CR}$ is such that $V_A = A|A|^\dagger$, and we call V_A the *polar factor*.

Symmetrically-normed ideals. We follow the classical book [38] (see [27, 42]). A *symmetrically-normed ideal* is a two-sided ideal $\mathfrak{S} \subseteq \mathcal{B}(\mathcal{H})$ endowed with a norm $\|\cdot\|_{\mathfrak{S}}$ satisfying $\|ABC\|_{\mathfrak{S}} \leq \|A\|\|B\|_{\mathfrak{S}}\|C\|$, for all $A, C \in \mathcal{B}(\mathcal{H})$ and $B \in \mathfrak{S}$; and $\|B\|_{\mathfrak{S}} = \|B\|$, for every rank-one operator B . We also assume that $(\mathfrak{S}, \|\cdot\|_{\mathfrak{S}})$ is a Banach space. The previous facts imply that $\|B\| \leq \|B\|_{\mathfrak{S}}$, for all $B \in \mathfrak{S}$ and that $\|B_1 B_2\|_{\mathfrak{S}} \leq \|B_1\|_{\mathfrak{S}}\|B_2\|_{\mathfrak{S}}$, for $B_1, B_2 \in \mathfrak{S}$, i.e. the norm of the ideal is submultiplicative. Recall that $\mathcal{F} \subseteq \mathfrak{S} \subseteq \mathcal{K}$, for every symmetrically-normed ideal, where $\mathcal{F} = \mathcal{F}(\mathcal{H})$ is the ideal of finite-rank operators and $\mathcal{K} = \mathcal{K}(\mathcal{H})$ is the ideal of compact operators. The p -Schatten \mathfrak{S}_p ($1 \leq p \leq \infty$) are well-known examples of symmetrically-normed ideals, whose norms are given by $\|A\|_p = \text{Tr}(|A|^p)^{1/p} = (\sum_{n \geq 1} s_n^p(A))^{1/p}$, $p \geq 1$; and for $p = \infty$, $\mathfrak{S}_\infty = \mathcal{K}$ endowed with the usual operator norm $\|A\|_\infty = \|A\| = s_1(A)$. Here $s(A) = \{s_n(A)\}_{n \geq 1}$ is the sequence of singular values of A arranged in non-increasing order and counting multiplicities. Other examples of symmetrically-normed ideals can be found in the aforementioned references.

Essential codimension. Next we recall the notion of essential codimension (see [1, 9, 15]). Let $P, Q \in \mathcal{B}(\mathcal{H})$ be two orthogonal projections such that the operator $QP|_{R(P)} : R(P) \rightarrow R(Q)$ is Fredholm. In this case, (P, Q) is known as a Fredholm pair and the index of this Fredholm operator

$$\begin{aligned} [P : Q] &:= \text{Ind}(QP|_{R(P)} : R(P) \rightarrow R(Q)) \\ &= \dim(N(Q) \cap R(P)) - \dim(R(Q) \cap N(P)) \end{aligned}$$

is called the *essential codimension* (or *Fredholm index of the pair*). We will often have two projections such that $P - Q \in \mathcal{K}$. In such a case, it is easy to see that (P, Q) is a Fredholm pair, and the essential codimension is well-defined. Among some elementary properties of the essential codimension that we will use frequently are the following: $[P : Q] = -[Q : P]$; if (P_i, Q_i) , $i = 1, 2$, are two Fredholm pairs such that $P_1 P_2 = 0$ and $Q_1 Q_2 = 0$, then $(P_1 + P_2, Q_1 + Q_2)$ is a Fredholm pair and $[P_1 + P_2 : Q_1 + Q_2] = [P_1 : Q_1] + [P_2 : Q_2]$; and if (P, Q) and (Q, R) are Fredholm pairs, and either $Q - R \in \mathcal{K}$ or $P - Q \in \mathcal{K}$, then (P, R) is a Fredholm pair and $[P : R] = [P : Q] + [Q : R]$.

Banach manifolds. We consider real analytic manifolds modeled on Banach spaces (see [10, 46]). Given M, N manifolds and a real analytic map $f : M \rightarrow N$, we denote by $T_p f : (TM)_p \rightarrow (TN)_{f(p)}$ the tangent map at $p \in M$, where $(TM)_p$ and $(TN)_{f(p)}$ are the tangent spaces of M at p and N at $f(p)$. A bijective map $f : M \rightarrow N$ is real bianalytic if f and f^{-1} are real analytic. A real analytic map $f : M \rightarrow N$ is called a submersion at $p \in M$ if $N(T_p f)$ is a closed complemented subspace of $(TN)_{f(p)}$ and $T_p f$ is surjective. If $f : M \rightarrow N$ is a submersion at every point $p \in M$, then f is called a submersion. A real Banach-Lie group is a real analytic Banach manifold G such that the group multiplication $G \times G \rightarrow G$, $(g, h) \mapsto gh$, and the inverse $G \rightarrow G$, $g \mapsto g^{-1}$, are real analytic maps. The construction of the Lie algebra $\mathfrak{g} \simeq (TG)_1$ and the exponential map $\exp_G : \mathfrak{g} \rightarrow G$ of a Banach-Lie group G can be carried out similarly to the case of finite-dimensional Lie groups. Also the exponential map $\exp_G : \mathfrak{g} \rightarrow G$ is a local bianalytic map. An action of a Banach-Lie group G on a manifold M is a map $L : G \times M \rightarrow M$, $L(g, p) = g \cdot p$, $g \in G$ and $p \in M$, such that $h \cdot (g \cdot p) = (hg) \cdot p$ and $1 \cdot p = p$, for all $h, g \in G$ and $p \in M$. The action is said to be real analytic if the map L is real analytic. A *real analytic homogeneous space of a Banach-Lie group* G is a manifold M such that G acts transitively and analytically on M , and there exists $p \in M$ such that the map $\pi_p : G \rightarrow M$, $\pi_p(g) = g \cdot p$, is a submersion.

Let M be a manifold, and $N \subseteq M$. A chart (ϕ, \mathcal{V}, E) at $p \in M$ consists in an open neighborhood \mathcal{V} of p , a Banach space E and a homeomorphism $\phi : \mathcal{V} \rightarrow \phi(\mathcal{V}) \subseteq E$. If for every $p \in N$ there exists a chart (ϕ, \mathcal{V}, E) at p , and a closed subspace F complemented in E satisfying $\phi(\mathcal{V} \cap N) = F \cap \phi(\mathcal{V})$, then N is called a *submanifold* of M . In this case, N turns out to be a manifold endowed with the topology inherited from M . If H is a subgroup of a Banach-Lie group G , then H is said to be a *Banach-Lie subgroup* of G when H is a submanifold of G .

Let M and N be two manifolds. A *real analytic fiber bundle* is a real analytic surjective map $f : M \rightarrow N$ such that for every $p \in N$ then $f^{-1}(p)$ is a manifold, and there exists an open neighborhood \mathcal{V} of p and a real bianalytic map $\Psi : f^{-1}(\mathcal{V}) \rightarrow \mathcal{V} \times f^{-1}(p)$ such that $\pi_1 \circ \Psi = f$, where $\pi_1 : \mathcal{V} \times f^{-1}(p) \rightarrow \mathcal{V}$ is the canonical projection. In such case, f turns out to be a submersion.

Restricted groups. Let $\mathcal{G}l(\mathcal{H})$ be the group of invertible operators on \mathcal{H} . For each symmetrically-normed ideal \mathfrak{S} there is associated the following group

$$\mathcal{G}l_{\mathfrak{S}} := \{G \in \mathcal{G}l(\mathcal{H}) : G - I \in \mathfrak{S}\}.$$

Also each symmetrically-normed ideal \mathfrak{S} gives raise to a subgroup of the full unitary group $\mathcal{U}(\mathcal{H})$ defined by

$$\mathcal{U}_{\mathfrak{S}} := \{U \in \mathcal{U}(\mathcal{H}) : U - I \in \mathfrak{S}\}.$$

For a standard reference for these groups in the case of the p -Schatten ideals see [26], meanwhile for the case of general symmetrically-normed ideals see [10].

Remark 2.1. We collect here several properties of the groups defined above. In what follows we let \mathfrak{S} denote a symmetrically-normed operator ideal.

i) If P, Q are orthogonal projections, then there is a unitary operator $U \in \mathcal{U}_{\mathfrak{S}}$ such that $Q = UPU^*$ if and only if $P - Q \in \mathfrak{S}$ and $[P : Q] = 0$ (see [17, Prop. 3.6], or more generally, [37, Prop 2.3]).

ii) $\mathcal{G}l_{\mathfrak{S}}$ is a real Banach-Lie group endowed with the metric $d_{\mathfrak{S}}(G, K) = \|G - K\|_{\mathfrak{S}}$, for $G, K \in \mathcal{G}l_{\mathfrak{S}}$, whose Lie algebra is \mathfrak{S} . Next, consider the unitalization $\tilde{\mathfrak{S}} = \{X + \lambda I : X \in \mathfrak{S}, \lambda \in \mathbb{C}\} \simeq \mathfrak{S} \oplus \mathbb{C}$. Each element $Z \in \tilde{\mathfrak{S}}$ is written as $Z = X + \lambda I$, for uniquely determined $X \in \mathfrak{S}$ and $\lambda \in \mathbb{C}$, and $\tilde{\mathfrak{S}}$ is equipped with the norm $\|X + \lambda I\|_{\tilde{\mathfrak{S}}} := \|X\|_{\mathfrak{S}} + |\lambda|$. In this case, $\tilde{\mathfrak{S}}$ is a unital Banach algebra. We can embed into $\mathcal{G}l_{\mathfrak{S}}$ in $\tilde{\mathfrak{S}}$ by the identification $T \mapsto (T - I) + I \in \tilde{\mathfrak{S}}$. Then $\mathcal{G}l_{\mathfrak{S}}$ is a Lie subgroup of the Banach-Lie group of invertible elements of $\tilde{\mathfrak{S}}$ having real codimension 2. On the other hand, $\mathcal{U}_{\mathfrak{S}}$ is a Banach-Lie subgroup of $\mathcal{G}l_{\mathfrak{S}}$, whose Lie algebra is $\mathfrak{S}_{ah} = \{X \in \mathfrak{S} : X^* = -X\}$, the anti-hermitian operators in \mathfrak{S} ([10, Prop 9.28]). The exponential maps of these Lie groups are given by $\exp_{\mathcal{G}l_{\mathfrak{S}}} : \mathfrak{S} \rightarrow \mathcal{G}l_{\mathfrak{S}}$, $\exp_{\mathcal{G}l_{\mathfrak{S}}}(X) = e^X = \sum_{n \geq 0} \frac{X^n}{n!}$ and $\exp_{\mathcal{U}_{\mathfrak{S}}} = \exp_{\mathcal{G}l_{\mathfrak{S}}} |_{\mathfrak{S}_{ah}}$.

iii) The exponential map of $\mathcal{G}l_{\mathfrak{S}}$ is surjective. We give a proof since we do not find references to this fact. For $G \in \mathcal{G}l_{\mathfrak{S}}$, $G - I \in \mathfrak{S} \subseteq \mathcal{K}$ yields that $\sigma(G)$ is a countable set having 1 as its unique limit point. Thus, there is a ray L from the origin such that $\sigma(G) \subseteq \mathbb{C} \setminus L$, and the analytic logarithm $\log : \mathbb{C} \setminus L \rightarrow \{z : \theta - 2\pi < \arg(z) \leq \theta\}$ is well-defined, where $\theta \in [0, 2\pi)$ is defined by the ray L . Denote by $\sigma_{\tilde{\mathfrak{S}}}(G)$ the spectrum of G in the Banach algebra $\tilde{\mathfrak{S}}$. Observe that $\sigma_{\tilde{\mathfrak{S}}}(G) = \sigma(G)$, so we can use the analytic functional calculus in $\tilde{\mathfrak{S}}$ to get $e^{\log(G)} = G$. That is, $\log(G) = X + \lambda I \in \tilde{\mathfrak{S}}$, $X \in \mathfrak{S}$ and $\lambda \in \mathbb{C}$, satisfies $G = e^{X + \lambda I} = e^X e^{\lambda I}$. But $G = (e^X - I)e^{\lambda I} + e^{\lambda I}$, with $e^X - I \in \mathfrak{S}$. Hence, the uniqueness of writing in $\tilde{\mathfrak{S}}$ gives $e^{\lambda I} = 1$, so $G = e^X$.

3 Moore-Penrose inverse

We first study the continuity of the Moore-Penrose inverse. Then we prove that the maximal sets in which it is continuous admit the structure of Banach manifolds. This is achieved by using the theory of Banach-Lie groups and their homogeneous spaces. We conclude that the Moore-Penrose inverse is a real bianalytic map between Banach manifolds.

3.1 Continuity of the Moore-Penrose inverse

We recall an estimate for matrices with equal rank and some elementary facts obtained in [22].

Lemma 3.1 ([47]). *Suppose that A, B are matrices such that $\text{rank}(A) = \text{rank}(B)$ and $\|A - B\| < \|A^\dagger\|^{-1}$, then*

$$\|B^\dagger\| \leq \frac{\|A^\dagger\|}{1 - \|A^\dagger\|\|A - B\|}.$$

Lemma 3.2 ([22]). *Let \mathfrak{S} be a symmetrically-normed ideal and take $A, B \in \mathcal{CR}$ be such that $A - B \in \mathfrak{S}$. Then $A^\dagger - B^\dagger \in \mathfrak{S}$, $P_{R(A)} - P_{R(B)} \in \mathfrak{S}$ and $P_{N(A)} - P_{N(B)} \in \mathfrak{S}$.*

Proof. We include a short proof of this result for the convenience of the reader. From Wedin's formula in Eq. (4) we get $A^\dagger - B^\dagger \in \mathfrak{S}$. The other assertions follow by using that $P_{N(A)^\perp} = A^\dagger A$ and $P_{R(A)} = AA^\dagger$. \square

We now present a generalization of Wedin's estimate in Lemma 3.1 in terms of the essential codimension.

Proposition 3.3. *Consider operators $A, B \in \mathcal{CR}$ satisfying $\|A - B\| < \|A^\dagger\|^{-1}$, $A - B \in \mathcal{K}$ and $[P_{N(A)} : P_{N(B)}] = 0$. Then,*

$$\|B^\dagger\| \leq \frac{\|A^\dagger\|}{1 - \|A^\dagger\|\|A - B\|}.$$

Proof. Since $A^* - B^* \in \mathcal{K}$, then $P_{N(A)^\perp} - P_{N(B)^\perp} \in \mathcal{K}$ by Lemma 3.2. From Remark 2.1 *i*) applied to the ideal of compact operators, we know that there exists a unitary $L \in \mathcal{U}_{\mathcal{K}}$ such that $LP_{N(A)^\perp}L^* = P_{N(B)^\perp}$. Next pick $\{E_n\}_{n \geq 1}$ a sequence of finite-rank projections such that $E_n \leq P_{N(A)^\perp}$ and $E_n \nearrow P_{N(A)^\perp}$ strongly. We set $F_n = LE_nL^*$, $B_n = BF_n$ and $A_n = AE_n$, for all $n \geq 1$. Notice that $\text{rank}(A_n) = \text{rank}(B_n)$, and also $A_n - B_n \rightarrow A - B$ strongly. Further, we observe that

$$A_n - B_n = AE_n - BLE_nL^* = (A - B)E_n - B(L - I)E_nL^* - BE_n(L^* - I),$$

where each term is multiplied by a compact operator. Thus, we get $\|A_n - B_n - (A - B)\| \rightarrow 0$ by a well-known result (see, e.g. [38, Thm. 6.3]). On the other hand, since $A_n^*A_n = E_nA^*AE_n \geq \gamma(A)E_nP_{N(A)^\perp}E_n = \gamma(A)E_n$ and $N(A_n)^\perp = R(E_n)$, then $\|A_n^\dagger\|^{-1} = \gamma(A_n) \geq \gamma(A) = \|A^\dagger\|^{-1}$.

Therefore for large n , $\|A_n - B_n\| < \|A^\dagger\|^{-1} \leq \|A_n^\dagger\|^{-1}$, so we can apply Lemma 3.1 to obtain

$$\|B_n^\dagger\| \leq \frac{\|A_n^\dagger\|}{1 - \|A_n^\dagger\|\|A_n - B_n\|} \leq \frac{\|A^\dagger\|}{1 - \|A^\dagger\|\|A_n - B_n\|}. \quad (5)$$

We claim that B_n^\dagger converges strongly to B^\dagger . This follows by using the formula in Eq. (4), which implies that for $f \in \mathcal{H}$ one has

$$\begin{aligned} \|(B_n^\dagger - B^\dagger)f\| &\leq \|B_n^\dagger\|\|(B_n - B)B^\dagger f\| + \|(B_n^*B_n)^\dagger\|\|(B_n^* - B^*)(I - BB^\dagger)f\| + \\ &\quad + \|I - B_n^\dagger B_n\|\|(B_n^* - B^*)(BB^*)^\dagger f\|. \end{aligned}$$

Here note that $\|(B_n^*B_n)^\dagger\| = \|B_n^\dagger(B_n^\dagger)^*\| = \|B_n^\dagger\|^2 \leq \|B^\dagger\|^2$, for all n , by a similar argument as before with A_n and A . Also observe that $\|I - B_n^\dagger B_n\| = 1$ and $B_n^* = F_n B^*$ converges strongly to B^* since $F_n \nearrow P_{N(B)^\perp} = P_{R(B^*)}$. This proves our claim.

Consider $\epsilon > 0$, and take a vector $f \in \mathcal{H}$, $\|f\| = 1$, such that $\|B^\dagger\| \leq \|B^\dagger f\| + \epsilon$. Using that B_n^\dagger converges strongly to B^\dagger we have $\|B^\dagger f\| \leq \|B_n^\dagger f\| + \epsilon \leq \|B_n^\dagger\| + \epsilon$ for all n large enough. This gives

$$\|B^\dagger\| \leq \|B_n^\dagger\| + 2\epsilon \leq \frac{\|A^\dagger\|}{1 - \|A^\dagger\|\|A_n - B_n\|} + 2\epsilon.$$

Letting $n \rightarrow \infty$ and noting that $\epsilon > 0$ is arbitrary, we find the desired estimate. \square

Now we can give our main result on the continuity of the Moore-Penrose inverse.

Theorem 3.4. *Let \mathfrak{S} be a symmetrically-normed ideal. Let $\{B_n\}_{n \geq 1}$ be a sequence in \mathcal{CR} such that $B_n - B \in \mathfrak{S}$ and $\|B_n - B\|_{\mathfrak{S}} \rightarrow 0$, for some $B \in \mathcal{CR}$. The following conditions are equivalent:*

- i)* $[P_{N(B_n)} : P_{N(B)}] = 0$ for all sufficiently large n ;
- ii)* $\sup_n \|B_n^\dagger\| < \infty$;
- iii)* $\|B_n^\dagger - B^\dagger\|_{\mathfrak{S}} \rightarrow 0$;
- iv)* $\|P_{N(B_n)} - P_{N(B)}\|_{\mathfrak{S}} < 1$ for all sufficiently large n ;
- v)* $\|P_{N(B_n)} - P_{N(B)}\| < 1$ for all sufficiently large n ;
- vi)* $N(B_n)^\perp \cap N(B) = \{0\}$ for all sufficiently large n .

Proof. *i) \rightarrow ii)* First, notice that Lemma 3.2 implies that $P_{N(B)} - P_{N(B_n)} \in \mathfrak{S} \subset \mathcal{K}$, so that the essential codimension in the statement above is well defined. Suppose that $[P_{N(B_n)} : P_{N(B)}] = 0$ for large n . Since $\|B_n - B\| \leq \|B_n - B\|_{\mathfrak{S}} \rightarrow 0$, we derive from Proposition 3.3 that $\|B_n^\dagger\| \leq \frac{\|B^\dagger\|}{1 - \|B_n^\dagger\| \|B - B_n\|}$ for sufficiently large n . Hence $\sup_n \|B_n^\dagger\| < \infty$.

ii) \rightarrow iii) The Moore-Penrose inverse of an operator $A \in \mathcal{CR}$ satisfies $(A^*A)^\dagger = A^\dagger(A^*)^\dagger$ and $(A^*)^\dagger = (A^\dagger)^*$. Using these facts in the identity in Eq. (4), we get

$$\begin{aligned} \|B_n^\dagger - B^\dagger\|_{\mathfrak{S}} &\leq \|B_n\| \|B_n - B\|_{\mathfrak{S}} \|B\| + \|B_n^\dagger\|^2 \|B_n - B\|_{\mathfrak{S}} \|I - BB^\dagger\| \\ &\quad + \|I - B_n^\dagger B_n\| \|B_n - B\|_{\mathfrak{S}} \|B^\dagger\|^2. \end{aligned} \tag{6}$$

Notice that $\|I - B_n^\dagger B_n\| = \|P_{N(B_n)}\| = 1$, and $\sup_n \|B_n\| < \infty$ because $\|B_n - B\| \leq \|B_n - B\|_{\mathfrak{S}} \rightarrow 0$. Thus, the assumption $\sup_n \|B_n^\dagger\| < \infty$ implies that $\|B_n^\dagger - B^\dagger\|_{\mathfrak{S}} \rightarrow 0$.

iii) \rightarrow iv) Recall that $P_{N(B_n)} = I - B_n^\dagger B_n$ and $P_{N(B)} = I - B^\dagger B$. Then

$$\|P_{N(B_n)} - P_{N(B)}\|_{\mathfrak{S}} \leq \|(B_n^\dagger - B^\dagger)B_n\|_{\mathfrak{S}} + \|B^\dagger(B_n - B)\|_{\mathfrak{S}}.$$

Thus, $\|P_{N(B_n)} - P_{N(B)}\|_{\mathfrak{S}}$ becomes arbitrarily small for sufficiently large n because $\|B_n - B\|_{\mathfrak{S}} \rightarrow 0$ so $\sup_n \|B_n\|_{\mathfrak{S}} < \infty$, $\|B_n^\dagger - B^\dagger\|_{\mathfrak{S}} \rightarrow 0$ and the norm of the ideal is submultiplicative.

iv) \rightarrow v) This follows again by the estimate $\|P_{N(B_n)} - P_{N(B)}\| \leq \|P_{N(B_n)} - P_{N(B)}\|_{\mathfrak{S}}$.

v) \rightarrow vi) Straightforward.

vi) \rightarrow i) Notice that $\|B_n - B\| \leq \|B_n - B\|_{\mathfrak{S}} \rightarrow 0$, so it holds $N(B)^\perp \cap N(B_n) = \{0\}$ for large n . Indeed, if there is a unit vector $f_n \in N(B)^\perp \cap N(B_n)$ for infinitely many $n \geq 1$, we find that $0 < \gamma(B) \leq \|Bf_n\| = \|(B_n - B)f_n\| \rightarrow 0$, a contradiction. Hence $[P_{N(B_n)} : P_{N(B)}] = 0$ for all sufficiently large n . \square

Remark 3.5. We can take the operator adjoint and use elementary properties of the essential codimension to state other equivalent conditions. Indeed, we can replace conditions *i)*, *iv)*, *v)* and *vi)* by: *i')* $[P_{R(B)} : P_{R(B_n)}] = 0$; *iv')* $\|P_{R(B_n)} - P_{R(B)}\|_{\mathfrak{S}} < 1$; *v')* $\|P_{R(B_n)} - P_{R(B)}\| < 1$; and *vi')* $R(B_n) \cap R(B)^\perp = \{0\}$, for all sufficiently large n (in each case).

Remark 3.6. Among the equivalent conditions of Theorem 3.4, we have considered $\sup_n \|B_n\| < \infty$ in a self-contained exposition, using properties of the essential codimension (Proposition 3.3). We point out that the following results in the literature on the convergence of the Moore-Penrose inverse can be used to give alternative proofs.

i) In the infinite dimensional setting Izumino [30, Lemma 2.2] proved that for operators $B, B_n \in \mathcal{CR}$ such that $\|B - B_n\| < \|B^\dagger\|^{-1}$ and $\|BB^\dagger - B_n B_n^\dagger\| < 1$, then

$$\|B_n^\dagger\| \leq \frac{2\|B^\dagger\|}{1 - \|B - B_n\|\|B^\dagger\|}.$$

This was adapted by Koliha [33, Thm. 1.5] to the context of C^* -algebras.

ii) In a work by Chen, Wei and Sue on the perturbation of the Moore-Penrose inverse in the operator norm, they proved the following estimate ([18, Thm. 3.2]):

$$\|B_n^\dagger\| \leq \frac{\|B^\dagger\|}{1 - \frac{1}{2}(3 + \sqrt{5})\|B^\dagger\|\|B_n - B\|},$$

whenever $R(B_n) \cap R(B)^\perp = \{0\}$ and $\|B_n - B\| \leq \frac{3 - \sqrt{5}}{2\|B^\dagger\|}$.

3.2 Geometric structure of maximal continuity sets

For $A \in \mathcal{CR}$ we begin by considering the set of perturbations of the form

$$\mathcal{CR} \cap (A + \mathfrak{S}) = \{B \in \mathcal{CR} : B - A \in \mathfrak{S}\}.$$

This set is endowed with the metric $d_{\mathfrak{S}}(B_1, B_2) = \|B_1 - B_2\|_{\mathfrak{S}}$, for $B_1, B_2 \in \mathfrak{S}$. As we will see below, the essential codimension provides a decomposition of $\mathcal{CR} \cap (A + \mathfrak{S})$ in connected sets where the Moore-Penrose inverse has nice continuity properties. It is worth mentioning that the essential codimension was used to give a parametrization of the connected components of infinite-dimensional Grassmannians or Stiefel manifolds (see [17] and Remark 4.13).

Remark 3.7. The following facts, which are [22, Thm 3.5 and 3.7], will be useful. Given $A, B \in \mathcal{CR}$, such that $B - A \in \mathfrak{S}$, then

- i) There exists $G \in \mathcal{G}\ell_{\mathfrak{S}}$ such that $B = GA$ if and only if $N(B) = N(A)$.
- ii) There exist $G, K \in \mathcal{G}\ell_{\mathfrak{S}}$ such that $B = GAK^{-1}$ if and only if $[P_{N(B)} : P_{N(A)}] = 0$.

Notation 3.8. For a fixed $A \in \mathcal{CR}$, set $n_1 = \dim(N(A))$, $n_2 = \dim(N(A)^\perp)$, $n_3 = \dim(R(A)^\perp)$ and $\mathbb{J}_A = \{k \in \mathbb{Z} : -\min\{n_1, n_3\} \leq k \leq n_2\}$. When $n_2 = \infty$ we mean \mathbb{J}_A contains all the positive integers; meanwhile when $n_1 = n_3 = \infty$ we have \mathbb{J}_A contains all the negative integers.

Theorem 3.9. *Let $A \in \mathcal{CR}$ and let \mathfrak{S} be a symmetrically-normed ideal. Then $\mathcal{CR} \cap (A + \mathfrak{S})$ can be decomposed as the following disjoint union*

$$\mathcal{CR} \cap (A + \mathfrak{S}) = \bigcup_{k \in \mathbb{J}_A} \mathcal{C}_k(A),$$

where

$$\begin{aligned} \mathcal{C}_k(A) &= \{B \in \mathcal{CR} : B - A \in \mathfrak{S}, [P_{N(B)} : P_{N(A)}] = k\} \\ &= \{B \in \mathcal{CR} : B - A \in \mathfrak{S}, [P_{R(B)} : P_{R(A)}] = -k\}. \end{aligned}$$

The set \mathbb{J}_A is infinite and $\mathcal{C}_k(A) \neq \emptyset$, for each $k \in \mathbb{J}_A$. Furthermore, the following assertions hold:

- i) Given $B \in \mathcal{C}_k(A)$, the action $(G, K) \cdot B = GBK^{-1}$, $G, K \in \mathcal{G}\ell_{\mathfrak{S}}$, is well defined and transitive on $\mathcal{C}_k(A)$.
- ii) $\mathcal{C}_k(A)$ is connected.

iii) For $\ell, k \in \mathbb{J}_A$ and $B \in \mathcal{C}_\ell(A)$ there is a sequence $\{B_n\}_{n \geq 1}$ in $\mathcal{C}_k(A)$ such that $\|B_n - B\|_{\mathfrak{S}} \rightarrow 0$. In particular, $d_{\mathfrak{S}}(\mathcal{C}_k(A), \mathcal{C}_\ell(A)) := \inf\{\|B_1 - B_2\|_{\mathfrak{S}} : B_1 \in \mathcal{C}_k(A), B_2 \in \mathcal{C}_\ell(A)\} = 0$, and every $\mathcal{C}_k(A)$ is dense in $\mathcal{CR} \cap (A + \mathfrak{S})$.

iv) The map $\mu : \mathcal{C}_k(A) \rightarrow \mathcal{CR}$, $\mu(B) = B^\dagger$, is locally Lipschitz.

v) Let $\mathcal{C}_k(A) \subseteq \mathcal{C} \subseteq \mathcal{CR} \cap (A + \mathfrak{S})$ endowed with the metric $d_{\mathfrak{S}}$ be such that the map $\mu : \mathcal{C} \rightarrow \mathcal{CR}$, $\mu(B) = B^\dagger$ is continuous. Then, $\mathcal{C} = \mathcal{C}_k(A)$.

Proof. In the proof we write $\mathcal{C}_k := \mathcal{C}_k(A)$. For $B \in \mathcal{CR}$, $B - A \in \mathfrak{S}$, notice that $[P_{N(B)} : P_{N(A)}] = 0$ if and only if $[P_{R(B)} : P_{R(A)}] = 0$. This follows easily from Remark 3.7 by taking the operator adjoint and elementary properties of the essential codimension. Next suppose $[P_{N(B)} : P_{N(A)}] = k \neq 0$. If $k > 0$, then the operators defined on $\mathcal{H} \oplus \mathbb{C}^k$ by $\tilde{A} = A \oplus 0_k$ and $\tilde{B} = B \oplus I_k$ now satisfy $0 = [P_{N(\tilde{B})} : P_{N(\tilde{A})}]$. Hence by the previous case, $0 = [P_{R(\tilde{B})} : P_{R(\tilde{A})}] = k + [P_{R(B)} : P_{R(A)}]$. The case $k < 0$ follows from the property $[P_{N(B)} : P_{N(A)}] = -[P_{N(A)} : P_{N(B)}]$. This proves the equivalence between the two conditions defining the sets \mathcal{C}_k .

In the forthcoming inequalities we use similar conventions to those of Notation 3.8 for the cases $n_1 = n_3 = \infty$, and $n_2 = \infty$. According to the definition of the essential codimension, it follows that $-\min\{n_1, n_3\} \leq [P_{N(B)} : P_{N(A)}] = -[P_{R(B)} : P_{R(A)}] \leq n_2$, for $B \in \mathcal{CR} \cap (A + \mathfrak{S})$. Therefore $\mathcal{CR} \cap (A + \mathfrak{S})$ can be expressed as the disjoint union in the statement. Moreover, note that for $-\min\{n_1, n_3\} \leq k \leq n_2$, one can construct operators B such that $A - B \in \mathfrak{S}$ and $[P_{N(B)} : P_{N(A)}] = k$. For instance, when $n_1 \leq n_3$, there is partial isometry X_l such that $N(X_l)^\perp \subseteq N(A)$, $R(X_l) \subseteq R(A)^\perp$ and $\dim(N(X_l)^\perp) = l$, for $0 \leq l \leq n_1$. Set $B = A + X_l$, which gives $[P_{N(B)} : P_{N(A)}] = -l$. Now for $0 < l \leq n_2$, take a subspace $\mathcal{S} \subseteq N(A)^\perp$ such that $\dim(N(A)^\perp \ominus \mathcal{S}) = l$. For $B = AP_{\mathcal{S}}$, it holds that $[P_{N(B)} : P_{N(A)}] = l$. The case $n_1 > n_3$ can be treated similarly. Hence $\mathcal{C}_k \neq \emptyset$ for $k \in \mathbb{J}_A$. Furthermore, \mathbb{J}_A is an infinite set because the underlying Hilbert space \mathcal{H} is infinite dimensional.

i) Given $B \in \mathcal{C}_k$ and $G, K \in \mathcal{GL}_{\mathfrak{S}}$ then $B_1 = (G, K) \cdot B = GBK^{-1}$ is such that $B_1 - B \in \mathfrak{S}$, $P_{N(B)} - P_{N(B_1)} \in \mathfrak{S}$ and $[P_{N(B)} : P_{N(B_1)}] = 0$ (see Remark 3.7). Hence, $A - B_1 = A - B + B - B_1 \in \mathfrak{S}$ and by the properties of the essential codimension (see Section 2) we have that $[P_{N(B_1)} : P_{N(A)}] = [P_{N(B_1)} : P_{N(B)}] + [P_{N(B)} : P_{N(A)}] = k + 0$, so $B_1 \in \mathcal{C}_k$. On the other hand, given $B, B_2 \in \mathcal{C}_k$ then $B - B_2 = (B - A) + (A - B_2) \in \mathfrak{S}$ and $[P_{N(B)} : P_{N(B_2)}] = [P_{N(B)} : P_{N(A)}] + [P_{N(A)} : P_{N(B_2)}] = k + (-k) = 0$. Again, by Remark 3.7 we get that there exists $G, K \in \mathcal{GL}_{\mathfrak{S}}$ such that $(G, K) \cdot B = B_2$ and the action is transitive on \mathcal{C}_k .

ii) From the previous item, every $B \in \mathcal{C}_k$ is written as $B = GB^{(k)}K^{-1}$, for a fixed $B^{(k)} \in \mathcal{C}_k$ and $G, K \in \mathcal{GL}_{\mathfrak{S}}$. According to Remark 2.1 iii) there exist $X, Y \in \mathfrak{S}$ such that $e^X = G$ and $e^Y = K$. Then $\gamma : [0, 1] \rightarrow \mathcal{C}_k$ defined by $\gamma(t) = e^{tX} B^{(k)} e^{-tY}$ is continuous and $\gamma(0) = B^{(k)}$ and $\gamma(1) = B$.

iii) Take $B_1 \in \mathcal{C}_k$ and $B_2 \in \mathcal{C}_l$, and suppose $k > l$. Therefore, $[P_{N(B_2)} : P_{N(B_1)}] = k - l$, so that $\dim(N(B_2) \cap N(B_1)^\perp) \geq k - l$. Let $\mathcal{S} \subseteq N(B_2) \cap N(B_1)^\perp$ be a subspace of dimension $k - l$, and for $\epsilon > 0$ let $B_1^\epsilon = B_2 + \frac{\epsilon}{k-l} P_{\mathcal{S}}$. One can verify that $B_1 - B_1^\epsilon \in \mathfrak{S}$ and $[P_{N(B_1^\epsilon)} : P_{N(B_1)}] = 0$; hence $B_1^\epsilon \in \mathcal{C}_k$ for $\epsilon > 0$ (see Remark 3.7 and item i) above). Since the singular values satisfy $s_j(B_1^\epsilon - B_2) = \frac{\epsilon}{k-l}$ for $j = 1, \dots, k - l$, and $s_j(B_1^\epsilon - B_2) = 0$ for $j > k - l$, it follows that $\|B_1^\epsilon - B_2\|_{\mathfrak{S}} \leq \epsilon$.

iv) Since $\mathcal{C}_k(A) = \mathcal{C}_0(B^{(k)})$ we assume, without loss of generality, that $k = 0$ and prove that μ is locally Lipschitz in a neighborhood of A . Indeed, take the open ball $\mathcal{V} := \{B \in \mathcal{C}_0 : \|B - A\|_{\mathfrak{S}} < \frac{1}{2\|A^\dagger\|}\}$. For $B_1, B_2 \in \mathcal{V}$, we consider the same estimate as in (6), i.e.

$$\begin{aligned} \|B_2^\dagger - B_1^\dagger\|_{\mathfrak{S}} &\leq \|B_2\| \|B_2 - B_1\|_{\mathfrak{S}} \|B_1\| + \|B_2^\dagger\|^2 \|B_2 - B_1\|_{\mathfrak{S}} \|I - B_1 B_1^\dagger\| \\ &\quad + \|I - B_2^\dagger B_2\| \|B_2 - B_1\|_{\mathfrak{S}} \|B_1^\dagger\|^2. \end{aligned}$$

Since $B_i \in \mathcal{V}$, $i = 1, 2$, we have $\|B_i - A\| \leq \frac{1}{2\|A^\dagger\|}$, so that $\|B_i\| \leq \|A\| + \frac{1}{2\|A^\dagger\|}$. By Proposition 3.3 it follows that $\|B_i^\dagger\| \leq \frac{\|A^\dagger\|}{1 - \|A^\dagger\|\|A - B_i\|} \leq 2\|A^\dagger\|$. Therefore, we get

$$\|B_2^\dagger - B_1^\dagger\|_{\mathfrak{S}} \leq \left[\left(\|A\| + \frac{1}{2\|A^\dagger\|} \right)^2 + 8\|A^\dagger\|^2 \right] \|B_2 - B_1\|_{\mathfrak{S}}.$$

v) If $\mathcal{C} \neq \mathcal{C}_k$, then there exists some $l \neq k$ with $B \in \mathcal{C}_l \cap \mathcal{C}$. By item iii) there is a sequence $\{B_n\}_{n \geq 1}$ in \mathcal{C}_k such that $\|B_n - B\|_{\mathfrak{S}} \rightarrow 0$. But this contradicts the continuity of $\mu : \mathcal{C} \rightarrow \mathcal{CR}$ by Theorem 3.4. \square

Remark 3.10. Regarding the Lipschitz condition in the context of infinite-dimensional Hilbert spaces, we recall the following result. Put $\mathcal{R}_k = \{B \in \mathcal{B}(\mathcal{H}) : \gamma(B) \geq \frac{1}{k}\}$. Then $\|B_1^\dagger - B_2^\dagger\| \leq 3k^2\|B_1 - B_2\|$, for all $B_1, B_2 \in \mathcal{R}_k$ ([24, Lemma 3.10]).

The following result shows that in case a sequence in $\mathcal{C}_k(A)$ approaches A in the norm of \mathfrak{S} , then it can be modified in a controlled way so that the Moore-Penrose of the modified sequence converges to A^\dagger in the norm of \mathfrak{S} .

Corollary 3.11. *Let $\{B_n\}_{n \geq 1}$ be a sequence in $\mathcal{C}_k(A)$ for some $0 \neq k \in \mathbb{J}_A$ such that $\|A - B_n\|_{\mathfrak{S}} \rightarrow 0$. Then, there exists a sequence $\{C_n\}_{n \geq 1}$ such that $\|C_n\|_{\mathfrak{S}} \rightarrow 0$, $\text{rank}(C_n) = |k|$ for $n \geq 1$, and $\|A^\dagger - (B_n + C_n)^\dagger\|_{\mathfrak{S}} \rightarrow 0$.*

Proof. Assume that $k < 0$, so that $\dim(N(A) \cap N(B_n)^\perp) \geq -k$, and let $\mathcal{S}_n \subset N(A) \cap N(B_n)^\perp$ be such that $\dim \mathcal{S}_n = -k$. If we let P_n denote the orthogonal projection onto \mathcal{S}_n then, by construction, $\|B_n P_n\|_{\mathfrak{S}} = \|(A - B_n)P_n\|_{\mathfrak{S}} \leq \|A - B_n\|_{\mathfrak{S}} \rightarrow 0$. Moreover, we also get that $[P_{N(B_n - B_n P_n)} : P_{N(A)}] = 0$. Thus, if we let $C_n = -B_n P_n$ then $\|A - (B_n + C_n)\|_{\mathfrak{S}} \rightarrow 0$ and $[P_{N(B_n + C_n)} : P_{N(A)}] = 0$. By the continuity of the Moore-Penrose inverse in $\mathcal{C}_0(A)$, we get that $\|A^\dagger - (B_n + C_n)^\dagger\|_{\mathfrak{S}} \rightarrow 0$.

Assume now that $k > 0$, so that $\dim(N(B_n) \cap N(A)^\perp) \geq k$, and let $\mathcal{S}_n \subset N(B_n) \cap N(A)^\perp$ be such that $\dim \mathcal{S}_n = k$. If we let P_n denote the orthogonal projection onto \mathcal{S}_n then, by construction, $\|A P_n\|_{\mathfrak{S}} = \|(A - B_n)P_n\|_{\mathfrak{S}} \leq \|A - B_n\|_{\mathfrak{S}} \rightarrow 0$. Moreover, we also get that $[P_{N(B_n + A P_n)} : P_{N(A)}] = 0$. Thus, if we let $C_n = A P_n$ then $\|A - (B_n + C_n)\|_{\mathfrak{S}} \rightarrow 0$ and $[P_{N(B_n + C_n)} : P_{N(A)}] = 0$. Again, by the continuity of the Moore-Penrose inverse in $\mathcal{C}_0(A)$, we obtain that $\|A^\dagger - (B_n + C_n)^\dagger\|_{\mathfrak{S}} \rightarrow 0$. \square

Notice that item i) of Theorem 3.9 says that $\mathcal{C}_k(A)$ is an orbit by the action of the (product) restricted group $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$, for every $k \in \mathbb{J}_A$. In the sequel, we may fix any $B^{(k)} \in \mathcal{C}_k(A)$ and write

$$\begin{aligned} \mathcal{C}_k(A) &= \{B \in \mathcal{CR} : B - A \in \mathfrak{S}, [P_{N(B)} : P_{N(A)}] = k\} \\ &= \{GB^{(k)}K^{-1} : G, K \in \mathcal{G}\ell_{\mathfrak{S}}\}. \end{aligned}$$

In particular, we can take $B^{(0)} = A$. From item iv), the operators in the sets $\mathcal{C}_k(A)$ can be considered as perturbations of the fixed operators $B^{(k)}$. Moreover, item v) may be interpreted as saying that the set $\mathcal{C}_0(A)$ is a maximal subset of $\mathcal{CR} \cap (A + \mathfrak{S})$ containing A in which the Moore-Penrose inverse is continuous.

To further study the structure of $\mathcal{C}_k(A)$, we now introduce the maps

$$\pi_k : \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}} \rightarrow \mathcal{C}_k(A), \quad \pi_k(G, K) = GB^{(k)}K^{-1}.$$

Recall that we consider $\mathcal{C}_k(A)$ endowed with the topology induced by metric $d_{\mathfrak{S}}(B, C) = \|B - C\|_{\mathfrak{S}}$.

Lemma 3.12. *The map π_k admits continuous local cross sections.*

Proof. We may assume that $k = 0$. We show that π_0 has continuous local cross sections at A . The arguments can be adapted to other points $B \in \mathcal{C}_0(A)$, because $\mathcal{C}_0(A)$ is an orbit by the continuous action π_0 of the product group $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$. Thus, we need to find an open neighborhood \mathcal{W} of $A \in \mathcal{C}_0(A)$ and a continuous map $\sigma : \mathcal{W} \rightarrow \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$ such that $(\pi_A \circ \sigma)(B) = B$, for all $B \in \mathcal{W}$. Observe that as an immediate consequence of Theorem 3.4, there exists an open set $\mathcal{W} \subset \mathcal{C}_0(A)$, where the map $\sigma : \mathcal{W} \rightarrow \mathcal{B}(\mathcal{H}) \times \mathcal{B}(\mathcal{H})$ defined by

$$\sigma(B) = (BA^\dagger + (I - P_{R(B)})(I - P_{R(A)}), P_{N(B)}P_{N(A)} + (I - P_{N(B)})(I - P_{N(A)}))$$

is continuous and it takes values on $\mathcal{G}\ell(\mathcal{H}) \times \mathcal{G}\ell(\mathcal{H})$. We remark that this section σ was first defined in [4, Prop. 1.1] in the context of generalized inverses in C^* -algebras (see also [24, Prop. 5.7]). The fact that this is indeed a section follows analogously in our setting. Finally, we may rewrite the first coordinate as $\sigma_1(B) = B(A^\dagger - B^\dagger) + P_{R(A)}(P_{R(B)} - P_{R(A)}) + I$; while the second coordinate may be written as $\sigma_2(B) = P_{N(B)}(P_{N(A)} - P_{N(B)}) + (P_{N(B)} - P_{N(A)})P_{N(A)} + I$. From these expressions together with Lemma 3.2 we find that $\sigma_i(B) \in \mathcal{G}\ell_{\mathfrak{S}}$, $i = 1, 2$. \square

To establish the homogeneous space structure of \mathcal{C}_k induced by the action of $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$ we first consider the isotropy subgroup of the action at $A \in \mathcal{C}_0(A)$.

Lemma 3.13. *Let $A \in \mathcal{C}\mathcal{R}$ and let \mathfrak{S} be a symmetrically-normed ideal. Then*

$$\mathcal{G}_A = \{(G, K) \in \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}} : GA = AK\}$$

is a Banach-Lie subgroup of $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$.

Proof. We write $\mathcal{G} := \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$ and $\mathfrak{g} = \mathfrak{S} \times \mathfrak{S}$ for its Lie algebra. Notice that \mathcal{G}_A is a closed subgroup of \mathcal{G} . Then according to [46, Prop 8.12] we must show that the closed subalgebra

$$\begin{aligned} (T\mathcal{G}_A)_{(I,I)} = \mathfrak{g}_A &:= \{(X, Y) \in \mathfrak{g} : (e^{tX}, e^{tY}) \in \mathcal{G}_A \text{ for all } t \in \mathbb{R}\} \\ &= \{(X, Y) \in \mathfrak{g} : XA = AY\} \end{aligned}$$

is a closed complemented subspace of \mathfrak{g} , and for every neighborhood \mathcal{T} of $(0, 0) \in \mathfrak{g}_A$, $\exp_{\mathcal{G}}(\mathcal{T})$ is a neighborhood of $(I, I) \in \mathcal{G}_A$.

We put $P = P_{R(A)}$ and $Q = P_{N(A)^\perp}$. From $XA = AY$ we see that $XP = AYA^\dagger$, and thus, $P^\perp XP = P^\perp AYA^\dagger = 0$. Similarly, $QYQ^\perp = 0$ and $QYQ = A^\dagger XPA = A^\dagger PXP$. We may represent the elements $(X, Y) \in \mathfrak{h}$ as follows

$$X = \begin{pmatrix} X_{11} & X_{12} \\ 0 & X_{22} \end{pmatrix}, \quad Y = \begin{pmatrix} A^\dagger X_{11} A & 0 \\ Y_{21} & Y_{22} \end{pmatrix},$$

where the first and second matrix representations are with respect to the decompositions $\mathcal{H} = R(P) \oplus N(P)$ and $\mathcal{H} = R(Q) \oplus N(Q)$, respectively. Here we identify the operator entries as $X_{11} \in P\mathfrak{S}P$, $X_{12} \in P\mathfrak{S}P^\perp$ and $X_{22} \in P^\perp\mathfrak{S}P^\perp$, and similarly for the operator entries corresponding to Y with respect to Q . From the above representations, it is now clear that \mathfrak{g}_A is closed in \mathfrak{g} . Moreover, a closed supplement of \mathfrak{g}_A in \mathfrak{g} is given by

$$\mathfrak{m} = \left\{ \left(\begin{pmatrix} X_{11} & 0 \\ X_{21} & 0 \end{pmatrix}, \begin{pmatrix} 0 & Y_{12} \\ 0 & 0 \end{pmatrix} \right) : X_{11} \in P\mathfrak{S}P, X_{21} \in P^\perp\mathfrak{S}P, Y_{12} \in Q\mathfrak{S}Q^\perp \right\}.$$

For the property of the exponential map, it suffices to show that there exist two open neighborhoods \mathcal{V} and \mathcal{W} of $(0, 0) \in \mathfrak{g}$ and $(I, I) \in \mathcal{G}$, respectively, such that $\exp_{\mathcal{G}} : \mathcal{V} \rightarrow \mathcal{W}$ is bianalytic, and $\exp_{\mathcal{G}}(\mathcal{V} \cap \mathfrak{g}_A) = \mathcal{W} \cap \mathcal{G}_A$. The nontrivial inclusion here can be formulated as follows. Given $(G, K) \in \mathcal{W} \cap \mathcal{G}_A$, we have to show that $G = e^X$ and $K = e^Y$, for some $(X, Y) \in \mathcal{V} \cap \mathfrak{g}_A$. Notice

that $GA = AK$, yields $(G - I)^n A = A(K - I)^n$, for all $n \geq 0$. Thus, the logarithm can be defined by the usual series if we take \mathcal{W} small enough; we further consider $\mathcal{V} = \exp_{\mathcal{G}}^{-1}(\mathcal{W})$. Therefore,

$$\log(G)A = \left(\sum_{n \geq 1} (-1)^{n+1} \frac{(G - I)^n}{n} \right) A = A \left(\sum_{n \geq 1} (-1)^{n+1} \frac{(K - I)^n}{n} \right) = A \log(K).$$

Hence we may take $X = \log(G)$ and $Y = \log(K)$ with $X, Y \in \mathcal{V} \cap \mathfrak{g}_A$. \square

Theorem 3.14. *Let $A \in \mathcal{CR}$ and let \mathfrak{S} be a symmetrically-normed ideal. Then $\mathcal{C}_k(A)$ is a real analytic homogeneous space of $\mathcal{G}_{\mathfrak{S}} \times \mathcal{G}_{\mathfrak{S}}$. Furthermore, $\mathcal{C}_k(A)$ is also a real analytic submanifold of $A + \mathfrak{S}$ with the same differential structure, whose tangent space at $B \in \mathcal{C}_k(A)$ is*

$$(T\mathcal{C}_k(A))_B = \{XB - BY : X, Y \in \mathfrak{S}\} \subset \mathfrak{S} = (T(A + \mathfrak{S}))_B.$$

Proof. We may suppose that $k = 0$ and $A = B^{(0)} = B$. Notice that the isotropy subgroup of the action π_0 is given by $\mathcal{G}_A = \{(G, K) \in \mathcal{G}_{\mathfrak{S}} \times \mathcal{G}_{\mathfrak{S}} : GA = AK\}$. From Lemma 3.13, \mathcal{G}_A is a Banach-Lie subgroup of $\mathcal{G} = \mathcal{G}_{\mathfrak{S}} \times \mathcal{G}_{\mathfrak{S}}$. Then according to [46, Thm. 8.19] we get that the quotient space $\mathcal{G}/\mathcal{G}_A$ has the structure of real analytic manifold such that $\pi_0 : \mathcal{G} \rightarrow \mathcal{C}_0(A)$ is a real analytic submersion. Furthermore, Lemma 3.12 implies that $\mathcal{C}_0(A)$ is homeomorphic to the quotient $\mathcal{G}/\mathcal{G}_A$, so that $\mathcal{C}_0(A)$ inherits the real analytic homogeneous space structure from $\mathcal{G}/\mathcal{G}_A$.

Now we show that $\mathcal{C}_0(A)$ is a submanifold of $A + \mathfrak{S}$. We first observe that from the previous facts the tangent space $(T\mathcal{C}_0(A))_A$ of $\mathcal{C}_0(A)$ at A can be identified as

$$(T\mathcal{C}_0(A))_A \simeq \mathfrak{g}/\mathfrak{g}_A \simeq \{XA - AY : X, Y \in \mathfrak{S}\},$$

where we have used that $(T\mathcal{G}_A)_A = \mathfrak{g}_A = \{(X, Y) \in \mathfrak{g} : XA - AY = 0\}$, so that the elements in the quotient space $\mathfrak{g}/\mathfrak{g}_A$ can be identified with $\{XA - AY : X, Y \in \mathfrak{S}\}$ as above. We put $P = P_{R(A)}$ and $Q = P_{N(A)^\perp}$. If we take a tangent vector $V = XA - AY$, then $PVQ = P(XA - AY)Q$, $PVQ^\perp = -PAYQ^\perp$, $P^\perp VQ = P^\perp XAQ$ and $P^\perp VQ^\perp = 0$. Notice that the identity $PVQ^\perp = -PAYQ^\perp$ is equivalent to $PVQ^\perp = -A(QYQ^\perp)$, meanwhile $P^\perp VQ = P^\perp XAQ$ is equivalent to $P^\perp VQ = (P^\perp XP)AQ$. Thus, the operators $PXAQ = (PXP)AQ$ and $PAYQ = A(QYQ)$, which appear in PVQ , are independent of the operators PVQ^\perp and $P^\perp VQ$ (since these last operators can be determined in terms of the independent blocks QYQ^\perp and $P^\perp XP$, respectively). Therefore, as an operator from $\mathcal{H} = R(Q) \oplus N(Q)$ to $\mathcal{H} = R(P) \oplus N(P)$ we get that V has the form

$$V = \begin{pmatrix} Z_{11} & Z_{12} \\ Z_{21} & 0 \end{pmatrix}.$$

Conversely, if V is an operator having such matrix representation, then we may take

$$X = \begin{pmatrix} 0 & 0 \\ Z_{21}A^\dagger & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} A^\dagger Z_{11} & A^\dagger Z_{12} \\ 0 & 0 \end{pmatrix},$$

which satisfy $V = XA - AY$. Then the tangent space is given by

$$(T\mathcal{C}_0(A))_A = \left\{ \begin{pmatrix} Z_{11} & Z_{12} \\ Z_{21} & 0 \end{pmatrix} : Z_{11} \in P\mathfrak{S}Q, Z_{12} \in P\mathfrak{S}Q^\perp, Z_{21} \in P^\perp\mathfrak{S}Q \right\}. \quad (7)$$

From this last representation is now evident that $(T\mathcal{C}_0(A))_A$ is closed in \mathfrak{S} (tangent space of $A + \mathfrak{S}$), and it has a closed supplement defined by

$$\mathfrak{n} = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & Z_{22} \end{pmatrix} : Z_{22} \in P^\perp\mathfrak{S}Q^\perp \right\}.$$

Thus the inclusion map $\iota : \mathcal{C}_0 \simeq \mathcal{G}/\mathcal{G}_A \rightarrow A + \mathfrak{S}$, which is analytic, satisfies that its tangent map has a (closed) complemented range at every point. This means that ι is an immersion. Now we recall that the quotient topology on $\mathcal{C}_0(A)$ is always stronger than the relative topology, but in this case both topologies coincide as a consequence of Lemma 3.12. Hence we can apply [46, Prop. 8.7] to conclude that $\mathcal{C}_0(A)$ is a real analytic submanifold of $A + \mathfrak{S}$. Furthermore, the manifold structure as a homogeneous space coincides with that as a submanifold of $A + \mathfrak{S}$. Notice that the tangent map of π_0 at (I, I) is given by $T_{(I, I)}\pi_0 : \mathfrak{S} \times \mathfrak{S} \rightarrow (T\mathcal{C}_0(A))_A$, $T_{(I, I)}(X, Y) = XA - AY$. Since this map is a submersion, it follows that $(T\mathcal{C}_0(A))_A = R(T_{(I, I)}\pi_0) = \{XA - AY : X, Y \in \mathfrak{S}\}$. \square

3.3 Real analyticity of the Moore-Penrose inverse

In this subsection we show that the Moore-Penrose inverse is a real analytic map between Banach manifolds. Hence, we consider the decomposition of $\mathcal{CR} \cap (A + \mathfrak{S})$ into the connected manifolds $\mathcal{C}_k(A)$, $k \in \mathbb{J}_A$, in which the Moore-Penrose inverse is a continuous map. We further consider each maximal continuity set $\mathcal{C}_k(A)$ endowed with its homogeneous space structure induced by the action of $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$, or equivalently, its submanifold structure.

Remark 3.15. Let E, F be complex Banach spaces, let $\Omega \subseteq E$ be an open set and let $f : \Omega \subseteq E \rightarrow F$ be a complex analytic function. Consider E_0, F_0 real closed subspaces of E and F , respectively, and suppose that $f(E_0) \subseteq F_0$. We claim that the function $f_0 : \Omega \cap E_0 \rightarrow F_0$, $f_0 = f|_{\Omega \cap E_0}$, is real analytic. Since f is complex analytic, for every $x_0 \in \Omega$, there exists a convergent power series $\sum_{n \geq 0} f_n$ such that $f(x) = \sum_{n \geq 0} f_n(x - x_0)$ locally at x_0 , where each f_n is a continuous n -homogeneous polynomial defined on E with values on F . Denote by \tilde{f}_n the continuous multilinear function associated with f_n . To prove our claim, it suffices to check that if f is restricted to $\Omega \cap E_0$, then the corresponding multilinear functions satisfy $\tilde{f}_n(E_0^n) \subseteq F_0$. The case $n = 0$, that is $f_0 \in F_0$, follows by the assumption $f(E_0) \subseteq F_0$. Next we use that f is complex analytic, so in particular f is a C^∞ function such that its derivatives satisfy $f^{(n)}(x_0) = n! \tilde{f}_n$ for all $n \geq 1$ (see [46, Corol. 1.8]). For $x_0, x \in E$, notice that the Gateux derivative at x_0 in the direction of x gives

$$\tilde{f}_1(x) = f'(x_0)x = \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \in F_0,$$

since $f(E_0) \subseteq F_0$ and F_0 is closed. Hence $\tilde{f}_1(E_0) \subseteq F_0$. We can now use again the Gateaux derivative of \tilde{f}_1 to get $\tilde{f}_2(E_0 \times E_0) \subseteq F_0$. Continuing with this argument we can obtain $\tilde{f}_n(E_0^n) \subseteq F_0$, for all $n \geq 1$.

Lemma 3.16. *Let $\mathcal{S} \subseteq \mathcal{H}$ be a closed subspace. Then there exists a real analytic map $\mathcal{G}\ell_{\mathfrak{S}} \rightarrow \mathcal{U}_{\mathfrak{S}}$, $T \mapsto U_T$, such that $P_{T(\mathcal{S})} = U_T P_{\mathcal{S}} U_T^*$, for $T \in \mathcal{G}\ell_{\mathfrak{S}}$.*

Proof. Let $T \in \mathcal{G}\ell_{\mathfrak{S}}$ and set $P = P_{T(\mathcal{S})}$, $Q = TP_{\mathcal{S}}T^{-1}$. Then,

$$T_0 = PQ + (I - P)(I - Q) = Q + (I - P)(I - Q)$$

satisfies $PT_0 = T_0Q = Q$, $T_0|_{R(Q)} = I|_{R(Q)}$ and $T_0|_{N(Q)} = (I - P)|_{N(Q)}$ is an isomorphism between $N(Q)$ and $N(P)$. Since Q is an oblique projection we get that \mathcal{H} is the (non-orthogonal) direct sum of $R(Q)$ and $N(Q)$; thus, T_0 is an invertible operator.

We now show $T_0 \in \mathcal{G}\ell_{\mathfrak{S}}$. Since $T \in \mathcal{G}\ell_{\mathfrak{S}}$ then, by [22, Thm. 3.5] we get that $P - P_{\mathcal{S}} = P_{R(TP_{\mathcal{S}})} - P_{R(P_{\mathcal{S}})} \in \mathfrak{S}$ (and furthermore, $[P : P_{\mathcal{S}}] = 0$). On the other hand, $Q - P_{\mathcal{S}} = TP_{\mathcal{S}}(T^{-1} - I) + (T - I)P_{\mathcal{S}} \in \mathfrak{S}$. These facts imply that $Q - P \in \mathfrak{S}$ and hence $T_0 - I = P(Q - P) \in \mathfrak{S}$.

Now set $T_1 = T_0T$, which satisfies $T_1 \in \mathcal{G}\ell_{\mathfrak{S}}$ and $T_1P_{\mathcal{S}}T_1^{-1} = P$. Then, $T_1P_{\mathcal{S}} = PT_1$ and $P_{\mathcal{S}}T_1^* = T_1^*P$ gives $T_1^*T_1P_{\mathcal{S}} = P_{\mathcal{S}}T_1^*T_1$. Thus, we get $|T_1|P_{\mathcal{S}} = P_{\mathcal{S}}|T_1|$. The unitary $U_T = T_1|T_1|^{-1}$ now can be seen to satisfy $U_T \in \mathcal{U}_{\mathfrak{S}}$ and $U_T P_{\mathcal{S}} U_T^* = P_{T(\mathcal{S})}$.

To obtain that the map $T \mapsto U_T$ is real analytic and thereby complete the proof, we express this map as a composition of real analytic maps. Recall that $\tilde{\mathfrak{S}} = \{X + \lambda I : X \in \mathfrak{S}, \lambda \in \mathbb{C}\}$ is the

unitalization of \mathfrak{S} , and consider its self-adjoint part, i.e. $\tilde{\mathfrak{S}}_{sa} := \{X + \lambda I : X = X^*, \lambda \in \mathbb{R}\}$ (see Remark 2.1).

Since T is invertible, $P_{\mathcal{S}}T^*TP_{\mathcal{S}} \geq cP_{\mathcal{S}}$, $c = \min_{\lambda \in \sigma(|T|)} \lambda^2$, so that $P_{\mathcal{S}}T^*T|_{\mathcal{S}}$ is invertible on \mathcal{S} and $P_{\mathcal{S}}(P_{\mathcal{S}}T^*TP_{\mathcal{S}})^{\dagger}P_{\mathcal{S}} = (P_{\mathcal{S}}T^*T|_{\mathcal{S}})^{-1}P_{\mathcal{S}}$. Then, the map $\mathcal{G}l_{\mathfrak{S}} \rightarrow \tilde{\mathfrak{S}}_{sa}$, $T \mapsto P_{T(\mathcal{S})} = TP_{\mathcal{S}}(P_{\mathcal{S}}T^*T|_{\mathcal{S}})^{-1}P_{\mathcal{S}}T^*$ is clearly real analytic since the operator adjoint, multiplication and inversion maps are real analytic. If $Q_T = TP_{\mathcal{S}}T^{-1}$, then the map $\mathcal{G}l_{\mathfrak{S}} \times \tilde{\mathfrak{S}}_{sa} \rightarrow \tilde{\mathfrak{S}}$, $(T, R) \mapsto Q_T + (I - R)(I - Q_T)$ is also real analytic. Therefore,

$$T_0 : \mathcal{G}l_{\mathfrak{S}} \rightarrow \tilde{\mathfrak{S}}, \quad T_0(T) = Q_T + (I - P_{T(\mathcal{S})})(I - Q_T),$$

is real analytic. Since $\mathcal{G}l_{\mathfrak{S}}$ is a submanifold of $\tilde{\mathfrak{S}}$ as we state in Remark 2.1 *ii*), and $T_0(T) \in \mathcal{G}l_{\mathfrak{S}}$ for every $T \in \mathcal{G}l_{\mathfrak{S}}$, it follows that $T_0 : \mathcal{G}l_{\mathfrak{S}} \rightarrow \mathcal{G}l_{\mathfrak{S}}$ turns out to be real analytic. We conclude that $T_1 : \mathcal{G}l_{\mathfrak{S}} \rightarrow \mathcal{G}l_{\mathfrak{S}}$, $T_1(T) = T_0(T)T$, is real analytic.

On the other hand, notice that $\sigma(Z) = \sigma_{\tilde{\mathfrak{S}}}(Z)$ for any $Z = (Z - I) + I \in \mathcal{G}l_{\mathfrak{S}} \subset \tilde{\mathfrak{S}}$, where the left-hand spectrum is the spectrum as an operator on \mathcal{H} and the right-hand spectrum is the spectrum in the Banach algebra $\tilde{\mathfrak{S}}$. Put $\mathbb{C}_+ := \{z \in \mathbb{C} : \Re(z) > 0\}$, and then take the open subset $\Omega = \{Z \in \tilde{\mathfrak{S}} : \sigma(Z) \subseteq \mathbb{C}_+\}$ of $\tilde{\mathfrak{S}}$. Now we apply Remark 3.15 with $E = F = \tilde{\mathfrak{S}}$, $E_0 = F_0 = \tilde{\mathfrak{S}}_{sa}$ and the map $f : \Omega \rightarrow \tilde{\mathfrak{S}}$, $f(Z) = Z^{1/2}$, which is certainly complex analytic by the holomorphic functional calculus in the Banach algebra $\tilde{\mathfrak{S}}$ (see, e.g., [46, p. 30]). We thus get that $f_0 : \Omega \cap \tilde{\mathfrak{S}}_{sa} \rightarrow \tilde{\mathfrak{S}}_{sa}$, $f_0(Z) = Z^{1/2}$, is real analytic. For $T \in \mathcal{G}l_{\mathfrak{S}}$ notice that $|T| - I = (|T| + I)^{-1}(|T|^2 - I) \in \mathfrak{S}$, whence $T^*T = |T|^2$, $|T| \in \mathcal{G}l_{\mathfrak{S}}$. Hence the maps $\mathcal{G}l_{\mathfrak{S}} \rightarrow \mathcal{G}l_{\mathfrak{S}}$, $T \mapsto |T| = f_0(T^*T)$ and $T \mapsto T|T|^{-1}$ are real analytic. Since $\mathcal{U}_{\mathfrak{S}}$ is a submanifold $\mathcal{G}l_{\mathfrak{S}}$, we can co-restrict the previous map and find that $\mathcal{G}l_{\mathfrak{S}} \rightarrow \mathcal{U}_{\mathfrak{S}}$, $T \mapsto T|T|^{-1}$, is real analytic. Finally, we obtain that the map $\mathcal{G}l_{\mathfrak{S}} \rightarrow \mathcal{U}_{\mathfrak{S}}$, $T \mapsto U_T$, where $U_T = T_1(T)|T_1(T)|^{-1}$ is real analytic. \square

Remark 3.17. For fixed $A \in \mathcal{CR}$ we may also express $\mathcal{CR} \cap (A^{\dagger} + \mathfrak{S})$ as follows

$$\mathcal{CR} \cap (A^{\dagger} + \mathfrak{S}) = \bigcup_{k \in \mathbb{J}_A} \mathcal{C}_k(A^{\dagger}),$$

where $\mathbb{J}_A = \mathbb{J}_{A^{\dagger}}$ is the set defined in Notation 3.8, and we set

$$\mathcal{C}_k(A^{\dagger}) := \{B \in \mathcal{CR} : B - A^{\dagger} \in \mathfrak{S}, [P_{N(B)} : P_{N(A^{\dagger})}] = k\}.$$

In particular, by Theorem 3.14, $\mathcal{C}_k(A^{\dagger})$ is also a real analytic homogeneous space of $\mathcal{G} = \mathcal{G}l_{\mathfrak{S}} \times \mathcal{G}l_{\mathfrak{S}}$ and a submanifold of $A^{\dagger} + \mathfrak{S}$. From Lemma 3.2, we know that $B - A \in \mathfrak{S}$ if and only if $B^{\dagger} - A^{\dagger} \in \mathfrak{S}$. Thus we can consider the bijection $\mu : \mathcal{CR} \cap (A + \mathfrak{S}) \rightarrow \mathcal{CR} \cap (A^{\dagger} + \mathfrak{S})$, $\mu(B) = B^{\dagger}$. Since $[P_{N(B^{\dagger})} : P_{N(A^{\dagger})}] = -[P_{N(B^{\dagger})^{\perp}} : P_{N(A^{\dagger})^{\perp}}] = -[P_{R(B)} : P_{R(A)}] = [P_{N(B)} : P_{N(A)}]$ (the last equality follows from Theorem 3.9), we get $\mu(\mathcal{C}_k(A)) = \mathcal{C}_k(A^{\dagger})$, for $k \in \mathbb{J}_A$. Hence $\mu : \mathcal{C}_k(A) \rightarrow \mathcal{C}_k(A^{\dagger})$, $\mu(B) = B^{\dagger}$, is a homeomorphism.

The previous remark concerns the map defined by the Moore-Penrose inverse at a topological level. The geometric structures we have constructed in Theorem 3.14 allow us to consider the Moore-Penrose inverse as a map between Banach manifolds. We can now give one of our main results about the Moore-Penrose inverse in this framework.

Theorem 3.18. *Let $A \in \mathcal{CR}$ and let \mathfrak{S} be a symmetrically-normed ideal. Then for every $k \in \mathbb{J}_A$ the map*

$$\mu : \mathcal{C}_k(A) \rightarrow \mathcal{C}_k(A^{\dagger}), \quad \mu(B) = B^{\dagger},$$

is real bianalytic between these Banach manifolds, and its tangent map is given by

$$(T_B\mu)(V) = -B^{\dagger}VB^{\dagger} + (B^*B)^{\dagger}V^*(I - BB^{\dagger}) + (I - B^{\dagger}B)V^*(BB^*)^{\dagger},$$

for $B \in \mathcal{C}_k(A)$ and $V \in (T\mathcal{C}_k(A))_B$.

Proof. We may assume that $k = 0$. As we observed in Remark 3.17 the map μ in the statement is a homeomorphism. Now we recall two properties of the Moore-Penrose inverse. For $C \in \mathcal{CR}$, and U, V unitary operators, it holds that $(UCV)^\dagger = V^*C^\dagger U^*$, and that C^\dagger is determined by the conditions $C^\dagger|_{R(C)^\perp} = 0$ and $C^\dagger|_{R(C)} = (C|_{R(C)})^{-1}$, where the inverse here is given by $(C|_{R(C)})^{-1} : R(C) \rightarrow N(C)^\perp$ satisfying $(C|_{R(C)})^{-1}C|_{N(C)^\perp} = I_{N(C)^\perp}$ and $C(C|_{R(C)})^{-1} = I_{R(C)}$. According to Lemma 3.16, for every $(G, K) \in \mathcal{GL}_\mathfrak{S} \times \mathcal{GL}_\mathfrak{S}$ there are unitary operators $U_K, U_G \in \mathcal{U}_\mathfrak{S}$ such that $U_K(N(A)^\perp) = K(N(A)^\perp)$, and $U_G(R(A)) = G(R(A))$. Then,

$$\begin{aligned} (GAK^{-1})^\dagger &= U_K(U_G^*(GAK^{-1})U_K)^\dagger U_G^* \\ &= U_K \begin{pmatrix} A^\dagger & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} [U_G^*(GAK^{-1})U_K A^\dagger]^{-1} & 0 \\ 0 & 0 \end{pmatrix} U_G^*, \end{aligned}$$

where the block matrix representation of factor to the right is with respect to the decomposition $\mathcal{H} = R(A) \oplus R(A)^\perp$ and the block matrix representation of the factor to the left is as an operator acting from $\mathcal{H} = R(A) \oplus R(A)^\perp$ into $\mathcal{H} = N(A)^\perp \oplus N(A)$. The operator obtained by restriction $U_G^*(GAK^{-1})U_K A^\dagger : R(A) \rightarrow R(A)$ is invertible, and using that $P_{R(A)} = AA^\dagger$ we further get that $U_G^*(GAK^{-1})U_K A^\dagger - P_{R(A)} \in \mathfrak{S}$. We write $\mathfrak{S}(R(A))$ for the corresponding symmetrically-normed ideal \mathfrak{S} on $R(A)$. By considering also the restricted invertible group on $R(A)$, i.e. $\mathcal{GL}_\mathfrak{S}(R(A)) := \{G \in \mathcal{GL}(R(A)) : G - I_{R(A)} \in \mathfrak{S}(R(A))\}$, we can use that the inversion map is real analytic on this Lie group. Since $\mathcal{U}_\mathfrak{S}$ is a Lie group, the inversion is real analytic, and using that $K \mapsto U_K$ and $G \mapsto U_G$ are real analytic by Lemma 3.16, we get from the above expression that the map $\tilde{\pi}_0 : \mathcal{GL}_\mathfrak{S} \times \mathcal{GL}_\mathfrak{S} \rightarrow \mathcal{C}_0(A^\dagger)$, $\tilde{\pi}_0(G, K) = (GAK^{-1})^\dagger$ is real analytic. By Theorem 3.14 we conclude that $\pi_0 : \mathcal{GL}_\mathfrak{S} \times \mathcal{GL}_\mathfrak{S} \rightarrow \mathcal{C}_0(A)$ is real analytic, and hence it has real analytic local cross sections ([46, Corol. 8.3]). Thus for every $B \in \mathcal{C}_0(A)$, there is an open set $B \in \mathcal{W} \subseteq \mathcal{C}_0(A)$, and a real analytic map $s : \mathcal{W} \rightarrow \mathcal{GL}_\mathfrak{S} \times \mathcal{GL}_\mathfrak{S}$ such that $\pi_0 \circ s = id|_{\mathcal{W}}$. Therefore, we can write locally $\mu = \tilde{\pi}_0 \circ s$. This shows that μ is real analytic. Clearly, μ becomes a bianalytic map between these manifolds.

Now we can compute the tangent map at $B \in \mathcal{C}_0(A)$ in the direction of a vector $V = XB - BY \in (T\mathcal{C}_0(A))_B$. Take the curve $\gamma(t) = e^{tX} B e^{-tY} \in \mathcal{C}_0(A)$, for some $X, Y \in \mathfrak{S}$, and $t \in (-1, 1)$, which satisfies $\gamma(0) = B$, $\dot{\gamma}(0) = V$. By Wedin's formula (4) and the continuity of μ in $\mathcal{C}_0(A)$ we get that

$$\begin{aligned} (T_B \mu)(V) &= \left. \frac{d}{dt} \right|_{t=0} \mu(\gamma(t)) \\ &= \lim_{t \rightarrow 0} \frac{1}{t} \{ -\gamma(t)^\dagger (\gamma(t) - \gamma(0)) \gamma(0)^\dagger + (\gamma(t)^* \gamma(t))^\dagger (\gamma(t)^* - \gamma(0)^*) (I - \gamma(0) \gamma(0)^\dagger) + \\ &\quad + (I - \gamma(t)^\dagger \gamma(t)) (\gamma(t)^* - \gamma(0)^*) (\gamma(0) \gamma(0)^*)^\dagger \} \\ &= -B^\dagger V B^\dagger + (B^* B)^\dagger V^* (I - B B^\dagger) + (I - B^\dagger B) V^* (B B^*)^\dagger. \end{aligned}$$

□

Remark 3.19. *i)* For $A \in \mathcal{CR}$, $G, K \in \mathcal{GL}_\mathfrak{S}$, we may compute $(GAK^{-1})^\dagger$ using the following formula proved in [29, Thm. 2]:

$$(GAK^{-1})^\dagger = (AK^{-1})^* [AK^{-1}(AK^{-1})^* + I - AA^\dagger]^{-1} A [(GA)^* GA + I - A^\dagger A]^{-1} (GA)^*.$$

A straightforward consequence is that $(G, K) \mapsto (GAK^{-1})^\dagger$ is a real analytic map. Thus, this gives another proof of the fact that μ is a real analytic map. However, the approach considered in the proof of Theorem 3.18, which is based on Lemma 3.16, is also needed in Section 4.

ii) Since $\mathcal{GL}_\mathfrak{S}$ is also a complex Lie group, the same proof of Theorem 3.14 also shows that \mathcal{C}_k , $k \in \mathbb{J}_A$, are complex analytic homogeneous spaces and submanifolds. The operator adjoint that shows up in the proofs of Lemma 3.16 and Theorem 3.18 implies that μ cannot be complex analytic. We refer to [36] for a study of other generalized inverses as complex analytic mappings operator valued mappings of a complex variable.

4 Polar decomposition

We first study the real analyticity of operator monotone functions defined on invertible positive perturbations by symmetrically-normed ideals. Then we develop a geometric study of the closed range positive operator perturbations by an ideal. This leads to a decomposition into connected sets, which are real analytic homogeneous spaces (congruence orbits) and submanifolds. In the next section we use these results, and also the ones from the previous sections, to show that the maps given by the operator modulus and the partial isometry in the polar decomposition are real analytic fiber bundles.

4.1 Real analyticity of operator monotone functional calculus

We revisit the Ando-van Hemmen theory on the perturbation problem for operator monotone functions [7]. We are interested in a rather different problem, related to the real analyticity of the operator monotone functional calculus with respect to operator ideal perturbations. Nevertheless, our approach is deeply influenced by the techniques from [7]. Recall that a function $f : [0, \infty) \rightarrow \mathbb{R}$ is said to be operator monotone if $C, D \in \mathcal{B}(\mathcal{H})^+$ are such that $C \leq D$, then we have that $f(C) \leq f(D)$. In this case f belongs to the Pick class. Hence, there exist a positive Borel measure ν on $(0, \infty)$ such that $\int_0^\infty (t^2 + 1)^{-1} d\nu(t)$ is finite, $\alpha \in \mathbb{R}$ and $\beta \in [0, \infty)$ such that

$$f(\lambda) = \alpha + \beta \lambda - \int_0^\infty \left(\frac{1}{t + \lambda} - \frac{t}{t^2 + 1} \right) d\nu(t) \quad \text{for } \lambda \in [0, \infty). \quad (8)$$

Moreover, notice that the expression for $f(\lambda)$ above is well defined for $\lambda \in \mathbb{C} \setminus (-\infty, 0)$. For example, if $f(\lambda) = \lambda^{1/2}$, for $\lambda \in [0, \infty)$, then

$$\lambda^{1/2} = \frac{1}{\sqrt{2}} - \int_0^\infty \left(\frac{1}{t + \lambda} - \frac{t}{t^2 + 1} \right) \frac{t^{1/2}}{\pi} dt \quad \text{for } \lambda \in [0, \infty). \quad (9)$$

Using the integral representation in Eq. (8) for the operator monotone function $f(\lambda)$, then given $C \geq 0$ we can represent the self-adjoint operator

$$f(C) = \alpha I + \beta C - \int_0^\infty \left((tI + C)^{-1} - \frac{t}{t^2 + 1} I \right) d\nu(t).$$

In what follows we use the notation $\mathcal{G}l^+ := \mathcal{G}l^+(\mathcal{H}) = \{G \in \mathcal{G}l(\mathcal{H}) : G > 0\}$. For a fixed $C \in \mathcal{G}l^+$ notice that $\mathcal{G}l^+ \cap (C + \mathfrak{S})$ is an open subset of the affine space $C + \mathfrak{S}$, where the topology is, as usual, defined by the metric $d_{\mathfrak{S}}(B_1, B_2) = \|B_1 - B_2\|_{\mathfrak{S}}$, $B_1, B_2 \in C + \mathfrak{S}$. We recall that \mathfrak{S}_{sa} denotes the set of self-adjoint operators in \mathfrak{S} .

Remark 4.1. Consider a monotone operator function represented as in Eq. (8) and let $C, D \in \mathcal{G}l^+$ be such that $D - C \in \mathfrak{S}$. In [7] Ando and van-Hemmen showed, in particular, that $f(D) - f(C)$ lies in the maximal ideal associated to the symmetric norming function induced by \mathfrak{S} ; for the notion of maximal ideal see [38]. In this context, they noticed that

$$\begin{aligned} f(D) - f(C) &= \beta(D - C) + \int_0^\infty \left((tI + C)^{-1} - (tI + D)^{-1} \right) d\nu(t) \\ &= \beta(D - C) + \int_0^\infty (tI + C)^{-1} (D - C) (tI + D)^{-1} d\nu(t) \end{aligned}$$

The previous identities suggest to consider the operator valued function $h : [0, \infty) \rightarrow \mathfrak{S}_{sa}$ given by

$$h(t) = (tI + C)^{-1} (D - C) (tI + D)^{-1} \quad \text{for } t \in [0, \infty).$$

In this case it is straightforward to check that:

1. $\|h(t)\|_{\mathfrak{S}} \leq q(t)$, where $q(t)$ is a bounded, continuous, positive and non-increasing function such that $\int_0^\infty q(t) d\nu(t) < \infty$ (e.g. $q(t) = \|D - C\|_{\mathfrak{S}}((t + \gamma_C)(t + \gamma_D))^{-1}$, $t \geq 0$);
2. If $\delta > 0$, then $\|h(t + \delta) - h(t)\|_{\mathfrak{S}} \leq \alpha \delta r(t)$, where $\alpha > 0$ and $r : [0, \infty) \rightarrow [0, \infty)$ is a continuous and non-increasing function, such that

$$\lim_{t \rightarrow \infty} r(t) = 0 \quad \text{and} \quad \int_0^\infty r(t) d\nu(t) < \infty.$$

We point out that in the case above we can take $r(t)$ to be

$$r(t) = ((t + \gamma_C)(t + \gamma_D))^{-1} \|D - C\|_{\mathfrak{S}} ((t + \gamma_C)^{-1} + (t + \gamma_D)^{-1}), \quad t \in [0, \infty).$$

The following lemma is an integral formulation of the fact that absolute convergence implies convergence in symmetrically-normed ideals.

Lemma 4.2. *Let \mathfrak{S} be symmetrically-normed ideal, let ν be a positive Borel measure on $(0, \infty)$ and let $h : [0, \infty) \rightarrow \mathfrak{S}_{sa}$ be a function satisfying items 1 and 2 in Remark 4.1. Then,*

$$\int_0^\infty h(t) d\nu(t) \in \mathfrak{S}_{sa} \quad \text{and} \quad \left\| \int_0^\infty h(t) d\nu(t) \right\|_{\mathfrak{S}} \leq \int_0^\infty \|h(t)\|_{\mathfrak{S}} d\nu(t).$$

Proof. See Section 5. □

Proposition 4.3. *Let $C, D \in \mathcal{G}\ell^+$ be such that $D - C \in \mathfrak{S}$, and let $f : [0, \infty) \rightarrow \mathbb{R}$ be an operator monotone function with integral representation as in Eq. (8). Then, we have that $f(D) - f(C) \in \mathfrak{S}_{sa}$ is such that*

$$\|f(D) - f(C)\|_{\mathfrak{S}} \leq \|D - C\|_{\mathfrak{S}} \left(\beta + \int_0^\infty \frac{1}{(t + \gamma_C)(t + \gamma_D)} d\nu(t) \right).$$

Proof. With the notation above, and arguing as in Remark 4.1 we get that

$$f(D) - f(C) = \beta(D - C) + \int_0^\infty h(t) d\nu(t) \quad \text{for} \quad h(t) = (tI + C)^{-1}(D - C)(tI + D)^{-1}. \quad (10)$$

Notice that $h(t)$ fulfills the conditions in items 1 and 2 in Remark 4.1 with $\|h(t)\|_{\mathfrak{S}} \leq \|D - C\|_{\mathfrak{S}}((t + \gamma_C)(t + \gamma_D))^{-1}$, for $t \geq 0$. Then, by Lemma 4.2 we see that

$$\int_0^\infty h(t) d\nu(t) \in \mathfrak{S}_{sa} \quad \text{and} \quad \left\| \int_0^\infty h(t) d\nu(t) \right\|_{\mathfrak{S}} \leq \int_0^\infty \frac{\|D - C\|_{\mathfrak{S}}}{(t + \gamma_C)(t + \gamma_D)} d\nu(t).$$

The result now follows from Eq. (10) and the last remarks. □

Next we study the continuity of the functional calculus induced by a (fixed) monotone operator function, with respect to certain perturbations of a positive closed range operator. To do this, we consider the following facts.

Remark 4.4. Let P, Q_n be orthogonal projections such that $P - Q_n \in \mathfrak{S}$, for all $n \geq 1$, and $\|P - Q_n\|_{\mathfrak{S}} \rightarrow 0$. Then, there exists a sequence $\{U_n\}_{n \geq 1}$ in $\mathcal{U}_{\mathfrak{S}}$ such that $U_n Q_n U_n^* = P$, for $n \geq 1$, and $\|U_n - I\|_{\mathfrak{S}} \rightarrow 0$. Indeed, this is a consequence of [5, Proposition 2.2.] for the ideal \mathfrak{S}_2 of Hilbert-Schmidt operators; there it is shown the existence of continuous local cross sections for the map $\mathcal{U}_{\mathfrak{S}_2} \ni U \mapsto U^* P U \subset P + (\mathfrak{S}_2)_{sa}$, where $\mathcal{U}_{\mathfrak{S}_2}$ and $P + (\mathfrak{S}_2)_{sa}$ are endowed with the Hilbert-Schmidt metric $d_2(C, D) = \|C - D\|_2$. The general case of a symmetrically-normed operator ideal follows with a straightforward adaption of the proof of the previous result.

In what follows we let $\mathcal{CR}^+ = \{C \in \mathcal{CR} : C \geq 0\}$.

Corollary 4.5. *Let $f : [0, \infty) \rightarrow \mathbb{R}$ be an operator monotone function and let \mathfrak{S} be a symmetrically-normed ideal. Fix $C \in \mathcal{CR}^+$ and let $\{D_n\}_{n \geq 1}$ be a sequence in \mathcal{CR}^+ such that $C - D_n \in \mathfrak{S}$, $[P_{N(C)} : P_{N(D_n)}] = 0$ for $n \geq 1$, and $\|D_n - C\|_{\mathfrak{S}} \rightarrow 0$. Then, there exists $n_0 \geq 1$ such that $f(D_n) - f(C) \in \mathfrak{S}$, for $n \geq n_0$, and $\|f(D_n) - f(C)\|_{\mathfrak{S}} \rightarrow 0$.*

Proof. Let $\{D_n\}_{n \geq 1}$ be as above and assume further that $N(D_n) = N(C)$ for $n \geq 1$. Then the restrictions $D_n|_{\mathcal{H}_0}, C|_{\mathcal{H}_0} \in \mathcal{B}(\mathcal{H}_0)$ are positive invertible operators acting on $\mathcal{H}_0 = R(C)$ such that $\|D_n|_{\mathcal{H}_0} - C|_{\mathcal{H}_0}\|_{\mathfrak{S}} \rightarrow 0$. Let $n_0 \geq 1$ be such that $\|D_n - C\|_{\mathfrak{S}} \leq \gamma_{C|_{\mathcal{H}_0}}/2$, so that $\gamma_{D_n|_{\mathcal{H}_0}} > \gamma_{C|_{\mathcal{H}_0}}/2$, for $n \geq n_0$. In this case

$$((t + \gamma_{D_n|_{\mathcal{H}_0}})(t + \gamma_{C|_{\mathcal{H}_0}}))^{-1} \leq 4(t + \gamma_{C|_{\mathcal{H}_0}})^{-2}, \quad t \geq 0.$$

By Proposition 4.3 we get that $f(D_n|_{\mathcal{H}_0}) - f(C|_{\mathcal{H}_0}) \in \mathfrak{S}(\mathcal{H}_0)$ for $n \geq n_0$, and that $\|f(D_n|_{\mathcal{H}_0}) - f(C|_{\mathcal{H}_0})\|_{\mathfrak{S}} \rightarrow 0$. Now notice that $f(C) = f(C|_{\mathcal{H}_0}) + f(0)(I - P_{\mathcal{H}_0})$ and similarly for $f(D_n)$, $n \geq 1$; thus, $f(D_n) - f(C) \in \mathfrak{S}$ for $n \geq n_0$ and $\|f(D_n) - f(C)\|_{\mathfrak{S}} = \|f(D_n|_{\mathcal{H}_0}) - f(C|_{\mathcal{H}_0})\|_{\mathfrak{S}} \rightarrow 0$.

Consider now a general sequence $\{D_n\}_{n \geq 1}$ as in the statement above. Notice that by hypothesis and Theorem 3.4, we get that $\|D_n^\dagger - C^\dagger\|_{\mathfrak{S}} \rightarrow 0$ and hence $\|P_{N(D_n)} - P_{N(C)}\|_{\mathfrak{S}} = \|D_n^\dagger D_n - C^\dagger C\|_{\mathfrak{S}} \rightarrow 0$. Let $\{U_n\}_{n \geq 1}$ be a sequence in $\mathcal{G}\ell_{\mathfrak{S}}$ such that $U_n P_{N(D_n)} U_n^* = P_{N(C)}$ and $\|U_n - I\|_{\mathfrak{S}} \rightarrow 0$ (see Remark 4.4). Hence, $U_n D_n U_n^* - C \in \mathfrak{S}$, $N(U_n D_n U_n^*) = N(C)$ and $\|U_n D_n U_n^* - C\|_{\mathfrak{S}} \rightarrow 0$. By the first part of the proof we conclude that $\|f(U_n D_n U_n^*) - f(C)\|_{\mathfrak{S}} \rightarrow 0$. On the other hand, $f(U_n D_n U_n^*) = U_n f(D_n) U_n^*$ so $\|f(U_n D_n U_n^*) - f(D_n)\|_{\mathfrak{S}} \rightarrow 0$. The previous facts imply that

$$\|f(D_n) - f(C)\|_{\mathfrak{S}} \leq \|f(D_n) - f(U_n D_n U_n^*)\|_{\mathfrak{S}} + \|f(U_n D_n U_n^*) - f(C)\|_{\mathfrak{S}} \rightarrow 0. \quad \square$$

In the proof of Lemma 3.16 we have used that the square root is a real analytic map in the set $\mathcal{G}\ell_{\mathfrak{S}} \cap \mathcal{G}\ell^+$, by considering the enveloping unital Banach algebra $\tilde{\mathfrak{S}}$ (see Remark 2.1). Thus, the argument in that proof cannot be repeated when the domain is changed to the set of perturbations $\{C + K \in \mathcal{G}\ell^+ : K \in \mathfrak{S}_{sa}\}$ endowed with the distance $d_{\mathfrak{S}}$, where $C \in \mathcal{G}\ell^+$ is some fixed positive invertible operator. Below we prove that the square root, and moreover any operator monotone function, is real analytic on these more general domains.

Theorem 4.6. *Let $f : [0, \infty) \rightarrow \mathbb{R}$ be an operator monotone function on $[0, \infty)$ with integral representation as in Eq. (8) and let $C \in \mathcal{G}\ell^+$. Consider $f : (C + \mathfrak{S}) \cap \mathcal{G}\ell^+ \rightarrow f(C) + \mathfrak{S}_{sa}$ given by $(C + \mathfrak{S}) \cap \mathcal{G}\ell^+ \ni D \mapsto f(D) \in f(C) + \mathfrak{S}_{sa}$. Then, f is a real analytic function. Moreover, for $\|D - C\|_{\mathfrak{S}} < \gamma_C$ we get a local series representation $f(D) = f(C) + \sum_{n=1}^{\infty} f_n(D - C)$, where*

$$f_1(D - C) = \beta(D - C) + \int_0^{\infty} (tI + C)^{-1} (D - C) (tI + C)^{-1} d\nu(t) \in \mathfrak{S}_{sa},$$

$$f_n(D - C) = (-1)^{n+1} \int_0^{\infty} ((tI + C)^{-1} (D - C))^n (tI + C)^{-1} d\nu(t) \in \mathfrak{S}_{sa}, \quad n \geq 2.$$

Proof. Given C as above we show that f admits a local power series around C , as in Remark 3.15. Indeed, assume that $D \in \mathcal{G}\ell^+$ is such that $D - C \in \mathfrak{S}$. By Proposition 4.3 we get that $f(D) \in f(C) + \mathfrak{S}_{sa}$. As noticed in [7] (see also Remark 4.1) we have that

$$f(D) - f(C) = \beta(D - C) + \int_0^{\infty} (tI + C)^{-1} (D - C) (tI + D)^{-1} d\nu(t). \quad (11)$$

Now, a closer look at the integrand reveals that for $t \in (0, \infty)$,

$$\begin{aligned} (tI + C)^{-1} (D - C) (tI + D)^{-1} &= (tI + C)^{-1} (D - C) (tI + C + (D - C))^{-1} \\ &= (tI + C)^{-1} (D - C) (I + (tI + C)^{-1} (D - C))^{-1} (tI + C)^{-1}. \end{aligned}$$

Since $\gamma_{tI+C} = t + \gamma_C \geq \gamma_C > 0$ then $\|(tI + C)^{-1}\| \leq \gamma_C^{-1}$, for $t \in (0, \infty)$. Hence, if $\|D - C\|_{\mathfrak{S}} < \gamma_C$, then we get that

$$\|(tI + C)^{-1}(D - C)\|_{\mathfrak{S}} \leq \|(tI + C)^{-1}\| \|D - C\|_{\mathfrak{S}} \leq \gamma_C^{-1} \|D - C\|_{\mathfrak{S}} < 1 \quad \text{for } t \in (0, \infty). \quad (12)$$

Thus, for $n \geq 1$ we have that

$$\|((tI + C)^{-1}(D - C))^n\| \leq \|((tI + C)^{-1}(D - C))^n\|_{\mathfrak{S}} \leq \|((tI + C)^{-1}(D - C))\|_{\mathfrak{S}}^n$$

since $\|\cdot\|_{\mathfrak{S}}$ is submultiplicative. The previous estimates show that the geometric series

$$\sum_{n=0}^{\infty} (-1)^n ((tI + C)^{-1}(D - C))^n = (I + (tI + C)^{-1}(D - C))^{-1} \in \mathcal{G}\ell_{\mathfrak{S}}$$

is $\|\cdot\|_{\mathfrak{S}}$ -absolute convergent and $\|\cdot\|_{\mathfrak{S}}$ -uniformly convergent for $t \in (0, \infty)$, by Eq. (12). In particular, the series is $\|\cdot\|$ -absolute convergent and $\|\cdot\|$ -uniformly convergent for $t \in (0, \infty)$. Therefore, we now see that

$$\begin{aligned} & \int_0^{\infty} ((tI + C)^{-1} - (tI + D)^{-1}) d\nu(t) = \\ & \int_0^{\infty} (tI + C)^{-1} (D - C) (I + (tI + C)^{-1}(D - C))^{-1} (tI + C)^{-1} d\nu(t) = \\ & \int_0^{\infty} (tI + C)^{-1} (D - C) \sum_{n=0}^{\infty} (-1)^n ((tI + C)^{-1}(D - C))^n (tI + C)^{-1} d\nu(t) = \\ & \sum_{n=0}^{\infty} (-1)^n \int_0^{\infty} ((tI + C)^{-1} (D - C))^{n+1} (tI + C)^{-1} d\nu(t) = \sum_{n=1}^{\infty} \tilde{f}_n(D - C) \end{aligned}$$

where, for $n \geq 1$ we let

$$\tilde{f}_n(D - C) = (-1)^{n+1} \int_0^{\infty} ((tI + C)^{-1}(D - C))^n (tI + C)^{-1} d\nu(t) \in \mathfrak{S}_{sa}$$

which is a $\|\cdot\|_{\mathfrak{S}}$ -continuous homogeneous polynomial of degree n (as a function of $D - C$) with values in \mathfrak{S}_{sa} , and the series is $\|\cdot\|_{\mathfrak{S}}$ -absolutely convergent. Indeed, to show $\tilde{f}_n(D - C) \in \mathfrak{S}_{sa}$ we argue as follows: for $n \geq 1$, let $h_n : [0, \infty) \rightarrow \mathfrak{S}_{sa}$ be given by

$$h_n(t) = ((tI + C)^{-1}(D - C))^n (tI + C)^{-1} \in \mathfrak{S}_{sa}, \quad t \geq 0.$$

In this case, $h(t)$ satisfies item 1 in Remark 4.1 since

$$\|h_n(t)\|_{\mathfrak{S}} \leq (t + \gamma_C)^{-2} (\gamma_C^{-1} \|D - C\|_{\mathfrak{S}})^{n-1} \|D - C\|_{\mathfrak{S}}, \quad t \geq 0, \quad (13)$$

$$(\gamma_C^{-1} \|D - C\|_{\mathfrak{S}})^{n-1} \|D - C\|_{\mathfrak{S}} \int_0^{\infty} (t + \gamma_C)^{-2} d\nu(t) < \infty,$$

where we have used that $\int_0^{\infty} (t^2 + 1)^{-1} d\nu(t)$ is finite. Furthermore, for $\delta > 0$ we have that

$$\|h(t + \delta) - h(t)\|_{\mathfrak{S}} \leq \delta (\gamma_C^{-1} \|D - C\|_{\mathfrak{S}})^{n-1} \|D - C\|_{\mathfrak{S}} (t + \gamma_C)^{-3} \quad \text{with} \quad \int_0^{\infty} (t + \gamma_C)^{-3} d\nu(t) < \infty.$$

Hence, $h(t)$ also satisfies item 2 in Remark 4.1. By Lemma 4.2 we conclude that

$$\tilde{f}_n(D - C) = (-1)^{n+1} \int_0^{\infty} h(t) d\nu(t) \in \mathfrak{S}_{sa}$$

and, using Eq. (13), we also get that

$$\|\tilde{f}_n(D - C)\|_{\mathfrak{S}} \leq (\gamma_C^{-1} \|D - C\|_{\mathfrak{S}})^{n-1} \|D - C\|_{\mathfrak{S}} \int_0^{\infty} (t + \gamma_C)^{-2} d\nu(t). \quad (14)$$

Hence, $\tilde{f}_n(D - C) \in \mathfrak{S}_{sa}$ is a $\|\cdot\|_{\mathfrak{S}}$ -continuous homogeneous polynomial of degree $n \geq 1$. Also, Eq. (14) shows that the series

$$\sum_{n=1}^{\infty} \tilde{f}_n(D - C) \in \mathfrak{S}_{sa}$$

is $\|\cdot\|_{\mathfrak{S}}$ -absolutely convergent for $\|D - C\|_{\mathfrak{S}} < \gamma_C$. The previous facts together with Eq. (11) show that for $D \in \mathcal{G}\ell^+$ with $\|D - C\|_{\mathfrak{S}} < \gamma_C$ we have the local expansion

$$f(D) = f(C) + \beta(D - C) + \sum_{n=1}^{\infty} \tilde{f}_n(D - C).$$

In particular, $f : (C + \mathfrak{S}) \cap \mathcal{G}\ell^+ \rightarrow f(C) + \mathfrak{S}_{sa}$ is a real analytic function. \square

Corollary 4.7. *Consider the notation in Theorem 4.6. For $D \in (C + \mathfrak{S}) \cap \mathcal{G}\ell^+$ with $\|D - C\|_{\mathfrak{S}} < \gamma_C$ we have that:*

$$\|f_1(D - C)\|_{\mathfrak{S}} \leq (\beta + \int_0^{\infty} (t + \gamma_C)^{-2} d\nu(t)) \|D - C\|_{\mathfrak{S}}$$

and for $n \geq 2$,

$$\begin{aligned} \|f_n(D - C)\|_{\mathfrak{S}} &\leq \int_0^{\infty} (t + \gamma_C)^{-(n+1)} d\nu(t) \|D - C\|_{\mathfrak{S}}^n \\ &\leq \int_0^{\infty} (t + \gamma_C)^{-2} d\nu(t) (\|D - C\|_{\mathfrak{S}} \gamma_C^{-1})^{n-1} \|D - C\|_{\mathfrak{S}} \end{aligned}$$

Proof. The result follows from a direct inspection of the proof of Theorem 4.6 above and Lemma 4.2. \square

Integral representations of (Fréchet) derivatives of the functional calculus induced by an operator monotone function have been considered before (see [16, 23, 41]). We remark that our previous result does not only provide integral representations of the derivatives (of all orders) of the functional calculus by operator monotone functions, but it also provides theoretical insights about the relevance of these derivatives for the computation of approximations of the function with respect to symmetrically-normed operator ideals.

Remark 4.8. Consider the notation in Theorem 4.6. The bounds in Corollary 4.7 allow to obtain simple estimates for the norm $\|\cdot\|_{\mathfrak{S}}$ of the error in the approximation (remainder) $f(D) \approx f(C) + \sum_{n=1}^m f_n(D - C)$, for $m \geq 1$. For example, if $f(\lambda) = \lambda^{1/2}$ for $\lambda \in [0, \infty)$ then Corollary 4.7 implies that

$$\|f_n(D - C)\|_{\mathfrak{S}} \leq \frac{1}{\pi} \int_0^{\infty} (t + \gamma_C)^{-(n+1)} t^{1/2} dt \|D - C\|_{\mathfrak{S}}, \quad n \geq 1,$$

where we have used the integral representation in Eq. (9) (so that $\beta = 0$). It is not difficult to obtain upper bounds for the integral above that in turn allows to obtain upper bounds for the remainder. Similar results have been obtained in [23] for this particular choice of f . Nevertheless, notice that the fact that the corresponding functional calculus is real analytic with respect to symmetrically-normed ideals seems to be new even in this case.

There are other important operator monotone functions whose Fréchet derivatives have been considered in the literature. Our results also imply some relevant information about the properties of the corresponding functional calculus and Taylor approximations; we will consider these results elsewhere.

4.2 Geometric structure of positive perturbations

We first give the following (algebraic) characterization of perturbations of a fixed positive operator by symmetrically-normed ideals.

Lemma 4.9. *Let $C \in \mathcal{CR}^+$ and \mathfrak{S} be a symmetrically-normed ideal. Then the following conditions are equivalent:*

- i) $D \in \mathcal{CR}^+$ satisfies $D - C \in \mathfrak{S}$ and $[P_{N(D)} : P_{N(C)}] = 0$;*
- ii) There exists $G \in \mathcal{G}\ell_{\mathfrak{S}}$ such that $D = GCG^*$.*

Proof. *i) \rightarrow ii)* From Remark 3.2 we know that $D - C \in \mathfrak{S}$ implies $P_{N(D)} - P_{N(C)} \in \mathfrak{S}$. By the assumption on the essential codimension of these projections and Remark 2.1, we have $UP_{N(D)}U^* = P_{N(C)}$ for some $U \in \mathcal{U}_{\mathfrak{S}}$. Notice that $UDU^* - C \in \mathfrak{S}$ and $N(UDU^*) = N(C)$; by Corollary 4.5 we conclude that $UD^{1/2}U^* - C^{1/2} \in \mathfrak{S}$. From the last condition, and noting that $N(UD^{1/2}U^*) = N(UDU^*) = N(C) = N(C^{1/2})$, it follows that there exists $G \in \mathcal{G}\ell_{\mathfrak{S}}$ such that $GC^{1/2} = UD^{1/2}U^*$ (see Remark 3.7). Hence, we find that $(U^*G)C(U^*G)^* = D$, where $U^*G \in \mathcal{G}\ell_{\mathfrak{S}}$.

ii) \rightarrow i) Clearly, we have $D \in \mathcal{CR}^+$. Using that $G \in \mathcal{G}\ell_{\mathfrak{S}}$, it follows that $D - C = GCG^* - C \in \mathfrak{S}$. From $D^* = GCG^*$ we have $G(N(C)^\perp) = G(R(C^*)) = R(D^*) = N(D)^\perp$. By Lemma 3.16 there is a unitary $U_G \in \mathcal{U}_{\mathfrak{S}}$ such that $U_G P_{N(C)} U_G^* = P_{N(D)}$. Then, again by Remark 2.1, we get $[P_{N(D)} : P_{N(C)}] = 0$ \square

For an operator $C \in \mathcal{CR}^+$, since $R(C)^\perp = N(C)$, the three dimensions used in Notation 3.8 reduce to two dimensions. That is, we have $\mathbb{J}_C = \{k \in \mathbb{Z} : -\dim(N(C)) \leq k \leq \dim(N(C)^\perp)\}$. The following result is an analog of Theorem 3.9 for positive perturbations.

Theorem 4.10. *Let $C \in \mathcal{CR}^+$ and \mathfrak{S} be a symmetrically-normed ideal. Then*

$$\mathcal{CR}^+ \cap (C + \mathfrak{S}) = \bigcup_{k \in \mathbb{J}_C} \mathcal{P}_k(C)$$

where

$$\mathcal{P}_k(C) = \{B \in \mathcal{CR}^+ : B - C \in \mathfrak{S}, [P_{N(B)} : P_{N(C)}] = k\}.$$

The following assertions hold:

- i) The action $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{CR}^+ \cap (C + \mathfrak{S}) \rightarrow \mathcal{CR}^+ \cap (C + \mathfrak{S})$, $G \cdot B = GBG^*$ restricted to $\mathcal{P}_k(C)$ is well defined and transitive. In other words, $\mathcal{P}_k(C) = \{GB^{(k)}G^* : G \in \mathcal{G}\ell_{\mathfrak{S}}\}$ for any $B^{(k)} \in \mathcal{P}_k(C)$.*
- ii) $\mathcal{P}_k(C)$ is a real analytic homogeneous space of $\mathcal{G}\ell_{\mathfrak{S}}$ and a submanifold of $C + \mathfrak{S}$, whose tangent space at $B \in \mathcal{P}_k(C)$ is $(T\mathcal{P}_k(C))_B = \{XB + BX^* : X \in \mathfrak{S}\} \subset \mathfrak{S} = (T(C + \mathfrak{S}))_B$.*
- iii) $\mathcal{P}_k(C)$ endowed with the previous differential structure is also a submanifold of $\mathcal{C}_k(C)$.*

Proof. We write $\mathcal{P}_k = \mathcal{P}_k(C)$ for short. Analogously to the proof of Theorem 3.9 one can see that $\mathcal{CR}^+ \cap (C + \mathfrak{S})$ is decomposed as the above disjoint union, and each $\mathcal{P}_k \neq \emptyset$ for $k \in \mathbb{J}_C$.

i) This follows by elementary properties of the essential codimension.

ii) We only treat the case $k = 0$ and take $B^{(0)} = C$. We first show that the isotropy of the action at C given by $\mathcal{G}_C := \{G \in \mathcal{G}\ell_{\mathfrak{S}} : GCG^* = C\}$ is a Banach-Lie subgroup of $\mathcal{G} := \mathcal{G}\ell_{\mathfrak{S}}$. Using a matrix representation with respect to $\mathcal{H} = R(C) \oplus R(C)^\perp$ we rewrite this group as

$$\mathcal{G}_C = \left\{ \begin{pmatrix} G_{11} & G_{12} \\ 0 & G_{22} \end{pmatrix} \in \mathcal{G}\ell_{\mathfrak{S}} : G_{11}C = C(G_{11}^*)^{-1} \right\},$$

which has its Lie algebra given by

$$\begin{aligned}\mathfrak{g}_C &= \{X \in \mathfrak{S} : XC = -CX^*\} \\ &= \left\{ \begin{pmatrix} X_{11} & X_{12} \\ 0 & X_{22} \end{pmatrix} \in \mathfrak{S} : X_{11}C = -CX_{11}^* \right\}.\end{aligned}$$

For every neighborhood \mathcal{T} of $0 \in \mathfrak{h}$, one can show that $\exp_{\mathcal{G}}(\mathcal{T})$ is a neighborhood of $I \in \mathcal{G}_C$ by using the logarithm series as in Lemma 3.13. To see that \mathfrak{g}_C is complemented in $\mathfrak{g} := \mathfrak{S}$, we put $P = P_{R(C)}$ and define the closed real subspaces

$$\mathfrak{n}^{\pm} = \{X \in P\mathfrak{S}P : X^* = \pm C^{\dagger}XC\}.$$

Since every $X \in P\mathfrak{S}P$ can be written as $X = X_+ + X_-$, where $X_+ = \frac{1}{2}(X^* + CXC^{\dagger})$ and $X_- = \frac{1}{2}(X^* - CXC^{\dagger})$, we have $P\mathfrak{S}P = \mathfrak{n}^+ \oplus \mathfrak{n}^-$. Therefore,

$$\mathfrak{m} = \left\{ \begin{pmatrix} X_{11} & 0 \\ Y & 0 \end{pmatrix} : X_{11} \in \mathfrak{n}^+, Y \in P^{\perp}\mathfrak{S}P \right\}.$$

is a closed supplement of \mathfrak{g}_C in \mathfrak{S} . This shows that \mathcal{G}_C is a Banach-Lie subgroup of \mathcal{G} , and consequently, $\mathcal{P}_0 \simeq \mathcal{G}/\mathcal{G}_C$ inherits the structure of real analytic homogeneous space of \mathcal{G} from the quotient space $\mathcal{G}/\mathcal{G}_C$. In particular, the tangent space $(T\mathcal{P}_0)_C$ can be identified with the quotient space $\mathfrak{g}/\mathfrak{g}_C = \{XC + CX^* : X \in \mathfrak{S}\}$.

Now we show that \mathcal{P}_0 with its homogeneous structure is also a real submanifold of $C + \mathfrak{S}$. We proceed as in Theorem 3.14. Notice that the map $\pi_0 : \mathcal{G}\ell_{\mathfrak{S}} \rightarrow \mathcal{P}_0$, $\pi_0(G) = GCG^*$ admits continuous local cross sections. Indeed, let $B \in \mathcal{P}_0$, $P = P_{R(C)}$, $Q = P_{R(B)}$; notice that $Q = Q(B)$ is a continuous function of $B \in \mathcal{P}_0$ since $Q = BB^{\dagger}$ and the continuity of the Moore-Penrose inverse on $\mathcal{C}_0(C)$ shown in Theorem 3.4 (notice that $\mathcal{P}_0 \subset \mathcal{C}_0(C)$ with the same distance function $d_{\mathfrak{S}}$). Set $S = QP + (I - Q)(I - P)$ and notice that $SP = QS$ so then $S(I - P) = (I - Q)S$. It can be checked that S is invertible when $\|C - B\|_{\mathfrak{S}} < \gamma$, for sufficiently small $\gamma > 0$; in this case, $SPS^{-1} = Q$ and $S(I - P)S^{-1} = (I - Q)$. Then, we define the map

$$\sigma(B) = B^{1/2}S(C^{\dagger})^{1/2} + (I - Q)S(I - P).$$

It follows by construction that $\sigma(B) \in \mathcal{G}\ell_{\mathfrak{S}}$ is such that $\pi_0(\sigma(B)) = B$ for $B \in \mathcal{P}_0$ such that $\|C - B\|_{\mathfrak{S}} < \gamma$. Moreover, σ is continuous by the continuity of the square root with respect to the metric $d_{\mathfrak{S}}$ (see Corollary 4.12).

It remains to show that the tangent space $(T\mathcal{P}_0)_C = \{XC + CX^* : X \in \mathfrak{S}\}$ is a closed complemented subspace in \mathfrak{S} . We may use again the above matrix representation,

$$XC + CX^* = \begin{pmatrix} X_{11}C + CX_{11}^* & CX_{21}^* \\ X_{21}C & 0 \end{pmatrix}.$$

From this expression, it suffices to show that $\Sigma = \{XC + CX^* : X \in P\mathfrak{S}P\}$ is closed and complemented in $P\mathfrak{S}P$. Consider the bounded and invertible operator $\mathcal{R}_T : P\mathfrak{S}P \rightarrow P\mathfrak{S}P$, $\mathcal{R}_T(Y) = YT$, for fixed $T \in \mathcal{B}(\mathcal{H})$ with $R(T) = N(T)^{\perp} = R(C)$. Since $\mathcal{R}_{C^{\dagger}}(\Sigma) = \mathfrak{n}^+$ we conclude that $\Sigma = \mathcal{R}_C(\mathfrak{n}^+)$ is closed and such that $\Sigma \oplus \mathcal{R}_C(\mathfrak{n}^-) = P\mathfrak{S}P$. This completes the proof.

iii) Again, we only consider the case $k = 0$ and set $\mathcal{C}_0 = \mathcal{C}_0(C)$. Notice that since both \mathcal{P}_0 and \mathcal{C}_0 are submanifolds of $C + \mathfrak{S}$, so their topologies coincide and the inclusion map $\iota : \mathcal{P}_0 \rightarrow \mathcal{C}_0$ is real analytic. We now show that this map is an immersion. For it is enough to see that the image under the tangent map $T_C\iota[(T\mathcal{P}_0)_C] = (T\mathcal{P}_0)_C$ is a closed complemented subspace of $(T\mathcal{C}_0)_C$. By inspection of the proof of Theorem 3.14 (see Eq. (7)) we have that

$$(T\mathcal{C}_0)_C = \left\{ \begin{pmatrix} Z_{11} & Z_{12} \\ Z_{21} & 0 \end{pmatrix} : Z_{11} \in P\mathfrak{S}P, Z_{12} \in P\mathfrak{S}P^{\perp}, Z_{21} \in P^{\perp}\mathfrak{S}P \right\},$$

where $P = P_{R(C)} = P_{N(C)^\perp}$ as before. Thus, from the proof of item *ii*) above we see that

$$\left\{ \begin{pmatrix} Z_{11} & 0 \\ 0 & 0 \end{pmatrix} : Z_{11} \in \mathcal{R}_C(\mathfrak{n}^-) \right\} \subset (TC_0)_C$$

is a complement for $(TP_0)_C$ inside $(TC_0)_C$. Thus, by [46, Prop. 8.7] we have that $\mathcal{P}_0 \subset \mathcal{C}_0$ is a real analytic submanifold. \square

Remark 4.11. We observe that in the context of C^* -algebras the constructions of continuous local cross sections for the action on congruence orbits can be given when $\dim(N(C)) < \infty$ or $\dim(R(C)) < \infty$ (see [25, Thm. 3.4]). The fact that the essential codimension is fixed is the key for the construction of continuous local cross sections in our previous proof.

Now that we have a manifold structure for $\mathcal{P}_k(C)$ we can complement Theorem 4.6 by showing the smoothness of the functional calculus by operator monotone functions for the more general domain $\mathcal{P}_k(C)$. This result will be needed to prove Theorem 4.17 below.

Corollary 4.12. *Let $f : [0, \infty) \rightarrow \mathbb{R}$ be an operator monotone function on $[0, \infty)$, let $C \in \mathcal{CR}^+$ and \mathfrak{S} be a symmetrically-normed ideal. Then, $f : \mathcal{P}_k(C) \rightarrow f(C) + \mathfrak{S}_{sa}$ is a real analytic function.*

Proof. Without loss of generality we can assume that $k = 0$ and check that f is real analytic in a neighborhood of $C \in \mathcal{P}_0 = \mathcal{P}_0(C)$. By Theorem 4.10 we get that the structure of \mathcal{P}_0 as a submanifold of $C + \mathfrak{S}$ coincides with its structure as an homogeneous space of $\mathcal{G}\ell$. Hence, it suffices to show that the map $\mathcal{G}\ell_{\mathfrak{S}} \ni G \mapsto f(GCG^*)$ is real analytic. By Lemma 3.16 there exists a real analytic function $\mathcal{G}\ell_{\mathfrak{S}} \ni G \mapsto U_G \in \mathcal{U}_{\mathfrak{S}}$ such that $U_G(R(C)) = G(R(C))$; thus, the map $\mathcal{G}\ell_{\mathfrak{S}} \ni G \mapsto U_G^*(GCG^*)U_G$ is real analytic and such that $R(U_G^*(GCG^*)U_G) = R(C)$, for $G \in \mathcal{G}\ell_{\mathfrak{S}}$. Hence, we can consider the matrix representation of $U_G^*(GCG^*)U_G$ with respect to the decomposition $\mathcal{H} = R(C) \oplus N(C)$ given by

$$\begin{pmatrix} U_G^*(GCG^*)U_G|_{R(C)} & 0 \\ 0 & 0 \end{pmatrix}.$$

Moreover, $U_G^*(GCG^*)U_G|_{R(C)} \in (C|_{R(C)} + \mathfrak{S}) \cap \mathcal{G}\ell(R(C))^+$, for $G \in \mathcal{G}\ell_{\mathfrak{S}}$. Therefore, by Theorem 4.6 we get that $f(U_G^*(GCG^*)U_G|_{R(C)}) \in f(C|_{R(C)}) + \mathfrak{S}_{sa}$ and that the map

$$\mathcal{G}\ell_{\mathfrak{S}} \ni G \mapsto f(U_G^*(GCG^*)U_G) = \begin{pmatrix} f(U_G^*(GCG^*)U_G|_{R(C)}) & 0 \\ 0 & f(0) \end{pmatrix} \in f(C) + \mathfrak{S}_{sa}$$

is real analytic, where we have also considered the decomposition $\mathcal{H} = R(C) \oplus N(C)$ above. Finally, notice that $\mathcal{G}\ell_{\mathfrak{S}} \ni G \mapsto f(GCG^*) = U_G f(U_G^*(GCG^*)U_G) U_G^*$ is real analytic, because it is the composition of real analytic maps. \square

4.3 Operator modulus and polar factor as real analytic fiber bundles

We apply the previous results to study the maps given by the polar factor and the operator modulus, defined on $\mathcal{C}_k(A)$, for $A \in \mathcal{CR}$. We remark the well-known fact that the essential codimension is also helpful for the analysis of perturbations of partial isometries by symmetrically-normed ideals; we collect several results from the literature in the following remark.

Remark 4.13. For a fixed $V \in \mathcal{PI} \subset \mathcal{CR}$ we consider below the set \mathbb{J}_V previously defined (Notation 3.8). The essential codimension can be used to write $\mathcal{PI} \cap (V + \mathfrak{S})$ as the union of connected components:

$$\mathcal{PI} \cap (V + \mathfrak{S}) = \bigcup_{k \in \mathbb{J}_V} \mathcal{V}_k(V)$$

where $\mathcal{V}_k(V) = \{X \in \mathcal{PI} : X - V \in \mathfrak{S}, [P_{N(X)} : P_{N(V)}] = k\} \neq \emptyset$. The action $(\mathcal{U}_{\mathfrak{S}} \times \mathcal{U}_{\mathfrak{S}}) \times \mathcal{PI} \rightarrow \mathcal{PI}$, $(U, W) \cdot V = UVW^*$ leaves invariant each $\mathcal{V}_k(V)$, and moreover, it holds

$$\mathcal{V}_k(V) = \{UV^{(k)}W^* : U, W \in \mathcal{U}_{\mathfrak{S}}\},$$

where $V^{(k)}$ is any partial isometry in $\mathcal{V}_k(V)$. These facts were proved in [20, Prop 3.5] for the ideal of Hilbert-Schmidt operators. The same proofs can be carried out for arbitrary symmetrically-normed ideals. In addition, we recall that $\mathcal{V}_k(V)$ are real analytic homogeneous spaces of the group $\mathcal{U}_{\mathfrak{S}} \times \mathcal{U}_{\mathfrak{S}}$, and submanifolds of $V + \mathfrak{S}$ (see [19]).

Lemma 4.14. *Let $A \in \mathcal{CR}$ with polar decomposition $A = V_A|A|$, $k \in \mathbb{J}_A$ and \mathfrak{S} be a symmetrically-normed ideal. The following assertions hold:*

i) $\alpha : \mathcal{C}_k(A) \rightarrow \mathcal{P}_k(|A|)$, $\alpha(B) = |B|$, is well defined and surjective.

ii) $v : \mathcal{C}_k(A) \rightarrow \mathcal{V}_k(V_A)$, $v(B) = V_B$, is well defined and surjective.

Proof. Without loss of generality we can assume that $k = 0$.

i) If $B \in \mathcal{C}_0(A)$, then $A - B \in \mathfrak{S}$ and $[P_{N(A)} : P_{N(B)}] = 0$. Since $A - B \in \mathfrak{S}$ then $A^*A - B^*B \in \mathfrak{S}$ and since $N(A^*A) = N(A)$ and $N(B^*B) = N(B)$ then $B^*B \in \mathcal{P}_0(A^*A)$. By Corollary 4.12 we see that $|B| - |A| = (B^*B)^{1/2} - (A^*A)^{1/2} \in \mathfrak{S}$, since $f(x) = x^{1/2}$ is an operator monotone function on $[0, \infty)$. Since $N(|B|) = N(B^*B)$ and $N(|A|) = N(A^*A)$ we get that $\alpha(B) = |B| \in \mathcal{P}_0(|A|)$. If $C \in \mathcal{P}_0(|A|)$ then, by item i) in Theorem 4.10 we see that there exists $G \in \mathcal{GL}_{\mathfrak{S}}$ such that $C = G|A|G^*$. By Lemma 3.16 there exists $U \in \mathcal{U}_{\mathfrak{S}}$ such that $U(R(|A|)) = G(R(|A|)) = R(G|A|)$. If we let $B = (V_A U)C = (V_A U^*)G|A|G^*$, then $V_B = V_A U$ and $|B| = G|A|G^* = C$ are the polar factor and the modulus corresponding to the polar decomposition of B . In particular, $V_B - V_A \in \mathfrak{S}$ and $|B| - |A| \in \mathfrak{S}$; then, $A - B = V_A|A| - V_B|B| = V_A(|A| - |B|) + (V_A - V_B)|B| \in \mathfrak{S}$ and moreover, $R(B) = R(A)$. Therefore, $B \in \mathcal{C}_0(A)$ is such that $\alpha(B) = C$.

ii) If $B \in \mathcal{C}_0(A)$, then by the first part of the proof, we have that $|B| \in \mathcal{P}_0(|A|)$. In particular, $|A| - |B| \in \mathfrak{S}$ and then $|A|^\dagger - |B|^\dagger \in \mathfrak{S}$, by Lemma 3.2. Hence, $V_A - V_B = A|A|^\dagger - B|B|^\dagger = A(|A|^\dagger - |B|^\dagger) + (A - B)|B|^\dagger \in \mathfrak{S}$. On the other hand, observe that $0 = [P_{N(A)} : P_{N(B)}] = [P_{N(V_A)} : P_{N(V_B)}]$ so $v(B) = V_B \in \mathcal{V}_0(V_A)$. If $V \in \mathcal{V}_0(V_A)$ then, by Remark 4.13, we see that there exist $U, W \in \mathcal{U}_{\mathfrak{S}}$ such that $V = UV_A W$. If we set $B = UV_A W(W^*|A|W)$, then $V_B = UV_A W = V$ and $|B| = W^*|A|W$ are the polar factor and the operator modulus corresponding to the polar decomposition of B . Since $\mathcal{U}_{\mathfrak{S}} \subset \mathcal{GL}_{\mathfrak{S}}$ then $B = UAW \in \mathcal{C}_0(A)$ is such that $v(B) = V_B = V$. \square

Remark 4.15. We remark that since $v(B) = V_B = B|B|^\dagger$ then, using similar arguments to those considered for the Moore-Penrose map and Corollary 4.5, we can show that $\mathcal{C}_k(A)$ are maximal subsets (of the metric space $(\mathcal{CR} \cap (A + \mathfrak{S}), d_{\mathfrak{S}})$) in which the polar factor becomes a continuous map, for $k \in \mathbb{J}_A$.

Under the same notation of Lemma 4.14 we now describe the structure of the fibers.

Lemma 4.16. *Given $C_0 \in \mathcal{P}_k(|A|)$ and $V_0 \in \mathcal{V}_k(V_A)$, then*

i) $\alpha^{-1}(C_0) := \{VC_0 \in \mathcal{C}_k : V \in \mathcal{V}_k(V_A), N(V) = N(C_0)\};$

ii) $v^{-1}(V_0) = \{V_0 C \in \mathcal{C}_k : C \in \mathcal{P}_k(|A|), N(C)^\perp = R(C) = N(V_0)^\perp\}.$

Furthermore, both fibers are submanifolds of $A + \mathfrak{S}$.

Proof. We may assume that $k = 0$.

i) Clearly, we have

$$\begin{aligned} \alpha^{-1}(C_0) &= \{VC_0 \in \mathcal{C}_0 : V \in \mathcal{V}_0(V_A), N(V) = N(C_0)\} \\ &\simeq \{V \in \mathcal{PI} : V - V_0 \in \mathfrak{S}, N(V) = N(V_0)\}, \end{aligned} \tag{15}$$

where V_0 is any partial isometry satisfying $N(V_0) = N(C_0)$ and $V_0 \in \mathcal{V}_0(V_A)$. The set in Eq. (15) is a submanifold of $V_A + \mathfrak{S}$ (see [19, Corol. 3.5]). The bijection above is given by $VC_0 \mapsto V$, and it induces a manifold structure on $\alpha^{-1}(C_0)$. We now prove that $\alpha^{-1}(C_0)$ with the aforementioned manifold structure is a submanifold of $A + \mathfrak{S}$. For we first observe that it is not difficult to see $\alpha^{-1}(C_0)$ has the topology defined by the metric $d_{\mathfrak{S}}$, and the inclusion map $\iota : \alpha^{-1}(C_0) \rightarrow A + \mathfrak{S}$ is real analytic. To prove that tangent spaces of $\alpha^{-1}(C_0)$ are closed and complemented in \mathfrak{S} , we identify the tangent space at VC_0 as

$$(T\alpha^{-1}(C_0))_{VC_0} = \{XVC_0 : X \in \mathfrak{S}_{ah}\}.$$

Without loss of generality, we now assume that $A = VC_0$. We can give the following alternative description of the tangent space

$$(T\alpha^{-1}(C_0))_A = \left\{ \begin{pmatrix} Z_{11} & 0 \\ Z_{21} & 0 \end{pmatrix} : Z_{11}A^\dagger \in P\mathfrak{S}_{ah}P, Z_{21} \in P^\perp\mathfrak{S}Q \right\},$$

where the above matrix decomposition is in terms of $Q = P_{N(A)^\perp}$ and $P = P_{R(A)}$ like in Eq. (7). So we only have to prove that $\Sigma := \{Z : ZA^\dagger \in P\mathfrak{S}_{ah}P\}$ is a closed complemented subspace of $P\mathfrak{S}Q$. Using the invertible map $\mathcal{R}_{A^\dagger} : P\mathfrak{S}Q \rightarrow P\mathfrak{S}P$, $\mathcal{R}_{A^\dagger}(Z) = ZA^\dagger$, this is equivalent to the fact that $P\mathfrak{S}_{ah}P$ is a (real) closed complemented subspace of $P\mathfrak{S}P$. Hence the inclusion map ι is an immersion, and $\alpha^{-1}(C_0)$ is a submanifold of $A + \mathfrak{S}$ (see [46, Prop. 8.7]).

ii) By Remark 4.13 there are unitaries $U, W \in \mathcal{U}_{\mathfrak{S}}$ such that $V_0 = UV_AW^*$. If $\mathcal{S} = N(V_0)^\perp$ and $E = P_{\mathcal{S}}$, then $W^*EW = P_{N(A)^\perp}$. Take the positive invertible operator $A_0 := W|A|W^*|_{\mathcal{S}} : \mathcal{S} \rightarrow \mathcal{S}$. Then the fiber can be computed as

$$v^{-1}(V_0) = \{V_0C \in \mathcal{C}_0(A) : C \in \mathcal{P}_0(|A|), N(C)^\perp = R(C) = \mathcal{S}\} \simeq \mathcal{P}_0(A_0). \quad (16)$$

Here we consider $\mathcal{P}_0(A_0)$ as a subset of the positive closed range operators acting on \mathcal{S} , and for its definition we use the ideal $\mathfrak{S}(\mathcal{S}) = \{EX|_{\mathcal{S}} : X \in \mathfrak{S}\} \simeq E\mathfrak{S}E$. The manifold structure of this fiber is induced by the bijection $v^{-1}(V_0) \simeq \mathcal{P}_0(A_0)$, which is given by the map $V_0C \mapsto C|_{\mathcal{S}}$. Using the characterization in Lemma 4.9 of $\mathcal{P}_0(|A|)$ and $\mathcal{P}_0(A_0)$ one can verify that this map is well defined, and it has the inverse $C_1 \mapsto V_0(C_1 \oplus 0)$. We follow similar steps to those of the previous item. Again the topology of $v^{-1}(V_0)$ is clearly given by the metric $d_{\mathfrak{S}}$, and the inclusion map $\iota : v^{-1}(V_0) \rightarrow A + \mathfrak{S}$ is real analytic. The tangent space at V_0C is given by

$$(Tv^{-1}(V_0))_{V_0C} = \{V_0(XC + CX^*) : X \in E\mathfrak{S}E\}.$$

For simplicity, we assume that $V_0C = A$. This implies that $V_0 = V_A$, $C = |A|$, $\mathcal{S} = N(A)^\perp$ and $E = Q = P_{N(A)^\perp}$. Next define the continuous invertible map $S_A : P\mathfrak{S}Q \rightarrow Q\mathfrak{S}Q$, $S_A(T) = V_A^*T|A|^\dagger$, which has the following property

$$S_A((Tv^{-1}(V_A))_A) = \{X + |A|X^*|A|^\dagger : X \in Q\mathfrak{S}Q\}.$$

The latter subspace we have already proved to be closed and complemented in $Q\mathfrak{S}Q$ (see the proof of Theorem 4.10 ii)). These facts show that the inclusion map is an immersion, and hence the fiber is a submanifold of $A + \mathfrak{S}$. \square

Next, we state our main result on the operator modulus and polar factor.

Theorem 4.17. *Let $A \in \mathcal{CR}$ with polar decomposition $A = V_A|A|$, $k \in \mathbb{J}_A$ and \mathfrak{S} be a symmetrically-normed ideal. The following assertions hold:*

i) $\alpha : \mathcal{C}_k(A) \rightarrow \mathcal{P}_k(|A|)$, $\alpha(B) = |B|$, is a real analytic fiber bundle.

ii) $v : \mathcal{C}_k(A) \rightarrow \mathcal{V}_k(V_A)$, $v(B) = V_B$, is a real analytic fiber bundle.

Proof. We may assume again that $k = 0$. Set $\mathcal{C}_0 = \mathcal{C}_0(A)$, $\mathcal{P}_0 = \mathcal{P}_0(|A|)$ and $\mathcal{V}_0 = \mathcal{V}_0(V_A)$.

i) In this case we show that α is a real analytic function in a neighborhood of $A \in \mathcal{C}_0(A)$. Since $\pi_0 : \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}} \rightarrow \mathcal{C}_0$ given by $\pi_0(G, K) = GAK^{-1}$ is a real analytic submersion, then it is enough to show that $\beta = \alpha \circ \pi_0 : \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}} \rightarrow \mathcal{P}_0$ is a real analytic function around the point $(I, I) \in \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}}$. We show that this is the case by expressing β as a composition of real analytic functions. Set $\mathcal{S} = N(A)^\perp$ and let $\mathcal{G}\ell_{\mathfrak{S}} \ni K \mapsto U_K \in \mathcal{U}_{\mathfrak{S}}$ be the real analytic map from Lemma 3.16, such that $U_K(N(A)^\perp) = K(N(A)^\perp)$. Then the map $\delta : \mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}} \rightarrow \mathcal{C}_0$ given by $\delta(G, K) = \pi_0(G, K)U_K = (GAK^{-1})U_K$ is a real analytic map. Notice that $\mathcal{G}\ell_{\mathfrak{S}} \times \mathcal{G}\ell_{\mathfrak{S}} \ni (G, K) \rightarrow |\delta(G, K)|^2$ is a real analytic map taking values in $\mathcal{P}_0(|A|^2)$. Indeed, $|(GAK^{-1})U_K|^2 = (K^{-1}U_K)^*|GA|^2K^{-1}U_K$ and $|GA|^2 = A^*|G|^2A$ show that $|GA|^2 - |A|^2 \in \mathfrak{S}$ and $N(|GA|) = N(|A|)$, so that by Lemma 4.9, we get $|(GAK^{-1})U_K|^2 \in \mathcal{P}_0(|GA|^2) = \mathcal{P}_0(|A|^2)$. By Corollary 4.12 applied to the function $f(\lambda) = \lambda^{1/2}$ we conclude that the map $(G, K) \mapsto |\delta(G, K)| = (U_K^*|GAK^{-1}|^2U_K)^{1/2} = U_K^*|GAK^{-1}|U_K$ is real analytic. Therefore, $\beta(G, K) = \alpha \circ \pi_0(G, K) = U_K|\delta(G, K)|U_K^*$ is a real analytic map. Hence, α is real analytic.

To prove that α is a real analytic fiber bundle, we also need to show here that v is a real analytic function. To this end notice that $v(B) = V_B = B|B|^\dagger$, for $B \in \mathcal{C}_0$. On the one hand, $\mathcal{C}_0 \ni B \mapsto |B| \in \mathcal{P}_0$ is real analytic by the first part of the proof. On the other hand, since $\mathcal{P}_0 \subset \mathcal{C}_0(|A|)$ is a submanifold by Theorem 4.10 then, $\mathcal{P}_0(|A|) \ni |B| \mapsto |B|^\dagger$ is a real analytic map, by Theorem 3.18. Hence, $v(B) = B|B|^\dagger$ is a real analytic map.

We now show that α is a real analytic fiber bundle. Recall that the fiber at $C_0 \in \mathcal{P}_0$ is given by $\alpha^{-1}(C_0) = \{VC_0 \in \mathcal{C}_0 : V \in \mathcal{V}_0, N(V) = N(C_0)\}$. We consider the action $\mathcal{G}\ell_{\mathfrak{S}} \ni G \mapsto GC_0G^* \in \mathcal{P}_0$ which, by Theorem 4.10, is a submersion. Hence, there exist an open set $\mathcal{W} \subset \mathcal{P}_0$ with $C_0 \in \mathcal{W}$, and an analytic map (cross-section) $\gamma : \mathcal{W} \rightarrow \mathcal{G}\ell_{\mathfrak{S}}$ such that $\gamma(C)C_0\gamma(C)^* = C$, for $C \in \mathcal{W}$ ([46, Corol. 8.3]). By Lemma 3.16 there exists an analytic map $\mathcal{W} \ni C \mapsto U_{\gamma(C)}$ such that $U_{\gamma(C)}(R(C_0)) = \gamma(C)(R(C_0)) = R(C)$. Since α and v are real analytic, and $\alpha^{-1}(C_0)$ is a submanifold of $A + \mathfrak{S}$ by Lemma 4.16, the map given by

$$\alpha^{-1}(\mathcal{W}) \rightarrow \mathcal{W} \times \alpha^{-1}(C_0), \quad B = V_B|B| \mapsto (|B|, (V_B U_{\gamma(|B|)})C_0)$$

is real analytic, and its inverse

$$\mathcal{W} \times \alpha^{-1}(C_0) \ni (C, VC_0) \mapsto VU_{\gamma(C)}^*C \in \alpha^{-1}(\mathcal{W})$$

is also real analytic. We have used that $\alpha((V_B U_{\gamma(|B|)})C_0) = C_0$ and $v((V_B U_{\gamma(|B|)})C_0) = V_B U_{\gamma(|B|)}$, for $B \in \alpha^{-1}(\mathcal{W})$.

ii) By the previous item, we already know that v is a real analytic map. Recall that for $V_0 \in \mathcal{V}_0(V_A)$, we have that $v^{-1}(V_0) = \{V_0C \in \mathcal{C}_0 : C \in \mathcal{P}_0, N(C)^\perp = R(C) = N(V_0)^\perp\}$. Consider the action $\mathcal{U}_{\mathfrak{S}} \times \mathcal{U}_{\mathfrak{S}} \ni (U, W) \mapsto UV_0W \in \mathcal{V}_0(V_A)$ which, by Remark 4.13, is a submersion. Then, there exist an open set $\mathcal{Y} \subset \mathcal{V}_0(V_A)$ with $V_0 \in \mathcal{Y}$ and a real analytic cross section $\eta : \mathcal{Y} \rightarrow \mathcal{U}_{\mathfrak{S}} \times \mathcal{U}_{\mathfrak{S}}$. So if $\eta(V) = (\eta_1(V), \eta_2(V))$, then $\eta_1(V)V_0\eta_2(V)^* = V$, for $V \in \mathcal{Y}$. Using that $v^{-1}(V_0)$ is a submanifold of $A + \mathfrak{S}$ by Lemma 4.16, the map given by

$$v^{-1}(\mathcal{Y}) \rightarrow \mathcal{Y} \times v^{-1}(V_0), \quad B = V_B|B| \mapsto (V_B, V_0(\eta_2(V_B)^*|B|\eta_2(V_B)))$$

is real analytic with inverse given by

$$\mathcal{Y} \times v^{-1}(V_0) \ni (V, B) \mapsto V\eta_2(V)|B|\eta_2(V)^* \in v^{-1}(\mathcal{Y})$$

which is also real analytic. □

Remark 4.18. We notice that the maps α and v in Theorem 4.17 satisfy some compatibility relations with respect to the Moore-Penrose inverse. Indeed, consider $A \in \mathcal{CR}$ and $k \in \mathbb{J}_A = -\mathbb{J}_{A^*}$. Let $\alpha_{A^*} : \mathcal{C}_{-k}(A^*) \rightarrow \mathcal{P}_{-k}(|A^*|)$ and $v_{A^*} : \mathcal{C}_{-k}(A^*) \rightarrow \mathcal{V}_{-k}(V_A^*)$ be defined as above, where $A^* = V_A^*|A^*|$ is the polar decomposition of A^* . We further consider $\alpha_{A^\dagger} : \mathcal{C}_k(A^\dagger) \rightarrow \mathcal{P}_k(|A^\dagger|)$ and $v_{A^\dagger} : \mathcal{C}_k(A^\dagger) \rightarrow \mathcal{V}_k(V_{A^\dagger})$. Let $\mu : \mathcal{C}_k(A) \rightarrow \mathcal{C}_k(A^\dagger)$ be given by $\mu(B) = B^\dagger$. Then, notice that for $B \in \mathcal{C}_k(A)$ we get that

$$\alpha_{A^\dagger}(\mu(B)) = |B^\dagger| = |B^*|^\dagger = \mu(\alpha_{A^*}(B^*)) \quad , \quad v_{A^\dagger}(\mu(B)) = V_B^* = \mu(v_{A^*}(B^*)) .$$

5 Appendix

In this section we present a proof of Lemma 4.2. Hence, we let \mathfrak{S} be a symmetrically-normed ideal, ν be a positive Borel measure on $(0, \infty)$ and we let $h : [0, \infty) \rightarrow \mathfrak{S}_{sa}$ be a function satisfying:

1. $\|h(t)\|_{\mathfrak{S}} \leq q(t)$, where $q(t)$ is a bounded, continuous, positive and non-increasing function such that $\int_0^\infty q(t) d\nu(t) < \infty$;
2. For $\delta > 0$ then $\|h(t + \delta) - h(t)\|_{\mathfrak{S}} \leq \alpha \delta r(t)$, where $\alpha > 0$ and $r : [0, \infty) \rightarrow [0, \infty)$ is a continuous and non-increasing function such that

$$\lim_{t \rightarrow \infty} r(t) = 0 \quad \text{and} \quad \int_0^\infty r(t) d\nu(t) < \infty .$$

Recall that the operator $\int_0^\infty h(t) d\nu(t) \in \mathcal{B}(\mathcal{H})_{sa}$ is determined by the identity

$$\left\langle \int_0^\infty h(t) d\nu(t) x, y \right\rangle = \int_0^\infty \langle h(t) x, y \rangle d\nu(t) \quad , \quad x, y \in \mathcal{H} .$$

Since $\|\cdot\| \leq \|\cdot\|_{\mathfrak{S}}$ then item 1 above shows that $\int_0^\infty \|h(t)\| d\nu(t) < \infty$. Then, the previous facts imply that for every $p \geq 1$,

$$\left\| \int_0^\infty h(t) d\nu(t) \right\| \leq \int_0^\infty \|h(t)\| d\nu(t) \quad , \quad \int_0^\infty h(t) d\nu(t) = \sum_{m=1}^\infty \int_{[0, \frac{1}{2^p})} h\left(\frac{m-1}{2^p} + t\right) d\nu(t) ,$$

where the series converges in the operator norm. We now introduce the sequence

$$R_p = \sum_{m=1}^\infty \nu \left(\left[\frac{m-1}{2^p}, \frac{m}{2^p} \right) \right) h\left(\frac{m}{2^p}\right) \in \mathfrak{S}_{sa} \quad , \quad p \geq 1 ,$$

where we have used that $\|h(t)\|_{\mathfrak{S}} \leq q(t)$ is a decreasing function,

$$\sum_{m=1}^\infty \nu \left(\left[\frac{m-1}{2^p}, \frac{m}{2^p} \right) \right) \|h\left(\frac{m}{2^p}\right)\|_{\mathfrak{S}} \leq \sum_{m=1}^\infty \nu \left(\left[\frac{m-1}{2^p}, \frac{m}{2^p} \right) \right) q\left(\frac{m}{2^p}\right) \leq \int_0^\infty q(t) d\nu(t) < \infty ,$$

so that the series defining R_p is absolutely convergent in \mathfrak{S} (and hence determines an element in \mathfrak{S}_{sa}). By item 2 above we get that for $m, p \geq 1$:

$$\left\| h\left(\frac{m}{2^p}\right) - h\left(\frac{m-1}{2^p} + t\right) \right\| \leq \left\| h\left(\frac{m}{2^p}\right) - h\left(\frac{m-1}{2^p} + t\right) \right\|_{\mathfrak{S}} \leq \frac{\alpha}{2^p} r\left(\frac{m-1}{2^p}\right) , \quad \text{for } t \in [0, \frac{1}{2^p}) , \quad (17)$$

where we have used that $\frac{m}{2^p} = \left(\frac{m-1}{2^p} + t\right) + \delta$, for some $\delta \in [0, 2^{-p})$, and that $r\left(\frac{m-1}{2^p} + t\right) \leq r\left(\frac{m-1}{2^p}\right)$, for $t \in [0, 2^{-p})$. By the same item we also get that there exists $\eta \geq 1$ such that for $p \geq 1$ we have that

$$\sum_{m=1}^\infty r\left(\frac{m-1}{2^p}\right) \nu\left(\left[\frac{m-1}{2^p}, \frac{m}{2^p} \right) \right) \leq \eta \int_0^\infty r(t) d\nu(t) . \quad (18)$$

Therefore, we get that

$$\left\| R_p - \int_0^\infty h(t) d\nu(t) \right\| \leq \frac{\alpha \cdot \eta}{2^p} \int_0^\infty r(t) d\nu(t) \xrightarrow{p \rightarrow \infty} 0. \quad (19)$$

We now show that $\{R_p\}_{p \geq 1}$ is a Cauchy sequence in \mathfrak{S} : indeed, for $p \geq 1$ we have that

$$R_p = \sum_{m=1}^{\infty} \left(\nu \left(\left[\frac{2m-2}{2^{p+1}}, \frac{2m-1}{2^{p+1}} \right) \right) + \nu \left(\left[\frac{2m-1}{2^{p+1}}, \frac{2m}{2^{p+1}} \right) \right) \right) h\left(\frac{2m}{2^{p+1}}\right)$$

and hence,

$$R_{p+1} - R_p = \sum_{m=1}^{\infty} \nu \left(\left[\frac{2m-2}{2^{p+1}}, \frac{2m-1}{2^{p+1}} \right) \right) \left(h\left(\frac{2m-1}{2^{p+1}}\right) - h\left(\frac{2m}{2^{p+1}}\right) \right).$$

The previous identity implies that

$$\|R_{p+1} - R_p\|_{\mathfrak{S}} \leq \sum_{m=1}^{\infty} \nu \left(\left[\frac{2m-2}{2^{p+1}}, \frac{2m-1}{2^{p+1}} \right) \right) \frac{\alpha}{2^{p+1}} r\left(\frac{2m-1}{2^{p+1}}\right) \leq \frac{\alpha \cdot \eta}{2^{p+1}} \int_0^\infty r(t) d\nu(t).$$

This last fact shows that $\{R_p\}_{p \geq 1}$ is a Cauchy sequence in \mathfrak{S} so that it converges to an operator $R \in \mathfrak{S}_{sa}$; but then, $\{R_p\}_{p \geq 1}$ also converges to R in the operator norm. The previous comments together with Eq. (19) imply that $R = \int_0^\infty h(t) d\nu(t) \in \mathfrak{S}_{sa}$. On the other hand, using Eq. (17) we have that

$$\left| \left\| h\left(\frac{m-1}{2^p} + \delta\right) \right\|_{\mathfrak{S}} - \left\| h\left(\frac{m}{2^p}\right) \right\|_{\mathfrak{S}} \right| \leq \left\| h\left(\frac{m-1}{2^p} + \delta\right) - h\left(\frac{m}{2^p}\right) \right\|_{\mathfrak{S}} \leq \frac{\alpha}{2^p} r\left(\frac{m-1}{2^p}\right), \quad \text{for } \delta \in \left[0, \frac{1}{2^p}\right).$$

Hence, using the previous fact and Eq. (18) we get that

$$\left| \int_0^\infty \|h(t)\| d\nu(t) - \sum_{m=1}^{\infty} \nu \left(\left[\frac{m-1}{2^p}, \frac{m}{2^p} \right) \right) \left\| h\left(\frac{m}{2^p}\right) \right\|_{\mathfrak{S}} \right| \leq \frac{\alpha \cdot \eta}{2^p} \int_0^\infty r(t) d\nu(t). \quad (20)$$

Since $\|R_p - \int_0^\infty h(t) d\nu(t)\|_{\mathfrak{S}} \rightarrow 0$ when $p \rightarrow \infty$ then, given $\epsilon > 0$ there exists $p_0 \geq 1$ such that if $p \geq p_0$ then $\|\int_0^\infty h(t) d\nu(t)\|_{\mathfrak{S}} \leq \|R_p\|_{\mathfrak{S}} + \epsilon$. On the other hand, Eq. (20) shows that there exists $p_1 \geq p_0$ such that for $p \geq p_1$ we get that

$$\left\| \int_0^\infty h(t) d\nu(t) \right\|_{\mathfrak{S}} \leq \|R_p\|_{\mathfrak{S}} + \epsilon \leq \sum_{m=1}^{\infty} \nu \left(\left[\frac{m-1}{2^p}, \frac{m}{2^p} \right) \right) \left\| h\left(\frac{m}{2^p}\right) \right\|_{\mathfrak{S}} + \epsilon \leq \int_0^\infty \|h(t)\|_{\mathfrak{S}} d\nu(t) + 2\epsilon.$$

Acknowledgment

This research was partially supported by CONICET (PIP 2021/2023 11220200103209CO), AN-PCyT (2015 1505/ 2017 0883) and FCE-UNLP (11X974).

References

- [1] W.O. Amrein, K.B. Sinha, On pairs of projections in a Hilbert space, *Linear Algebra Appl.* 208/209 (1994), 425-435.
- [2] E. Andruchow, Pairs of Projections: Geodesics, Fredholm and Compact Pairs. *Complex Anal. Oper. Theory* 8 (2014), 1435-1453.
- [3] E. Andruchow, E. Chiumiento, G. Larotonda, Geometric significance of Toeplitz kernels, *J. Funct. Anal.* (2018), 329-355.

- [4] E. Andruchow, G. Corach, M. Mbekhta, On the geometry of generalized inverses, *Math. Nachr.* 278 (2005), 756-770.
- [5] E. Andruchow, G. Larotonda, Hopf-Rinow theorem in the Sato Grassmannian. *J. Funct. Anal.* 255 (2008), no. 7, 1692-1712.
- [6] E. Andruchow, G. Larotonda, L. Recht, Finsler geometry and actions of the p -Schatten unitary groups, *Trans. Amer. Math. Soc.* 362 (2010), 319-344.
- [7] T. Ando, J.L. van Hemmen, An inequality for trace ideals, *Comm. Math. Phys.* 76 (1980), 143-148.
- [8] C. Apostol, The reduced minimum modulus. *Michigan Math. J.* 32 (1985), no. 3, 279-294.
- [9] J. Avron, R. Seiler, B. Simon, The index of a pair of projections, *J. Funct. Anal.* 120 (1994), no. 1, 220-237.
- [10] D. Beltiță, Smooth homogeneous structures in operator theory, Chapman and Hall/CRC, Monographs and Surveys in Pure and Applied Mathematics 137, Boca Raton, 2006.
- [11] D. Beltiță, G. Larotonda, Unitary Group Orbits Versus Groupoid Orbits of Normal Operators, *J. Geom. Anal.* 33 (2023), 95.
- [12] D. Beltiță, T. Goliński, G. Jakimowicz, F. Pelletier, Banach-Lie groupoids and generalized inversion. *J. Funct. Anal.* 276 (2019), no. 5, 1528-1574.
- [13] D. Beltiță, T.S. Ratiu, Symplectic leaves in real Banach Lie-Poisson spaces. *Geom. Funct. Anal.* 15 (2005), no. 4, 753-779.
- [14] E. Boasso, On the Moore-Penrose inverse in C^* -algebras. *Extracta Math.* 21 (2006), no. 2, 93-106.
- [15] L.G. Brown, R.G. Douglas, and P. A. Fillmore, Unitary equivalence modulo the compact operators and extensions of C^* -algebras, in: P. A. Fillmore (Ed.) *Lect. Not. Math.* 345 (Apr. 13, 1973) 58-128.
- [16] J.R. Cardoso, Computation of the matrix p -th root and its Fréchet derivative by integrals. *Electron. Trans. Numer. Anal.* 39 (2012), 414-436.
- [17] A.L. Carey, Some homogeneous spaces and representations of the Hilbert Lie group $U(H)_2$, *Rev. Roum. Math. Pures Appl.* 30 (1985), no. 7, 505-520.
- [18] G. Chen, M. Wei, Y. Xue, Perturbation Analysis of the Least Squares Solution in Hilbert Spaces, *Linear Algebra Appl.* 244 (1996), 69-80.
- [19] E. Chiumiento, Geometry of \mathcal{J} -Stiefel manifolds, *Proc. Amer. Math. Soc.* 138 (2010), no. 1, 341-353.
- [20] E. Chiumiento, Global symmetric approximation of frames, *J. Fourier Anal. Appl.* 25 (2019), no. 4, 1395-1423.
- [21] E. Chiumiento, P. Massey, On restricted diagonalization, *J. Funct. Anal.* 282 (2021), 109342.
- [22] E. Chiumiento, P. Massey, Restricted orbits of closed range operators and equivalences of frames for subspaces, submitted.
- [23] P. Del Moral, A. Niclas, A Taylor expansion of the square root matrix function. *J. Math. Anal. Appl.* 465 (2018), no. 1, 259-266.
- [24] G. Corach, A. Maestripieri, M. Mbekhta, Metric and homogeneous structure of closed range operators, *J. Operator Theory* 61 (2009), no. 1, 171-190.
- [25] G. Corach, A. Maestripieri, D. Stojanoff, Orbits of positive operators from a differentiable viewpoint, *Positivity* 8 (2004), no. 1, 31-48.
- [26] P. de la Harpe, Classical Banach-Lie Algebras and Banach-Lie Groups of Operators in Hilbert Space, *Lecture Notes in Mathematics*, Vol. 285. Springer-Verlag, Berlin-New York, 1972.
- [27] K. Dykema, T. Figiel, G. Weiss, M. Wodzicki, Commutator structure of operator ideals, *Adv. Math.* 185 (2004), no. 1, 1-79.
- [28] G.H. Golub, V. Pereyra, The differentiation of pseudo-inverses and nonlinear least squares problems whose variables separate. *SIAM J. Numer. Anal.* 10 (1973), 413-432.

- [29] M.C. Gouveia, R. Puystjens, About the group inverse and Moore-Penrose inverse of a product, *Linear Algebra Appl.* 150 (1991), 361-369.
- [30] S. Izumino, Convergence of generalized splines and splines projections, *J. Approx. Theory* 38 (1983), 269-278.
- [31] B. Jia, Compact perturbations of Moore-Penrose invertible operators. *J. Math. Anal. Appl.* 523 (2023), no. 2, Paper No. 127048, 10 pp.
- [32] V. Kaftal, J. Loreaux, Kadison's Pythagorean theorem and essential codimension, *Integral Equations Operator Theory* 87 (2017), no. 4, 565-580.
- [33] J.J. Koliha, Continuity and differentiability of the Moore-Penrose inverse in C^* -algebras, *Math. Scand.* 88 (2001), 154-160.
- [34] J.-Ph. Labrousse, M. Mbekhta, Point-of-continuity operators for the conorm and the Moore-Penrose inverse. *Houston J. Math.* 18 (1992), no. 1, 7-23.
- [35] G. Larotonda, Metric geometry of infinite dimensional Lie groups and their homogeneous spaces, *Forum Math.* 31 (2019) 1567-1605.
- [36] J. Leiterer, L. Rodman, Smoothness of generalized inverses. *Indag. Math. (N.S.)* 23 (2012), no. 3, 487-521.
- [37] J. Loreaux, Restricted diagonalization of finite spectrum normal operators and a theorem of Arveson, *J. Operator Theory* 81 (2019), no. 2, 257-272.
- [38] I.C. Gohberg, M.G. Krein, *Introduction to the Theory of Linear Non-Self-Adjoint Operators*, Amer. Math. Soc., Providence, RI, 1960.
- [39] K.-H. Neeb, Holomorphic highest weight representations of infinite dimensional complex classical groups, *J. Reine Angew. Math.* 497 (1998), 171-222.
- [40] K.-H. Neeb, *Infinite-dimensional groups and their representations. Lectures at the European School in Group Theory, SDU-Odense Univ., August 2000.*
- [41] T. Sano, Fréchet derivatives for operator monotone functions. *Linear Algebra Appl.* 456 (2014), 88-92.
- [42] B. Simon, *Trace Ideals and their Applications*, in: London Mathematical Society Lecture Note Series, vol. 35, Cambridge University Press, Cambridge, 1979.
- [43] B. Simon, Unitaries permuting two orthogonal projections, *Linear Algebra Appl.* 528 (2017), 436-441.
- [44] S. Strătilă, D. Voiculescu, On a class of KMS states for the group $U(\infty)$, *Math. Ann.* 235 (1978), 87-110.
- [45] G.W. Stewart, On the perturbation of pseudo-inverses, projections, and linear least squares problems, *SIAM Rev.* 19 (1977), 634-662.
- [46] H. Upmeyer, *Symmetric Banach Manifolds and Jordan C^* -Algebras*, North-Holland Math. Stud., vol. 104, Notas de Matemática, vol. 96, North-Holland, Amsterdam 1985.
- [47] P. Å. Wedin, Perturbation theory for pseudo-inverses, *BIT* 13 (1973), 217-232.

(EDUARDO CHIUMENTO) Departamento de Matemática & Centro de Matemática La Plata, FCE-UNLP, Calles 50 y 115, (1900) La Plata, Argentina and Instituto Argentino de Matemática, 'Alberto P. Calderón', CONICET, Saavedra 15, 3er. piso, (1083) Buenos Aires, Argentina.
e-mail: eduardo@mate.unlp.edu.ar

(PEDRO MASSEY) Departamento de Matemática & Centro de Matemática La Plata, FCE-UNLP, Calles 50 y 115, (1900) La Plata, Argentina and Instituto Argentino de Matemática, 'Alberto P. Calderón', CONICET, Saavedra 15, 3er. piso, (1083) Buenos Aires, Argentina.
e-mail: massey@mate.unlp.edu.ar