



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa


Abstract splines in Krein spaces

 J.I. Giribet^{a,c,1}, A. Maestriperi^{a,c,2}, F. Martínez Pería^{b,c,*,2}
^a Departamento de Matemática, FI-UBA, Paseo Colón 850, (1063) Buenos Aires, Argentina

^b Departamento de Matemática, FCE-UNLP, CC 172, (1900) La Plata, Argentina

^c Instituto Argentino de Matemática "Alberto Calderón" – CONICET, Saavedra 15, Piso 3, (1083) Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 13 October 2009

Available online 16 March 2010

Submitted by J.A. Ball

Keywords:

Krein spaces

Abstract splines

Oblique projections

ABSTRACT

We present generalizations to Krein spaces of the abstract interpolation and smoothing problems proposed by Atteia in Hilbert spaces: given a Krein space \mathcal{K} and Hilbert spaces \mathcal{H} and \mathcal{E} (bounded) surjective operators $T : \mathcal{H} \rightarrow \mathcal{K}$ and $V : \mathcal{H} \rightarrow \mathcal{E}$, $\rho > 0$ and a fixed $z_0 \in \mathcal{E}$, we study the existence of solutions of the problems $\operatorname{argmin}\{[Tx, Tx]_{\mathcal{K}} : Vx = z_0\}$ and $\operatorname{argmin}\{[Tx, Tx]_{\mathcal{K}} + \rho \|Vx - z_0\|_{\mathcal{E}}^2 : x \in \mathcal{H}\}$.

© 2010 Elsevier Inc. All rights reserved.

1. Introduction

Since I.J. Schoenberg introduced the spline functions [1], they have become an important notion in several branches of mathematics such as approximation theory, statistics, numerical analysis and partial differential equations, among others. Moreover, they have been useful to solve some practical issues in signal and image processing [2–5], computer graphics [6–8], learning theory [9,10] and other applications.

In the sixties, a Hilbert space formulation of spline functions, known as abstract splines, was introduced by M. Atteia [11] and developed by several authors, see for instance [12–15]. Given Hilbert spaces \mathcal{H} , \mathcal{K} and \mathcal{E} , consider (bounded) surjective operators $T : \mathcal{H} \rightarrow \mathcal{K}$ and $V : \mathcal{H} \rightarrow \mathcal{E}$. The abstract interpolation problem in Hilbert spaces can be stated as follows: for a fixed $z_0 \in \mathcal{E}$, find $x_0 \in \mathcal{H}$ such that $Vx_0 = z_0$ and

$$\|Tx_0\|_{\mathcal{K}}^2 = \min\{\|Tx\|_{\mathcal{K}}^2 : Vx = z_0\}. \quad (1)$$

Observe that $x_0 \in V^{-1}(\{z_0\})$ is an abstract interpolating spline (i.e. x_0 satisfies Eq. (1)) if and only if Tx_0 realizes the distance between $TV^\dagger z_0$ and the subspace $T(N(V))$, where V^\dagger stands for the Moore–Penrose inverse of V . So, the existence of x_0 depends on the existence of a suitable (contractive) projection of $TV^\dagger z_0$ onto $T(N(V))$. Then, if $T(N(V))$ is a closed subspace of \mathcal{K} , the existence of x_0 is guaranteed because the selfadjoint projection onto $T(N(V))$ is always contractive.

On the other hand, the abstract smoothing problem introduces a new parameter $\rho > 0$ in order to balance the amounts $\|Tx\|_{\mathcal{K}}^2$ and $\|Vx - z_0\|_{\mathcal{E}}^2$. Formally, given $\rho > 0$ and a fixed $z_0 \in \mathcal{E}$, it consists in minimizing the function $F_\rho : \mathcal{H} \rightarrow \mathbb{R}^+$ defined by

$$F_\rho(x) = \|Tx\|_{\mathcal{K}}^2 + \rho \|Vx - z_0\|_{\mathcal{E}}^2. \quad (2)$$

* Corresponding author at: Instituto Argentino de Matemática "Alberto Calderón", Saavedra 15, Piso 3, (1083) Buenos Aires, Argentina.

E-mail addresses: jgiribet@fi.uba.ar (J.I. Giribet), amaestri@fi.uba.ar (A. Maestriperi), francisco@mate.unlp.edu.ar (F.M. Pería).

¹ Partially supported by ANPCyT 1728 PICT06, PUNQ 0530/07, FONCYT 808.

² Partially supported by FONCYT 808.

This problem can be reduced to a least squares problem. In fact,

$$F_\rho(x) = \|Lx - (0, z_0)\|_\rho^2,$$

where $\|\cdot\|_\rho$ is the norm associated to the inner product on $\mathcal{K} \times \mathcal{E}$ defined by $\langle (y, z), (y', z') \rangle_\rho = \langle y, y' \rangle_{\mathcal{K}} + \rho \langle z, z' \rangle_{\mathcal{E}}$ if $(y, z), (y', z') \in \mathcal{K} \times \mathcal{E}$, and L is an auxiliary operator from \mathcal{H} into $\mathcal{K} \times \mathcal{E}$. Therefore, the abstract smoothing problem is also related to the existence of a selfadjoint (contractive) projection onto $R(L)$.

There is also a variational problem which mixes both abstract interpolation and smoothing problems. In the abstract mixed problem (as it is known) the “measurement operator” $V : \mathcal{H} \rightarrow \mathcal{E}$ splits up into two surjective operators. The technique used by A. Rozhenko [16] to solve this problem is similar to the one mentioned above to solve the abstract smoothing problem. So, the existence of “abstract mixed splines” also depends on the existence of a suitable contractive projection.

For a complete exposition on these subjects see the books by Atteia [17], A. Bezhaev and V. Vasilenko [18], and the survey by R. Champion et al. [19].

In this work, mainly motivated by the ideas exposed above, we present generalizations of the abstract interpolation, smoothing and mixed problems to Krein spaces. As we have mentioned before, the techniques used to solve these problems in the Hilbert space setting, involved contractive projections onto some subspaces. So, they (or their complementary subspaces) are asked to be closed. In order to reproduce this geometrical approach for Krein spaces, the hypothesis on the subspaces has to be modified. Recall that to guarantee the existence of a selfadjoint projection onto a subspace of a Krein space, it has to be regular. Moreover, if the projection has to be contractive then its nullspace has to be uniformly J -positive, where J stands for the fundamental symmetry of the Krein space (see [20]).

First, we study the indefinite abstract interpolation problem. Specifically, if \mathcal{K} is a Krein space and \mathcal{E} and \mathcal{H} are Hilbert spaces, given (bounded) surjective operators $T : \mathcal{H} \rightarrow \mathcal{K}$ and $V : \mathcal{H} \rightarrow \mathcal{E}$ and a fixed $z_0 \in \mathcal{E}$, we are interested in characterizing (if there is any) those $x_0 \in \mathcal{H}$ such that $Vx_0 = z_0$ and

$$[Tx_0, Tx_0]_{\mathcal{K}} = \min\{[Tx, Tx]_{\mathcal{K}} : Vx = z_0\}.$$

Using a similar argument as in the definite interpolation problem, it can be shown that the existence of x_0 depends on the existence of a suitable (contractive) projection of $TV^\dagger z_0$ onto the J -orthogonal companion of $T(N(V))$ in \mathcal{K} . Then, if $T(N(V))$ is a closed uniformly J -positive subspace of \mathcal{K} , the existence of x_0 is guaranteed.

On the other hand, in the indefinite abstract smoothing problem, we look for the minimizers of the function $F_\rho : \mathcal{H} \rightarrow \mathbb{R}$ defined by

$$F_\rho(x) = [Tx, Tx]_{\mathcal{K}} + \rho \|Vx - z_0\|_{\mathcal{E}}^2, \quad x \in \mathcal{H}. \tag{3}$$

This problem can be no longer restated as a least squares problem, but as an indefinite least squares problem. The technique used to describe its solutions is similar to the one used in the definite smoothing problem, but a particular orthogonal decomposition of the range of a given operator is needed.

The last problem we consider is the indefinite abstract mixed problem. If \mathcal{K} is a Krein space and $\mathcal{E}_1, \mathcal{E}_2$ and \mathcal{H} are Hilbert spaces, consider (bounded) surjective operators $T : \mathcal{H} \rightarrow \mathcal{K}$, $V_1 : \mathcal{H} \rightarrow \mathcal{E}_1$ and $V_2 : \mathcal{H} \rightarrow \mathcal{E}_2$. Given $\rho > 0$ and a fixed $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$, we look for those $x_0 \in \mathcal{H}$ such that $V_1 x_0 = z_1$ which are minimizers of the function

$$G_\rho(x) = [Tx, Tx]_{\mathcal{K}} + \rho \|V_2 x - z_2\|_{\mathcal{E}_2}^2, \quad x \in V_1^{-1}(\{z_1\}). \tag{4}$$

Spline functions in indefinite metric spaces have already been studied in [9] to solve numerical aspects related to learning theory problems. Although the problems presented there are different from those studied in this work, they are closely related. In [10] another version of the abstract indefinite smoothing problem is studied: given $z_0 \in \mathcal{E}$, instead of finding the minimum of the function F_ρ given in Eq. (3), the authors are interested in stabilizing it.

The paper is organized as follows: Section 2 contains the preliminaries. In Section 3 we study the indefinite abstract interpolation problem, we give necessary and sufficient conditions for the existence (and uniqueness) of solutions of this problem, and characterize them. Also, given a frame $\{f_n\}_{n \in \mathbb{N}}$ for the Hilbert space \mathcal{E} , we give conditions to obtain different frames for subspaces of \mathcal{H} composed by interpolating splines corresponding to the family $\{f_n\}_{n \in \mathbb{N}}$.

Section 4 is devoted to the study of the indefinite abstract smoothing problem: after characterizing its set of solutions (for a fixed ρ), we show that it is related to the set of solutions of an indefinite interpolation problem for a certain $z_\rho \in \mathcal{E}$. Then, as it was studied by Atteia in Hilbert spaces, we analyze the convergence of the solutions of the indefinite smoothing problem to the solutions of the indefinite interpolation problem as ρ goes to infinity.

In Section 5 the abstract mixed problem studied by A. Rozhenko and V. Vasilenko [16,21,22], is extended to Krein spaces.

2. Preliminaries

Along this work \mathcal{E} denotes a complex (separable) Hilbert space. If \mathcal{F} is another Hilbert space then $L(\mathcal{E}, \mathcal{F})$ is the algebra of bounded linear operators from \mathcal{E} into \mathcal{F} , $L(\mathcal{E}) = L(\mathcal{E}, \mathcal{E})$ and denote by \mathcal{Q} the set of (oblique) projections, i.e. $\mathcal{Q} = \{Q \in L(\mathcal{E}) : Q^2 = Q\}$. If $T \in L(\mathcal{E}, \mathcal{F})$ then $T^* \in L(\mathcal{F}, \mathcal{E})$ denotes the adjoint operator of T , $R(T)$ stands for its range and $N(T)$ for its nullspace. Also, if $T \in L(\mathcal{E}, \mathcal{F})$ has closed range, T^\dagger denotes the Moore–Penrose inverse of T .

If \mathcal{S} and \mathcal{T} are two (closed) subspaces of \mathcal{E} , denote by $\mathcal{S} \dot{+} \mathcal{T}$ the direct sum of \mathcal{S} and \mathcal{T} , $\mathcal{S} \oplus \mathcal{T}$ the (direct) orthogonal sum of them and $\mathcal{S} \ominus \mathcal{T} := \mathcal{S} \cap (\mathcal{S} \cap \mathcal{T})^\perp$. If $\mathcal{E} = \mathcal{S} \dot{+} \mathcal{T}$, the oblique projection onto \mathcal{S} along \mathcal{T} , $P_{\mathcal{S} // \mathcal{T}}$, is the unique $Q \in \mathcal{Q}$ with $R(P_{\mathcal{S} // \mathcal{T}}) = \mathcal{S}$ and $N(P_{\mathcal{S} // \mathcal{T}}) = \mathcal{T}$. In particular, $P_{\mathcal{S}} := P_{\mathcal{S} // \mathcal{S}^\perp}$ is the orthogonal projection onto \mathcal{S} .

2.1. Krein spaces

In what follows we present the standard notation and some basic results on Krein spaces. For a complete exposition on the subject (and the proofs of the results below) see the books by J. Bognár [23] and T.Ya. Azizov and I.S. Iokhvidov [24], the monographs by T. Ando [25] and by M. Dritschel and J. Rovnyak [26] and the paper by J. Rovnyak [27].

Given a Krein space $(\mathcal{K}, [\cdot, \cdot])$ with a *fundamental decomposition* $\mathcal{K} = \mathcal{K}_+ \dot{+} \mathcal{K}_-$, the direct (orthogonal) sum of the Hilbert spaces $(\mathcal{K}_+, [\cdot, \cdot])$ and $(\mathcal{K}_-, -[\cdot, \cdot])$ is denoted by $(\mathcal{K}, \langle \cdot, \cdot \rangle)$. Sometimes we use the notation $[\cdot, \cdot]_{\mathcal{K}}$ instead of $[\cdot, \cdot]$ to emphasize the Krein space considered.

Observe that the indefinite metric and the inner product of \mathcal{K} are related by means of a *fundamental symmetry*, i.e. a unitary selfadjoint operator $J \in L(\mathcal{K})$ which satisfies:

$$[x, y] = \langle Jx, y \rangle, \quad x, y \in \mathcal{K}.$$

If \mathcal{H} is another Krein space, given $T \in L(\mathcal{H}, \mathcal{K})$ the J -adjoint operator of T is defined by $T^+ = J_{\mathcal{H}} T^* J_{\mathcal{K}}$, where $J_{\mathcal{H}}$ and $J_{\mathcal{K}}$ are the fundamental symmetries associated to \mathcal{H} and \mathcal{K} , respectively. An operator $T \in L(\mathcal{K})$ is said to be J -selfadjoint if $T = T^+$.

Given a subspace \mathcal{S} of a Krein space \mathcal{K} , the J -orthogonal companion to \mathcal{S} is defined by

$$\mathcal{S}^{[\perp]} = \{x \in \mathcal{K} : [x, s] = 0, \text{ for every } s \in \mathcal{S}\}.$$

Notice that if \mathcal{H} is a Hilbert space, $T \in L(\mathcal{H}, \mathcal{K})$ and \mathcal{S} is a closed subspace of \mathcal{K} then

$$T^+(\mathcal{S})^\perp = T^{-1}(\mathcal{S}^{[\perp]_{\mathcal{K}}}). \tag{5}$$

A subspace \mathcal{S} of \mathcal{K} is non-degenerated if $\mathcal{S} \cap \mathcal{S}^{[\perp]} = \{0\}$. A vector $x \in \mathcal{K}$ is J -positive if $[x, x] > 0$. A subspace \mathcal{S} of \mathcal{K} is J -positive if every $x \in \mathcal{S}$, $x \neq 0$, is a J -positive vector. Moreover, it is said to be *uniformly J -positive* if there exists $\alpha > 0$ such that

$$[x, x] \geq \alpha \|x\|^2, \quad \text{for every } x \in \mathcal{S},$$

where $\|\cdot\|$ stands for the norm of the associated Hilbert space $(\mathcal{K}, \langle \cdot, \cdot \rangle)$. J -nonnegative, J -neutral, J -negative and J -nonpositive vectors (and subspaces) are defined analogously. Notice that if \mathcal{S} is a J -definite subspace of \mathcal{K} then it is non-degenerated.

Definition 2.1. Let $(\mathcal{K}, [\cdot, \cdot])$ be a Krein space with fundamental symmetry J . A subspace \mathcal{S} of \mathcal{K} is called *regular* if $(\mathcal{S}, [\cdot, \cdot])$ is also a Krein space, or equivalently, \mathcal{S} is the range of a J -selfadjoint projection.

Proposition 2.2. (See [24, Corollary 7.17].) Let \mathcal{K} be a Krein space with fundamental symmetry J and \mathcal{S} a J -nonnegative closed subspace of \mathcal{K} . Then, \mathcal{S} is regular if and only if \mathcal{S} is uniformly J -positive.

Corollary 2.3. (See [23, Theorem 8.4].) Let \mathcal{K} be a Krein space with fundamental symmetry J and \mathcal{S} a closed uniformly J -positive subspace of \mathcal{K} . If Q is the J -selfadjoint projection onto \mathcal{S} then, given $x \in \mathcal{K}$,

$$[x - Qx, x - Qx] = \min_{y \in \mathcal{S}} [x - y, x - y].$$

2.2. Angles between subspaces and reduced minimum modulus

Definition 2.4. Let \mathcal{S} and \mathcal{T} be two closed subspaces of a Hilbert space \mathcal{E} . The cosine of the *Friedrichs angle* between \mathcal{S} and \mathcal{T} is defined by

$$c(\mathcal{S}, \mathcal{T}) = \sup \{ |\langle x, y \rangle| : x \in \mathcal{S} \ominus \mathcal{T}, \|x\| = 1, y \in \mathcal{T} \ominus \mathcal{S}, \|y\| = 1 \}.$$

It is well known that

$$c(\mathcal{S}, \mathcal{T}) < 1 \Leftrightarrow \mathcal{S} + \mathcal{T} \text{ is closed} \Leftrightarrow \mathcal{S}^\perp + \mathcal{T}^\perp \text{ is closed} \Leftrightarrow c(\mathcal{S}^\perp, \mathcal{T}^\perp) < 1.$$

Furthermore, if $P_{\mathcal{S}}$ and $P_{\mathcal{T}}$ are the orthogonal projections onto \mathcal{S} and \mathcal{T} , respectively, then $c(\mathcal{S}, \mathcal{T}) < 1$ if and only if $(I - P_{\mathcal{S}})P_{\mathcal{T}}$ has closed range, or equivalently, $(I - P_{\mathcal{T}})P_{\mathcal{S}}$ has closed range. See [28] for further details.

Proposition 2.5. (See [29,30].) Given a Hilbert space \mathcal{E} , let $A, B \in L(\mathcal{E})$ be closed range operators. Then, AB has closed range if and only if $c(R(B), N(A)) < 1$.

The next definition is due to T. Kato, see [31, Ch. IV, §5] for a complete exposition on this subject.

Definition 2.6. The reduced minimum modulus $\gamma(T)$ of an operator $T \in L(\mathcal{E})$ is defined by

$$\gamma(T) = \inf\{\|Tx\|: \|x\| = 1; x \in N(T)^\perp\}.$$

It is well known that $\gamma(T) = \gamma(T^*) = \gamma(T^*T)^{1/2}$. Also, it can be shown that an operator $T \neq 0$ has closed range if and only if $\gamma(T) > 0$. In this case, $\gamma(T) = \|T^\dagger\|^{-1}$.

3. Indefinite abstract splines: definitions and basic results

Recently, some interpolation methods in Reproducing Kernel Hilbert Spaces (RKHS) have shown to be useful to deal with machine learning problems. Given a data set $X = \{x_1, \dots, x_m\} \subseteq \mathcal{X}$ and labels $Y = \{y_1, \dots, y_m\} \subset \mathbb{R}$, it is necessary to estimate the minimal norm function $f \in \mathcal{H}$ such that $f(x_i) = y_i$, where \mathcal{H} is a RKHS with kernel

$$k: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}.$$

If $E: \mathcal{H} \rightarrow \mathbb{R}^m$ is the evaluation map given by $Ef = (f(x_1), \dots, f(x_m))$, the above interpolation problem consists in finding $f \in \mathcal{H}$ such that

$$Ef = (y_1, \dots, y_m) = \mathbf{y} \quad \text{and} \quad \|f\|^2 = \min_{g \in E^{-1}(\mathbf{y})} \|g\|^2.$$

Notice that the adjoint operator $E^*: \mathbb{R}^m \rightarrow \mathcal{H}$ is given by $E^*\alpha = \sum_{i=1}^m \alpha_i k(x_i, \cdot)$ where $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{R}^m$. Then, it follows that $K = EE^*$ is the Gram matrix associated to the kernel k , i.e. $K_{ij} = k(x_i, x_j)$.

Since $\mathcal{H} = R(E^*) \oplus N(E)$, it is easy to see that there is a solution to the above problem if and only if there exists $f \in R(E^*)$ such that $Ef = \mathbf{y}$, or equivalently, there exist $\alpha \in \mathbb{R}^m$ such that $K\alpha = \mathbf{y}$ (in this case the minimizing function is reconstructed as $f(x) = E^*\alpha = \sum_{i=1}^m \alpha_i k(x_i, x)$). So, the interpolation spline can be defined without using the norm of the RKHS but only its kernel.

In order to admit indefinite kernels to study machine learning problems, S. Canu et al. provided a definition of interpolating splines in a Reproducing Kernel Krein Space (RKKS):

Definition 3.1 (S. Canu et al.). Let \mathcal{K} be a RKKS with kernel $k: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ and suppose that $N(E)$ is a regular subspace of \mathcal{K} . Given $\mathbf{y} \in \mathbb{R}^m$, the interpolation spline f is given by

$$f(x) = \sum_{i=1}^m \alpha_i k(x_i, x),$$

where α satisfies $K\alpha = EE^+\alpha = \mathbf{y}$, see [10, Definition 3.3].

Notice that, if $N(E)$ is a uniformly J -positive subspace of \mathcal{K} , then the interpolation spline f defined by S. Canu et al. satisfies

$$Ef = \mathbf{y} \quad \text{and} \quad [f, f] = \min_{g \in E^{-1}(\mathbf{y})} [g, g], \tag{6}$$

because the projection onto $R(E^+)$ along $N(E)$ is J -contractive and $f = E^+\alpha$ is the unique vector in $E^{-1}(\mathbf{y})$ which coincides with its projection.

The aim of this section is to consider a general indefinite version of the abstract interpolation problem considered by M. Atteia (see Eq. (1)).

Throughout this work, \mathcal{K} is a Krein space with fundamental symmetry J , \mathcal{H} and \mathcal{E} are Hilbert spaces and the operators $T \in L(\mathcal{H}, \mathcal{K})$ and $V \in L(\mathcal{H}, \mathcal{E})$ are surjective. Consider the following generalization of the abstract interpolation problem [11]:

Problem 3.2. Given $z_0 \in \mathcal{E}$, find $x_0 \in V^{-1}(\{z_0\})$ such that

$$[Tx_0, Tx_0]_{\mathcal{K}} = \min\{[Tx, Tx]_{\mathcal{K}}: Vx = z_0\}. \tag{7}$$

Definition 3.3. Any element $x_0 \in V^{-1}(\{z_0\})$ satisfying Eq. (7) is called an *indefinite abstract spline* or, more specifically, a (T, V) -interpolant to $z_0 \in \mathcal{E}$. The set of (T, V) -interpolants to z_0 is denoted by $sp(T, V, z_0)$.

Considering the Moore–Penrose inverse of V , the above problem can be restated as: For a fixed $z_0 \in \mathcal{E}$, find $u_0 \in N(V)$ such that

$$[T(V^\dagger z_0 + u_0), T(V^\dagger z_0 + u_0)]_{\mathcal{K}} = \min\{[T(V^\dagger z_0 + u), T(V^\dagger z_0 + u)]_{\mathcal{K}} : u \in N(V)\}. \quad (8)$$

As it was mentioned in the introduction, the following lemma shows under which conditions indefinite abstract splines do exist.

Lemma 3.4. *Given $z_0 \in \mathcal{E}$, $x_0 \in V^{-1}(\{z_0\})$ is a (T, V) -interpolant to z_0 if and only if $T(N(V))$ is a J -nonnegative subspace of \mathcal{K} and $Tx_0 \in T(N(V))^{\perp\perp}$.*

Proof. Suppose that $x_0 \in \mathcal{H}$ is a (T, V) -interpolant to z_0 . Then, for every $u \in N(V)$ and $\alpha \in \mathbb{R}$,

$$[Tx_0, Tx_0] \leq [T(x_0 + \alpha u), T(x_0 + \alpha u)] = [Tx_0, Tx_0] + 2\alpha \operatorname{Re}[Tx_0, Tu] + \alpha^2[Tu, Tu].$$

Therefore, $2\alpha \operatorname{Re}[Tx_0, Tu] + \alpha^2[Tu, Tu] \geq 0$ for every $\alpha \in \mathbb{R}$, and a standard argument shows that $\operatorname{Re}[Tx_0, Tu] = 0$. Analogously, if $\beta = i\alpha$, $\alpha \in \mathbb{R}$, it follows that $\operatorname{Im}[Tx_0, Tu] = 0$. Then, $[Tx_0, Tu] = 0$ and $[Tu, Tu] \geq 0$ for every $u \in N(V)$.

Conversely, suppose that $T(N(V))$ is a J -nonnegative subspace of \mathcal{K} and there exists $x_0 \in V^{-1}(\{z_0\})$ such that $Tx_0 \in T(N(V))^{\perp\perp}$. If $u_0 = x_0 - V^\dagger z_0 \in N(V)$ then, for every $u \in N(V)$,

$$[T(V^\dagger z_0 + u), T(V^\dagger z_0 + u)] = [T(V^\dagger z_0 + u_0), T(V^\dagger z_0 + u_0)] + [T(u - u_0), T(u - u_0)] \geq [Tx_0, Tx_0].$$

Therefore, x_0 is a (T, V) -interpolant to z_0 . \square

Remark 3.5. Notice that the framework considered by S. Canu et al. to define interpolating splines in RKKS is a particular case of ours. If \mathcal{K} is a RKKS consider the identity operator as T and the evaluation map E as V . Then, the hypothesis mentioned before Eq. (6) to obtain the variational characterization of the interpolating spline is the same as in Lemma 3.4.

As a consequence of Eq. (5), $sp(T, V, z_0)$ can be characterized as the intersection of a subspace and an affine manifold of \mathcal{H} .

Corollary 3.6. *Suppose that $T(N(V))$ is a J -nonnegative subspace of \mathcal{K} and let $z_0 \in \mathcal{E}$. Then,*

$$sp(T, V, z_0) = (V^\dagger z_0 + N(V)) \cap T^+T(N(V))^{\perp}.$$

Proof. Given $z_0 \in \mathcal{E}$, suppose that $x_0 \in \mathcal{H}$ is a (T, V) -interpolant to z_0 . Then, $u_0 = x_0 - V^\dagger z_0 \in N(V)$ and by the above lemma, $Tx_0 \in T(N(V))^{\perp\perp}$, or equivalently by Eq. (5), $x_0 \in T^+T(N(V))^{\perp}$. Therefore, $x_0 \in (V^\dagger z_0 + N(V)) \cap T^+T(N(V))^{\perp}$.

On the other hand, if $u \in N(V)$ is such that $x = V^\dagger z_0 + u \in T^+T(N(V))^{\perp}$, then $Tx \in T(N(V))^{\perp\perp}$ and $Vx = z_0$. So, applying Lemma 3.4, it follows that $x \in sp(T, V, z_0)$. \square

The following lemma shows how regularity conditions on $T(N(V))$ determine relationships between the subspaces $N(T)$ and $T^+T(N(V))^{\perp}$.

Lemma 3.7.

- (i) *If $T(N(V))$ is non-degenerated, then $N(V) \cap T^+T(N(V))^{\perp} = N(V) \cap N(T)$.*
- (ii) *If $T(N(V))$ is regular, then $\mathcal{H} = N(V) + T^+T(N(V))^{\perp}$.*

Proof. (i) By Eq. (5), the inclusion $N(T) \cap N(V) \subseteq N(V) \cap T^+T(N(V))^{\perp}$ is straightforward. On the other hand, if $x \in N(V) \cap T^+T(N(V))^{\perp}$ then $Tx \in T(N(V)) \cap T(N(V))^{\perp\perp} = \{0\}$. Thus, $x \in N(V) \cap N(T)$.

(ii) If $T(N(V))$ is a regular subspace of \mathcal{K} then $\mathcal{K} = T(N(V)) + T(N(V))^{\perp\perp}$. Therefore,

$$\mathcal{H} = T^{-1}(T(N(V))) + T^{-1}(T(N(V))^{\perp\perp}) = N(V) + T^+T(N(V))^{\perp}$$

(see Eq. (5)). \square

As mentioned above, if $T(N(V))$ is a regular subspace of \mathcal{K} then $\mathcal{H} = N(V) + T^+T(N(V))^{\perp}$. But this may not be a direct sum. Therefore, there is a family of closed subspaces of $T^+T(N(V))^{\perp}$ which are complementary to $N(V)$. Along this work, if $T(N(V))$ is a regular subspace of \mathcal{K} we will consider the following projection:

$$Q_0 = P_{N(V)/T^+T(N(V))^{\perp} \oplus N(V)}. \quad (9)$$

Proposition 3.8. *Suppose that $T(N(V))$ is a closed subspace of \mathcal{K} . Then, the set $sp(T, V, z) \neq \emptyset$ for every $z \in \mathcal{E}$ if and only if $T(N(V))$ is uniformly J -positive. In this case, $sp(T, V, z)$ is an affine manifold parallel to $N(V) \cap N(T)$.*

Proof. Suppose that $T(N(V))$ is a closed uniformly J -positive subspace of \mathcal{K} . Then, by Proposition 2.2, $T(N(V))$ is a regular subspace of \mathcal{K} , and $Q_0 \in \mathcal{Q}$ (see Lemma 3.7). For a fixed $z \in \mathcal{E}$, let $x = (I - Q_0)V^\dagger z \in \mathcal{H}$. Then, $Vx = z$ and $Tx \in T(N(V))^{\perp\perp}$. So, by Lemma 3.4, $x \in sp(T, V, z)$, i.e. $sp(T, V, z) \neq \emptyset$ for every $z \in \mathcal{E}$.

Conversely, suppose that $sp(T, V, z) \neq \emptyset$ for every $z \in \mathcal{E}$. Then, as a consequence of Lemma 3.4, $T(N(V))$ is a J -nonnegative subspace of \mathcal{K} . Furthermore, for each $z \in \mathcal{E}$, there exists a vector $x_z \in \mathcal{H}$ such that $Vx_z = z$ and $Tx_z \in T(N(V))^{\perp\perp}$. Since $V^\dagger z = (V^\dagger z - x_z) + x_z$ and $V(V^\dagger z - x_z) = 0$ for every $z \in \mathcal{E}$, it is easy to see that $N(V)^\perp \subseteq N(V) + T^+T(N(V))^\perp$. Therefore, $\mathcal{H} = N(V) + T^+T(N(V))^\perp$ and $\mathcal{K} = T(N(V)) + T(N(V))^{\perp\perp}$. So, $T(N(V))$ is a regular J -nonnegative subspace of \mathcal{K} , i.e. $T(N(V))$ is a uniformly J -positive subspace of \mathcal{K} (see Proposition 2.2).

Assuming that $T(N(V))$ is uniformly J -positive, if $x_1, x_2 \in sp(T, V, z)$ then, by Lemma 3.7,

$$x_1 - x_2 \in N(V) \cap T^+T(N(V))^\perp = N(V) \cap N(T). \quad \square$$

Corollary 3.9. *Suppose that $T(N(V))$ is a closed uniformly J -positive subspace of \mathcal{K} and $N(T) \cap N(V) = \{0\}$. Then, given $z \in \mathcal{E}$, $sp(T, V, z)$ is a singleton. More precisely,*

$$sp(T, V, z) = \{P_{T^+T(N(V))^\perp // N(V)} V^\dagger z\}.$$

Example 3.10. In signal processing applications it is frequently assumed that the mathematical model, describing the physical phenomena under study, satisfies the following equation:

$$y = Hx + \eta,$$

where $x \in \mathbb{C}^n$ is the quantity that needs to be estimated and $H \in \mathbb{C}^{m \times n}$ is known.

Sometimes, due to physical restrictions, it is not possible to measure x , but the measurement y may be available. This measurement is corrupted by some noise η . According to the known information on the measurement noise, different estimation techniques can be used to approximate x . For instance, when no statistical information about the measurement noise is available, the \mathcal{H}^∞ -estimation technique has been proved to be a good solution for different engineering problems. Given $\gamma > 0$, the \mathcal{H}^∞ -estimation technique consists in finding an estimation \hat{x} of the vector x , such that:

$$\max_{x \in \mathbb{C}^n} \frac{\|x - \hat{x}\|^2}{\|y - Hx\|^2} \leq \gamma^2, \tag{10}$$

or equivalently,

$$\min_{x \in \mathbb{C}^n} \|y - Hx\|^2 - \frac{1}{\gamma^2} \|x - \hat{x}\|^2 \geq 0. \tag{11}$$

In what follows we show that the \mathcal{H}^∞ -estimation technique can be formulated as an indefinite abstract spline problem. This connection has been previously considered, see for instance [32] and references therein.

If $w = x - \hat{x}$, then Eq. (11) can be written in matrix form as

$$\min_{w \in \mathbb{C}^n} \left[\begin{pmatrix} y - H\hat{x} \\ 0 \end{pmatrix} - \begin{pmatrix} H \\ \gamma^{-1}I_n \end{pmatrix} w \right]^* \begin{bmatrix} I_m & 0 \\ 0 & -I_n \end{bmatrix} \left[\begin{pmatrix} y - H\hat{x} \\ 0 \end{pmatrix} - \begin{pmatrix} H \\ \gamma^{-1}I_n \end{pmatrix} w \right] \geq 0. \tag{12}$$

So, if $\mathcal{K} = \mathbb{C}^{m+n}$, define the symmetry $J \in L(\mathcal{K})$ as $J = \begin{pmatrix} I_m & 0 \\ 0 & -I_n \end{pmatrix}$ and let $[x, y] = x^* J y$, for every $x, y \in \mathcal{K}$. Then, the \mathcal{H}^∞ -estimation technique can be rewritten as: for a fixed $\gamma > 0$, find $z_0 \in \mathbb{C}^{m+n}$ such that $Vz_0 = y_1$ and

$$[z_0, z_0] = \min_{Vz=y_1} [z, z] \geq 0, \tag{13}$$

where $I - V$ is the orthogonal projection onto the range of the matrix $\begin{pmatrix} H \\ \gamma^{-1}I_n \end{pmatrix}$ and $y_1 = V \begin{pmatrix} y - H\hat{x} \\ 0 \end{pmatrix}$.

Observe that Eq. (13) is more than an indefinite splines problem, because the solution z_0 also has to satisfy $[z_0, z_0] \geq 0$. This last condition depends on the chosen parameter $\gamma > 0$.

In what follows, for a fixed $z_0 \in \mathcal{E}$, it is shown that $sp(T, V, z_0)$ can be parametrized by means of a family of projections onto $N(V)$.

Proposition 3.11. *Suppose that $T(N(V))$ is a closed uniformly J -positive subspace of \mathcal{K} . Given $z_0 \in \mathcal{E}$, $x \in sp(T, V, z_0)$ if and only if there exists $Q \in \mathcal{Q}$ with $R(Q) = N(V)$ and $N(Q) \subseteq T^+T(N(V))^\perp$ such that $x = (I - Q)V^\dagger z_0$.*

To prove the above proposition, we need the following lemma.

Lemma 3.12. *Let $Q \in \mathcal{Q}$ and suppose that $T(N(V))$ is a regular subspace of \mathcal{K} . Then, $R(Q) = N(V)$ and $N(Q) \subseteq T^+T(N(V))^\perp$ if and only if $Q = Q_0 + Z$, where $Z \in L(\mathcal{H})$ is such that $N(V) \subseteq N(Z)$ and $R(Z) \subseteq N(V) \cap N(T)$.*

Proof. If $Q \in L(\mathcal{H})$ is a projection with $R(Q) = N(V)$ and $N(Q) \subseteq T^+T(N(V))^\perp$, let $Z = Q - Q_0$. Since $R(Q) = R(Q_0) = N(V)$ it is trivial that $N(V) \subseteq N(Z)$. On the other hand, consider $y = Zx \in R(Z)$: $y = Qx - Q_0x \in N(V)$ and $y = (I - Q_0)x - (I - Q)x \in T^+T(N(V))^\perp$. Then $y \in N(V) \cap T^+T(N(V))^\perp = N(V) \cap N(T)$.

Conversely, given $Z \in L(\mathcal{H})$ with $N(V) \subseteq N(Z)$ and $R(Z) \subseteq N(V) \cap N(T)$, consider $Q = Q_0 + Z$. Then, $Q^2 = Q$ because $Z^2 = 0$, $Q_0Z = Z$ and $ZQ_0 = 0$. It is easy to see that $R(Q) \subseteq N(V)$ and, if $x \in N(V)$ then $Qx = Q_0x = x$. Therefore, $R(Q) = N(V)$. Finally, observe that if $x \in N(Q)$ then $x = (I - Q)x = (I - Q_0)x - Zx \in T^+T(N(V))^\perp$, because $N(Q_0) + R(Z) \subseteq T^+T(N(V))^\perp$. \square

Proof of Proposition 3.11. If $x = (I - Q)V^\dagger z_0$, where $Q \in \mathcal{Q}$ with $R(Q) = N(V)$ and $N(Q) \subseteq T^+T(N(V))^\perp$, it is easy to see that $Vx = z_0$ and $Tx \in T(N(V))^{\perp\perp}$. Then, by Lemma 3.4, $x \in sp(T, V, z_0)$.

Conversely, as a consequence of Proposition 3.8, $sp(T, V, z_0) = (I - Q_0)V^\dagger z_0 + N(V) \cap N(T)$ because $(I - Q_0)V^\dagger z_0 \in sp(T, V, z_0)$. Then, if $x \in sp(T, V, z_0)$ there exists $u \in N(V) \cap N(T)$ such that $x = (I - Q_0)V^\dagger z_0 + u$. So, consider $Z \in L(\mathcal{H})$ such that $Z(V^\dagger z_0) = -u$ and $Zy = 0$ if $y \perp V^\dagger z_0$. Then,

$$x = (I - Q_0)V^\dagger z_0 - ZV^\dagger z_0 = (I - (Q_0 + Z))V^\dagger z_0,$$

$N(V) \subseteq N(Z)$ and $R(Z) \subseteq N(V) \cap N(T)$. Therefore, by the above lemma, $Q = Q_0 + Z \in \mathcal{Q}$ with $R(Q) = N(V)$ and $N(Q) \subseteq T^+T(N(V))^\perp$. \square

3.1. Frames of indefinite abstract splines

Recall that a sequence $\{f_n\}_{n \in \mathbb{N}}$ in a Banach space X is called a *Schauder basis* of X if for every $x \in X$ there is a unique sequence of scalars $\{c_n\}_{n \in \mathbb{N}}$ such that $x = \sum_{n=1}^\infty c_n f_n$, where the series converges in the norm topology. A vector sequence $\{f_n\}_{n \in \mathbb{N}}$ in X is a *Riesz basis* if there exist constants $0 < A < B$ such that

$$A \sum_{n=1}^m |c_n|^2 \leq \left\| \sum_{n=1}^m c_n f_n \right\|^2 \leq B \sum_{n=1}^m |c_n|^2, \tag{14}$$

for all finite sequences c_1, \dots, c_m .

On the other hand, given a Hilbert space \mathcal{E} , a sequence $\{f_n\}_{n \in \mathbb{N}}$ in \mathcal{E} is a *frame* for \mathcal{E} if there exist constants $0 < A < B$ such that

$$A \|z\|^2 \leq \sum_{n=1}^\infty |\langle z, f_n \rangle|^2 \leq B \|z\|^2, \quad \text{for every } z \in \mathcal{E}. \tag{15}$$

Observe that, if \mathcal{E} is a Hilbert space, $\{f_n\}_{n \in \mathbb{N}}$ is a Riesz basis of \mathcal{E} if and only if $\{f_n\}_{n \in \mathbb{N}}$ is a frame for \mathcal{E} such that, if $\sum_{n=1}^\infty c_n f_n = 0$, then $c_n = 0$ for every $n \in \mathbb{N}$. See [33,34] for further details on this subject.

In what follows, recall that $T \in L(\mathcal{H}, \mathcal{K})$ and $V \in L(\mathcal{H}, \mathcal{E})$ are surjective operators and suppose that $T(N(V))$ is a closed uniformly J -positive subspace of \mathcal{K} .

Proposition 3.13. *Given a sequence $\{f_n\}_{n \in \mathbb{N}}$ in \mathcal{E} , suppose that there exists a frame $\{g_n\}_{n \in \mathbb{N}}$ for $\mathcal{W} = T^+T(N(V))^\perp$ such that $g_n \in sp(T, V, f_n)$ for every $n \in \mathbb{N}$. Then, $\{f_n\}_{n \in \mathbb{N}}$ is a frame for \mathcal{E} .*

Proof. If $g_n \in sp(T, V, f_n)$ then, by Proposition 3.11, there exists $Q_n \in \mathcal{Q}$ with $R(Q_n) = N(V)$ and $N(Q_n) \subseteq \mathcal{W}$, such that $g_n = (I - Q_n)V^\dagger f_n$. Since $V(I - Q_n)V^\dagger = I_{\mathcal{E}}$ for every $n \in \mathbb{N}$, it is easy to see that

$$\sum_{n=1}^\infty |\langle z, f_n \rangle|^2 = \sum_{n=1}^\infty \|(V^*z, (I - Q_n)V^\dagger f_n)\|^2 = \sum_{n=1}^\infty |(P_{\mathcal{W}}V^*z, g_n)|^2, \quad \text{for every } z \in \mathcal{E},$$

since $P_{\mathcal{W}}(I - Q_n) = (I - Q_n)$. Therefore, if $\{g_n\}_{n \in \mathbb{N}}$ is a frame for \mathcal{W} with frame bounds $0 < A < B$,

$$A \|P_{\mathcal{W}}V^*z\|^2 \leq \sum_{n=1}^\infty |\langle z, f_n \rangle|^2 \leq B \|P_{\mathcal{W}}V^*z\|^2 \leq B \|V\|^2 \|z\|^2,$$

for every $z \in \mathcal{E}$. But $\|P_{\mathcal{W}}V^*z\|^2 \geq \gamma(P_{\mathcal{W}}V^*)^2 \|z\|^2 = \gamma(V P_{\mathcal{W}})^2 \|z\|^2$. Since $c(\mathcal{W}, N(V)) < 1$ it follows by Proposition 2.5 that $V P_{\mathcal{W}}$ has closed range, so $\gamma(V P_{\mathcal{W}}) > 0$. Then, $\{f_n\}_{n \in \mathbb{N}}$ is a frame for \mathcal{E} , with frame bounds $0 < A\gamma(V P_{\mathcal{W}})^2 < B\|V\|^2$. \square

The next result shows that, given a frame $\{f_n\}_{n \in \mathbb{N}}$ for \mathcal{E} , it is possible to obtain frames of splines for any complement of $N(V)$ contained in $T^+T(N(V))^\perp$.

Proposition 3.14. *Given a sequence $\{f_n\}_{n \in \mathbb{N}}$ in \mathcal{E} , consider $g_n = (I - Q)V^\dagger f_n \in sp(T, V, f_n)$, $n \in \mathbb{N}$, where $Q \in L(\mathcal{H})$ is any fixed projection such that $R(Q) = N(V)$ and $N(Q) \subseteq T^+T(N(V))^\perp$. Then,*

- (i) $\{f_n\}_{n \in \mathbb{N}}$ is a frame for \mathcal{E} if and only if $\{g_n\}_{n \in \mathbb{N}}$ is a frame for $N(Q)$.
- (ii) $\{f_n\}_{n \in \mathbb{N}}$ is a Riesz basis of \mathcal{E} if and only if $\{g_n\}_{n \in \mathbb{N}}$ is a Riesz basis of $N(Q)$.
- (iii) $\{f_n\}_{n \in \mathbb{N}}$ is a (Schauder) basis of \mathcal{E} if and only if $\{g_n\}_{n \in \mathbb{N}}$ is a (Schauder) basis of $N(Q)$.

Proof. Observe that, if $W = (I - Q)V^\dagger$, then $R(W) = R(I - Q) = N(Q)$ is closed. Then, $\gamma(W) > 0$.

(i) Suppose that $\{f_n\}_{n \in \mathbb{N}}$ is a frame for \mathcal{E} . Notice that

$$\sum_{n=1}^{\infty} |\langle x, g_n \rangle|^2 = \sum_{n=1}^{\infty} |\langle x, Wf_n \rangle|^2 = \sum_{n=1}^{\infty} |\langle W^*x, f_n \rangle|^2 \quad \text{for every } x \in \mathcal{H}.$$

So, if $0 < A < B$ are frame bounds for $\{f_n\}_{n \in \mathbb{N}}$ then

$$A\gamma(W)^2 \|x\|^2 = A\gamma(W^*)^2 \|x\|^2 \leq A \|W^*x\|^2 \leq \sum_{n=1}^{\infty} |\langle x, g_n \rangle|^2 \leq B \|W^*x\|^2 \leq B \|W\|^2 \|x\|^2,$$

for every $x \in N(W^*)^\perp = N(Q)$. Therefore, $\{g_n\}_{n \in \mathbb{N}}$ is a frame for $N(Q)$. The other implication is a consequence of Proposition 3.13.

(ii) Suppose that $\{f_n\}_{n \in \mathbb{N}}$ is a Riesz basis of \mathcal{E} . Then it is also a frame for \mathcal{E} and, by item (i), the sequence $\{g_n\}_{n \in \mathbb{N}}$ is a frame for $N(Q)$. Furthermore, if there exists a sequence $(\alpha_k)_{k \in \mathbb{N}}$ such that $\sum_{k=1}^{\infty} \alpha_k g_k = 0$, then applying V to both sides of the equation we obtain that $\sum_{k=1}^{\infty} \alpha_k f_k = 0$. So, $\alpha_k = 0$ for every $k \in \mathbb{N}$ because $\{f_n\}_{n \in \mathbb{N}}$ is a Riesz basis of \mathcal{E} . Therefore, $\{g_n\}_{n \in \mathbb{N}}$ is a Riesz basis of $N(Q)$. The other implication follows in the same way.

(iii) It is analogous to the proof of [17, Ch. III, Proposition 1.1]. \square

Given a sequence $\{f_n\}_{n \in \mathbb{N}}$ in \mathcal{E} , if $N(T) \cap N(V) = \{0\}$ it is easy to see that $\{f_n\}_{n \in \mathbb{N}}$ is a frame for \mathcal{E} if and only if $\{g_n\}_{n \in \mathbb{N}}$ is a frame for $T^+T(N(V))^\perp$, where g_n is the (unique) (T, V) -interpolant to f_n (see Proposition 3.14). However, the following example shows that, if $N(T) \cap N(V) \neq \{0\}$, given a frame $\{f_n\}_{n \in \mathbb{N}}$ for \mathcal{E} it is easy to construct $g_n \in sp(T, V, f_n)$ (for every $n \in \mathbb{N}$) such that $\{g_n\}_{n \in \mathbb{N}}$ is not a frame.

Example 3.15. Observe that if $\{f_n\}_{n \in \mathbb{N}}$ is a frame with frame bounds $0 < A < B$ then $\|f_n\|^2 \leq B$. Given $u \in N(T) \cap N(V)$ with $\|u\| = 1$, define

$$Z_n(x) = \begin{cases} n\alpha u & \text{if } x = \alpha V^\dagger f_n, \alpha \in \mathbb{C}; \\ 0 & \text{if } x \perp V^\dagger f_n. \end{cases}$$

Then, $Z_n \in L(\mathcal{H})$ and satisfies $N(V) \subseteq N(Z_n)$ and $R(Z_n) \subseteq N(T) \cap N(V)$. Furthermore, by Lemma 3.12, $Q_n = Q_0 + Z_n$ is a projection with $R(Q_n) = N(V)$ and $N(Q_n) \subseteq T^+T(N(V))^\perp$. Therefore, $g_n = (I - Q_n)V^\dagger f_n \in sp(T, V, f_n)$ for every $n \in \mathbb{N}$.

But observe that $\{g_n\}_{n \in \mathbb{N}}$ cannot be a frame because $\|g_n\| \rightarrow +\infty$ as $n \rightarrow \infty$. Indeed, it is easy to see that

$$\|g_n\| \geq \|Z_n V^\dagger f_n\| - \|(I - Q_0)V^\dagger f_n\| \geq n - \|I - Q_0\| \|V^\dagger\| B^{1/2} \rightarrow +\infty \quad \text{as } n \rightarrow \infty.$$

4. Indefinite abstract smoothing splines

Let \mathcal{K} be a Krein space with fundamental symmetry $J_{\mathcal{K}}$, and consider two Hilbert spaces \mathcal{H} and \mathcal{E} . Given surjective operators $T \in L(\mathcal{H}, \mathcal{K})$ and $V \in L(\mathcal{H}, \mathcal{E})$, consider the following generalization of the abstract smoothing problem [17]:

Problem 4.1. Given $\rho > 0$ and a fixed $z_0 \in \mathcal{E}$, find $x_0 \in \mathcal{H}$ such that

$$[Tx_0, Tx_0]_{\mathcal{K}} + \rho \|Vx_0 - z_0\|_{\mathcal{E}}^2 = \min_{x \in \mathcal{H}} ([Tx, Tx]_{\mathcal{K}} + \rho \|Vx - z_0\|_{\mathcal{E}}^2). \tag{16}$$

Definition 4.2. Any element $x_0 \in \mathcal{H}$ satisfying Eq. (16) is called a (T, V, ρ) -smoothing spline to $z_0 \in \mathcal{E}$. The set of (T, V, ρ) -smoothing splines to z_0 is denoted by $sm(T, V, \rho, z_0)$.

To study this problem consider the indefinite metric defined on $\mathcal{K} \times \mathcal{E}$ by:

$$[(y, z), (y', z')]_{\rho} = [y, y']_{\mathcal{K}} + \rho [z, z']_{\mathcal{E}}, \quad (y, z), (y', z') \in \mathcal{K} \times \mathcal{E}. \tag{17}$$

Notice that $\mathcal{K} \times \mathcal{E}$ is a Krein space with the indefinite metric defined above. In fact, considering the fundamental symmetry $J_{\mathcal{K}}$ of \mathcal{K} and the inner product $\langle \cdot, \cdot \rangle_{\rho}$ in $\mathcal{K} \times \mathcal{E}$ given by $\langle (y, z), (y', z') \rangle_{\rho} = \langle y, y' \rangle_{\mathcal{K}} + \rho \langle z, z' \rangle_{\mathcal{E}}$ where $(y, z), (y', z') \in \mathcal{K} \times \mathcal{E}$, the operator $J_{\rho} \in L(\mathcal{K} \times \mathcal{E})$ defined as

$$J_{\rho}(y, z) = (J_{\mathcal{K}}y, z), \quad (y, z) \in \mathcal{K} \times \mathcal{E},$$

is a fundamental symmetry associated to $(\mathcal{K} \times \mathcal{E}, [\cdot, \cdot]_{\rho})$. Also, considering the operator $L : \mathcal{H} \rightarrow \mathcal{K} \times \mathcal{E}$ defined by

$$Lx = (Tx, Vx), \quad x \in \mathcal{H},$$

observe that Problem 4.1 can be restated as the following indefinite least squares problem: given $\rho > 0$ and a fixed $z_0 \in \mathcal{E}$, find $x_0 \in \mathcal{H}$ such that

$$[Lx_0 - (0, z_0), Lx_0 - (0, z_0)]_{\rho} = \min_{x \in \mathcal{H}} [Lx - (0, z_0), Lx - (0, z_0)]_{\rho}. \tag{18}$$

Using the formulation given above, the next results characterize the solutions of the indefinite abstract smoothing problem.

Lemma 4.3. *Given $z_0 \in \mathcal{E}, x_0 \in \mathcal{H}$ is a solution of Problem 4.1 if and only if $R(L)$ is J_{ρ} -nonnegative and x_0 is a solution of the equation:*

$$(T^+T + \rho V^*V)x = \rho V^*z_0.$$

Proof. Following the same arguments as in Lemma 3.4, it is easy to see that $x_0 \in \mathcal{H}$ satisfies Eq. (18) if and only if $R(L)$ is J_{ρ} -nonnegative and

$$[Lx_0 - (0, z_0), Lx]_{\rho} = 0, \quad \text{for every } x \in \mathcal{H},$$

or equivalently, $L^+(Lx_0 - (0, z_0)) = 0$. Since $L^+ \in L(\mathcal{K} \times \mathcal{E}, \mathcal{H})$ is given by $L^+(y, z) = T^+y + \rho V^*z$, $(y, z) \in \mathcal{K} \times \mathcal{E}$, it follows that $(T^+T + \rho V^*V)x_0 = \rho V^*z_0$. \square

In order to obtain some alternative characterizations for the solutions of Problem 4.1, it is necessary to consider the particular case of a closed range operator L . The next lemma gives a condition between the operators V and T that guarantees that L has closed range. The proof is similar to the one given in [17, Ch. III, Lemma 2.1] for the Hilbert space case.

Lemma 4.4. *If $T(N(V))$ is a closed subspace of \mathcal{K} then $R(L)$ is a closed subspace of $\mathcal{K} \times \mathcal{E}$.*

Proof. Given $(y, z) \in \mathcal{K} \times \mathcal{E}$, suppose that $\{x_n\}_{n \geq 1} \subseteq N(L)^{\perp}$ is such that $Lx_n \rightarrow (y, z)$. If $v_n = V^{\dagger}Vx_n \in N(V)^{\perp} \subseteq N(L)^{\perp}$, then $v_n \rightarrow V^{\dagger}z \in \mathcal{H}$ and $u_n = x_n - v_n \in N(V) \cap N(L)^{\perp}$. Therefore, $Vv_n = Vx_n \rightarrow z$ and $Tu_n \rightarrow y - TV^{\dagger}z$.

Since $T(N(V))$ is a closed subspace of \mathcal{K} , the operator $W = T|_{N(V)} : N(V) \rightarrow \mathcal{K}$ has closed range and, for every $n \geq 1$, $u_n = W^{\dagger}Tu_n$ because $u_n \in N(V) \cap N(L)^{\perp} = N(W)^{\perp}$. Thus, $x_n = v_n + u_n = v_n + W^{\dagger}Tu_n \rightarrow V^{\dagger}z + W^{\dagger}(y - TV^{\dagger}z)$. Furthermore, if $x = V^{\dagger}z + W^{\dagger}(y - TV^{\dagger}z)$, it follows that $Tx = y$ and $Vx = z$ because $y - TV^{\dagger}z \in T(N(V))$. Therefore, $R(L)$ is a closed subspace of $\mathcal{K} \times \mathcal{E}$. \square

As a consequence of Corollary 2.3, if there exists $\arg \min_{x \in \mathcal{H}} [Lx - (y, z), Lx - (y, z)]_{\rho}$ for every $(y, z) \in \mathcal{K} \times \mathcal{E}$, then $R(L)$ is a regular subspace of $\mathcal{K} \times \mathcal{E}$. The following proposition shows that this assertion also holds considering the proper subspace of $\mathcal{K} \times \mathcal{E}$ obtained by embedding \mathcal{E} into $\mathcal{K} \times \mathcal{E}$.

Proposition 4.5. *Problem 4.1 admits a solution for every $z \in \mathcal{E}$ if and only if $R(L)$ is a closed uniformly J_{ρ} -positive subspace of $\mathcal{K} \times \mathcal{E}$.*

Proof. Suppose that, Problem 4.1 admits a solution for every $z \in \mathcal{E}$. Applying Lemma 4.3, it follows that $R(L)$ is J_{ρ} -nonnegative. Given $(y, z) \in \mathcal{K} \times \mathcal{E}$, consider $w = u + T^{\dagger}y$, where $u \in \mathcal{H}$ satisfies

$$[Lu - (0, z - VT^{\dagger}y), Lu - (0, z - VT^{\dagger}y)]_{\rho} = \min_{x \in \mathcal{H}} [Lx - (0, z - VT^{\dagger}y), Lx - (0, z - VT^{\dagger}y)]_{\rho}.$$

Then, for every $x \in \mathcal{H}$,

$$\begin{aligned} [Lw - (y, z), Lw - (y, z)]_{\rho} &= [Lu + (y, VT^{\dagger}y) - (y, z), Lu + (y, VT^{\dagger}y) - (y, z)]_{\rho} \\ &= [Lu - (0, z - VT^{\dagger}y), Lu - (0, z - VT^{\dagger}y)]_{\rho} \\ &\leq [L(x - T^{\dagger}y) - (0, z - VT^{\dagger}y), L(x - T^{\dagger}y) - (0, z - VT^{\dagger}y)]_{\rho} \\ &= [Lx - (y, z), Lx - (y, z)]_{\rho}. \end{aligned}$$

Therefore, for every $(y, z) \in \mathcal{K} \times \mathcal{E}$, there exists $w \in \mathcal{H}$ such that

$$[Lw - (y, z), Lw - (y, z)]_\rho = \min_{x \in \mathcal{H}} [Lx - (y, z), Lx - (y, z)]_\rho.$$

Then, as in Lemma 3.4, it is easy to see that for every $(y, z) \in \mathcal{K} \times \mathcal{E}$ there exists $w \in \mathcal{H}$ such that $Lw - (y, z) \in R(L)^{\perp\perp}$. So, $\mathcal{K} \times \mathcal{E} = R(L) + R(L)^{\perp\perp}$, i.e. $R(L)$ is a regular subspace of $\mathcal{K} \times \mathcal{E}$. Thus, by Proposition 2.2, $R(L)$ is a closed uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}$.

The converse implication follows from Corollary 2.3, considering the J_ρ -selfadjoint projection $Q \in L(\mathcal{K} \times \mathcal{E})$ onto $R(L)$. \square

4.1. Every indefinite smoothing spline is an indefinite interpolating spline

This subsection is devoted to show that $sm(T, V, \rho, z_0) = sp(T, V, z')$ for a suitable $z' \in \mathcal{E}$. In order to do so, a particular decomposition of $R(L)$ is needed. If $T(N(V))$ is a regular subspace of \mathcal{K} and Q_0 is the projection considered in Eq. (9), consider the (bounded) operator $U : \mathcal{E} \rightarrow \mathcal{K} \times \mathcal{E}$ given by

$$Uz = (T(I - Q_0)V^\dagger z, z), \quad z \in \mathcal{E}.$$

Observe that $N(U) = \{0\}$ and $R(U)$ is closed (because it is isometrically isomorphic to the graph of the bounded operator $T(I - Q_0)V^\dagger$).

Lemma 4.6. *If $T(N(V))$ is a regular subspace of \mathcal{K} then*

$$R(L) = (T(N(V)) \times \{0\}) \dot{+} R(U),$$

and this decomposition of $R(L)$ is orthogonal in the Krein space $(\mathcal{K} \times \mathcal{E}, [\cdot, \cdot]_\rho)$.

Proof. Since $R(Q_0) = N(V)$, observe that $R(L) = L(N(V)) + L(N(Q_0))$ and $L(N(V)) = T(N(V)) \times \{0\}$. In order to compute $L(N(Q_0))$, observe that $I - Q_0 = (I - Q_0)P_{N(V)^\perp} = (I - Q_0)V^\dagger V$ because $N(I - Q_0) = N(P_{N(V)^\perp}) = N(V)$. Therefore, if $x \in N(Q_0)$,

$$Lx = (Tx, Vx) = (T(I - Q_0)x, Vx) = (T(I - Q_0)V^\dagger Vx, Vx) = (T(I - Q_0)V^\dagger z, z) = Uz,$$

where $z = Vx$. Since $V(N(Q_0)) = \mathcal{E}$, it follows that $L(N(Q_0)) = \{(T(I - Q_0)V^\dagger z, z) : z \in \mathcal{E}\} = R(U)$. Finally, since $T(N(Q_0)) \subseteq T(N(V))^{\perp\perp}$, it follows that $L(N(V)) \perp_\rho L(N(Q_0))$. \square

The next theorem shows the existence of a vector $z' \in \mathcal{E}$ such that $sm(T, V, \rho, z_0) = sp(T, V, z')$. Also, along the proof, an expression of such z' is given in terms of the J_ρ -selfadjoint projection onto one of the subspaces of $R(L)$ presented in the above decomposition.

Theorem 4.7. *Suppose that $T(N(V))$ is a closed subspace of \mathcal{K} and $R(L)$ is a uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}$. Then, given $z_0 \in \mathcal{E}$, $sm(T, V, \rho, z_0) = sp(T, V, z')$, where z' is an adequate vector in \mathcal{E} .*

Proof. If $z_0 = 0$ then $sm(T, V, \rho, 0) = N(L) = N(T) \cap N(V) = sp(T, V, 0)$. On the other hand, notice that $R(L)$ is closed (see Lemma 4.4). Then, by Proposition 2.2, $R(L)$ and $T(N(V))$ are regular subspaces of $\mathcal{K} \times \mathcal{E}$ and \mathcal{K} , respectively. So, the projection considered in Eq. (9) is bounded. Given $x \in \mathcal{H}$, it can be decomposed as

$$x = Q_0x + (I - Q_0)x = Q_0x + (I - Q_0)P_{N(V)^\perp}x = v + (I - Q_0)V^\dagger z,$$

where $v = Q_0x \in N(V)$ and $z = Vx \in \mathcal{E}$. Observe that, by Lemma 4.6,

$$[Lx - (0, z_0), Lx - (0, z_0)]_\rho = [(Tv, 0), (Tv, 0)]_\rho + [Uz - (0, z_0), Uz - (0, z_0)]_\rho.$$

Then, $x_0 \in sm(T, V, \rho, z_0)$ if and only if $[TQ_0x_0, TQ_0x_0]_\mathcal{K} = \min_{u \in N(V)} [Tu, Tu]_\mathcal{K}$ and $z_1 = Vx_0$ satisfies

$$[Uz_1 - (0, z_0), Uz_1 - (0, z_0)]_\rho = \min_{z \in \mathcal{E}} [Uz - (0, z_0), Uz - (0, z_0)]_\rho.$$

Notice that $\min_{u \in N(V)} [Tu, Tu]_\mathcal{K}$ is attained at every $u \in N(T) \cap N(V)$, because $T(N(V))$ is uniformly $J_\mathcal{K}$ -positive. Therefore, $Q_0x_0 \in N(T) \cap N(V)$.

On the other hand, since $R(U)$ is a regular subspace of $R(L)$ (see Lemma 4.6), $R(U)$ is a (closed) uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}$. Thus, by Corollary 2.3, z_1 satisfies the above equation if and only if $Uz_1 = P(0, z_0)$, where P is the J_ρ -selfadjoint projection onto $R(U)$.

If $S : \mathcal{K} \times \mathcal{E} \rightarrow \mathcal{E}$ is defined as $S(y, z) = z$ then $SU = I_{\mathcal{E}}$ and $z_1 = SUz_1 = SP(0, z_0)$. So, $(I - Q_0)V^\dagger z_1 = (I - Q_0)V^\dagger SP(0, z_0)$. Therefore, $x_0 \in sm(T, V, \rho, z_0)$ if and only if $x_0 \in (I - Q_0)V^\dagger SP(0, z_0) + N(T) \cap N(V)$, i.e.

$$sm(T, V, \rho, z_0) = sp(T, V, SP(0, z_0)).$$

4.2. The smoothing splines converge to the interpolating spline

In the following paragraph we show that, given $z_0 \in \mathcal{E}$, if $\{x_\rho\}_{\rho \geq 1}$ is a net in \mathcal{H} such that $x_\rho \in sm(T, V, \rho, z_0)$, then it converges to an interpolating spline $x_0 \in sp(T, V, z_0)$ as $\rho \rightarrow \infty$. The proof of this result is analogous to [17, Ch. III, Proposition 2.2].

Proposition 4.8. *Given a fixed vector $z_0 \in \mathcal{E}$, suppose that $T(N(V))$ is a closed subspace of \mathcal{K} , $N(T) \cap N(V) = \{0\}$ and $R(L)$ is a uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}$. Let $x_\rho \in sm(T, V, \rho, z_0)$ for every $\rho \geq 1$. Then, there exists $x_0 \in sp(T, V, z_0)$ such that*

$$\lim_{\rho \rightarrow \infty} \|x_\rho - x_0\| = 0.$$

Proof. First, observe that if $x_\rho \in sm(T, V, \rho, z_0)$ then $\{[Tx_\rho, Tx_\rho]\}_{\rho \geq 1}$ is an increasing net in \mathbb{R} with an upper bound, and $\|Vx_\rho - z_0\| \rightarrow 0$ as $\rho \rightarrow \infty$. Indeed, given $\rho_1, \rho_2 \geq 1$, notice that $[Tx_{\rho_1}, Tx_{\rho_1}] + \rho_1 \|Vx_{\rho_1} - z_0\|^2 \leq [Tx_{\rho_j}, Tx_{\rho_j}] + \rho_j \|Vx_{\rho_j} - z_0\|^2$, if $i \neq j$. Then, if $\rho_1 < \rho_2$ it follows that $\|Vx_{\rho_1} - z_0\|^2 - \|Vx_{\rho_2} - z_0\|^2 \geq 0$ and

$$[Tx_{\rho_2}, Tx_{\rho_2}] - [Tx_{\rho_1}, Tx_{\rho_1}] \geq \rho_1 (\|Vx_{\rho_1} - z_0\|^2 - \|Vx_{\rho_2} - z_0\|^2) \geq 0.$$

Furthermore, if $x \in sp(T, V, z_0)$ for every $\rho \geq 1$, $[Tx_\rho, Tx_\rho] + \rho \|Vx_\rho - z_0\|^2 \leq [Tx, Tx] + \rho \|Vx - z_0\|^2 = [Tx, Tx]$. So, $[Tx, Tx] - [Tx_\rho, Tx_\rho] \geq \rho \|Vx_\rho - z_0\|^2 \geq 0$ for every $\rho \geq 1$, and this inequality implies that

$$\lim_{\rho \rightarrow \infty} \|Vx_\rho - z_0\| = 0.$$

The next step is to prove that $\lim_{\rho \rightarrow \infty} \|x_\rho - x_0\| = 0$, where $x_0 = V^\dagger z_0 + u$ for some $u \in N(V)$. Let $y_\rho = P_{N(V)^\perp} x_\rho$ and observe that $y_\rho = V^\dagger Vx_\rho \rightarrow V^\dagger z_0$ as $\rho \rightarrow \infty$.

If $u_\rho = x_\rho - y_\rho = P_{N(V)} x_\rho \in N(V)$, then $\{u_\rho\}_{\rho \geq 1}$ converges to some $u \in N(V)$. To prove this assertion, consider the closed range operator $W = T|_{N(V)} : N(V) \rightarrow \mathcal{K}$ (see Lemma 4.4). If Q is the $J_{\mathcal{K}}$ -selfadjoint projection onto $T(N(V))$, let $W' = W^\dagger Q$. Then, W' satisfies $WW'W = W$, $W'W'W' = W'$ and $N(W') = T(N(V))^{\perp\perp}$. By Theorem 4.7, $x_\rho \in sp(T, V, z_\rho)$ for a suitable $z_\rho \in \mathcal{E}$; then, it follows that $Tx_\rho \in T(N(V))^{\perp\perp}$ (see Lemma 3.4). Therefore, $W'Tx_\rho = 0$ for every $\rho \geq 1$, and

$$W'Tu_\rho = -W'Ty_\rho \rightarrow -W'TV^\dagger z_0 = u \in R(W') \subseteq N(V) \quad \text{as } \rho \rightarrow \infty. \quad \square$$

5. The indefinite abstract mixed problem

Given Hilbert spaces \mathcal{H} , \mathcal{E}_1 and \mathcal{E}_2 , and a Krein space \mathcal{K} with fundamental symmetry $J_{\mathcal{K}}$, let $T \in L(\mathcal{H}, \mathcal{K})$, $V_1 \in L(\mathcal{H}, \mathcal{E}_1)$ and $V_2 \in L(\mathcal{H}, \mathcal{E}_2)$ be surjective operators. Then, consider the following problem:

Problem 5.1. Let $\rho > 0$. For a fixed $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$, find $x_0 \in \mathcal{H}$ such that $V_1 x_0 = z_1$ and

$$([Tx_0, Tx_0]_{\mathcal{K}} + \rho \|V_2 x_0 - z_2\|_{\mathcal{E}_2}^2) = \min_{V_1 x = z_1} ([Tx, Tx]_{\mathcal{K}} + \rho \|V_2 x - z_2\|_{\mathcal{E}_2}^2). \tag{19}$$

This is a generalization to Krein spaces of the mixed problem in Hilbert spaces proposed by A.I. Rozhenko in [16] (see also [21,22]).

It is clear that the indefinite abstract and smoothing problems are the partial cases of the indefinite abstract mixed problem corresponding to $\mathcal{E}_2 = \{0\}$, $V_2 = 0$ and $\mathcal{E}_1 = \{0\}$, $V_1 = 0$, respectively. Thus, it is expected that similar results to those given in the previous sections, can be stated with some additional restrictions. We prefer to introduce the indefinite abstract mixed problem after studying the other problems in order to motivate it.

As in the previous section, $\mathcal{K} \times \mathcal{E}_2$ is a Krein space with the indefinite metric defined in Eq. (17) and its fundamental symmetry $J_\rho \in L(\mathcal{K} \times \mathcal{E}_2)$ is given by $J_\rho(y, z) = (J_{\mathcal{K}}y, z)$, where $(y, z) \in \mathcal{K} \times \mathcal{E}_2$. Also, consider the operators $L \in L(\mathcal{H}, \mathcal{K} \times \mathcal{E}_2)$ given by

$$Lx = (Tx, V_2x), \quad x \in \mathcal{H},$$

and $L_1 = LP_{N(V_1)} \in L(\mathcal{H}, \mathcal{K} \times \mathcal{E}_2)$. Then, Problem 5.1 can be restated as: given $\rho > 0$ and a fixed $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$, find $x_0 \in \mathcal{H}$ such that

$$[L_1x_0 - (w_1, w_2), L_1x_0 - (w_1, w_2)]_\rho = \min_{x \in \mathcal{H}} [L_1x - (w_1, w_2), L_1x - (w_1, w_2)]_\rho, \tag{20}$$

where $w_1 = -TV_1^\dagger z_1$ and $w_2 = z_2 - V_2V_1^\dagger z_1$.

Lemma 5.2. *Given $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$, $x_0 \in \mathcal{H}$ is a solution of Problem 5.1 if and only if $R(L_1)$ is J_ρ -nonnegative and x_0 is a solution of the equation:*

$$P_{N(V_1)}(T^+T + \rho V_2^*V_2)P_{N(V_1)}x_0 = P_{N(V_1)}(T^+w_1 + \rho V_2^*w_2).$$

Proof. It is analogous to the proof of Lemma 4.3. Notice that, in this case, $L_1^+ \in L(\mathcal{K} \times \mathcal{E}_2, \mathcal{H})$ is given by $L_1^+(y, z) = P_{N(V_1)}L^+(y, z) = P_{N(V_1)}(T^+y + \rho V_2^*z)$, $(y, z) \in \mathcal{K} \times \mathcal{E}_2$. \square

Proposition 5.3. *Problem 5.1 admits a solution for every $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$ if and only if $R(L_1)$ is a closed uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}_2$.*

Proof. Suppose that, Problem 5.1 admits a solution for every $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$. Given $(y, z) \in \mathcal{K} \times \mathcal{E}_2$, let $z_1 = -V_1T^\dagger y$ and $z_2 = z - V_2T^\dagger y$. Consider $x_0 = u + T^\dagger y$, where $u \in \mathcal{H}$ satisfies

$$[L_1u - (w_1, w_2), L_1u - (w_1, w_2)]_\rho = \min_{x \in \mathcal{H}} [L_1x - (w_1, w_2), L_1x - (w_1, w_2)]_\rho,$$

for this particular pair $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$.

Observe that $L_1x_0 - (y, z) = L_1u + (TP_{N(V_1)}T^\dagger y, V_2P_{N(V_1)}T^\dagger y) - (y, z) = L_1u - (-TV_1^\dagger z_1, z_2 - V_2V_1^\dagger z_1) = L_1u - (w_1, w_2)$. Then, for every $x \in \mathcal{H}$,

$$\begin{aligned} [L_1x_0 - (y, z), L_1x_0 - (y, z)]_\rho &= [L_1u - (w_1, w_2), L_1u - (w_1, w_2)]_\rho \\ &\leq [L_1(x - T^\dagger y) - (w_1, w_2), L_1(x - T^\dagger y) - (w_1, w_2)]_\rho \\ &= [L_1x - (y, z), L_1x - (y, z)]_\rho, \end{aligned}$$

because $L_1x - (y, z) = L_1(x - T^\dagger y) - (w_1, w_2)$. Therefore, for every $(y, z) \in \mathcal{K} \times \mathcal{E}_2$, there exists $x_0 \in \mathcal{H}$ such that

$$[L_1x_0 - (y, z), L_1x_0 - (y, z)]_\rho = \min_{x \in \mathcal{H}} [L_1x - (y, z), L_1x - (y, z)]_\rho.$$

Following the same arguments as in the proof of Proposition 4.5, it is easy to see that the above condition holds if and only if $R(L_1)$ is a closed uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}_2$. \square

5.1. Parametrization of the set of solutions of the indefinite abstract mixed problem

The following paragraphs follow analogous ideas to those presented in the previous section to show that every smoothing spline is an interpolating spline.

Consider the operator $V \in L(\mathcal{H}, \mathcal{E}_1 \times \mathcal{E}_2)$ given by $Vx = (V_1x, V_2x)$, $x \in \mathcal{H}$, and notice that $N(V) = N(V_1) \cap N(V_2)$ but V is not surjective. However, Lemma 3.7 also holds in this case. So, if $T(N(V))$ is a regular subspace of \mathcal{K} then, denoting $\mathcal{W} = T^+T(N(V))^\perp \ominus N(V)$, the projection $Q_0 = P_{N(V)/\mathcal{W}}$ is bounded. Before stating the main theorem, we need the following key lemma.

Lemma 5.4. *Suppose that $T(N(V))$ is a regular subspace of \mathcal{K} and $N(V_1) + N(V_2)$ is closed in \mathcal{H} . Then,*

- (i) $\mathcal{M}_1 = (I - Q_0)(N(V_1))$ and $\mathcal{M}_2 = V_2(N(V_1))$ are closed subspaces of \mathcal{H} and \mathcal{E}_2 , respectively.
- (ii) $V_2|_{\mathcal{M}_1} : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ is an isomorphism.
- (iii) $R(L_1) = (T(N(V)) \times \{0\}) \dot{+} L(\mathcal{M}_1)$. Furthermore, $L(\mathcal{M}_1)$ is closed in $\mathcal{K} \times \mathcal{E}_2$ and the decomposition is orthogonal in the Krein space $(\mathcal{K} \times \mathcal{E}_2, [\cdot, \cdot]_\rho)$.

Proof. (i) First of all, notice that $\mathcal{M}_1 = R(I - Q_0) \cap N(V_1)$. Therefore, it is closed and $N(V_1) = N(V) \dot{+} \mathcal{M}_1$ because $Q_0(N(V_1)) = N(V)$. On the other hand, by Proposition 2.5, $\mathcal{M}_2 = R(V_2P_{N(V_1)})$ is closed if and only if $c(N(V_2), N(V_1)) < 1$, or equivalently, $N(V_1) + N(V_2)$ is closed. Therefore, \mathcal{M}_2 is closed.

(ii) To show that $V_2|_{\mathcal{M}_1} : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ is an isomorphism observe that $V_2(\mathcal{M}_1) = V_2(\mathcal{M}_1 + N(V)) = V_2(N(V_1)) = \mathcal{M}_2$, so it only remains to prove that $V_2|_{\mathcal{M}_1}$ is injective. But, if $x \in \mathcal{M}_1$ and $V_2x = 0$ then $x \in N(V_2) \cap \mathcal{M}_1 = N(V) \cap R(I - Q_0) = \{0\}$.

(iii) Observe that $R(L_1) = L(N(V_1)) = L(N(V)) + L(\mathcal{M}_1)$ because $N(V_1) = N(V) \dot{+} \mathcal{M}_1$. Furthermore, if $x \in N(V_1)$ then $Q_0x \in N(V)$ and $(I - Q_0)x \in \mathcal{M}_1$. So, $Lx = (TQ_0x, 0) + L(I - Q_0)x$. Therefore, $R(L_1) = L(N(V)) + L(\mathcal{M}_1) = (T(N(V)) \times \{0\}) \dot{+} L(\mathcal{M}_1)$.

If $(y, 0) \in (T(N(V)) \times \{0\}) \cap L(\mathcal{M}_1)$, there exists $m \in \mathcal{M}_1$ such that $Tm = y$ and $V_2m = 0$. Then, $m = 0$ because $V_2|_{\mathcal{M}_1}$ is an isomorphism. So, $y = Tm = 0$ and $R(L_1) = L(N(V)) \dot{+} L(\mathcal{M}_1)$. As in Lemma 4.6, it is easy to see that this decomposition is orthogonal respect to the indefinite metric defined on $\mathcal{K} \times \mathcal{E}_2$.

It only remains to prove that $L(\mathcal{M}_1)$ is a closed subspace of $\mathcal{K} \times \mathcal{E}_2$. Given $(y, z) \in \overline{L(\mathcal{M}_1)}$ consider $\{m_k\}_{k \geq 1} \subseteq \mathcal{M}_1$ such that $Tm_k \rightarrow y$ and $V_2m_k \rightarrow z$ as $k \rightarrow \infty$. Notice that $m_k = (V_2|_{\mathcal{M}_1})^{-1}V_2m_k$, because $V_2|_{\mathcal{M}_1} : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ is an isomorphism. Therefore, $m_k \rightarrow (V_2|_{\mathcal{M}_1})^{-1}z \in \mathcal{M}_1$ and $(y, z) = L((V_2|_{\mathcal{M}_1})^{-1}z)$. \square

Corollary 5.5. *If $T(N(V))$ is a regular subspace of \mathcal{K} and $N(V_1) + N(V_2)$ is closed in \mathcal{H} then $R(L_1)$ is closed in $\mathcal{K} \times \mathcal{E}_2$.*

The next theorem shows that every mixed spline is an interpolating spline.

Theorem 5.6. *Suppose that $N(V_1) + N(V_2)$ is closed in \mathcal{K} , $T(N(V))$ is a closed subspace of \mathcal{K} and $R(L_1)$ is a (closed) uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}_2$. Then, given $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$, an element $x_0 \in \mathcal{H}$ is a solution of Problem 5.1 if and only if*

$$x_0 \in sp(T, V, (e_1, e_2)),$$

where (e_1, e_2) is a suitable vector in $\mathcal{E}_1 \times \mathcal{E}_2$.

Proof. Given $(z_1, z_2) \in \mathcal{E}_1 \times \mathcal{E}_2$, recall that if $x_0 \in \mathcal{H}$ is a solution of Problem 5.1 then $V_1x_0 = z_1$, or equivalently, $P_{N(V_1)^\perp}x_0 = V_1^\dagger z_1$. Assuming that $T(N(V))$ is a regular subspace of \mathcal{K} , $V_1^\dagger z_1$ can be decomposed as $V_1^\dagger z_1 = u_1 + v_1$, where $u_1 = Q_0V_1^\dagger z_1 \in N(V)$ and $v_1 = (I - Q_0)V_1^\dagger z_1 \in \mathcal{W}$. Then, the pair (w_1, w_2) considered in Eq. (20) satisfies

$$-w_1 = TV_1^\dagger z_1 = Tu_1 + Tv_1 \in T(N(V)) \dot{+} T(N(V))^{\perp\perp} \quad \text{and} \quad w_2 = z_2 - V_2v_1.$$

If $N(V_1) + N(V_2)$ is a closed subspace of \mathcal{H} , given $x \in \mathcal{H}$ there exist (unique) $u \in N(V)$ and $m \in \mathcal{M}_1$ such that $P_{N(V_1)^\perp}x = u + m$ (see Lemma 5.4). Thus, $x = u + m + P_{N(V)^\perp}x$ and

$$L_1x - (w_1, w_2) = (T(u + u_1), 0) + Lm - (-Tv_1, w_2).$$

Observe that $Lm - (-Tv_1, w_2) = L(m + v_1) - (0, z_2) \in (T(N(V)) \times \{0\})^{\perp\perp}$ because $m + v_1 \in N(Q_0)$. Then,

$$[L_1x - (w_1, w_2), L_1x - (w_1, w_2)]_\rho = [T(u + u_1), T(u + u_1)]_{\mathcal{K}} + [Lm - (-Tv_1, w_2), Lm - (-Tv_1, w_2)]_\rho.$$

Therefore, x_0 is a solution to Problem 5.1 if and only if $P_{N(V_1)^\perp}x_0 = u_0 + m_0$, with $u_0 \in N(V)$ and $m_0 \in \mathcal{M}_1$ satisfying $[T(u_0 + u_1), T(u_0 + u_1)]_{\mathcal{K}} = \min_{u \in N(V)} [T(u + u_1), T(u + u_1)]_{\mathcal{K}}$ and

$$[Lm_0 - (-Tv_1, w_2), Lm_0 - (-Tv_1, w_2)]_\rho = \min_{m \in \mathcal{M}_1} [Lm - (-Tv_1, w_2), Lm - (-Tv_1, w_2)]_\rho.$$

Notice that, if $R(L_1)$ is a closed uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}_2$, then $T(N(V))$ is a closed uniformly $J_{\mathcal{K}}$ -positive subspace of \mathcal{K} and $\min_{u \in N(V)} [T(u + u_1), T(u + u_1)]_{\mathcal{K}}$ is attained at every $y \in -u_1 + N(V) \cap N(T)$.

On the other hand, consider the bounded operator $U : \mathcal{M}_2 \rightarrow \mathcal{K} \times \mathcal{E}_2$ defined by

$$Uz = (T(V_2|_{\mathcal{M}_1})^{-1}z, z).$$

Observe that U has closed range, because it is isometrically isomorphic to the graph of the bounded operator $T(V_2|_{\mathcal{M}_1})^{-1}$, and

$$\min_{m \in \mathcal{M}_1} [Lm - (-Tv_1, w_2), Lm - (-Tv_1, w_2)]_\rho = \min_{z \in \mathcal{M}_2} [Uz - (-Tv_1, w_2), Uz - (-Tv_1, w_2)]_\rho.$$

Thus, following the same argument as in Theorem 4.7 and observing that $R(U) = L(\mathcal{M}_1)$ is a closed uniformly J_ρ -positive subspace of $\mathcal{K} \times \mathcal{E}_2$, this last problem admits a (unique) solution given by $z_0 = V_2m_0 = SP(-Tv_1, w_2)$, where P is the J_ρ -selfadjoint projection onto $L(\mathcal{M}_1)$ and $S : \mathcal{K} \times \mathcal{E}_2 \rightarrow \mathcal{E}_2$ is defined by $S(y, z) = z$. So, $x_0 \in \mathcal{H}$ is a solution to Problem 5.1 if and only if

$$x_0 = V_1^\dagger z_1 + P_{N(V_1)^\perp}x_0 = u_1 + v_1 + u_0 + m_0 \in (v_1 + (V_2|_{\mathcal{M}_1})^{-1}SP(-Tv_1, w_2)) + N(T) \cap N(V).$$

Therefore, $x_0 \in \mathcal{H}$ is a solution to Problem 5.1 if and only if $x_0 \in sp(T, V, (e_1, e_2))$, where

$$e_1 = z_1 + V_1(V_2|_{\mathcal{M}_1})^{-1}SP(-Tv_1, w_2) \in \mathcal{E}_1 \quad \text{and} \quad e_2 = V_2V_1^\dagger z_1 + SP(-Tv_1, w_2) \in \mathcal{E}_2. \quad \square$$

Acknowledgments

The authors would like to acknowledge Professor Gustavo Corach who brought into their attention the mixed problem for abstract splines in Hilbert spaces. Also they would like to thank the referee for his useful comments.

Francisco Martínez Pería would like to thank Centre de Recerca Matemàtica staff for their hospitality while he held a Lluís Santaló visiting position in Barcelona, Spain, and to Fundació Carolina for supporting him during this visit with a Formació Permanente fellowship.

References

- [1] I.J. Schoenberg, Contributions to the problem of approximation of equidistant data by analytic functions. Parts A, B, Q. *J. Math.* 4 (1946) 45–99, 112–141.
- [2] H.S. Hou, H.C. Andrews, Cubic splines for image interpolation and digital filtering, *IEEE Trans. ASSP* 26 (1978) 508–517.
- [3] K. Torachi, S. Yang, M. Kamada, R. Mori, Two dimensional spline interpolation for image reconstruction, *Patt. Recognition* 21 (1998) 275–284.
- [4] M. Unser, A. Aldroubi, M. Eden, Fast B-spline transform for continuous image reconstruction and interpolation, *IEEE Trans. Patt. Anal. Machine Intell.* 13 (1991) 277–285.
- [5] J.I. Giribet, M. España, C. Miranda, Synthetic data for validation of navigation systems, *Acta Astronaut.* 60 (2) (2007) 88–95.
- [6] R.H. Bartels, J.C. Beatty, B.A. Barsky, *An Introduction to Splines for Use in Computer Graphics & Geometric Modeling*, Morgan Kaufmann Publishers Inc., San Francisco, 1987.
- [7] D.S. Meek, D.J. Walton, A note on planar minimax arc splines, *Comput. & Graphics* 16 (1992) 431–433.
- [8] D.S. Meek, D.J. Walton, Approximating smooth planar curves by arc splines, *J. Comput. Appl. Math.* 59 (1995) 221–231.
- [9] S. Canu, C.S. Ong, X. Mary, A. Smola, Learning with non-positive kernels, in: *Proceedings of the 21st International Conference on Machine Learning*, 2004, pp. 639–646.
- [10] S. Canu, C.S. Ong, X. Mary, Splines with non positive kernels, in: *Proceedings of the 5th International ISAAC Congress*, 2005, pp. 1–10.
- [11] M. Atteia, Généralization de la définition et des propriétés des “splines fonctions”, *C. R. Sci. Paris* 260 (1965) 3550–3553.
- [12] P.M. Anselone, P.J. Laurent, A general method for the construction of interpolating or smoothing spline-functions, *Numer. Math.* 12 (1968) 66–82.
- [13] C. de Boor, Convergence of abstract splines, *J. Approx. Theory* 31 (1981) 80–89.
- [14] P.J. Laurent, *Approximation et optimisation*, Hermann, Paris, 1972.
- [15] A. Sard, Optimal approximation, *J. Funct. Anal.* 1 (1967) 222–244; *J. Funct. Anal.* 2 (1968) 368–369 (addendum).
- [16] A.I. Rozhenko, Mixed spline approximation, *Bull. Novosibirsk Comput. Center Ser.: Numer. Anal.* 5 (1994) 67–86.
- [17] M. Atteia, *Hilbertian Kernels and Spline Functions*, North-Holland Publishing Co., Amsterdam, 1992.
- [18] A.Yu. Bezhaev, V.A. Vasilenko, *Variational Theory of Splines*, Kluwer Academic/Plenum Publishers, New York, 2001.
- [19] R. Champion, C.T. Lenard, T.M. Mills, An introduction to abstract splines, *Math. Sci.* 21 (1996) 8–26.
- [20] S. Hassi, K. Nordström, On projections in a space with an indefinite metric, *Linear Algebra Appl.* 208/209 (1994) 401–417.
- [21] A.I. Rozhenko, The conditions of unique solvability of mixed spline approximation problem, Preprint N1023, *Comp. Center USSR Acad. Sci. Press*, Novosibirsk, 1994 (in Russian).
- [22] A.I. Rozhenko, V.A. Vasilenko, Variational approach in abstract splines: achievements and open problems, *East J. Approx.* 1 (1995) 277–308.
- [23] J. Bognár, *Indefinite Inner Product Spaces*, Springer-Verlag, 1974.
- [24] I.S. Iokhvidov, T.Ya. Azizov, *Linear Operators in Spaces with an Indefinite Metric*, John Wiley & Sons, 1989.
- [25] T. Ando, *Linear Operators on Krein Spaces*, Hokkaido University, Sapporo, Japan, 1979.
- [26] M.A. Dritschel, J. Rovnyak, Operators on indefinite inner product spaces, in: Peter Lancaster (Ed.), *Fields Inst. Monogr.*, vol. 3, Amer. Math. Soc., 1996, pp. 141–232.
- [27] J. Rovnyak, Methods on Krein space operator theory, in: *Interpolation Theory, Systems Theory and Related Topics*, Tel Aviv/Rehovot, 1999, in: *Oper. Theory Adv. Appl.*, vol. 134, 2002, pp. 31–66.
- [28] F. Deutsch, The angle between subspaces of a Hilbert space, in: *Approximation Theory, Wavelets and Applications*, Maratea, 1994, in: *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, vol. 454, Kluwer Acad. Publ., Dordrecht, 1995, pp. 107–130.
- [29] R. Bouldin, The product of operators with closed range, *Tohoku Math. J.* 25 (1973) 359–363.
- [30] S. Izumino, The product of operators with closed range and an extension of the reverse order law, *Tohoku Math. J.* 34 (1982) 43–52.
- [31] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, New York, 1966.
- [32] H. Patel, Solving the indefinite least squares problem, PhD thesis, Univ. of Manchester, 2002.
- [33] P. Casazza, O. Christensen, Frames containing a Riesz basis and preservation of this property under perturbations, *SIAM J. Math. Anal.* 29 (1) (1998) 266–278.
- [34] D.F. Walnut, *An Introduction to Wavelets Analysis*, Springer-Verlag, 2004.