

Therefore, comparative genomic analysis of different plasmids carrying *bla*_{CTX-M-2} might be useful to fully understand their evolution, plasticity and spread. While *bla*_{CTX-M-2} has frequently been detected in human pathogens,¹ this is the first report of *bla*_{CTX-M-2}-producing *E. coli* isolated from diseased animals and, more specifically, horses. The emergence of *bla*_{CTX-M-2} is therefore a clear cause for concern.

Acknowledgements

We thank Mrs Nathalie Van Ryselberghe and Isabelle Monchaux for their skilled technical assistance.

Funding

This work was supported by internal funding.

Transparency declarations

None to declare.

Supplementary data

Table S1 and Figure S1 are available as Supplementary data at JAC Online (<http://jac.oxfordjournals.org/>).

References

- 1 Smet A, Martel A, Persoons D *et al.* Broad-spectrum β -lactamases among Enterobacteriaceae of animal origin: molecular aspects, mobility and impact on public health. *FEMS Microbiol Rev* 2010; **34**: 295–316.
- 2 Poupard MC, Chanal C, Sirot D *et al.* Identification of CTX-2, a novel cefotaximase from a *Salmonella mbandaka* isolate. *Antimicrob Agents Chemother* 1991; **35**: 1498–500.
- 3 Bertrand S, Weill FX, Cloeckaert A *et al.* Clonal emergence of extended-spectrum β -lactamase (CTX-M-2)-producing *Salmonella enterica* serovar Virchow isolates with reduced susceptibilities to ciprofloxacin among poultry and humans in Belgium and France (2000 to 2003). *J Clin Microbiol* 2006; **44**: 2897–903.
- 4 Karczmarczyk M, Abbott Y, Walsh C *et al.* Characterization of multidrug-resistant *Escherichia coli* from animals presenting at a university veterinary hospital. *Appl Environ Microbiol* 2011; **77**: 7104–12.
- 5 Smet A, Martel A, Persoons D *et al.* Comparative analysis of extended-spectrum- β -lactamase-carrying plasmids from different members of Enterobacteriaceae isolated from poultry, pigs and humans: evidence for a shared β -lactam resistance gene pool? *J Antimicrob Chemother* 2009; **63**: 1286–8.
- 6 Members of the SFM Antibiogram Committee. Comité de l'Antibiogramme de la Société Française de Microbiologie report 2003. *Int J Antimicrob Agents* 2003; **21**: 364–91.
- 7 Carattoli A, Bertini A, Villa L *et al.* Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005; **63**: 219–28.
- 8 Lévesque C, Piché L, Larose C *et al.* PCR mapping of integrons reveals several novel combinations of resistance genes. *Antimicrob Agents Chemother* 1995; **39**: 185–91.

J Antimicrob Chemother 2012

doi:10.1093/jac/dks032

Advance Access publication 20 February 2012

Detection of an international multiresistant clone belonging to sequence type 654 involved in the dissemination of KPC-producing *Pseudomonas aeruginosa* in Argentina

Fernando Pasteran¹, Diego Faccione¹, Sonia Gomez¹, Sabrina De Bunder², Federico Spinelli^{3,4}, Melina Rapoport¹, Alejandro Petroni¹, Marcelo Galas¹ and Alejandra Corso^{1*} on behalf of the *Pseudomonas aeruginosa* KPC Group†

¹Servicio Antimicrobianos, Departamento Bacteriología, Instituto Nacional de Enfermedades Infecciosas (INEI), ANLIS 'Dr Carlos G. Malbrán', Ciudad Autónoma de Buenos Aires, Argentina;

²Bacteriología, Hospital Zonal de Bariloche 'Ramón Carillo', Río Negro, Argentina; ³Infectología, Hospital Privado Regional del Sur, Río Negro, Argentina; ⁴Infectología, Sanatorio del Sol, Bariloche, Río Negro, Argentina

*Corresponding author. Tel/Fax: +54-11-4303-2812;

E-mail: acorso@anlis.gov.ar

†Members of the *Pseudomonas aeruginosa* KPC Group are given in the Acknowledgements section.

Keywords: clone, multidrug resistance, outbreak

Sir,

The emergence of *Klebsiella pneumoniae* carbapenemase (KPC) has now become a global concern. KPC producers are mostly Enterobacteriaceae, but *Pseudomonas aeruginosa* have also been reported and mostly identified in the American continent.^{1–4} However, it is unknown if this is due to the spread of epidemic strains, since the multilocus sequence type (ST) has not been provided in most of those reports. The aim of this work was to characterize KPC-producing *P. aeruginosa* isolated in Argentina from 2006 to June 2011.

Since 2005, we designed an algorithm to detect carbapenemases (metallo- β -lactamases, KPC, etc.) in *P. aeruginosa* at the level of the clinical microbiology laboratory, which was implemented among 432 hospitals. All *P. aeruginosa* were screened through that algorithm, and KPC production was suspected in isolates with high-level resistance to carbapenems and aztreonam (absence of disc zones) and a negative synergy test result between the carbapenems and EDTA, a phenotype consistent with the reported patterns of KPC-producing *P. aeruginosa*.^{1,4}

As a result, 65 isolates were suspected to be KPC producers (Table 1). Strains were isolated from nine hospitals (seven cities, five provinces), six of which were located in the Patagonia region. The first KPC producer was detected in a hospital from Bariloche in 2006. Dissemination to other locations (except

Table 1. Epidemiological and molecular characteristics of KPC-positive *P. aeruginosa* isolates

Year	City, Province	Hospital(s) ^a	No. of isolates suspected of KPC/total no. of <i>P. aeruginosa</i> isolates (%) ^b	No. of isolates submitted to the INEI	PFGE pattern (ST)	Susceptibility to non-β-lactam agents ^c
Pre-epidemic period						
2005 ^d	Argentina (24 provinces)	WHONET AR	0/8846 (0)	0	NA ^e	NA
Epidemic period						
2006	Bariloche, Rio Negro	SSC	3/28 (11)	2	A (654)	S: CST, AMK, GEN
2008	Bariloche, Rio Negro	SSC, HPR, BAR	16/26 (62)	7	A	S: CST, AMK, GEN
	Lago Puelo, Chubut	HZLP	1 ^f /2	1	A	S: CST, AMK, GEN
2009	Bariloche, Rio Negro	BAR, HPR	12/48 (25)	4	A	S: CST, AMK, GEN
	General Roca, Rio Negro	CGR	3 ^f /43 (7)	1	A	S: CST, AMK, GEN
	Esquel, Chubut	HZE	2 ^f /26 (8)	2	A	S: CST, AMK, GEN
	Córdoba, Córdoba	HPC	1 ^f /27 (4)	1	A	S: CST, AMK, GEN
2010	Bariloche, Rio Negro	BAR, HPR	9/29 (31)	4	A	S: CST, AMK, GEN (3) S: CST (1)
	General Roca, Rio Negro	CGR	3/54 (6)	1	A	S: CST, AMK, GEN
	Esquel, Chubut	HZE	1/22 (4)	1	A	S: CST, AMK, GEN
	Ciudad de Buenos Aires	UDA	1/not available	1	A	S: CST, AMK, GEN
	Resistencia, Chaco	HJP	1/167 (0.6)	1	B (162)	S: CST, AMK, GEN, CIP
2011 ^g	Bariloche, Rio Negro	SSC	1/6 (17)	1	A	S: CST, AMK, GEN
	General Roca, Rio Negro	CGR	11/36 (31)	3	A	S: CST, AMK, GEN
total	7 cities/5 Provinces	9 hospitals	65/514 (12)	30	A (654), B (162)	S: CST 100%, AMK 97%, GEN 97%, CIP 3%

INEI, Instituto Nacional de Enfermedades Infecciosas.

^aWHONET AR, WHONET Argentina Network (90 Hospitals, 24 Provinces); SSC, Sanatorio San Carlos; HPR, Hospital Privado Regional del Sur y Sanatorio del Sol; BAR, Hospital Zonal de Bariloche; HZLP, Hospital Zonal de Lago Puelo; CGR, Laboratorio Roca, Clínica Roca; HZE, Hospital Zonal de Esquel; HPC, Hospital Privado Centro Modelo de Córdoba; UDA, Hospital Municipal de Gastroenterología Udaondo; HJP, Hospital Julio Perrando.

^bIsolates suspected of producing KPC defined with the phenotypic algorithm indicated in the text. The denominator represents the total *P. aeruginosa* isolates recovered in the indicated hospital(s) of the corresponding row.

^cAs defined by agar dilution according to CLSI. S, susceptible; CST, colistin; AMK, amikacin; GEN, gentamicin; CIP, ciprofloxacin. The number of isolates is shown in parentheses.

^dBaseline obtained from the WHONET Argentina Network.

^eNA, not applicable.

^fThe respective index patients of these locations were previously hospitalized in Bariloche.

^gJanuary–June 2011.

Chaco) was associated with patients previously hospitalized in Bariloche (Table 1).

A subset ($n=30$) of suspected isolates was submitted to the National Reference Laboratory (Malbrán Institute), where *bla*_{KPC-2} was confirmed in all strains by PCR/sequencing. XbaI PFGE analysis, using previously described criteria,⁵ revealed that all but one of the isolates belonged to a single pulsotype (Table 1). In Argentina, these PFGE patterns had not previously been observed. Multilocus sequence typing of one strain of each PFGE type revealed that the dominant clone (PFGE type A) belonged to ST654, while the unique strain of PFGE type B (Chaco), belonged to ST162 (<http://pubmlst.org/paeruginosa>) (Table 1).

To investigate the genetic organization of *bla*_{KPC-2} in *P. aeruginosa*, we used PCR primer pairs located in the Tn4401 structure and in the flanking sequences found in Enterobacteriaceae with KPC in our country, called Variant 1a (ISKpn8 and ISKpn6-like).⁶ All isolates belonging to ST654 harboured *bla*_{KPC-2}

in Tn4401b, which was confirmed by sequencing. The ST162 isolate harboured *bla*_{KPC-2} in Variant 1a.⁶ By plasmid content analysis and Southern blotting with a *bla*_{KPC-2} probe,⁶ we found that ST654 contains *bla*_{KPC} in a plasmid of 50 kb, while the strain ST162 contains several plasmids but only one of 47 kb hybridized with KPC. Plasmid incompatibility (Inc) groups were further analysed using the PCR-based replicon-typing protocol described for Enterobacteriaceae.⁷ Both ST162 and ST654 gave negative results for all the Inc groups tested.

The strains were highly resistant (MICs >128 mg/L) to aztreonam, cefepime, imipenem, meropenem and piperacillin/tazobactam by agar dilution. Colistin (MICs 1 mg/L), gentamicin (MIC_{90S} 2 mg/L) and amikacin (MIC_{90S} 4 mg/L) were the most active drugs. ST162 was susceptible to ciprofloxacin (MIC 0.5 mg/L) (Table 1).

The origin of *bla*_{KPC-2}-possessing ST654 remains uncertain. The mobilization of *bla*_{KPC} from Enterobacteriaceae was ruled

out: first, these strains were not detected in Bariloche until 2010 and, second, they harboured *bla*_{KPC-2} in the Variant 1a.⁶ ST162 was isolated from a patient with no recent history of travel. Interestingly, this patient shared the same ward simultaneously with two other patients undergoing infections due to KPC-producing *Enterobacter cloacae* and *K. pneumoniae*. In these isolates, the *bla*_{KPC-2} genetic environment matched that of ST162. However, *bla*_{KPC-2} in *E. cloacae* and *K. pneumoniae* was associated with plasmids of different sizes (40 and >150 kb, respectively) and different Inc groups (N and FIIS, respectively)⁶ from that detected in ST162. Thus, we speculated that the surge of ST162 could be due to *bla*_{KPC-2} mobilization among different plasmids (i.e. transposition).

Clonal complexes CC111 and CC235 have been reported as the major clones involved in the global dissemination of extended-spectrum and metallo- β -lactamases in *P. aeruginosa*.⁸ However, we observed that the main dissemination of KPC was mediated by a new clone (ST654) not related to both CC111 and CC235. ST654 is endemic in Singapore,⁹ where it has been associated with IMP-1 and IMP-26; it has also been reported in Sweden as a VIM-2 producer, although the isolate was imported from Tunisia.¹⁰ Unlike other Latin American regions, such as Puerto Rico and Colombia, where different PFGE types and STs have been identified,^{1,4} the dissemination of KPC-producing *P. aeruginosa* in Argentina is mainly associated with a single clone. These findings confirm that ST654 plays an important role in the global spread of carbapenemases, either metallo- β -lactamases or KPC. Thus, the worldwide dissemination of KPC-producing *P. aeruginosa* of ST654 might be expected and should be monitored.

Acknowledgements

Part of this work was presented at the Fifty-first Interscience Conference on Antimicrobial Agents and Chemotherapy, Chicago, IL, 2011 (Abstract no. C2-1700).

Pseudomonas aeruginosa KPC Group

Sanatorio San Carlos, Bariloche, Rio Negro, Argentina: Barbara Bavdaz; Hospital Privado Regional del Sur y Sanatorio del Sol, Bariloche, Rio Negro, Argentina: Gabriela Rubinstein; Hospital Zonal de Bariloche, Bariloche, Rio Negro, Argentina: Nestor Blazquez; Hospital Zonal de Lago Puelo: Pablo Nouveau; Laboratorio Roca, Clínica Roca, General Roca, Rio Negro, Argentina: Silvana Cecchi; Hospital Zonal de Esquel, Chubut, Argentina: Omar Daher; Hospital Privado Centro Modelo de Córdoba, Córdoba: Mario Vilaró; Hospital Municipal de Gastroenterología 'B. Udaondo', Ciudad Autónoma de Buenos Aires, Argentina: Claudia Brotto; and Hospital Julio Perrando, Resistencia, Chaco, Argentina: Isabel Marques.

Funding

This work was supported by the regular federal budget of the Ministry of Health of Argentina.

Transparency declarations

None to declare.

References

- 1 Cuzon G, Naas T, Villegas MV *et al.* Wide dissemination of *Pseudomonas* producing β -lactamase *bla*_{KPC-2} gene in Colombia. *Antimicrob Agents Chemother* 2011; **55**: 5350–3.
- 2 Akpaka PE, Swanston WH, Ihemere HN *et al.* Emergence of KPC-producing *Pseudomonas aeruginosa* in Trinidad and Tobago. *J Clin Microbiol* 2009; **47**: 2670–1.
- 3 Poirel L, Nordmann P, Lagrutta E *et al.* Emergence of KPC-producing *Pseudomonas aeruginosa* in the United States. *Antimicrob Agents Chemother* 2010; **54**: 3072.
- 4 Wolter DJ, Khalaf N, Robledo IE *et al.* Surveillance of carbapenem-resistant *Pseudomonas aeruginosa* isolates from Puerto Rican medical center hospitals: dissemination of KPC and IMP-18 β -lactamases. *Antimicrob Agents Chemother* 2011; **55**: 2968–70.
- 5 Tenover FC, Arbeit RD, Goering RV *et al.* Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995; **33**: 2233–9.
- 6 Gomez S, Pasteran F, Faccione D *et al.* Clonal dissemination of *Klebsiella pneumoniae* ST258 harbouring KPC-2. *Clin Microbiol Infect* 2011; **17**: 1520–4.
- 7 Carattoli A, Bertini A, Villa L *et al.* Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005; **63**: 219–28.
- 8 Woodford N, Turton JF, Livermore DM. Multiresistant Gram-negative bacteria: the role of high-risk clones in the dissemination of antibiotic resistance. *FEMS Microbiol Rev* 2011; **35**: 736–55.
- 9 Koh TH, Khoo CT, Tan TT *et al.* Multilocus sequence types of carbapenem-resistant *Pseudomonas aeruginosa* in Singapore carrying metallo- β -lactamase genes, including the novel *bla*_{IMP-26} gene. *J Clin Microbiol* 2010; **48**: 2563–4.
- 10 Samuelson O, Toleman MA, Sundsfjord A *et al.* Molecular epidemiology of metallo- β -lactamase-producing *Pseudomonas aeruginosa* isolates from Norway and Sweden shows import of international clones and local clonal expansion. *Antimicrob Agents Chemother* 2010; **54**: 346–52.

J Antimicrob Chemother 2012

doi:10.1093/jac/dkr593

Advance Access publication 18 January 2012

***In vitro* activity of tigecycline against multidrug-resistant Gram-negative blood culture isolates from critically ill patients**

Soham Gupta, Chowdappa Aruna, Savitha Nagaraj, Mary Dias and Sethumadhavan Muralidharan*

Department of Microbiology, St John's Medical College and Hospital, Bangalore-560034, India

*Corresponding author. Tel: +91-80-22065052; Email: dr.murali.2009@gmail.com

Keywords: MIC, VITEK 2C, intensive care unit

Sir,
With increasing resistance to currently available antibiotics and decreasing numbers of newer antimicrobial agents, tigecycline