

# *Journal of Threatened Taxa*

*Building evidence for conservation globally*



*Open Access*

[10.11609/jott.2021.13.11.19431-19674](https://doi.org/10.11609/jott.2021.13.11.19431-19674)  
[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

26 September 2021 (Online & Print)  
Vol. 13 | No. 11 | Pages: 19431–19674

ISSN 0974-7907 (Online)  
ISSN 0974-7893 (Print)



## Publisher

Wildlife Information Liaison Development Society

www.wild.zooreach.org

Host

Zoo Outreach Organization

www.zooreach.org

No. 12, Thiruvannamalai Nagar, Saravanampatti - Kalapatti Road, Saravanampatti,

Coimbatore, Tamil Nadu 641035, India

Ph: +91 9385339863 | [www.threatenedtaxa.org](http://www.threatenedtaxa.org)

Email: sanjay@threatenedtaxa.org

## EDITORS

## Founder &amp; Chief Editor

Dr. Sanjay Molur

Wildlife Information Liaison Development (WILD) Society & Zoo Outreach Organization (ZOO),  
12 Thiruvannamalai Nagar, Saravanampatti, Coimbatore, Tamil Nadu 641035, India

## Deputy Chief Editor

Dr. Neelesh Dahanukar

Noida, Uttar Pradesh, India

## Managing Editor

Mr. B. Ravichandran, WILD/ZOO, Coimbatore, India

## Associate Editors

Dr. Mandar Paingankar, Government Science College Gadchiroli, Maharashtra 442605, India  
Dr. Ulrike Streicher, Wildlife Veterinarian, Eugene, Oregon, USA  
Ms. Priyanka Iyer, ZOO/WILD, Coimbatore, Tamil Nadu 641035, India  
Dr. B.A. Daniel, ZOO/WILD, Coimbatore, Tamil Nadu 641035, India

## Editorial Board

Dr. Russel Mittermeier

Executive Vice Chair, Conservation International, Arlington, Virginia 22202, USA

## Prof. Mewa Singh Ph.D., FASc, FNA, FNAsc, FNAPsy

Ramanna Fellow and Life-Long Distinguished Professor, Biopsychology Laboratory, and  
Institute of Excellence, University of Mysore, Mysuru, Karnataka 570006, India; Honorary  
Professor, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore; and Adjunct  
Professor, National Institute of Advanced Studies, Bangalore

## Stephen D. Nash

Scientific Illustrator, Conservation International, Dept. of Anatomical Sciences, Health Sciences  
Center, T-8, Room 045, Stony Brook University, Stony Brook, NY 11794-8081, USA

## Dr. Fred Pluthero

Toronto, Canada

## Dr. Priya Davidar

Sigur Nature Trust, Chadapatti, Mavinahalli PO, Nilgiris, Tamil Nadu 643223, India

## Dr. Martin Fisher

Senior Associate Professor, Battcock Centre for Experimental Astrophysics, Cavendish  
Laboratory, JJ Thomson Avenue, Cambridge CB3 0HE, UK

## Dr. John Fellowes

Honorary Assistant Professor, The Kadoorie Institute, 8/F, T.T. Tsui Building, The University of  
Hong Kong, Pokfulam Road, Hong Kong

## Prof. Dr. Mirco Solé

Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas, Vice-coordenador  
do Programa de Pós-Graduação em Zoologia, Rodovia Ilhéus/Itabuna, Km 16 (45662-000)  
Salobrinho, Ilhéus - Bahia - Brasil

## Dr. Rajeev Raghavan

Professor of Taxonomy, Kerala University of Fisheries &amp; Ocean Studies, Kochi, Kerala, India

## English Editors

Mrs. Mira Bhojwani, Pune, India

Dr. Fred Pluthero, Toronto, Canada

Mr. P. Ilangovan, Chennai, India

## Web Maintenance

Mrs. Latha G. Ravikumar, ZOO/WILD, Coimbatore, India

## Typesetting

Mr. Arul Jagadish, ZOO, Coimbatore, India

Mrs. Radhika, ZOO, Coimbatore, India

Mrs. Geetha, ZOO, Coimbatore India

## Fundraising/Communications

Mrs. Payal B. Molur, Coimbatore, India

Subject Editors 2018–2020

## Fungi

Dr. B. Shivaraju, Bengaluru, Karnataka, India

Dr. R.K. Verma, Tropical Forest Research Institute, Jabalpur, India

Dr. Vatsavaya S. Raju, Kakatiya University, Warangal, Andhra Pradesh, India

Dr. M. Krishnappa, Jnana Sahyadri, Kuvenpura University, Shimoga, Karnataka, India

Dr. K.R. Sridhar, Mangalore University, Mangalagangotri, Mangalore, Karnataka, India

Dr. Gunjan Biswas, Vidyasagar University, Midnapore, West Bengal, India

## Plants

Dr. G.P. Sinha, Botanical Survey of India, Allahabad, India

Dr. N.P. Balakrishnan, Ret. Joint Director, BSI, Coimbatore, India

Dr. Shonil Bhagwat, Open University and University of Oxford, UK

Prof. D.J. Bhat, Retd. Professor, Goa University, Goa, India

Dr. Ferdinando Boero, Università del Salento, Lecce, Italy

Dr. Dale R. Calder, Royal Ontario Museum, Toronto, Ontario, Canada

Dr. Cleofas Cervancia, Univ. of Philippines Los Baños College Laguna, Philippines

Dr. F.B. Vincent Florens, University of Mauritius, Mauritius

Dr. Merlin Franco, Curtin University, Malaysia

Dr. V. Irudayaraj, St. Xavier's College, Palayamkottai, Tamil Nadu, India

Dr. B.S. Kholia, Botanical Survey of India, Gangtok, Sikkim, India

Dr. Pankaj Kumar, Kadoorie Farm and Botanic Garden Corporation, Hong Kong S.A.R., China

Dr. V. Sampath Kumar, Botanical Survey of India, Howrah, West Bengal, India

Dr. A.J. Solomon Raju, Andhra University, Visakhapatnam, India

Dr. Vijaysankar Raman, University of Mississippi, USA

Dr. B. Ravi Prasad Rao, Sri Krishnadevaraya University, Anantapur, India

Dr. K. Ravikumar, FRLHT, Bengaluru, Karnataka, India

Dr. Aparna Watve, Pune, Maharashtra, India

Dr. Qiang Liu, Xishuangbanna Tropical Botanical Garden, Yunnan, China

Dr. Noor Azhar Mohamed Shazili, Universiti Malaysia Terengganu, Kuala Terengganu, Malaysia

Dr. M.K. Vasudeva Rao, Shiv Ranjan Housing Society, Pune, Maharashtra, India

Prof. A.J. Solomon Raju, Andhra University, Visakhapatnam, India

Dr. Mandar Datar, Agharkar Research Institute, Pune, Maharashtra, India

Dr. M.K. Janarthanan, Goa University, Goa, India

Dr. K. Karthigeyan, Botanical Survey of India, India

Dr. Errol Vela, University of Montpellier, Montpellier, France

Dr. P. Lakshminarasimhan, Botanical Survey of India, Howrah, India

Dr. Larry R. Noblick, Montgomery Botanical Center, Miami, USA

Dr. K. Haridasan, Pallavur, Palakkad District, Kerala, India

Dr. Analinda Manila-Fajard, University of the Philippines Los Baños, Laguna, Philippines

Dr. P.A. Sinu, Central University of Kerala, Kasaragod, Kerala, India

Dr. Afroz Alam, Banasthali Vidyapith (accredited A grade by NAAC), Rajasthan, India

Dr. K.P. Rajesh, Zamorin's Guruvayurappan College, GA College PO, Kozhikode, Kerala, India

Dr. David E. Boufford, Harvard University Herbaria, Cambridge, MA 02138-2020, USA

Dr. Ritesh Kumar Choudhary, Agharkar Research Institute, Pune, Maharashtra, India

Dr. Navendu Page, Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand, India

## Invertebrates

Dr. R.K. Avasthi, Rohtak University, Haryana, India

Dr. D.B. Bastawade, Maharashtra, India

Dr. Partha Pratim Bhattacharjee, Tripura University, Suryamaninagar, India

Dr. Kailash Chandra, Zoological Survey of India, Jabalpur, Madhya Pradesh, India

Dr. Ansie Dippenaar-Schoeman, University of Pretoria, Queenswood, South Africa

Dr. Rory Dow, National Museum of Natural History Naturalis, The Netherlands

Dr. Brian Fisher, California Academy of Sciences, USA

Dr. Richard Gallon, Llandudno, North Wales, LL30 1UP

Dr. Hemanth V. Ghate, Modern College, Pune, India

Dr. M. Monwar Hossain, Jahangirnagar University, Dhaka, Bangladesh

Mr. Jatishwor Singh Irungbam, Biology Centre CAS, Brno, Czech Republic.

Dr. Ian J. Kitching, Natural History Museum, Cromwell Road, UK

Dr. George Mathew, Kerala Forest Research Institute, Peechi, India

Dr. John Noyes, Natural History Museum, London, UK

For Focus, Scope, Aims, and Policies, visit [https://threatenedtaxa.org/index.php/JoTT/aims\\_scope](https://threatenedtaxa.org/index.php/JoTT/aims_scope)For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions>For Policies against Scientific Misconduct, visit [https://threatenedtaxa.org/index.php/JoTT/policies\\_various](https://threatenedtaxa.org/index.php/JoTT/policies_various)

continued on the back inside cover

Caption: Malabar Slender Loris *Loris lydekkerianus malabaricus* © Dileep Anthikkad.



## Argentinian odonates (dragonflies and damselflies): current and future distribution and discussion of their conservation

A. Nava-Bolaños<sup>1</sup> ID, D.E. Vrech<sup>2</sup> ID, A.V. Peretti<sup>3</sup> ID & A. Córdoba-Aguilar<sup>4</sup> ID

<sup>1,4</sup> Departamento de Ecología Evolutiva, Instituto de Ecología, Universidad Nacional Autónoma de México, Apdo. Postal 70-275, Ciudad Universitaria, México, D.F. 04510, México.

<sup>1</sup> Biodiversity Institute, University of Kansas, Lawrence, KS, USA.

<sup>1</sup> Museo de Zoología, Departamento de Biología Evolutiva, Facultad de Ciencias, Universidad Nacional Autónoma de México, Apartado Postal 70-399, 04510 Ciudad de México, México.

<sup>2,3</sup> Instituto de Diversidad y Ecología Animal, CONICET - Universidad Nacional de Córdoba, Vélez Sarsfield 299 (5000), Córdoba, Argentina.

<sup>1</sup> anb@ciencias.unam.mx, <sup>2</sup> dvrech@unc.edu.ar,

<sup>3</sup> aperetti@unc.edu.ar (corresponding author), <sup>4</sup> acordoba@iecolologia.unam.mx (corresponding author)

**Abstract:** In terms of conservation, Argentinian odonates have not been assessed using a quantitative approach. One way to achieve this is by modelling their distribution to gather the extent of occurrence. Thus, we modelled the current and future (projected year, 2050) potential distribution of 44 odonate species that occur in Argentina as well as in neighboring countries. Our models of current times indicate a fairly wide distribution for most species but one exception is relevant for conservation purposes: *Lestes dichrostigma* has less than 30,000 km<sup>2</sup> and falls in the 'Near Threatened' category according to the IUCN Red List. Another seven species have less than or close to 100,000 km<sup>2</sup>: *Elasmothemis cannacioides*, *Erythemis credula*, *E. paraguayensis*, *Heteragrion angustipenne*, *H. inca*, *Lestes forficula*, and *Mecistogaster linearis*. Future distribution estimates suggest that: a) 12 species will lose or gain around 10%, four species will increase their distribution beyond 10% (up to 2,346%), and 28 species will lose more than 10% (up to 99%). Although current protected areas embrace most odonate species in Argentina, it is still premature to conclude whether this situation will remain in the future given the physiological tolerance and dispersal abilities of the study species among other drivers of distribution.

**Keywords:** Argentina, global change, IUCN, Odonata, potential distribution, status.

**Resumen:** En términos de conservación, los odonatos argentinos no han sido evaluados usando un enfoque cuantitativo. Una manera de hacer esto es modelando su distribución para obtener la extensión de la ocurrencia. En este trabajo modelamos la distribución actual y futura (año proyectado, 2050) de 44 especies de odonatos que se distribuyen en Argentina y países vecinos. Los modelos actuales indican una distribución amplia para la mayoría de especies aunque existe una excepción para propósitos de conservación: *Lestes dichrostigma* con menos de 30,000 km<sup>2</sup> y que cae en la categoría de "cercana a la amenaza" según la lista roja de la UICN. Otras siete especies tienen menos o cerca de 100,000 km<sup>2</sup>: *Elasmothemis cannacioides*, *Erythemis credula*, *E. paraguayensis*, *Heteragrion angustipenne*, *H. inca*, *Lestes forficula* y *Mecistogaster linearis*. Las estimas futuras sugieren que: a) 12 especies perderán o ganarán alrededor de 10% de área, cuatro especies incrementarán su distribución por más de 10% (hasta 2346 %), y 28 especies perderán más del 10% (hasta 99%). Aunque las áreas naturales protegidas actuales albergan la mayoría de especies en Argentina, es aún prematuro concluir que esta situación prevalecerá en el futuro dada la tolerancia fisiológica y capacidad de dispersión de las especies incluidas en este estudio así como otros efectores de su distribución.

**Editor:** Anonymity requested.

**Date of publication:** 26 September 2021 (online & print)

**Citation:** Nava-Bolaños, A., D.E. Vrech, A.V. Peretti & A. Córdoba-Aguilar (2021). Argentinian odonates (dragonflies and damselflies): current and future distribution and discussion of their conservation. *Journal of Threatened Taxa* 13(11): 19448–19465. <https://doi.org/10.11609/jott.7166.13.11.19448-19465>

**Copyright:** © Nava-Bolaños et al. 2021. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

**Funding:** To PAPIIT-UNAM grants IN 203115 and IN206618 to ACA. To a CONACyT-CONICET grant 190552 to ACA and AVP. To Secretaría de Educación, Ciencia, Tecnología e Innovación de la Ciudad de México (SECTEI) for the support to ANB.

**Competing interests:** The authors declare no competing interests.

**Author details, Author contributions and Acknowledgements:** See end of this article.



## INTRODUCTION

Given their analytical strength, species distribution models have been widely used to assess the potential area where a species occurs as predicted by environmental variables (Peterson 2006). Odonates have not been an exception to this practice with at least 30 different studies in distinct world regions (reviewed by Collins & McIntyre 2015). Such interest is partly understood on the basis of the intrinsic threat that humankind has posed to freshwater bodies (e.g. Sala et al. 2000) related to the direct dependence of odonates on these bodies. Furthermore, a more recent analysis indicated that odonates can be used as the indicators of global change given their practicality as study models (i.e. large body size), well-described macro-ecological responses, key role as predators in aquatic and terrestrial habitats and their trend of becoming field-animal models for temperature-mediated responses (Hassall 2015). Paradoxically, our current knowledge of the extinction risk for most odonates is extremely limited. For example, the IUCN (2018) shows a shortage of species with strong geographical biases, with country-based assessments frequently lacking firm quantitative-supporting data (see for example, Paulson 2004). One case is that of Argentina: 86 species are listed of which one is 'Endangered', one is 'Vulnerable', two are 'Near Threatened', four are 'Data Deficient', and 78 are 'Least Concern' (IUCN 2018). This implies that a proper assessment is badly needed for this country.

Distribution models of odonates have provided clues of how current distribution will be affected by increases in temperature (reviewed by Collins & McIntyre 2015). These studies have covered up to 25% of the total world odonate diversity, and have shown that in general there will be shifts in distribution, with lotic species and narrow-distribution species (e.g., endemic) showing a tendency to have their areas reduced (reviewed by Collins & McIntyre 2015). In this paper, we have carried out an exercise of calculating current and future distribution models for Argentinian odonates to supplement current studies of distribution gathered from provincial records (e.g. Muzón et al. 2014, 2015; von Ellenrieder & Muzón 1999, 2008; von Ellenrieder 2009, 2010). Our analysis is based on a fraction of the 271 species currently known to occur in Argentina (Muzón & von Ellenrieder 1999; von Ellenrieder & Muzón 2008). Our aim is to use our assessment to guide the current IUCN risk categories for Argentinian odonates based on criteria A and B, that define extent of occurrence.

## MATERIAL AND METHODS

### Occurrence data of species

Presence of odonate species was compiled from literature records, GBIF records ([www.gbif.org](http://www.gbif.org) as of 20 December 2017; GBIF Occurrence Download <http://doi.org/10.15468/dl.mf6nh7>), and odonate specialists (Rosser Garrison, Natalia von Ellenrieder, and Dennis Paulson). All data were checked carefully for geographic accuracy by removing duplicates and records with inconsistent georeferencing, for example coordinates on the sea, or missing as recommended in the literature of data cleaning (Chapman 2005). Most records were gathered by odonate experts, so we are confident that identification bias should be minimal. Niche models were built only when more than 10 records per species were available. Thus, the final data set included 1,734 unique presences of 44 species (see Table 1) which were those species with enough collecting data (range 11–158, see Table 1). The database of records is available upon request.

### Study area, background and environmental predictors

We have modeled the potential distribution of Argentinian species including cases outside the country's boundaries. Our study area included land between latitudes -55.08 and -21.55S, and longitudes -75.30 to -53.13W. As bioclimatic variables, we used the WorldClim 1.4 ([www.worldclim.org](http://www.worldclim.org)) data set (Hijmans et al. 2005) at 0.04166669 cell size. To establish a background and a set of uncorrelated climatic variables, we intersected the variables with target group points, and with 10,000 points randomly selected in the extension of the study area (M). We eliminated some variables with an exploratory data analysis and Pearson correlation analysis (values >0.7). Thus, we selected variables with low correlation and high contribution to reduce the parametrization of the models. After this, the final data set included uncorrelated variables which had more biological importance for our study species, and contributed the most according to the jackknife analysis. Variables were: mean diurnal range (bio 02), isothermality (bio 03), temperature seasonality (bio 04), mean temperature of driest quarter (bio 09), mean temperature of warmest quarter (bio 10), precipitation of wettest month (bio 13), precipitation seasonality (bio 15), precipitation of driest quarter (bio 17), precipitation of warmest quarter (bio 18), and precipitation of the coldest quarter (bio 19).

**Table 1.** Argentinian odonates species modeled, number of records, potential distribution of species in km<sup>2</sup>, TSS values and current and proposed IUCN categories.

Species	Records	Current area (km <sup>2</sup> )	TSS	Current IUCN status	Suggested IUCN status
<i>Acanthagrion aepiolum</i> Tennessen, 2004	23	206259	0.90	N/A	LC
<i>A. cuyabae</i> Calvert, 1909	55	1136583	0.86	LC	LC
<i>A. floridense</i> Fraser, 1946	47	166257	0.89	N/A	LC
<i>A. gracile</i> (Rambur, 1842)	43	865415	0.85	N/A	LC
<i>A. hidegarda</i> Gloger, 1967	27	112352	0.90	N/A	LC
<i>A. lancea</i> Selys, 1876	48	645339	0.87	N/A	LC
<i>Elasmothemis cannacioides</i> (Calvert, 1906)	12	79208	0.83	N/A	LC
<i>Erythemis attala</i> (Selys in Sagra, 1857)	70	368120	0.89	LC	LC
<i>E. credula</i> (Hagen, 1861)	16	67990	0.86	N/A	LC
<i>E. peruviana</i> (Rambur, 1842)	72	1056558	0.86	LC	LC
<i>E. plebeja</i> (Burmeister, 1839)	94	1523637	0.84	LC	LC
<i>E. vesiculosa</i> (Fabricius, 1775)	132	2228200	0.81	LC	LC
<i>Erythrodiplax fusca</i> (Rambur, 1842)	22	173798	0.90	LC	LC
<i>E. paraguayensis</i> (Förster, 1905)	11	40995	0.80	LC	LC
<i>E. umbrata</i> (Linnaeus, 1758)	59	184811	0.90	LC	LC
<i>Heteragrion angustipenne</i> Selys, 1886	14	74209	0.84	N/A	LC
<i>H. inca</i> Calvert, 1909	13	102730	0.82	N/A	LC
<i>Ischnura capreolus</i> (Hagen, 1861)	139	734839	0.88	N/A	LC
<i>I. fluviatilis</i> Selys, 1876	158	1714797	0.83	LC	LC
<i>I. ultima</i> Ris, 1908	34	11808573	0.90	N/A	LC
<i>Lestes dichrostigma</i> Calvert, 1909	11	28823	0.80	LC	NT
<i>L. forficula</i> Rambur, 1842	14	72423	0.83	N/A	LC
<i>L. spatula</i> Fraser, 1946	30	504657	0.88	N/A	LC
<i>L. undulatus</i> Say, 1840	34	195329	0.89	LC	LC
<i>Mecistogaster linearis</i> (Fabricius, 1777)	13	71030	0.82	N/A	LC
<i>Miathyria marcella</i> (Selys in Sagra, 1857)	44	4166276	0.87	LC	LC
<i>Micrathyria hesperis</i> Ris, 1911	19	7900041	0.87	N/A	LC
<i>M. hypodidyma</i> Calvert, 1906	33	653996	0.88	N/A	LC
<i>M. longifasciata</i> Calvert, 1909	48	416857	0.89	LC	LC
<i>M. tibialis</i> Kirby, 1897	11	184013	0.80	LC	LC
<i>Orthemis ferruginea</i> (Fabricius, 1775)	13	1401215	0.79	LC	LC
<i>Pantala flavescens</i> (Fabricius, 1798)	17	387339	0.85	LC	LC
<i>Perithemis mooma</i> Kirby, 1889	15	829042	0.83	N/A	LC
<i>Rhionaeschna absoluta</i> (Calvert, 1952)	133	934413	0.86	N/A	LC
<i>R. bonariesis</i> (Rambur, 1842)	158	1417407	0.84	N/A	LC
<i>R. confusa</i> (Rambur, 1842)	52	261179	0.88	N/A	LC
<i>R. diffinis</i> (Rambur, 1842)	40	226574	0.89	LC	LC
<i>R. pallipes</i> (Fraser, 1947)	26	142412	0.89	N/A	LC
<i>R. planaltica</i> (Calvert, 1952)	51	163524	0.89	LC	LC
<i>R. variegata</i> (Fabricius, 1775)	41	365158	0.88	N/A	LC
<i>R. viginpunctata</i> (Ris, 1918)	47	155497	0.90	N/A	LC
<i>Tramea darwini</i> Kirby, 1889	16	321819	0.85	LC	LC
<i>Uracis fastigiata</i> (Burmeister, 1839)	17	760515	0.85	N/A	LC
<i>U. imbuta</i> (Burmeister, 1839)	22	830556	0.84	N/A	LC

### Background selection

To choose the best background, preliminary species distribution models were generated with Maxent 3.3.3k (Phillips et al. 2006) with target group points (with 10,000 points randomly selected in the extension of the study area, M), and with a special extent delineating M for each particular species with ecoregions (World Wildlife Fund; [www.worldwildlife.org/](http://www.worldwildlife.org/) date accessed 20 January 2018). Models were constructed by setting several parameters to default ('Auto features', convergence= 10-5, maximum number of iterations= 500). However, we used random seed (with a 30 test percentage), 10 replicates, removed duplicate records, ran bootstrap replicated type, with no extrapolation and no clamping. All this to find which combination of settings and variables generated the best outcomes (highest area under the curve, or AUC) while minimizing the number of model parameters, as well as producing 'closed', bell-shaped response curves guaranteeing model calibration (Elith et al. 2010). The best background by the preliminary analyses was 10,000 points randomly selected in the extension of the study area.

### Training ecological niche models

Final models were built with BIOMOD (Biodiversity Modelling) package in R software. This package is a platform for predicting species' distribution, including the ability to model the distribution using various techniques and test patterns (Thuiller et al. 2009). We trained models using four widely used algorithms: maximum entropy (Maxent), random forest (RF), generalized boosting methods (GBM), and multivariate adaptive regression splines (MARS). These models have shown good performance in terms of predictive power (Broennimann et al. 2012; Pliscoff & Fuentes-Castillo 2011; Reiss et al. 2011). From individual models obtained with these different algorithms, we generated a 'consensus model'. Such model combination is the best logistic compromise to avoid either overfitting and overpredicting (Merow et al. 2014). In other words, this reduces biases and limitations of using only individual models. Seventy percent of data was used for training, and 30% for validation with 10 replicates. Final model validation was performed with TSS (True Skill Statistics), average net rate of successful prediction for sites of presence and absence (Liu et al. 2009), ranging from -1 to 1, where the more positive values indicate a higher degree of accuracy and discrimination model (Allouche et al. 2006) (Table 1). Notice that the result of these models is not the area that species occupy absolutely, because they do not consider population dynamics, dispersibility,

interactions with other species, and human impacts. However, these models predict where species can be potentially found given their environmental conditions. This assumes that the distribution known of each species provides enough information to characterize its environmental requirements.

A total of 224 models were generated, whose performance was assessed by means of the AUC and TSS statistics (Table 1), while minimizing the number of model parameters, and the best presence/absence models using the '10 percentile-training presence' are shown. This threshold was used because we prefer to err in the side of caution accepting that a 10% of our presences could be problematic (for a similar rationale, see Sánchez-Guillén et al. 2013). The best models of current climatic conditions of species were used to generate projections.

### Future projections

The best models of current climatic conditions of species were used to generate projections for the 2050 year assuming climatic change scenarios. The data for future projections were: Global Climate Models (GCM) (CNRM-CM5, HadGEM2-ES, and MPI-ESM-LR) in WorldClim (<http://worldclim.org/CMIP5v1/>; date accessed 12 December 2017), these climate projections were gathered from the Fifth Assessment (CMIP5) (<http://cmip-pcmdi.llnl.gov/cmip5/> date accessed 19/7/2017) report of The Intergovernmental Panel on Climate Change (IPPC) (<http://www.ipcc.ch/>). The representative concentration pathways used (RCP) were 4.5 and 8.5, for year 2050. A RCP 8.5 is considered a pessimistic scenario, where CO<sub>2</sub> emissions would continue to rise while a RCP 4.5 is considered a more optimistic situation.

We estimated areas of potential distribution of odonate species occurring within Argentinian borders in km<sup>2</sup>, and calculated the percentage of loss or gain of geographic areas with respect to current potential distribution. 2050 distribution was represented by a consensus model where only pixels-predicted-present by all models were considered as representing the presence of the species. We estimated areas with a function with stringr and raster packages in R (R Core Team 2017).

## RESULTS

Table 1 shows the potential current distribution (in km<sup>2</sup>) for each species, and the summary of the performance of the best models (with TSS). This table also shows the current IUCN Red List categories (as of 28 January 2018) and the new categories we suggest based on our analysis of distribution area. From these data, only *Lestes dichrostigma* Calvert, 1909 appears as 'Near Threatened' as its estimated distribution area is 28,823 km<sup>2</sup> (Figure 1). This as well as other seven species deserve some attention given that their distribution is less than- or close to 100,000 km<sup>2</sup> (Figure 1): *Elasmothemis cannacioides* (Calvert, 1906), *Erythemis credula* (Hagen, 1861), *Erythrodiplax paraguayensis* (Förster, 1905), *Heteragrion angustipenne* Selys, 1886, *H. inca* Calvert, 1909, *Lestes forficula* Rambur, 1842 and *Mecistogaster linearis* (Fabricius, 1777). Distributions of all species are included in supplementary material Figure 1.

In regard to climate change projections for the year 2050 the RCP 8.5 estimated the following: 12 species would maintain their distribution with loss or gain of only around 10% of change of their current distribution, four species would increase their distribution beyond 10%, and 28 species would lose their area of their distribution for more than 10% (Table 2). These changes, in general, were fairly consistent with the scenario RCP 4.5 with three species keeping their distribution for around 10% of change, 11 species increasing their distribution beyond 10%, and 30 species losing their distribution for more than 10% (Table 2). These coincidences for both scenarios include, for example, *Micrathyria tibialis* Kirby, 1897 and *Heteragrion angustipenne* Selys, 1886 which represent the extremes in terms of gaining and losing area, respectively.

## DISCUSSION

One benefit species distribution models can bring about is the conservation aspects. In this extent, our results suggest that although most Argentinian species have relatively large distributions, a few species deserve some attention. According to the current IUCN Red List (IUCN 2018), the following species face some risk: *Andinagrion garrisoni* von Ellenrieder & Muzón, 2006 and *Progomphus kimminsi* Belle, 1973 (Near Threatened), *Phyllogomphoides joaquinii* Rodrigues Capítulo, 1992 (Vulnerable) and *Staurophlebia bosqi* Navás, 1927 (Endangered). The remaining 82 are categorized as Data Deficient (4 species) or Least Concern (78 species). The

threatened four species were classified as such given the paucity of collecting records and their restricted areas of distribution. We were not able to locate enough collecting points for any of these four species. However, our work suggests that *Lestes dichrostigma* Calvert, 1909 deserves some attention, as its area is above but close to 20,000 km<sup>2</sup>. Although the remaining 43 species can be categorized as least concern, another five have less than 100,000 km<sup>2</sup> so we suggest their populations should be also monitored: *Elasmothemis cannacioides* (Calvert, 1906), *Erythemis credula* (Hagen, 1861), *Erythrodiplax paraguayensis* (Förster, 1905), *Heteragrion angustipenne* Selys, 1886, *H. inca* Calvert, 1909, *L. forficula* Rambur, 1842, and *Mecistogaster linearis* (Fabricius, 1777). Of course, several other population parameters should be gathered to complement IUCN categorization for all species, for example to detect the population reduction or less of variability. Notice that future projections would not help most species we modelled as 28–30 species would reduce their distribution dramatically in some cases. According to this, some other species not in danger currently would face threat according to these future scenarios: *Acanthagrion hidegarda* Gloger, 1967, *Heteragrion angustipenne* Selys, 1886, *Lestes dichrostigma* Calvert, 1909, *Mecistogaster linearis* (Fabricius, 1777), and *Rhionaeschna virginica* (Ris, 1918). These five species may reduce their area to less than 20,000 km<sup>2</sup>.

Essential to our present estimates of area is the fact that 70% of Argentinian species are currently present in protected areas (Muzón & von Ellenrieder 1999). However, given that global change will lead to shifts in current distribution (Sánchez-Guillén et al. 2016), a necessary step is to define whether current Argentinian protected areas will still embrace future odonate geographical distributions. A key issue here is to carry out more intensive collections to construct models for the remaining 227 odonate species that occur within Argentinian boundaries (von Ellenrieder & Muzón 2008). Moreover, research should pay attention to answer whether dispersal abilities can allow odonates catch up with different habitats located at different temperature regimes (Bush et al. 2014).

Related to global change scenarios, it is not surprising to find an inter-specific variation in projected responses to raising temperatures in odonates. Our explanations for this are incomplete yet but may have to do with odonate physiological abilities that affect themoregulatory responses (e.g., Corbet & May 2008) and development (especially at egg and larval stages; Pritchard & Leggot 1987). Given this, it is also not surprising that the largest




Figure 1. Potential distribution of a subset of Argentinian odonate species as predicted by ecological niche models: *Elasmothemis cannaeoides*, *Erythemis credula*, *E. paraguayensis*, *Heteragrion angustipenne*, *H. incia*, *Lestes dichrostigma*, *L. forficula*, and *Mecistogaster linearis*.

**Table 2. Absolute (in km<sup>2</sup>) and relative changes in suitable area per Argentinian odonate species according to different climatic changes scenarios. Losses are shown as negative values while gains are shown as positive values.**

Species	2050 (km <sup>2</sup> ) RCP4.5	2050 (km <sup>2</sup> ) RCP8.5	2050 (%) RCP4.5	2050 (%) RCP8.5
<i>Acanthagrion aepiolum</i> Tennessen, 2004	95025	77268	-53.93	-62.54
<i>A. cuyabae</i> Calvert, 1909	1085251	1128738	-4.52	-0.69
<i>A. floridense</i> Fraser, 1946	124121	148521	-25.34	-10.67
<i>A. gracile</i> (Rambur, 1842)	511056	459049	-40.95	-46.96
<i>A. hidegarda</i> Gloger, 1967	7430	7418	-93.39	-93.40
<i>A. lancea</i> Selys, 1876	334559	328591	-48.16	-49.08
<i>Elasmothemis cannacioides</i> (Calvert, 1906)	26652	20123	-66.35	-74.59
<i>Erythemis attala</i> (Selys in Sagra, 1857)	1040509	1672709	182.65	354.39
<i>E. credula</i> (Hagen, 1861)	104121	181602	53.14	167.10
<i>E. peruviana</i> (Rambur, 1842)	3475030	3977046	228.90	276.42
<i>E. plebeja</i> (Burmeister, 1839)	2875597	3578859	88.73	134.89
<i>E. vesiculosa</i> (Fabricius, 1775)	6394736	8249237	186.99	270.22
<i>Erythrodiplax fusca</i> (Rambur, 1842)	203928	185469	17.34	6.72
<i>E. paraguayensis</i> (Förster, 1905)	29488	30549	-28.07	-25.48
<i>E. umbrata</i> (Linnaeus, 1758)	1107621	1462042	499.33	691.10
<i>Heteragrion angustipenne</i> Selys, 1886	2709	566	-96.35	-99.24
<i>H. inca</i> Calvert, 1909	27552	18234	-73.18	-82.25
<i>Ischnura capreolus</i> (Hagen, 1861)	5444382	6676849	640.89	808.61
<i>I. fluviatilis</i> Selys, 1876	1087445	1034849	-36.58	-39.65
<i>I. ultima</i> Ris, 1908	1438693	1637097	-87.82	-86.14
<i>Lestes dichrostigma</i> Calvert, 1909	1497	1456	-94.81	-94.95
<i>L. forficula</i> Rambur, 1842	61821	78055	-14.64	7.78
<i>L. spatula</i> Fraser, 1946	297025	323398	-41.14	-35.92
<i>L. undulatus</i> Say, 1840	177025	181143	-9.37	-7.26
<i>Mecistogaster linearis</i> (Fabricius, 1777)	5896	2538	-91.70	-96.43
<i>Miathyria marcella</i> (Selys in Sagra, 1857)	8903701	9675724	113.71	132.24
<i>Micrathyria hesperis</i> Ris, 1911	1325471	1539839	-83.22	-80.51
<i>M. hypodidyma</i> Calvert, 1906	360230	360273	-44.92	-44.91
<i>M. longifasciata</i> Calvert, 1909	301298	304006	-27.72	-27.07
<i>M. tibialis</i> Kirby, 1897	3288689	4500751	1687.21	2345.89
<i>Orthemis ferruginea</i> (Fabricius, 1775)	856545	573823	-38.87	-59.05
<i>Pantala flavescens</i> (Fabricius, 1798)	345606	358468	-10.77	-7.45
<i>Perithemis mooma</i> Kirby, 1889	586843	671876	-29.21	-18.96
<i>Rhionaeschna absoluta</i> (Calvert, 1952)	775879	740279	-16.97	-20.78
<i>R. bonariesis</i> (Rambur, 1842)	713468	711143	-49.66	-49.83
<i>R. confusa</i> (Rambur, 1842)	211253	216912	-19.12	-16.95
<i>R. diffinis</i> (Rambur, 1842)	262980	259209	16.07	14.40
<i>R. pallipes</i> (Fraser, 1947)	70805	75227	-50.28	-47.18
<i>R. planaltica</i> (Calvert, 1952)	45782	44497	-72.00	-72.79
<i>R. variegata</i> (Fabricius, 1775)	295227	300756	-19.15	-17.64
<i>R. viginpunctata</i> (Ris, 1918)	89497	89484	-42.44	-42.45
<i>Tramea darwini</i> Kirby, 1889	343101	337055	6.61	4.73
<i>Uracis fastigiata</i> (Burmeister, 1839)	223876	175053	-70.56	-76.98
<i>U. imbuta</i> (Burmeister, 1839)	416894	126006	-49.81	-84.83



species turnover will occur at intermediate altitudes where drastic changes in temperature currently occur (Maes et al. 2010). The case of Argentina is actually very relevant to this altitude phenomenon given its sharp changes in elevation. Thus, special attention should be given to these areas. Given the small number of records for most species, we are far from ensuring a well-known distribution for a large number of Argentine species, where field work, as well as the digitization of records, is advisable to document regions that are poorly explored. One tool to help in this regard is the use of repositories of citizen science photographs.

Apart from North America (Canada and USA; Hassall 2012; Rangel-Sánchez et al. 2018) and Brazil (Nóbrega & De Marco 2011), our study adds a substantially high number of odonate species with projected distributions for America. Considering that there exist around 5,680 described odonate species, of which 25% had been modelled (Collins & McIntyre 2015), our study makes a valuable global contribution for the Southern Hemisphere. This importance can be seen not only in terms of conservation as discussed above, but also in terms of biogeography given the southerly location of our study species (currently, the southern extreme was Brazil with mainly tropical species; De Marco et al. 2015; Nóbrega & De Marco 2011). Thus our results can be used to understand biogeographical patterns based on odonate ecology (e.g., preference for lentic and lotic waters and global distribution; Hof et al. 2006).

## REFERENCES

- Allouche, O., A. Tsoar, & R. Kadmon (2006). Assessing the accuracy of species distribution models: Prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology* 43(6): 1223–1232. <https://doi.org/10.1111/j.1365-2664.2006.01214.x>
- Broennimann, O., M.C. Fitzpatrick, P.B. Pearman, B. Petitpierre, L. Pellissier, N.G. Yoccoz, W. Thuiller, M.-J. Fortin, C. Randin, N.E. Zimmermann, C.H. Graham & A. Guisan (2012). Measuring ecological niche overlap from occurrence and spatial environmental data. *Global Ecology and Biogeography* 21(4): 481–497. <https://doi.org/10.1111/j.1466-8238.2011.00698.x>
- Bush, A.A., D.A. Nipperess, D.E. Duursma, G. Theischinger, E. Turak & L. Hughes (2014). Continental-scale assessment of risk to the Australian Odonata from climate change. *PLoS ONE* 9(2). <https://doi.org/10.1371/journal.pone.0088958>
- Chapman, A.D. (2005). *Principles and Methods of Data Cleaning*. GBIF, Copenhagen, 75pp.
- Collins, S.D. & N.E. McIntyre (2015). Modeling the distribution of odonates: A review. *Freshwater Science* 34(3): 1144–1158. <https://doi.org/10.1086/682688>
- Corbet, P.S. & M.L. May (2008). Fliers and perchers among Odonata: Dichotomy or multidimensional continuum? A provisional reappraisal. *International Journal of Odonatology* 11(2): 155–171. <https://doi.org/10.1080/13887890.2008.9748320>
- De Marco Júnior, P., C.C. Nóbrega, R.A. De Souza, & U.G. Neiss (2015). Modeling the distribution of a rare Amazonian odonate in relation to future deforestation. *Freshwater Science* 34(3): 1123–1132. <https://doi.org/10.1086/682707>
- Elith, J., K. Michael & P. Steven (2010). The art of modelling range-shifting species. *Methods in Ecology and Evolution* 1(4): 330–342. <https://doi.org/10.1111/j.2041-210X.2010.00036.x>
- Hassall, C. (2012). Predicting the distributions of under-recorded Odonata using species distribution models. *Insect Conservation and Diversity* 5(3): 192–201. <https://doi.org/10.1111/j.1752-4598.2011.00150.x>
- Hassall, C. (2015). Odonata as candidate macroecological barometers for global climate change. *Freshwater Science* 34(3): 1040–1049. <https://doi.org/10.1086/682210>
- Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones & A. Jarvis (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25(15): 1965–1978. <https://doi.org/10.1002/joc.1276>
- Hof, C., M. Brändle & R. Brandl (2006). Lentic odonates have larger and more northern ranges than lotic species. *Journal of Biogeography* 33(1): 63–70. <https://doi.org/10.1111/j.1365-2699.2005.01358.x>
- IUCN (2018). The IUCN Red List of Threatened Specieswww.threddedlist.org accessed 25 August 2016.
- Liu, C., M. White, & G. Newell (2009). Measuring the accuracy of species distribution models: A review. *18th World IMACS Congress and MODSIM 2009 - International Congress on Modelling and Simulation: Interfacing Modelling and Simulation with Mathematical and Computational Sciences, Proceedings* (January 2009): 4241–4247.
- Maes, D., N. Titeux, J. Hortal, A. Anselin, K. Decleer, G. de Knijf, ... M. Luoto (2010). Predicted insect diversity declines under climate change in an already impoverished region. *Journal of Insect Conservation* 14(5): 485–498. <https://doi.org/10.1007/s10841-010-9277-3>
- Merow, C., M.J. Smith, T.C. Edwards, A. Guisan, S.M. Mcmahon, S. Normand, W. Thuiller, R.O. Wüest, N.E. Zimmermann & Elith (2014). What do we gain from simplicity versus complexity in species distribution models? *Ecography* 37(12): 1267–1281. <https://doi.org/10.1111/ecog.00845>
- Muzón, J., F. Lozano, A. del Palacio, L.S. Ramos, & A. Lutz (2015). Odonata from the Lower Delta of the Paraná River, Argentina. *Agrion* 20(2): 68–72.
- Muzón, J., P. Pessacq & F. Lozano (2014). The Odonata (Insecta) of Patagonia: A synopsis of their current status with illustrated keys for their identification. *Zootaxa* 3784(4): 346–388. <https://doi.org/10.11646/zootaxa.3784.4.2>
- Muzón, J. & N. von Ellenrieder (1999). Status and distribution of Odonata (Insecta) within natural protected areas in Argentina. *Biogeographica* 75(3): 119–128.
- Nóbrega, C.C. & P. de Marco (2011). Unprotecting the rare species: A niche-based gap analysis for odonates in a core Cerrado area. *Diversity and Distributions* 17(3): 491–505. <https://doi.org/10.1111/j.1472-4642.2011.00749.x>
- Paulson, D.R. (2004). Critical species of odonata in the neotropics. *International Journal of Odonatology* 7(2): 163–188. <https://doi.org/10.1080/13887890.2004.9748208>
- Peterson, A.T. (2006). Uses and Requirements of Ecological Niche Models and Related Distributional Models. *Biodiversity Informatics*, 3(0): 59–72.
- Phillips, S., R. Anderson & R. Schapire (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190: 231–259. <https://doi.org/10.1016/j.ecolmodel.2005.03.026>
- Pliscoff, P. & T. Fuentes-Castillo (2011). Modelación de la distribución de especies y ecosistemas en el tiempo y en el espacio: una revisión de las nuevas herramientas y enfoques disponibles 1. *Revista de Geografía Norte Grande* 48: 61–79. <https://doi.org/http://dx.doi.org/10.4067/S0718-34022011000100005>
- Pritchard, G., M.A. Leggott, G. Pritchard & M.A. Leggott (1987). Temperature, Incubation Rates and Origins of Dragonflies. *Odonatology* 3(1): 121–126.

- R Core Team (2017).** R: A language and environment for statistical computing. In: R Found Stat Comput Vienna, Austria.
- Rangel-Sánchez, L., A. Nava-Bolaños, F. Palacino-Rodríguez & A. Córdoba-Aguilar (2018).** Estimating distribution area in six Argia damselflies (Insecta: Odonata: Coenagrionidae) including *A. garrisoni*, a threatened species. *Revista Mexicana de Biodiversidad* 89(3): 921–926. <http://dx.doi.org/10.22201/ib.20078706e.2018.3.2469>
- Reiss, H., H. Cunze, K. König, K. Neumann & I. Kröncke (2011).** Species distribution modelling of marine benthos: A North Sea case study. *Marine Ecology Progress Series* 442: 71–86. <https://doi.org/10.3354/meps09391>
- Sala, O.E., F.S. Chapin, J.J. Armesto, E. Berlow, J. Bloomfield, R. Dirzo, E. Huber-Sanwald, L.F. Huenneke, R.B. Jackson, A. Kinzig, R. Leemans, D.M. Lodge, H.A. Mooney, M. Oesterheld, N.L. Poff, M.T. Sykes, B.H. Walker, M. Walker & D.H. Wall (2000).** Global biodiversity scenarios for the year 2100. *Science* 287(5459): 1770–1774. <https://doi.org/10.1126/science.287.5459.1770>
- Sánchez-Guillén, R.A., A. Córdoba-Aguilar, B. Hansson, J. Ott & M. Wellenreuther (2016).** Evolutionary consequences of climate-induced range shifts in insects. *Biological Reviews* 91(4): 1050–1064. <https://doi.org/10.1111/brv.12204>
- Sánchez-Guillén, R.A., J. Muñoz, G. Rodríguez-Tapia, T.P.F. Arroyo & A. Córdoba-Aguilar (2013).** Climate-induced range shifts and possible hybridisation consequences in insects. *PLoS ONE* 8(11): 1–10. <https://doi.org/10.1371/journal.pone.0080531>
- Thuiller, W., B. Lafourcade, R. Engler, & M.B. Araújo (2009).** BIOMOD - A platform for ensemble forecasting of species distributions. *Ecography* 32(3): 369–373. <https://doi.org/10.1111/j.1600-0587.2008.05742.x>
- Von Ellenrieder, N. (2009).** Odonata of the Argentine Yungas cloud forest: Distribution patterns and conservation status. *Odonatologica* 38(1): 39–53.
- Von Ellenrieder, N. (2010).** Odonata biodiversity of the argentine chaco biome. *International Journal of Odonatology* 13(1): 1–25. <https://doi.org/10.1080/13887890.2010.974835>
- Von Ellenrieder, N. & J. Muzón (2008).** An updated checklist of the Odonata from Argentina. *Odonatologica* 37(1): 55–68.
- Von Ellenrieder, N. & J. Muzón (1999).** The Argentinean species of the genus *Perithemis* Hagen (Anisoptera: Libellulidae). *Odonatologica* 28(4): 385–398.

**Author details:** A. NAVA-BOLAÑOS is a postdoctoral researcher at the Museum of Zoology (Universidad Nacional Autónoma de México) conducting global biodiversity analysis of pollinator species. Her research topics include hybridization, climate change, reproductive isolation barriers, species distribution, ecological niche models and insect conservation. D.E. VRECH is a researcher at the Instituto de Diversidad y Ecología Animal (Universidad Nacional de Córdoba, Argentina). His main research topics are behavioral ecology of arthropods, sperm biology, evolutionary biology, and reproductive behavior. A.V. PERETTI is a researcher at the Instituto de Diversidad y Ecología Animal (Universidad Nacional de Córdoba, Argentina). His research lines include the study of the reproductive biology of arthropods, and topics associated with ecology, functional morphology, physiology and genetics. He has worked with scorpions, spiders, and insects, mainly odonates. A. CÓRDOBA-AGUILAR is a researcher at the Instituto de Ecología (Universidad Nacional Autónoma de México). His research topics are insect vector control and insect conservation.

**Author contributions:** ANB, DV, AVP, and ACA planned the paper, ANB and DV executed all analyses and all authors wrote and revised the paper. ANB and DV contributed equally to the paper.

**Acknowledgements:** To PAPIIT-UNAM grants IN 203115 and IN206618 to ACA. To a CONACyT-CONICET grant 190552 to ACA and AVP. To Secretaría de Educación, Ciencia, Tecnología e Innovación de la Ciudad de México (SECTEI) for the support to ANB.

Supplementary material figure. Current potential distribution of Argentinian odonate species as predicted by ecological niche models. Predictions of suitable area appear in black.



*Acanthagrion aepiolum*



*Acanthagrion cuyabae*



*Acanthagrion floridense*



*Acanthagrion gracile*



*Acanthagrion hidegarda*



*Acanthagrion lancea*



*Elasmothemis cannacioides*



*Erythemis attala*



*Erythemis credula*



*Erythemis plebeja*



*Erythemis peruviana*



*Erythemis vesiculosa*



*Erythrodiplax umbrata*



*Erythrodiplax fusca*



*Ischnura capreolus*



*Ischnura fluviatilis*



*Ischnura ultima*



*Lestes undulates*



*Lestes spatula*



*Mecistogaster linearis*



*Miathyria marcella*



*Micrathyria hypodidyma*



*Micrathyria hesperis*



*Micrathyria longifasciata*



*Rhionaeschna absoluta*



*Rhionaeschna bonariensis*



*Rhionaeschna pallipes*



*Rhionaeschna planaltica*



*Rhionaeschna variegata*



*Rhionaeschna virginpunctata*



*Tramea darwini*



*Uracis fastigiate*



*Uracis imbuta*



Dr. Albert G. Orr, Griffith University, Nathan, Australia  
Dr. Sameer Padhye, Katholieke Universiteit Leuven, Belgium  
Dr. Nancy van der Poorten, Toronto, Canada  
Dr. Karen Schnabel, NIWA, Wellington, New Zealand  
Dr. R.M. Sharma, (Retd.) Scientist, Zoological Survey of India, Pune, India  
Dr. Manju Siliwal, WILD, Coimbatore, Tamil Nadu, India  
Dr. G.P. Sinha, Botanical Survey of India, Allahabad, India  
Dr. K.A. Subramanian, Zoological Survey of India, New Alipore, Kolkata, India  
Dr. P.M. Sureshan, Zoological Survey of India, Kozhikode, Kerala, India  
Dr. R. Varatharajan, Manipur University, Imphal, Manipur, India  
Dr. Eduard Vives, Museu de Ciències Naturals de Barcelona, Terrassa, Spain  
Dr. James Young, Hong Kong Lepidopterists' Society, Hong Kong  
Dr. R. Sundararaj, Institute of Wood Science & Technology, Bengaluru, India  
Dr. M. Nithyanandan, Environmental Department, La Al Al Kuwait Real Estate. Co. K.S.C., Kuwait  
Dr. Himender Bharti, Panjab University, Punjab, India  
Mr. Purnendu Roy, London, UK  
Dr. Saito Motoki, The Butterfly Society of Japan, Tokyo, Japan  
Dr. Sanjay Sondhi, TITLI TRUST, Kalpavriksh, Dehradun, India  
Dr. Nguyen Thi Phuong Lien, Vietnam Academy of Science and Technology, Hanoi, Vietnam  
Dr. Nitin Kulkarni, Tropical Research Institute, Jabalpur, India  
Dr. Robin Wen Jiang Ngiam, National Parks Board, Singapore  
Dr. Lionel Monod, Natural History Museum of Geneva, Genève, Switzerland.  
Dr. Asheesh Shivam, Nehru Gram Bharti University, Allahabad, India  
Dr. Rosana Moreira da Rocha, Universidade Federal do Paraná, Curitiba, Brasil  
Dr. Kurt R. Arnold, North Dakota State University, Saxony, Germany  
Dr. James M. Carpenter, American Museum of Natural History, New York, USA  
Dr. David M. Claborn, Missouri State University, Springfield, USA  
Dr. Karen Schnabel, Marine Biologist, Wellington, New Zealand  
Dr. Amazonas Chagas Júnior, Universidade Federal de Mato Grosso, Cuiabá, Brasil  
Mr. Monsoon Jyoti Gogoi, Assam University, Silchar, Assam, India  
Dr. Heo Chong Chin, Universiti Teknologi MARA (UiTM), Selangor, Malaysia  
Dr. R.J. Shiel, University of Adelaide, SA 5005, Australia  
Dr. Siddharth Kulkarni, The George Washington University, Washington, USA  
Dr. Priyadarshan Dharma Rajan, ATREE, Bengaluru, India  
Dr. Phil Alderslade, CSIRO Marine And Atmospheric Research, Hobart, Australia  
Dr. John E.N. Veron, Coral Reef Research, Townsville, Australia  
Dr. Daniel Whitmore, State Museum of Natural History Stuttgart, Rosenstein, Germany.  
Dr. Yu-Feng Hsu, National Taiwan Normal University, Taipei City, Taiwan  
Dr. Keith V. Wolfe, Antioch, California, USA  
Dr. Siddharth Kulkarni, The Hormiga Lab, The George Washington University, Washington, D.C., USA  
Dr. Tomas Ditrich, Faculty of Education, University of South Bohemia in Ceske Budejovice, Czech Republic  
Dr. Mihaly Foldvari, Natural History Museum, University of Oslo, Norway  
Dr. V.P. Uniyal, Wildlife Institute of India, Dehradun, Uttarakhand 248001, India  
Dr. John T.D. Caleb, Zoological Survey of India, Kolkata, West Bengal, India  
Dr. Priyadarshan Dharma Rajan, Ashoka Trust for Research in Ecology and the Environment (ATREE), Royal Enclave, Bangalore, Karnataka, India

#### Fishes

Dr. Neelish Dahanukar, IISER, Pune, Maharashtra, India  
Dr. Topiltzin Contreras MacBeath, Universidad Autónoma del estado de Morelos, México  
Dr. Heok Hee Ng, National University of Singapore, Science Drive, Singapore  
Dr. Rajeev Raghavan, St. Albert's College, Kochi, Kerala, India  
Dr. Robert D. Sluka, Chiltern Gateway Project, A Rocha UK, Southall, Middlesex, UK  
Dr. E. Vivekanandan, Central Marine Fisheries Research Institute, Chennai, India  
Dr. Davor Zanella, University of Zagreb, Zagreb, Croatia  
Dr. A. Biju Kumar, University of Kerala, Thiruvananthapuram, Kerala, India  
Dr. Akhilesh K.V., ICAR-Central Marine Fisheries Research Institute, Mumbai Research Centre, Mumbai, Maharashtra, India  
Dr. J.A. Johnson, Wildlife Institute of India, Dehradun, Uttarakhand, India

#### Amphibians

Dr. Sushil K. Dutta, Indian Institute of Science, Bengaluru, Karnataka, India  
Dr. Annemarie Ohler, Muséum national d'Histoire naturelle, Paris, France

#### Reptiles

Dr. Gernot Vogel, Heidelberg, Germany  
Dr. Raju Vyas, Vadodara, Gujarat, India  
Dr. Pritpal S. Soorae, Environment Agency, Abu Dhabi, UAE.  
Prof. Dr. Wayne J. Fuller, Near East University, Mersin, Turkey  
Prof. Chandrashekher U. Rironker, Goa University, Taleigao Plateau, Goa, India  
Dr. S.R. Ganesh, Chennai Snake Park, Chennai, Tamil Nadu, India  
Dr. Himansu Sekhar Das, Terrestrial & Marine Biodiversity, Abu Dhabi, UAE

#### Birds

Dr. Hem Sagar Baral, Charles Sturt University, NSW Australia  
Dr. Chris Bowden, Royal Society for the Protection of Birds, Sandy, UK  
Dr. Priya Davidar, Pondicherry University, Kalapet, Puducherry, India  
Dr. J.W. Duckworth, IUCN SSC, Bath, UK  
Dr. Rajah Jayopal, SACON, Coimbatore, Tamil Nadu, India  
Dr. Rajiv S. Kalsi, M.L.N. College, Yamuna Nagar, Haryana, India  
Dr. V. Santharam, Rishi Valley Education Centre, Chittoor Dt., Andhra Pradesh, India  
Dr. S. Balachandran, Bombay Natural History Society, Mumbai, India  
Mr. J. Praveen, Bengaluru, India  
Dr. C. Srinivasulu, Osmania University, Hyderabad, India  
Dr. K.S. Gopi Sundar, International Crane Foundation, Baraboo, USA  
Dr. Gombobaataa Sundev, Professor of Ornithology, Ulaanbaatar, Mongolia  
Prof. Reuven Yosef, International Birding & Research Centre, Eilat, Israel  
Dr. Taej Mundkur, Wetlands International, Wageningen, The Netherlands  
Dr. Carol Inskip, Bishop Auckland Co., Durham, UK  
Dr. Tim Inskip, Bishop Auckland Co., Durham, UK  
Dr. V. Gokula, National College, Tiruchirappalli, Tamil Nadu, India  
Dr. Arkady Lelej, Russian Academy of Sciences, Vladivostok, Russia  
Dr. Simon Dowell, Science Director, Chester Zoo, UK  
Dr. Mário Gabriel Santiago dos Santos, Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, Vila Real, Portugal  
Dr. Grant Connette, Smithsonian Institution, Royal, VA, USA  
Dr. M. Zafar-ul Islam, Prince Saud Al Faisal Wildlife Research Center, Taif, Saudi Arabia

#### Mammals

Dr. Giovanni Amori, CNR - Institute of Ecosystem Studies, Rome, Italy  
Dr. Anwaruddin Chowdhury, Guwahati, India  
Dr. David Mallon, Zoological Society of London, UK  
Dr. Shomita Mukherjee, SACON, Coimbatore, Tamil Nadu, India  
Dr. Angie Appel, Wild Cat Network, Germany  
Dr. P.O. Nameer, Kerala Agricultural University, Thrissur, Kerala, India  
Dr. Ian Redmond, UNEP Convention on Migratory Species, Lansdown, UK  
Dr. Heidi S. Riddle, Riddle's Elephant and Wildlife Sanctuary, Arkansas, USA  
Dr. Karin Schwartz, George Mason University, Fairfax, Virginia.  
Dr. Lala A.K. Singh, Bhubaneswar, Orissa, India  
Dr. Mewa Singh, Mysore University, Mysore, India  
Dr. Paul Racey, University of Exeter, Devon, UK  
Dr. Honnallini N. Kumara, SACON, Anaikatty P.O., Coimbatore, Tamil Nadu, India  
Dr. Nishith Dharaiya, HNG University, Patan, Gujarat, India  
Dr. Spartaco Gippoliti, Socio Onorario Società Italiana per la Storia della Fauna "Giuseppe Altobello", Rome, Italy  
Dr. Justus Joshua, Green Future Foundation, Tiruchirappalli, Tamil Nadu, India  
Dr. H. Raghuram, The American College, Madurai, Tamil Nadu, India  
Dr. Paul Bates, Harison Institute, Kent, UK  
Dr. Jim Sanderson, Small Wild Cat Conservation Foundation, Hartford, USA  
Dr. Dan Challender, University of Kent, Canterbury, UK  
Dr. David Mallon, Manchester Metropolitan University, Derbyshire, UK  
Dr. Brian L. Cypher, California State University-Stanislaus, Bakersfield, CA  
Dr. S.S. Talmale, Zoological Survey of India, Pune, Maharashtra, India  
Prof. Karan Bahadur Shah, Budhanilakantha Municipality, Kathmandu, Nepal  
Dr. Susan Cheyne, Borneo Nature Foundation International, Palangkaraya, Indonesia  
Dr. Hemanta Kafley, Wildlife Sciences, Tarleton State University, Texas, USA

#### Other Disciplines

Dr. Aniruddha Belsare, Columbia MO 65203, USA (Veterinary)  
Dr. Mandar S. Paingankar, University of Pune, Pune, Maharashtra, India (Molecular)  
Dr. Jack Tordoff, Critical Ecosystem Partnership Fund, Arlington, USA (Communities)  
Dr. Ulrike Streicher, University of Oregon, Eugene, USA (Veterinary)  
Dr. Hari Balasubramanian, EcoAdvisors, Nova Scotia, Canada (Communities)  
Dr. Rayanna Hellem Santos Bezerra, Universidade Federal de Sergipe, São Cristóvão, Brazil  
Dr. Jamie R. Wood, Landcare Research, Canterbury, New Zealand  
Dr. Wendy Collinson-Jonker, Endangered Wildlife Trust, Gauteng, South Africa  
Dr. Rajeshkumar G. Jani, Anand Agricultural University, Anand, Gujarat, India  
Dr. O.N. Tiwari, Senior Scientist, ICAR-Indian Agricultural Research Institute (IARI), New Delhi, India  
Dr. L.D. Singla, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana, India  
Dr. Rupika S. Rajakaruna, University of Peradeniya, Peradeniya, Sri Lanka  
Dr. Bahar Baviskar, Wild-CER, Nagpur, Maharashtra 440013, India

#### Reviewers 2018–2020

Due to paucity of space, the list of reviewers for 2018–2020 is available online.

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Print copies of the Journal are available at cost. Write to:  
The Managing Editor, JoTT,  
c/o Wildlife Information Liaison Development Society,  
No. 12, Thiruvannamalai Nagar, Saravanampatti - Kalapatti Road,  
Saravanampatti, Coimbatore, Tamil Nadu 641035, India  
ravi@threatenedtaxa.org

**Journal of Threatened Taxa** is indexed/abstracted in Bibliography of Systematic Mycology, Biological Abstracts, BIOSIS Previews, CAB Abstracts, EBSCO, Google Scholar, Index Copernicus, Index Fungorum, JournalSeek, National Academy of Agricultural Sciences, NewJour, OCLC WorldCat, SCOPUS, Stanford University Libraries, Virtual Library of Biology, Zoological Records.

NAAS rating (India) 5.64



OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](#) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

September 2021 | Vol. 13 | No. 11 | Pages: 19431-19674

Date of Publication: 26 September 2021 (Online & Print)

DOI: [10.11609/jott.2021.13.11.19431-19674](https://doi.org/10.11609/jott.2021.13.11.19431-19674)

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

#### Articles

##### Understanding human-flying fox interactions in the Agusan Marsh Wildlife Sanctuary as basis for conservation policy interventions

– Sherry L. Paz & Juan Carlos T. Gonzalez, Pp. 19431–19447

##### Argentinian odonates (dragonflies and damselflies): current and future distribution and discussion of their conservation

– A. Nava-Bolaños, D.E. Vrech, A.V. Peretti & A. Córdoba-Aguilar, Pp. 19448–19465

#### Communications

##### The diel activity pattern of small carnivores of Western Ghats, India: a case study at Nelliampathies in Kerala, India

– Devika Sanghamithra & P.O. Nameer, Pp. 19466–19474

##### Distribution and threats to Smooth-Coated Otters *Lutrogale perspicillata* (Mammalia: Carnivora: Mustelidae) in Shuklaphanta National Park, Nepal

– Gopi Krishna Joshi, Rajeev Joshi & Bishow Poudel, Pp. 19475–19483

##### Wildlife hunting practices of the Santal and Oraon communities in Rajshahi, Bangladesh

– Azizul Islam Barkat, Fahmida Tasnim Liza, Sumaiya Akter, Ashikur Rahman Shome & M. Fazle Rabbe, Pp. 19484–19491

##### Ethnozoological use of primates in northeastern India

– Deborah Daolagupu, Nazimur Rahman Talukdar & Parthakar Choudhury, Pp. 19492–19499

##### Factors influencing the flush response and flight initiation distance of three owl species in the Andaman Islands

– Shanmugavel Sureshmarimuthu, Santhanakrishnan Babu, Honnavalli Nagaraj Kumara & Nagaraj Rajeshkumar, Pp. 19500–19508

##### Birds of Barandabhar Corridor Forest, Chitwan, Nepal

– Saneer Lamichhane, Babu Ram Lamichhane, Kapil Pokharel, Pramod Raj Regmi, Tulasi Prasad Dahal, Santosh Bhattacharai, Chiranjibi Prasad Pokhrel, Pabitra Gotame, Trishna Rayamajhi, Ram Chandra Kandel & Aashish Gurung, Pp. 19509–19526

##### On some additions to the amphibians of Gunung Inas Forest Reserve, Kedah, Peninsular Malaysia

– Shahriza Shahrudin, Pp. 19527–19539

#### Reviews

##### A review of research on the distribution, ecology, behaviour, and conservation of the Slender Loris *Loris lydekkerianus* (Mammalia: Primates: Lorisidae) in India

– Mewa Singh, Mridula Singh, Honnavalli N. Kumara, Shanthala Kumar, Smitha D. Gnanaolivu & Ramamoorthy Sasi, Pp. 19540–19552

##### Bivalves (Mollusca: Bivalvia) in Malaysian Borneo: status and threats

– Abdulla-Al-Asif, Hadi Hamli, Abu Hena Mustafa Kamal, Mohd Hanafi Idris, Geoffrey James Gerusu, Johan Ismail & Muyassar H. Abualrehesh, Pp. 19553–19565

##### Disentangling earthworm taxonomic stumbling blocks using molecular markers

– Azhar Rashid Lone, Samrendra Singh Thakur, Nalini Tiwari, Olusola B. Sokefun & Shweta Yadav, Pp. 19566–19579

##### A reference of identification keys to plant-parasitic nematodes (Nematoda: Tylenchida\ Tylenchomorpha)

– Reza Ghaderi, Manouchehr Hosseinvand & Ali Eskandari, Pp. 19580–19602

#### Short Communications

##### Catalogue of herpetological specimens from Meghalaya, India at the Salim Ali Centre for Ornithology and Natural History

– S.R. Chandramouli, R.S. Naveen, S. Sureshmarimuthu, S. Babu, P.V. Karunakaran & Honnavalli N. Kumara, Pp. 19603–19610

##### A preliminary assessment of odonate diversity along the river Tirthan, Great Himalayan National Park Conservation Area, India with reference to the impact of climate change

– Amar Paul Singh, Kritish De, Virendra Prasad Uniyal & Sambandam Sathyakumar, Pp. 19611–19615

##### A checklist of orthopteran fauna (Insecta: Orthoptera) with some new records in the cold arid region of Ladakh, India

– M. Ali, M. Kamil Usmani, Hira Naz, Tajamul Hassan Baba & Mohsin Ali, Pp. 19616–19625

##### New distribution records of two *Begonias* to the flora of Bhutan

– Phub Gyeltshen & Sherab Jamtsho, Pp. 19626–19631

##### Rediscovery of *Aponogeton lakhonensis* A. Camus (Aponogetonaceae): a long-lost aquatic plant of India

– Debolina Dey, Shrirang Ramchandra Yadav & Nilakshee Devi, Pp. 19632–19635

##### *Glypochloa acuminata* (Hack.) Clayton var. *laevis* (Poaceae): a new variety from central Western Ghats of Karnataka, India

– H.U. Abhijit & Y.L. Krishnamurthy, Pp. 19636–19639

##### A cytomorphological investigation of three species of the genus *Sonchus* L. (Asterales: Asteraceae) from Punjab, India

– M.C. Sidhu & Rai Singh, Pp. 19640–19644

##### *Dryopteris lunanensis* (Dryopteridaceae) - an addition to the pteridophytic diversity of India

– Chhandam Chanda, Christopher Roy Fraser-Jenkins & Vineet Kumar Rawat, Pp. 19645–19648

#### Notes

##### First record of Spotted Linsang *Prionodon pardicolor* (Mammalia: Carnivora: Prionodontidae) with photographic evidence in Meghalaya, India

– Papori Khatonier & Adrian Wansaindor Lyngdoh, Pp. 19649–19651

##### First record of the Eastern Cat Snake *Boiga gocoal* (Gray, 1835) (Squamata: Colubridae) from Tripura, India

– Sumit Nath, Biswajit Singh, Chiranjib Debnath & Joydeb Majumder, Pp. 19652–19656

##### First record of the genus *Tibetanja* (Lepidoptera: Eupterotidae: Janinae) from India

– Alka Vaidya & H. Sankararaman, Pp. 19657–19659

##### *Austroborus cordillerae* (Mollusca: Gastropoda) from central Argentina: a rare, little-known land snail

– Sandra Gordillo, Pp. 19660–19662

##### Intestinal coccidiosis (Apicomplexa: Eimeriidae) in a Himalayan Griffon Vulture *Gyps himalayensis*

– Vimalraj Padayatchiar Govindan, Parag Madhukar Dhakate & Ayush Uniyal, Pp. 19663–19664

##### Two new additions to the orchid flora of Assam, India

– Sanswrang Basumatary, Sanjib Baruah & Lal Ji Singh, Pp. 19665–19670

##### Wildlife art and illustration – combining black and white ink drawings with colour: some experiments in Auroville, India

– M. Eric Ramanujam & Joss Brooks, Pp. 19671–19674

#### Publisher & Host

