

THE DESCRIPTIVE COMPLEXITY OF THE SET OF POISSON GENERIC NUMBERS

VERÓNICA BECHER, STEPHEN JACKSON, DOMINIK KWIETNIAK, AND BILL MANCE

ABSTRACT. Let $b \geq 2$ be an integer. We show that the set of real numbers that are Poisson generic in base b is Π_3^0 -complete in the Borel hierarchy of subsets of the real line. Furthermore, the set of real numbers that are Borel normal in base b and not Poisson generic in base b is complete for the class given by the differences between Π_3^0 sets. We also show that the effective versions of these results hold in the effective Borel hierarchy.

Keywords: Poisson generic numbers; normal numbers; descriptive set theory.

MSC Classification: 03E15; 11U99; 11K16.

1. INTRODUCTION AND STATEMENT OF RESULTS

Years ago Zeev Rudnick introduced Poisson generic real numbers: a real number x is Poisson generic in an integer base $b \geq 2$, if the counts of number of occurrences of words of length k over the alphabet $\{0, 1, \dots, b-1\}$ appearing in the initial segments of the base b expansion of x tends to the Poisson distribution with parameter λ as $k \rightarrow \infty$ for every $\lambda > 0$. That is, the fraction of k words appearing a given number of times among the first digits tends in distribution to the Poisson distribution with parameter λ as $k \rightarrow \infty$. Peres and Weiss [11] proved that Lebesgue almost all real numbers are Poisson generic. Their proof is presented in [3, Theorem 1]. Poisson genericity implies (Borel) normality.

For the rest of the paper, given an integer $b \geq 2$, we identify real numbers in the unit interval $[0, 1)$ with their base b expansions, that is, we identify each $x \in [0, 1)$ with a sequence $x_1x_2x_3\dots$ with values in $\{0, 1, \dots, b-1\}$ such that

$$(1) \quad x = \sum_{j=1}^{\infty} \frac{x_j}{b^j}$$

and $x_j \neq b-1$ for infinitely many $j \geq 1$. Every real number in $[0, 1)$ has a unique such expansion, and for all, but countably many real numbers the base b expansion is the unique representation of x as in (1) (the countable set of exceptions has another representation as in (1) with $x_j = b-1$ for all sufficiently large j 's).

In the sequel we consider an integer $b \geq 2$ that we take as the given base. For a real number x and an interval $A = [q, r]$ of real numbers (respectively $A = [q, r)$), where $1 \leq q < r$ we write $x \upharpoonright A$ to denote the segment of the base- b expansion of x corresponding to positive integers in the interval A . Many times instead of writing $x \upharpoonright [1, r]$ for some $r > 1$ we write we write $x \upharpoonright r$ to denote the initial segment of the base b expansion of x up to position $\lfloor r \rfloor$.

Since Poisson genericity in base b is a property that depends only of the tail of the base b representation of that real number, the integer part of the number is irrelevant. Thus, we

present our results just for the real numbers in the unit interval, but they also hold when the unit interval is replaced by the real line.

Definition (Poisson generic number). Let λ be positive real number. A real number $x \in [0, 1)$ is λ -Poisson generic in base b if for every non-negative integer j we have

$$\lim_{k \rightarrow \infty} Z_{j,k}^\lambda(x) = e^{-\lambda} \frac{\lambda^j}{j!},$$

where

$$Z_{j,k}^\lambda(x) = \frac{1}{b^k} |\{w \in \{0, \dots, (b-1)\}^k : w \text{ occurs } j \text{ times in } x \upharpoonright \lambda b^k + k\}|.$$

A real number x is Poisson generic in base b if it is λ -Poisson generic in base b for every positive real λ .

Let \mathcal{P}_b be the set of real numbers that are Poisson generic in base b . It is easy to see that \mathcal{P}_b is a Borel set. Our goal is to give the descriptive complexity of \mathcal{P}_b . In other words, we would like to locate the exact position of \mathcal{P}_b in the Borel hierarchy (both, lightface and boldface).

Recall that the Borel hierarchy for subsets of the real numbers is the stratification of the σ -algebra generated by the open sets with the usual topology. For references see Kechris's textbook [9].

A set A is Σ_1^0 if and only if A is open and A is Π_1^0 if and only if A is closed. We say that A is Σ_{n+1}^0 for $n \geq 1$ if and only if it is a countable union of Π_n^0 sets, and A is Π_{n+1}^0 if and only if it is a countable intersection of Σ_n^0 sets.

A set A is hard for a Borel class if and only if every set in the class is reducible to A by a continuous map. A set A is complete in a class if it is hard for this class and belongs to the class. By Wadge's celebrated theorem, in spaces like the real numbers with the usual interval topology, a Σ_n^0 set is Σ_n^0 -complete if and only if it is not Π_n^0 .

When we restrict to intervals with rational endpoints and computable countable unions and intersections, we obtain the effective or lightface Borel hierarchy. One way to present the finite levels of the effective Borel hierarchy is by means of the arithmetical hierarchy of formulas in the language of second-order arithmetic. Atomic formulas in this language assert algebraic identities between integers or membership of real numbers in intervals with rational endpoints. A formula in the arithmetic hierarchy involves only quantification over integers. A formula is Π_0^0 and Σ_0^0 if all its quantifiers are bounded. It is Σ_{n+1}^0 if it has the form $\exists x \theta$ where θ is Π_n^0 , and it is Π_{n+1}^0 if it has the form $\forall x \theta$ where θ is Σ_n^0 .

A set A of real numbers is Σ_n^0 (respectively Π_n^0) in the effective Borel hierarchy if and only if membership in that set is definable by a formula which is Σ_n^0 (respectively Π_n^0). Notice that every Σ_n^0 set is Σ_n^0 and every Π_n^0 set is Π_n^0 . In fact, for every set A in Σ_n^0 there is a Σ_n^0 formula and real parameter such that membership in A is defined by that Σ_n^0 formula relative to that real parameter.

A set A is hard for an effective Borel class if and only if every set in the class is reducible to A by a computable map. As before, A is complete in an effective class if it is hard for this class and belongs to the class. Since computable maps are continuous, proofs of hardness in the effective hierarchy often yield proofs of hardness in general by relativization.

The difference hierarchy over a pointclass is generated by taking differences of sets. In the sequel we are just interested in the class $D_2\text{-}\Pi_3^0$ which consists of all the sets that are difference between two sets in Π_3^0 . The class $D_2\text{-}\Pi_3^0$ is the effective counterpart.

Although the definition of Poisson genericity in a given base b asks for λ -Poisson genericity in base b for every positive real λ , it suffices to consider λ -Poisson genericity in base b for every positive rational λ . This is proved in Lemma 3. Then, by the form of its definition, the set \mathcal{P}_b is a Π_3^0 property, hence \mathcal{P}_b is a Borel set appearing as Π_3^0 set in the Borel hierarchy. We shall prove completeness. We first prove the boldface case, and then we add the needed subtleties to prove the lightface case. We start with the following result.

Theorem 1. \mathcal{P}_b is Π_3^0 -complete.

Definition (Borel normal number). Let an integer $b \geq 2$. A real number x is Borel normal in base b if for every block w of digits in $\{0, \dots, (b-1)\}$,

$$\lim_{n \rightarrow \infty} \frac{\text{the number of occurrences of } w \text{ in } x \upharpoonright n}{n} = b^{-|w|}.$$

The set \mathcal{P}_b of real numbers that are Borel normal in base b is Π_3^0 -complete [6, 10]. Every real Poisson generic in base b is Borel normal in base b , see [11] or [5, Theorem 2]. We study the descriptive complexity of the difference set. Let \mathcal{N}_b be the set of real numbers that Borel normal in base b .

Theorem 2. $\mathcal{N}_b \setminus \mathcal{P}_b$ is D_2 - Π_3^0 -complete.

The next two results are the lightface improvements of Theorems 1 and 2.

Theorem 3. \mathcal{P}_b is Π_3^0 -complete.

Theorem 4. $\mathcal{N}_b \setminus \mathcal{P}_b$ is D_2 - Π_3^0 -complete.

Similarly to previous consequences of differences sets of normal numbers for Cantor series expansions being D_2 - Π_3^0 -complete [2], Theorem 2 imposes limitations on the relationship between \mathcal{N}_b and \mathcal{P}_b . An immediate consequence of Theorem 2 is that the set $\mathcal{N}_b \setminus \mathcal{P}_b$ is uncountable. Also, since $\mathcal{N}_b \setminus \mathcal{P}_b$ is D_2 - Π_3^0 -complete, there cannot be a Σ_3^0 set A such that $A \cap \mathcal{N}_b = \mathcal{P}_b$ (as otherwise, we would have $\mathcal{N}_b \setminus \mathcal{P}_b = \mathcal{N}_b \setminus A \in \Pi_0^3$, a contradiction). Thus, no Σ_3^0 condition can be added to normality to give Poisson genericity. Equivalently, any time a Σ_3^0 set contains \mathcal{P}_b , it must contain elements of $\mathcal{N}_b \setminus \mathcal{P}_b$. As an application, consider the following definition of weakly Poisson generic:

Definition (Weakly-Poisson generic number). Say $x \in [0, 1)$ with base b expansion (x_j) is weakly Poisson generic in base b if for every $\epsilon > 0$, every rational λ , and non-negative integer j , we have that for infinitely many k that $|Z_{j,k}^\lambda(x) - e^{-\lambda} \frac{\lambda^j}{j!}| < \epsilon$.

Note that being Poisson generic in base b implies being weakly-Poisson generic. However, being weakly-Poisson generic is a Π_2^0 condition. So, from Theorem 2 we get the following:

Corollary 1. For every base b there is a base- b normal number which is weakly Poisson generic but not Poisson generic.

As another application, consider the following version of discrepancy. Suppose f is a function assigning to each word $w \in b^{<\omega}$ and each positive integer n a positive real number $f(w, n)$. Given $x \in [0, 1)$ with base b expansion (b_j) , say the (w, n) -discrepancy is $D(x, w, n) = |\frac{n}{b^{|w|}} - W(x \upharpoonright n, w)|$, where $W(u, w)$ is the number of occurrences of w in u . We say a real number x has base b f -large discrepancy if for all w and all n we have that $D(x, w, n) > f(w, n)$. The set of x with f -large discrepancy, for any fixed f , is easily a Π_1^0 set. The set of numbers that are Borel normal to base b are exactly those for which the

discrepancy of their initial segments of their expansion in base b goes to zero. We conjecture that the Poisson generic numbers in base b can not have very low discrepancy of their initial segments (for instance, the infinite de Bruijn sequences exist in bases $b \geq 3$, they satisfy that $Z_{1,k}^1 = 1$ for every k , hence they do not correspond to Poisson generic numbers, and they have low discrepancy.) However, we have the following, which states that the Poisson generic reals cannot be characterized as the set of normal numbers satisfying a large discrepancy condition.

Corollary 2. For every function f , the set of base- b Poisson generic reals is not equal to the set of normal numbers with f -large discrepancy.

There are also many other naturally occurring sets of real numbers defined by conditions which make them Σ_3^0 . Examples include countable sets, co-countable sets, the class BA of *badly approximable* numbers (which is a Σ_2^0 set), the Liouville numbers (which is a Π_2^0 set), and the set of $x \in [0, 1]$ where a particular continuous function $f: [0, 1] \rightarrow \mathbb{R}$ is not differentiable. In all these cases, the theorem implies that either the set omits some Poisson generic number, or else contains a number which is normal but not Poisson generic. Of course, many of these statements are easy to see directly, but the point is that they all follow immediately from the general complexity result, Theorem 2.

The set of real numbers whose expansion in *every* integer base is Poisson generic is of course Π_3^0 , but we do not know yet how to prove that this set is Π_3^0 -complete. The set of real numbers whose expansion in one base is λ' -Poisson generic but not λ -Poisson generic, for different positive real numbers λ and λ' , is D_2 - Π_3^0 but we do not know if it is complete.

The result in the present note contribute to the corpus of work on the descriptive complexity of properties of real numbers that started with the questions of Kechris on the descriptive complexity of the set of Borel normal numbers. He conjectured that set of absolutely normal numbers (normal to all integer bases) is $\Pi_3^0(\mathbb{R})$ -complete. Ki and Linton [10] gave the first result towards solving the conjecture by showing that the set of numbers that are normal to base 2 is Π_3^0 -complete. Then V. Becher, P. A. Heiber, and T. A. Slaman [4] settled Kechris' conjecture. Furthermore, V. Becher and T. A. Slaman [6] proved that the set of numbers normal in at least one base is $\Sigma_4^0(\mathbb{R})$ -complete. In another direction, D. Airey, S. Jackson, D. Kwietniak, and B. Mance [1] and, more generally K. Deka, S. Jackson, D. Kwietniak, and B. Mance in [8] showed that for any dynamical system with a weak form of the specification property, the set of generic points for any invariant measure of the system is Π_3^0 -complete. This result generalizes the Ki-Linton result to many numeration systems other than the standard base b one. In general, the Cantor series expansions are not covered even in the generality of [8], so D. Airey, S. Jackson, and B. Mance [2] determined the descriptive complexity of various sets of normal numbers in these numeration systems.

2. BOLDFACE

We write μ for the Lebesgue measure on the real numbers. From Peres and Weiss metric theorem [11, 3] asserting that μ -almost all real numbers in the unit interval are Poisson generic in each integer base b , we have the following.

For μ almost all real numbers x in the unit interval the following holds. Fix an integer base $b \geq 2$ and any $\alpha \in (0, 1)$. Then for any non negative integer j , and any $\epsilon > 0$, for all large enough k we have that

$$\left| Z_{j,k}^{(1-\alpha)}(x) - e^{-(1-\alpha)} \frac{(1-\alpha)^j}{j!} \right| < \epsilon.$$

Proof of Theorem 1. Let $\mathcal{C} = \{z \in (\omega \setminus \{0, 1\})^\omega : \lim_i z(i) = \infty\}$. So, \mathcal{C} is Π_3^0 -complete. We define a continuous map $f: \omega^\omega \rightarrow (0, 1)$ which reduces \mathcal{C} to \mathcal{P}_b , that is, $f(z) \in \mathcal{P}_b$ if and only if $z \in \mathcal{C}$. Fix $z \in (\omega \setminus \{0, 1\})^\omega$. At step i we define $f(z) \upharpoonright [b^{k_{i-1}}, b^{k_i})$, where $\{k_i\}$ is a sufficiently fast-growing sequence of positive integers. Let

$$B_i := [b^{k_{i-1}}, b^{k_i}),$$

$$B'_i := \left[b^{k_{i-1}}, \left(1 - \frac{1}{z(i)}\right)b^{k_i} \right).$$

The set B'_i is non-empty as we may assume $k_i > 2k_{i-1}$. Let $x \in (0, 1)$ be Poisson generic in base b . We set

$$f(z) \upharpoonright B'_i = x \upharpoonright B'_i \text{ and } f(z) \upharpoonright B_i \setminus B'_i = 0.$$

First suppose $z \notin \mathcal{C}$, and fix $p \in \omega$ such that for infinitely many i we have $z(i) = p$. Consider step i in the construction of $f(z)$ for such an i . For any $\epsilon > 0$, if i is large enough then the number of words w of length k_i which occur in $x \upharpoonright [1, (1 - \frac{1}{z(i)})b^{k_i}]$ is at most

$$b^{k_i} (1 - e^{-(1 - \frac{1}{z(i)})}) + \epsilon).$$

So, the number Z_i of words w of length k_i which occur in $f(z) \upharpoonright b^{k_i}$ is at most

$$b^{k_i} (1 - e^{-(1 - \frac{1}{z(i)})}) + \epsilon + b^{k_{i-1}}.$$

So,

$$\frac{1}{b^{k_i}} Z_i \leq (1 - e^{-(1 - \frac{1}{p})}) + 2\epsilon$$

if i is large enough using the fact that the k_i grow sufficiently fast. On the other hand, the Poisson estimate for the proportion of words of length k_i occurring in a Poisson generic sequence of length b^{k_i} is $1 - 1/e$. Since p is fixed, as i gets large we have a contradiction. So, $f(z)$ is not 1-Poisson generic.

Next suppose that $z \in \mathcal{C}$. We show that $f(z)$ is Poisson generic in base b . Fix $\lambda > 0$ and $\ell \in \omega$. Fix also $\epsilon > 0$. Consider $k \in \omega$, and let i be such that $k_{i-1} \leq k < k_i$. We show that for k (and hence i) sufficiently large that $|Z_{\ell, k}^\lambda(f(z)) - e^{-\lambda} \frac{\lambda^\ell}{\ell!}| < \epsilon$. Assume i is large enough so that $\frac{1}{z(j)} < \epsilon$ for all $j \geq i - 1$. First consider the case $\lambda \leq 1$. Note that, as $z(i) \geq 2$,

$$b^k \leq \frac{1}{b} b^{k_i} \leq b^{k_i} \left(1 - \frac{1}{z(i)}\right).$$

We have that

$$(2) \quad \left| \frac{1}{b^k} Z_{\ell, k}^\lambda(f(z)) - \frac{1}{b^k} Z_{\ell, k}^\lambda(x) \right| \leq \frac{1}{b^k} \left(b^{k_{i-1}} \frac{1}{z(i-1)} + 6k + b^{k_{i-2}} \right)$$

$$\leq \frac{1}{z(i-1)} + \epsilon$$

$$\leq 2\epsilon.$$

for i large enough. We have used here the fact that

$$|Z_{\ell, k}^\lambda(f(z)) - Z_{\ell, k}^\lambda(x)|$$

is at most the number of words of length k which appear in one of $f(z) \upharpoonright b^k$, $x \upharpoonright b^k$ at a position which overlaps the block $[b^{k_{i-1}}(1 - \frac{1}{z(i-1)}), b^{k_{i-1}})$, or else overlaps the block $[1, b^{k_{i-2}}]$, which gives the above estimate.

Consider now the case $\lambda > 1$. If $\lambda b^k < b^{k_i}(1 - \frac{1}{z(i)})$, then the same estimate above works. So, suppose $b^k \geq \frac{1}{\lambda} b^{k_i}(1 - \frac{1}{z(i)})$. We may assume that

$$\lambda b^k < \frac{1}{2} b^{k_{i+1}} \leq (1 - \frac{1}{z(i+1)}) b^{k_{i+1}}$$

since λ is fixed and the k_i grow sufficiently fast (in particular $\frac{b^{k_{i+1}}}{b^{k_i}} \rightarrow \infty$). In this case we also count the number of words w of length k which might overlap the block of 0s in $f(z) \upharpoonright [b^{k_i}(1 - \frac{1}{z(i)}), b^{k_i}]$. We then get

$$\begin{aligned} \left| \frac{1}{b^k} Z_{\ell,k}^\lambda(f(z)) - \frac{1}{b^k} Z_{\ell,k}^\lambda(x) \right| &\leq \frac{1}{b^k} \left(b^{k_{i-1}} \frac{1}{z(i-1)} + b^{k_i} \frac{1}{z(i)} + 10k + b^{k_{i-2}} \right) \\ &\leq \frac{1}{z(i-1)} + \frac{b^{k_i}}{b^k} \frac{1}{z(i)} + \epsilon \\ &\leq \frac{1}{z(i-1)} + \lambda \frac{1}{1 - \frac{1}{z(i)}} \frac{1}{z(i)} + \epsilon \\ &\leq 2\epsilon. \end{aligned}$$

if i is sufficiently large, since λ is fixed and $z(i) \rightarrow \infty$. □

For the proof of Theorem 2 we require the following two lemmas.

Lemma 1. Fix an integer $b \geq 2$. Almost all real numbers in $(0, 1)$ have the property that for any α of the form $\alpha = \frac{1}{2^\ell}$ we have

$$\lim_{i \rightarrow \infty} \frac{1}{b^{k_i}} |H_i| = (1 - e^{-\alpha})(e^{-(1-\alpha)}),$$

where H_i is the set of words of length k_i which occur in the base- b expansion of x with a starting position $[(1 - \alpha)b^{k_i}, b^{k_i})$, but do not occur with a starting position in $[b^{k_{i-1}}, (1 - \alpha)b^{k_i}]$.

In fact, this claim holds for any x which is Poisson generic in base b .

Proof. Let $x \in (0, 1)$ be Poisson generic in base b and fix α a negative power of 2. Let

- A_i be the set of words of length k_i occurring in $[b^{k_{i-1}}, b^{k_i})$.
- C_i be the set of words of length k_i occurring in $[b^{k_{i-1}}, (1 - \alpha)b^{k_i})$.

Clearly $C_i \subseteq A_i$. The words which occur in $[(1 - \alpha)b^{k_i}, b^{k_i})$ but not in $[b^{k_{i-1}}, (1 - \alpha)b^{k_i})$ are exactly the words which occur in A_i but not C_i .

Let

- A'_i be the set of words that occur in $[1, b^{k_i})$
- C'_i be the set of words that occur in $[1, (1 - \alpha)b^{k_i})$.

Then

$$||A_i \setminus C_i| - |A'_i \setminus C'_i|| \leq b^{k_{i-1}}.$$

Since x is Poisson generic in base b , for any $\epsilon > 0$ we have that for all large enough i that

$$\left| \frac{1}{b^{k_i}} |A'_i| - (1 - \frac{1}{e}) \right| < \epsilon.$$

Similarly, as x is Poisson generic in base b , and using $\lambda = 1 - \alpha$, we have that

$$\left| \frac{1}{b^{k_i}} |C'_i| - (1 - e^{-(1-\alpha)}) \right| < \epsilon.$$

By these, Lemma 1 readily follows. \square

Assume x lies in the measure one set of Lemma 1 and that the k_i grow fast enough, then

$$\left| \frac{1}{b^{k_i}} |H_i| - (1 - e^{-\alpha})(e^{-(1-\alpha)}) \right| < \frac{1}{2^i}.$$

A standard probability computation shows the following.

Lemma 2. There is a function $g: \omega \rightarrow \omega$ such that the following holds. Suppose $k_0 < k_1 < \dots$ are such that $b^{k_i} - b^{k_{i-1}} > g(i-1)$ for all i . Then μ -almost all $x \in (0, 1)$ satisfy the following: for any $j \in \omega$, any $w \in b^j$ and any $\epsilon > 0$, for all large enough i , and any $n > g(i-1)$

$$\left| \frac{1}{n} W(x \upharpoonright [b^{k_{i-1}}, b^{k_{i-1}} + n), w) - \frac{1}{b^j} \right| < \epsilon$$

where $W(s, w)$ is the number of occurrences of the word w in s .

Proof. We can take $g(n) = n$. Fix j and $w \in b^j$, and fix $\epsilon > 0$. It suffices to show that for almost all x that for all large enough i and any $n > g(i-1) = i-1$ that

$$\left| \frac{1}{n} W(x \upharpoonright [b^{k_{i-1}}, b^{k_{i-1}} + n), w) - \frac{1}{b^j} \right| < \epsilon.$$

Bernstein's inequality in Probability theory bounds the probability that the sum of random variables deviates from its mean. Applying it we obtain that there are constants $\alpha, \beta > 0$ such that for all n , the probability that a string $s \in b^n$ violates the inequality $|\frac{1}{n} W(s, w) - \frac{1}{b^j}| < \epsilon$ is less than $\alpha e^{-\beta n}$. Many works on normal numbers include a complete proof of this result, see for instance [7, Lemma 8].

So, the probability that an $x \in (0, 1)$ violates $|\frac{1}{n} W(x \upharpoonright [b^{k_{i-1}}, b^{k_{i-1}} + n), w) - \frac{1}{b^j}| < \epsilon$ for some $i \geq i_0$ and $n \geq i$ is at most

$$\sum_{i \geq i_0} \sum_{n \geq i} \alpha e^{-\beta n} \leq \sum_{i \geq i_0} \alpha \frac{e^{-\beta i}}{1 - e^{-\beta}} = \frac{\alpha e^{-\beta i_0}}{(1 - e^{-\beta})^2}.$$

Since this tends to 0 with i_0 , the result follows. \square

We can now give the proof of the $D_2\text{-}\Pi_3^0$ completeness of the difference set $\mathcal{N}_b \setminus \mathcal{P}_b$.

Proof of Theorem 2. We fix a sufficiently fast growing sequence $k_0 < k_1 < \dots$ as in Lemma 2, and then fix $x \in (0, 1)$ to be Poisson generic in base b (so that Lemma 1 holds) and also to be in the measure one set where Lemma 2 holds for this sequence $(k_i)_{i \geq 0}$.

We let $C = \{z \in \omega^\omega : z(2n) \rightarrow \infty\}$, and $D = \{z \in \omega^\omega : z(2n+1) \rightarrow \infty\}$. We define a continuous map $f: \omega^\omega \rightarrow (0, 1)$ which reduces $C \setminus D$ to $\mathcal{N}_b \setminus \mathcal{P}_b$. The idea to define f so that for $z \in \omega^\omega$, the even digits $z(2i)$ will control whether $f(z) \in \mathcal{N}_b$ and the odd digits $z(2i+1)$ will control whether $f(z) \in \mathcal{P}_b$. When we wish to violate Poisson genericity, we will do so for $\lambda = 1$ and $j = 0$. We may assume without loss of generality that all $z(i)$ and all k_i are positive powers of 2.

As before, at step i we define $f(z) \upharpoonright B_i$, where $B_i = [b^{k_{i-1}}, b^{k_i})$. Let

$$\begin{aligned}
B_i^1 &:= [b^{k_{i-1}}, (1 - \frac{1}{z(2i)} - \frac{1}{z(2i+1)})b^{k_i}) \\
B_i^2 &:= [(1 - \frac{1}{z(2i)} - \frac{1}{z(2i+1)})b^{k_i}, (1 - \frac{1}{z(2i+1)})b^{k_i}) \\
B_i^3 &:= [(1 - \frac{1}{z(2i+1)})b^{k_i}, b^{k_i})
\end{aligned}$$

So,

$$\begin{aligned}
|B_i^2| &= \frac{1}{z(2i)}b^{k_i}, \\
|B_i^3| &= \frac{1}{z(2i+1)}b^{k_i}.
\end{aligned}$$

We let

$$\begin{aligned}
f(z) \upharpoonright B_i^1 &:= x \upharpoonright B_i^1, \\
f(z) \upharpoonright B_i^2 &:= 0, \\
f(z) \upharpoonright B_i^3 &:= x \upharpoonright [b^{k_{i-1}}, b^{k_{i-1}} + |B_i^3|) = x \upharpoonright \left[b^{k_{i-1}}, b^{k_{i-1}} + \frac{1}{z(2i+1)}b^{k_i} \right).
\end{aligned}$$

We show that f is a reduction from $C \setminus D$ to $\mathcal{N}_b \setminus \mathcal{P}_b$.

First assume $z \notin C$, that is $z(2i)$ does not tend to ∞ . Fix ℓ such that $z(2i) = \ell$ for infinitely many i . We easily have that $f(z) \notin \mathcal{N}_b$. For example, if the digit 0 occurs with approximately the right frequency $\frac{1}{b}$ in

$$f(z) \upharpoonright [1, b^{k_{i-1}} + |B_i^1|) = \left[0, b^{k_i} \left(1 - \frac{1}{z(2i)} - \frac{1}{z(2i+1)} \right) \right),$$

then 0 will occur with too large a frequency in

$$f(z) \upharpoonright [1, b^{k_{i-1}} + |B_i^1| + |B_i^2|) = \left[0, b^{k_{i-1}} + |B_i^1| + \frac{1}{\ell}b^{k_i} \right).$$

This is because $f(z) \upharpoonright B_i^2 = 0$ and $|B_i^2| = \frac{1}{\ell}b^{k_i}$ for such i .

So we may henceforth assume that $z \in C$, so $\frac{1}{b^{k_i}}|B_i^2| = \frac{1}{z(2i)} \rightarrow 0$. We observe that this implies that $f(z) \in \mathcal{N}_b$. This follows from Lemma 2 and that we may assume $\lim_i \frac{g(i-1)}{b^{k_{i-1}}} = 0$.

Now assume that $z \in D$, so $z \notin C \setminus D$. We show $f(z) \in \mathcal{P}_b$, and so $f(z) \notin \mathcal{N}_b \setminus \mathcal{P}_b$. Since we are assuming $z \in C$ also, we have $\lim_{i \rightarrow \infty} z(i) = \infty$. So, $\lim_{i \rightarrow \infty} \frac{1}{b^{k_i}}(|B_i^2| + |B_i^3|) = 0$. It then follows exactly as in Equation 2 in the proof of Theorem 1 that $f(z) \in \mathcal{P}_b$.

Assume next that $z \notin D$ (but $z \in C$ still). We show that $f(z) \notin \mathcal{P}_b$, which shows $f(z) \in \mathcal{N}_b \setminus \mathcal{P}_b$. Fix m so that for infinitely many i we have $z(2i+1) = m$, and we may assume m is of the form $m = 2^\ell$. Recall $\frac{1}{b^{k_i}}|B_i^3| = \frac{1}{z(2i+1)} = \frac{1}{2^\ell}$ for such i . We restrict our attention to this set of i in the following argument. If $f(z)$ were Poisson generic, then from Lemma 1 we would have that for large enough i in our set that

$$\frac{1}{b^{k_i}}|H_i| \approx (1 - e^{-\alpha})(e^{-(1-\alpha)}),$$

where H_i is the set of words of length k_i which occur in $f(z)$ with a starting position in $[(1-\alpha)b^{k_i}, b^{k_i})$, but do not occur in $f(z)$ with a starting position in $[b^{k_{i-1}}, (1-\alpha)b^{k_i})$. However,

by the construction of $f(z)$ we have that every word which occurs in $[(1-\alpha)b^{k_i}, b^{k_i})$ also occurs in $[b^{k_{i-1}}, (1-\alpha)b^{k_i})$, and so $|H_i| = 0$. \square

3. LIGHTFACE REFINEMENTS

The existence of computable Poisson generic real number was proved in [3, Theorem 2]. We show how to compute an instance of a Poisson generic real number in base b . We start by proving that to obtain a Poisson generic real it suffices to construct an instance of a λ -Poisson generic real for every positive rational λ .

Lemma 3 (adapted from [3]). Let integer $b \geq 2$. If $x \in (0, 1)$ is λ -Poisson generic in base b for all positive rational λ then x is Poisson generic in base b .

Proof. For each $x \in (0, 1)$ and for each $k \in \mathbb{N}$, on the space of words of length k with uniform measure define the integer-valued random measure $M_k^x = M_k^x(v)$ on the real half-line $\mathbb{R}^+ = [0, +\infty)$ by setting for all Borel sets $S \subseteq \mathbb{R}^+$,

$$M_k^x(S)(v) := \sum_{p \in \mathbb{N} \cap b^k S} I_p(x, v),$$

where I_p is the indicator function that v occurs in x at position p and $\mathbb{N} \cap b^k S$ denotes the set of integer values in $\{b^k s : s \in S\}$. Then, $M_k^x(\cdot)$ is a point process on \mathbb{R}^+ . The function $Z_{j,k}^\lambda(x)$ can be formulated in terms of $M_k^x(S)$ for the sets $S = (0, \lambda]$, as follows:

$$Z_{j,k}^\lambda(x) = \frac{1}{b^k} \# \{v \in \{0, \dots, (b-1)\}^k : M_k^x((0, \lambda])(v) = j\}.$$

Observe that for every pair of positive reals λ, λ' , with $\lambda < \lambda'$,

$$M_k^x((0, \lambda'])(v) - M_k^x((0, \lambda])(v) = \sum_{p \in \mathbb{N} \cap b^k [\lambda, \lambda')} I_p(x, v).$$

The classical total variation distance d_{TV} between two probability measures P and Q on a σ -algebra \mathcal{F} is defined via

$$d_{TV}(P, Q) := \sup_{A \in \mathcal{F}} |P(A) - Q(A)|.$$

For a random variable X taking values in \mathbb{R} , the distribution of X is the probability measure μ_X on \mathbb{R} defined as the push-forward of the probability measure on the sample space of X . The total variation distance between two random variables X and Y is simply $d_{TV}(X, Y) = d_{TV}(\mu_X, \mu_Y)$. Hence, the total distance variation

$$d_{TV}(M_k^x((0, \lambda']), M_k^x((0, \lambda])) \leq \frac{1}{b^k} \#(\mathbb{N} \cap b^k [\lambda, \lambda')) = \lambda' - \lambda + O(b^{-k}).$$

Let $Po(\lambda)$ to denote the Poisson distribution with parameter $\lambda > 0$ (if X is a discrete random variable, the probability that X is equal to j , for each $j = 0, 1, 2, \dots$, is $\lambda^j e^{-\lambda} / j!$). Observe that $d_{TV}(Po(\lambda'), Po(\lambda)) \rightarrow 0$ as $\lambda \rightarrow \lambda'$. From these two observations and the fact that the rational numbers are a dense subset of the real numbers we conclude that if x is λ -Poisson generic for every positive rational λ then x is Poisson generic. \square

Definition (Values N_n and sets E_n). For each $n \geq 1$ define

$$N_n := b^{2^n},$$

$$E_n := (0, 1) \setminus \bigcup_{N_n \leq k < N_{n+1}} \text{Bad}_k,$$

where

$$\begin{aligned} \text{Bad}_k &:= \bigcup_{j \in J_k} \bigcup_{\lambda \in L_k} \text{Bad}(\lambda, k, j, 1/k), \\ J_k &:= \{0, \dots, b^k - 1\}, \\ L_k &:= \{p/q : q \in \{1, \dots, k\}, p/q < k\}, \\ \text{Bad}(\lambda, k, j, \varepsilon) &:= \left\{ x \in (0, 1) : \left| Z_{j,k}^\lambda(x) - \frac{e^{-\lambda} \lambda^j}{j!} \right| > \varepsilon \right\}. \end{aligned}$$

Observe that each set Bad_k is a finite union of intervals with rational endpoints. Also each set E_n is a finite union of intervals with rational endpoints.

Fact 1. There is n_0 such that for every n greater than n_0 , $\mu(E_n) > 1 - \frac{1}{N_n^2}$.

Proof. By [3, Proof of Theorem 2] there is k_0 such that for every $k \geq k_0$, for every $j \geq 0$,

$$\mu(\text{Bad}(\lambda, k, j, 1/k)) < 2e^{-\frac{b^k}{2\lambda k^4}}$$

and

$$\mu(\text{Bad}_k) = \mu\left(\bigcup_{j \in J_k} \bigcup_{\lambda \in L_k} \text{Bad}(\lambda, k, j, 1/k)\right) < 2b^k k^3 e^{-b^k/(2k^5)}.$$

Since $\mu(\text{Bad}_k)$ has double exponential decay as k increases, for $N_n = b^{2^n}$ with n sufficiently large,

$$\mu(\text{Bad}_{N_n}) < \frac{1}{2N_n^2}.$$

Since $E_n = (0, 1) \setminus \bigcup_{N_n \leq k < N_{n+1}} \text{Bad}_k$, there is n_0 such that for every $n > n_0$,

$$\mu(E_n) \geq 1 - 2\mu(\text{Bad}_{N_n}) > 1 - \frac{1}{N_n^2}. \quad \square$$

Fact 1 ensures that the set $\bigcap_{n \geq n_0} E_n$ has positive measure. Let see that $\bigcap_{n \geq n_0} E_n$ consists entirely of Poisson generic real numbers for base b . Suppose that x is not Poisson generic for base b . By Lemma 3 x is not λ -Poisson generic in base b for some positive rational λ . Then, there is a positive ε and a non-negative integer j such that for infinitely many n 's,

$$\left| Z_{j,n}^\lambda(x) - \frac{e^{-\lambda} \lambda^j}{j!} \right| > \varepsilon.$$

Let $n_1 = n_1(\lambda, \varepsilon, j)$ be the smallest such that $\lambda \in L_{n_1}, j \in J_{n_1}, \varepsilon \geq 1/n_1$. Since sets J_n and L_n are subset increasing in n , for every $n \geq n_1$ we have $\lambda \in L_n$ and $j \in J_n$. And since $\varepsilon > 1/n_1$ we have $\varepsilon > 1/n$, for every $n > n_1$. Then, for infinitely many values of n greater than or equal to n_1 , $x \in \text{Bad}_n$. Hence, for infinitely many values of n , $x \notin E_n$, and thus $x \notin \bigcap_{n \geq n_0} E_n$.

The following algorithm is an adaptation of Turing's algorithm for computing an absolutely normal number (see [7]). We modified it to obtain a real that is Poisson generic in base b .

Algorithm. Let n_0 be determined by Fact 1. Let $I_{n_0} := (0, 1)$.
 At each step $n > n_0$, divide I_{n-1} in b equal parts $I_{n-1}^0, I_{n-1}^1, \dots, I_{n-1}^{b-1}$.
 Let v be the smallest in $\{0, \dots, (b-1)\}$ such that $\mu(I_{n-1}^v \cap E_n) > \frac{1}{N_n}$.
 $I_n := I_{n-1}^v$.
 The n -digit in the base- b expansion of x is the digit v .

Remark. Observe that the number x computed by the algorithm ensures that for each $n \geq n_0$, $x \in I_n \cap E_n$. Since the intervals I_n are nested, we have

$$x \in I_n \cap \left(\bigcap_{n_0 \leq m \leq n} E_m \right),$$

where $E_n = (0, 1) \setminus \bigcup_{N_n \leq k \leq N_{n+1}} \text{Bad}_k$ with $N_n = b^{2n}$. Thus, to define $x \upharpoonright n$ the algorithm looks at all the possible continuations up to $x \upharpoonright b^{N_{n+1}}$.

We prove that the number x produced by the algorithm is indeed Poisson generic for base b . The algorithm defines a sequence of intervals $(I_n)_{n \geq n_0}$ such that $I_n = (\frac{a}{b^n}, \frac{a+1}{b^n})$ for some $a \in \{0, \dots, b^n - 1\}$, $I_{n+1} \subseteq I_n$ and $\mu(I_n) = b^{-n}$. The number x defined is the unique element in $\bigcap_{n \geq n_0} I_n$. We first prove that for every $n \geq n_0$,

$$\mu\left(I_n \cap \bigcap_{i=n_0}^n E_i\right) > 0.$$

To show this we prove by induction on n ,

$$\mu\left(I_n \cap \bigcap_{i=n_0}^n E_n\right) > \frac{1}{N_n}.$$

Base case. For n_0 it is immediate because $I_{n_0} = (0, 1)$, so $\mu(I_{n_0}) = 1$ and

$$\mu(E_{n_0}) > 1 - \frac{1}{N_{n_0}^2} > \frac{1}{N_{n_0}}.$$

Inductive case. Assume the inductive hypothesis

$$\mu\left(I_n \cap \bigcap_{i=n_0}^n E_i\right) > \frac{1}{N_n}.$$

Let's see it holds for $n+1$. Using the inductive hypothesis and Fact 1, we have

$$\begin{aligned} \mu\left(I_n \cap \bigcap_{i=n_0}^{n+1} E_i\right) &= \mu\left(\left(I_n \cap \bigcap_{i=n_0}^n E_i\right) \cap E_{n+1}\right) \\ &> \mu\left(I_n \cap \bigcap_{i=n_0}^n E_i\right) - \mu((0, 1) - E_{n+1}) \\ &> \frac{1}{N_n} - \frac{1}{N_n^2} \\ &> \frac{b}{N_{n+1}}. \end{aligned}$$

By this, there is at least one $v \in \{0, \dots, (b-1)\}$ such that

$$\mu\left(I_n^v \cap \bigcap_{i=n_0}^{n+1} E_i\right) > \frac{1}{N_{n+1}}.$$

Since the algorithm sets I_{n+1} to be the leftmost I_n^v with this property, we have

$$I_{n+1} \cap \bigcap_{i=n_0}^{n+1} E_i > \frac{1}{N_{n+1}}.$$

We conclude that $x \in \bigcap_{n \geq n_0} E_n$. So x is λ -Poisson generic in base b for all positive rational λ .

The proofs of Theorems 3 and 4 are very similar to those of Theorems 1 and 2. However we now include what is needed to prove the lightface results. We now start with a computable Poisson generic number in base b that we obtain with the Algorithm above, and we computably determine the sequence of values $(k_i)_{i \geq 1}$ using the input sequence $z \in \omega^\omega$.

Proof of Theorem 3. Let $\mathcal{C} = \{z \in \omega^\omega : \lim_{i \rightarrow \infty} z(i) = \infty\}$. So, \mathcal{C} is Π_3^0 -complete. We define a computable map $f: \omega^\omega \rightarrow (0, 1)$ which reduces \mathcal{C} to \mathcal{P}_b . Fix $z \in \omega^\omega$. At step i , let k_i be the least integer such that $k_i > k_{i-1}$, and $k_i > z(i)$. Fix $k_0 = 0$. For $i > 0$, we define $f(z) \upharpoonright [b^{k_{i-1}}, b^{k_i})$ as follows. Let

$$\begin{aligned} B_i &:= [b^{k_{i-1}}, b^{k_i}) \\ B'_i &:= \left[b^{k_{i-1}}, \left(1 - \frac{1}{z(i)}\right)b^{k_i}\right). \end{aligned}$$

We set

$$f(z) \upharpoonright B'_i := x \upharpoonright B'_i, \text{ and } f(z) \upharpoonright B_i \setminus B'_i := 0.$$

First suppose $z \notin \mathcal{C}$, and fix $\ell \in \omega$ such that for infinitely many i we have $z(i) = \ell$. Consider step i in the construction of $f(z)$ for such an i . For any $\epsilon > 0$, if i is large enough then the number of words w of length k_i which occur in $x \upharpoonright [1, (1 - \frac{1}{z(i)})b^{k_i}]$ is at most

$$b^{k_i}(1 - e^{-(1 - \frac{1}{z(i)})} + \epsilon).$$

Then, the number Z_i of words w of length k_i which occur in $f(z) \upharpoonright b^{k_i}$ is at most

$$b^{k_i}(1 - e^{-(1 - \frac{1}{z(i)})} + \epsilon).$$

So,

$$\frac{1}{b^{k_i}} Z_i \leq (1 - e^{-(1 - \frac{1}{\ell})} + 2\epsilon).$$

On the other hand, the Poisson estimate for the proportion of words of length k_i occurring in an initial segment of length b^{k_i} is $1 - 1/e$. Since ℓ is fixed, as i gets large we have a contradiction. So, $f(z)$ is not 1-Poisson generic in base b .

Next suppose that $z \in \mathcal{C}$. We show that $f(z)$ is Poisson generic in base b . Fix a positive rational λ and $\epsilon > 0$. Consider any $k \in \omega$, large enough so that the following holds: Let i be such that $k_{i-1} \leq k < k_i$,

- if $\lambda = \frac{p}{q}$ then $k_{i-1} \geq q$,
- $k_i > \frac{1}{\epsilon}$,
- for all $s \geq i - 1$ we have $\frac{1}{z(s)} < \epsilon$.

We show that for any such k , and for every non negative j less than b^k , $|Z_{j,k}^\lambda(f(z)) - e^{-\lambda} \frac{\lambda^j}{j!}| < \epsilon$. First consider the case $\lambda \leq 1$. Fix j . We have that

$$\begin{aligned} \left| \frac{1}{b^k} Z_{j,k}^\lambda(f(z)) - \frac{1}{b^k} Z_{j,k}^\lambda(x) \right| &\leq \frac{1}{b^k} \left(b^{k_{i-1}} \frac{1}{z(i-1)} + 2k + \sum_{m < i-1} b^{k_m} \right) \\ &\leq \frac{1}{z(i-1)} + \epsilon \\ &\leq 2\epsilon \end{aligned}$$

for i large enough. We have used here the fact that $|Z_{j,k}^\lambda(f(z)) - Z_{j,k}^\lambda(x)|$ is at most the number of words of length k which appear in one of $f(z) \upharpoonright b^k$, $x \upharpoonright b^k$ but not the other. Such a word must overlap the block of 0s in $f(z) \upharpoonright [b^{k_{i-1}}(1 - \frac{1}{z(i-1)}), b^{k_{i-1}}]$, or else overlap $[1, b^{k_{i-2}}]$, which gives the above estimate.

Consider now the case $\lambda > 1$. If $\lambda b^k < b^{k_i}(1 - \frac{1}{z(i)})$, then the same estimate above works. So, suppose $b^k \geq \frac{1}{\lambda} b^{k_i}(1 - \frac{1}{z(i)})$. In this case we also count the number of words w of length k which might overlap the block of 0s in $f(z) \upharpoonright [b^{k_i}(1 - \frac{1}{z(i)}), b^{k_i}]$. We then get

$$\begin{aligned} \left| \frac{1}{b^k} Z_{j,k}^\lambda(f(z)) - \frac{1}{b^k} Z_{j,k}^\lambda(x) \right| &\leq \frac{1}{b^k} \left(b^{k_{i-1}} \frac{1}{z(i-1)} + b^{k_i} \frac{1}{z(i)} + 3k + \sum_{s < i-1} b^{k_s} \right) \\ &\leq \frac{1}{z(i-1)} + \frac{b^{k_i}}{b^k} \frac{1}{z(i)} + \epsilon \\ &\leq \frac{1}{z(i-1)} + \lambda \frac{1}{1 - \frac{1}{z(i)}} \frac{1}{z(i)} + \epsilon \\ &\leq 2\epsilon \end{aligned}$$

if i is sufficiently large, since λ is fixed and $z(i) \rightarrow \infty$. □

We can now prove the D_2 - Π_3^0 -completeness of the difference set $\mathcal{N}_b \setminus \mathcal{P}_b$.

Proof of Theorem 4. The proof is exactly as that of Theorem 2 except that we start with a computable real x and we determine the sequence $(k_i)_{i \geq 1}$ using the input sequence $z \in \omega^\omega$. Let x be the number obtained by the Algorithm.

Let $C := \{z \in \omega^\omega : z(2n) \rightarrow \infty\}$ and $D := \{z \in \omega^\omega : z(2n+1) \rightarrow \infty\}$. We assume without loss of generality that all $z(i)$ are powers of 2. We define a computable map $f: \omega^\omega \rightarrow (0, 1)$ which reduces $C \setminus D$ to $\mathcal{N}_b \setminus \mathcal{P}_b$. Fix $z \in \omega^\omega$. At step i , let k_i be the least power of 2 such that $k_i > k_{i-1}$, and $k_i > z(i)$. We define f so that for $z \in \omega^\omega$, the even digits $z(2i)$ will control whether $f(z) \in \mathcal{N}_b$ and the odd digits $z(2i+1)$ control whether $f(z) \in \mathcal{P}_b$. When we wish to violate Poisson genericity, we do so for $\lambda = 1$ and $j = 0$.

As in the proof of Theorem 2, at step i we define $f(z) \upharpoonright B_i$, where $B_i = [b^{k_{i-1}}, b^{k_i})$. Let

$$\begin{aligned} f(z) \upharpoonright B_i^1 &:= x \upharpoonright B_i^1 \\ f(z) \upharpoonright B_i^2 &:= 0 \\ f(z) \upharpoonright B_i^3 &:= x \upharpoonright \left[b^{k_{i-1}}, b^{k_{i-1}} + \frac{1}{z(2i+1)} b^{k_i} \right) \end{aligned}$$

where

$$\begin{aligned} B_i^1 &:= \left[b^{k_{i-1}}, \left(1 - \frac{1}{z(2i)} - \frac{1}{z(2i+1)}\right) b^{k_i} \right) \\ B_i^2 &:= \left[\left(1 - \frac{1}{z(2i)} - \frac{1}{z(2i+1)}\right) b^{k_i}, \left(1 - \frac{1}{z(2i+1)}\right) b^{k_i} \right) \\ B_i^3 &:= \left[\left(1 - \frac{1}{z(2i+1)}\right) b^{k_i}, b^{k_i} \right). \end{aligned}$$

Notice that $|B_i^2| = \frac{1}{z(2i)} b^{k_i}$, and $|B_i^3| = \frac{1}{z(2i+1)} b^{k_i}$.

We show that f is a reduction from $C \setminus D$ to $\mathcal{N}_b \setminus \mathcal{P}_b$. First assume $z \notin C$, that is $z(2i)$ does not tend to infinity when i goes to infinity. Fix ℓ such that $z(2i) = \ell$ for infinitely many i . We easily have that $f(z) \notin \mathcal{N}_b$. For example, if the digit 0 occurs with approximately the right frequency $\frac{1}{b}$ in $f(z) \upharpoonright [0, b^{k_{i-1}} + |B_i^1|) = [0, b^{k_i} (1 - \frac{1}{z(2i)} - \frac{1}{z(2i+1)})]$, then 0 will occur with too large a frequency in

$$f(z) \upharpoonright \left[1, b^{k_{i-1}} + |B_i^1| + |B_i^2| \right) = [0, b^{k_{i-1}} + |B_i^1| + \frac{1}{\ell} b^{k_i}).$$

We use here that $\frac{1}{b^{k_i}} \sum_{k < i} b_k \rightarrow 0$. This is because $f(z) \upharpoonright B_2^i = 0$ and $|B_2^i| = \frac{1}{\ell} b^{k_i}$ for such i .

Now assume that $z \in C$, so $\frac{1}{b^{k_i}} |B_i^2| = \frac{1}{z(2i)} \rightarrow 0$. Then, we have $f(z) \in \mathcal{N}_b$. This follows from the the definition of f and the fact that by that x is Borel normal to base b , see [5, Theorem 2].

Assume first that $z \in D$, so $z \notin C \setminus D$. We show $f(z) \in \mathcal{P}_b$, and so $f(z) \notin \mathcal{N}_b \setminus \mathcal{P}_b$. Since we are assuming $z \in C$ also, we have $\lim_{i \rightarrow \infty} z(i) = \infty$. So, $\lim_{i \rightarrow \infty} \frac{1}{b^{k_i}} (|B_i^2| + |B_i^3|) = 0$. It then follows exactly as in the proof of Theorem 3 that $f(z) \in \mathcal{P}_b$.

Assume next that $z \notin D$ (but $z \in C$ still). We show that $f(z) \notin \mathcal{P}_b$, which shows $f(z) \in \mathcal{N}_b \setminus \mathcal{P}_b$. Fix m so that for infinitely many i we have $z(2i+1) = m$, and m is of the form $m = 2^\ell$. Recall $\frac{1}{b^{k_i}} |B_i^3| = \frac{1}{z(2i+1)} = \frac{1}{2^\ell}$ for such i . We restrict our attention to this set of i in the following argument: If $f(z)$ were Poisson generic, then from Lemma 1 we would have that for large enough i in our set that

$$\frac{1}{b^{k_i}} |H_i| = (1 - e^{-\alpha})(e^{-(1-\alpha)}),$$

where H_i is the set of words of length k_i which occur in $f(z)$ with a starting position in $[(1-\alpha)b^{k_i}, b^{k_i})$, but do not occur in x with a starting position in $[b^{k_{i-1}}, (1-\alpha)b^{k_i})$. However, by the construction of $f(z)$ we have that every word which occurs in $[(1-\alpha)b^{k_i}, b^{k_i})$ also occurs in $[b^{k_{i-1}}, (1-\alpha)b^{k_i})$, and so $|H_i| = 0$. This completes the proof of Theorem 4. \square

Acknowledgements. The authors extend their appreciation to an anonymous referee for their thorough review. V. Becher is supported by grant PICT 2018-2315 of Agencia Nacional de Promoción Científica y Tecnológica de Argentina. S. Jackson is supported by NSF grant DMS-1800323. W. Mance is supported by grant 2019/34/E/ST1/00082 for the project ‘‘Set theoretic methods in dynamics and number theory,’’ NCN (The National Science Centre of Poland). D. Kwietniak is supported by NCN (the National Science Centre, Poland) Preludium Bis project no. 2019/35/O/ST1/02266.

REFERENCES

- [1] Dylan Airey, Steve Jackson, Dominik Kwietniak, and Bill Mance. Borel complexity of sets of normal numbers via generic points in subshifts with specification. *Trans. Amer. Math. Soc.*, 373(7):4561–4584, 2020.
- [2] Dylan Airey, Steve Jackson, and Bill Mance. Descriptive complexity of sets of normal numbers for the Cantor series expansions. *J. Symb. Log.*, 74(1):170–198, 2022.
- [3] Nicolás Álvarez and Verónica Becher and Martín Mereb. Poisson generic sequences. *International Mathematics Research Notices*, rnac234, 2022.
- [4] Verónica Becher, Pablo Ariel Heiber, and Theodore Slaman. Normal numbers and the Borel hierarchy. *Fundamenta Mathematicae*, 226(1):63–78, 2014.
- [5] Verónica Becher and Gabriel Sac Himelfarb. Construction of a λ -Poisson generic sequence. *Mathematics of Computation*, 92:1453–1466, 2023.
- [6] Verónica Becher and Theodore Slaman. On the normality of numbers to different bases. *Journal of the London Mathematical Society*, 90(2):472–494, 2014.
- [7] Verónica Becher, Santiago Figueira, and Rafael Picchi. Turing’s unpublished algorithm for normal numbers. *Theoretical Computer Science*, 377(1):126–138, 2007.
- [8] Konrad Deka, Steve Jackson, Dominik Kwietniak, and Bill Mance. Borel complexity of sets of points with prescribed Birkhoff averages in Polish dynamical systems with a specification property. arXiv:2210.08937.
- [9] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [10] Haseo Ki and Tom Linton. Normal numbers and subsets of \mathbb{N} with given densities. *Fundamenta Mathematica*, 144(2):163–179, 1994.
- [11] Benjamin Weiss. Poisson generic points, 23-27 November 2020. Jean-Morlet Chair conference on Diophantine Problems, Determinism and Randomness. Centre International de Rencontres Mathématiques. Audio-visual resource: doi:10.24350/CIRM.V.19690103.

Verónica Becher

Departamento de Computación, Facultad de Ciencias Exactas y Naturales & ICC
 Universidad de Buenos Aires & CONICET Argentina
 vbecher@dc.uba.ar

Stephen Jackson

Department of Mathematics
 University of North Texas, General Academics Building 435
 1155 Union Circle, #311430, Denton, TX 76203-5017, USA
 Stephen.Jackson@unt.edu

Dominik Kwietniak

Faculty of Mathematics and Computer Science
 Jagiellonian University in Kraków, ul. Łojasiewicza 6, 30-348 Kraków, Poland
 dominik.kwietniak@uj.edu.pl

Bill Mance

Faculty of Mathematics and Computer Science
 Adam Mickiewicz University, ul. Umultowska 87, 61-614 Poznań, Poland
 william.mance@amu.edu.pl