arXiv:2401.06917v2 [quant-ph] 14 Sep 2024

Bipartite representations and many-body entanglement of pure states of N indistinguishable particles

J.A. Cianciulli,¹ R. Rossignoli,^{1, 2} M. Di Tullio,¹ N. Gigena,¹ and F. Petrovich¹

¹IFLP/CONICET-Departamento de Física, Universidad Nacional de La Plata, C.C. 67, La Plata (1900), Argentina

² Comisión de Investigaciones Científicas (CIC), La Plata (1900), Argentina

We analyze a general bipartite-like representation of arbitrary pure states of N indistinguishable particles, valid for both bosons and fermions, based on M- and (N-M)-particle states. It leads to exact (M, N-M) Schmidt-like expansions of the state for any M < N and is directly related to the isospectral reduced M- and (N-M)-body density matrices $\rho^{(M)}$ and $\rho^{(N-M)}$. The formalism also allows for reduced yet still exact Schmidt-like decompositions associated with blocks of these densities, in systems having a fixed fraction of the particles in some single particle subspace. Monotonicity of the ensuing M-body entanglement under a certain set of quantum operations is also discussed. Illustrative examples in fermionic and bosonic systems with pairing correlations are provided, which show that in the presence of dominant eigenvalues in $\rho^{(M)}$, approximations based on a few terms of the pertinent Schmidt expansion can provide a reliable description of the state. The associated one- and two-body entanglement spectrum and entropies are also analyzed.

I. INTRODUCTION

Quantum entanglement and particle indistinguishability are undoubtedly among the most fundamental features of quantum mechanics. Yet the extension of the concept of entanglement to systems of indistinguishable particles is not straightforward [1]. The standard theory of entanglement [2, 3] was originally devised for systems of distinguishable components, where the pertinent Hilbert space has a tensor product structure which plays an essential role already in the basic definition: separable states, i.e. those that can be generated by local operations and classical communication, are just product states (or convex mixtures of product states in the mixed case), all remaining states being entangled.

In systems of indistinguishable components all states are, however, necessarily symmetrized (bosons) or antisymmetrized (fermions), preventing in principle a direct extension of previous scheme. The definition of entanglement in these systems has then followed different approaches, starting from mode entanglement [4–9], where each side has access to different orthogonal (and hence distinguishable) single particle (sp) modes and entanglement is then defined in the standard form, although it becomes thus dependent on the choice of sp modes. A distinct approach is the so-called *particle entanglement* or entanglement beyond symmetrization [10–21], which is independent of the choice of basis and just basic independent particle states (i.e., Slater determinants (SDs) for fermions) are nonentangled. Other proposals based on correlations between observables or measurements have also been discussed [22–26], including the consideration of symmetrization correlations as entanglement [27–31]. Connections between these distinct forms of entanglement and their behavior in different contexts have been analyzed by several authors [1, 32-43].

In particular, in [17] we focused on the one-body entanglement in fermion systems, which is determined by the one-body reduced density matrix (DM) $\rho^{(1)}$ and vanishes just for SDs (or quasiparticle vacua when extended to states with no fixed particle number). It also represents the *minimum* mode entanglement associated with a sp basis [17] and is connected to the minimum bipartite mode entanglement in four-level systems [35]. In [20] we examined its interpretation as a quantum resource [44, 45] in fermion systems, showing through a general majorization relation that it cannot decrease under a class of sp measurements, and also identified its relation with a bipartite-like (1, N-1) representation of a general N-fermion state. In [21] we extended the previous scheme to general *M*-body entanglement in *N*-fermion states (M < N), determined by the *M*-body DM $\rho^{(M)}$. Interest on many-body DMs and their relation with entanglement and characterization of correlations have recently increased in different areas [32, 46–49].

The aim of this work is first to extend the formalism of [21] to the bosonic case, developing a unified secondquantized formalism valid for both bosons and fermions. The formalism still conserves, nonetheless, some of the features of the standard distinguishable case: Any pure state of N indistinguishable particles is shown to admit. for $1 \leq M \leq N-1$, a bipartite-like (M, N-M) representation and Schmidt-like decomposition, whose coefficients are independent of the choice of sp basis and are just the square roots of the eigenvalues of the reduced DMs $\rho^{(M)}$ and $\rho^{(N-M)}$, isospectral in any pure state. The ensuing (M, N-M) entanglement determined by the mixedness of these eigenvalues is shown, through a general majorization relation, to be nonincreasing under a certain set of L-particle operations (and to stay invariant under unitary sp transformations), thus opening the way to a basic mode-independent (M, N-M) quantum resource theory. Besides, for M > 1 the *M*-body DMs may exhibit a few large *dominant* eigenvalues in correlated states, enabling a reliable approximation of the state based on just a few terms of the associated Schmidt expansion.

In addition, we also consider here the case of states having a fixed fraction of the total number of particles within some sp subspace. This is a common situation, arising e.g. in eigenstates of interacting Hamiltonians conserving the number of particles within certain sp subspaces, like a set S and a partner set \overline{S} of time-reversed sp states in pairing-type Hamiltonians or Hubbard models, as discussed in sections III and IV. It also emerges, of course, in any bosonic or fermionic entangled state having fixed number N_i of particles in orthogonal subsets S_i of modes, like e.g. distinct sites. In such cases the M-body DMs exhibit a blocked structure, and it will be shown that reduced but still exact (M, N-M) bipartitelike expansions and Schmidt decompositions associated with each of these blocks are also possible. Moreover, the standard distinguishable bipartite case naturally emerges here as a particular instance in this general formulation.

The general formalism is presented in section II, while the special case of blocked DMs and reduced exact expansions are treated in section III. Illustrative examples in finite systems with pairing correlations are provided in section IV for both fermions and bosons. They include analytical results and bounds for the one and two-body entanglement spectrum in some typical paired states, as well as numerical results for the latter and the associated entanglement entropy in the exact GS of a finite pairing Hamiltonian. These results show the presence of a characteristic large dominant eigenvalue in the twobody DM of both fermionic and bosonic paired states, together with a highly mixed one-body DM, which provides a clear signature of such states. It is also shown that through such eigenvalue and the associated eigenvector, a good approximation to the exact GS of the previous Hamiltonian for *all* values of the coupling strength can be here achieved with just very few terms of the pertinent (2, N-2) Schmidt expansion. Conclusions are finally provided in section V.

II. FORMALISM

A. *N* particle states in boson and fermion systems

Let us consider a single particle (sp) space \mathcal{H} of finite dimension d and a set of particle creation and annihilation operators c_i^{\dagger} , c_i , $i = 1, \ldots, d$, satisfying

$$[c_i, c_j]_{\pm} = [c_i^{\dagger}, c_j^{\dagger}]_{\pm} = 0, \ [c_i, c_j^{\dagger}]_{\pm} = \delta_{ij}, \qquad (1)$$

for bosons (-) or fermions (+), where $[a, b]_{\pm} = ab \pm ba$. We define the *M*-particle creation operators

$$C_{\alpha}^{(M)\dagger} = \frac{c_1^{\dagger n_1}}{\sqrt{n_1!}} \frac{c_2^{\dagger n_2}}{\sqrt{n_2!}} \dots \frac{c_d^{\dagger n_d}}{\sqrt{n_d!}}, \quad \sum_{i=1}^d n_i = M, \qquad (2)$$

where $n_i = 0, 1, 2, ...$ for bosons and $n_i = 0, 1$ for fermions, while $\boldsymbol{\alpha} = (n_1, ..., n_d)$. When applied to the vacuum $|0\rangle$, they create normalized orthogonal *M*particle states $C_{\boldsymbol{\alpha}}^{(M)\dagger}|0\rangle$ having n_i particles in sp state *i*. For $M \geq 0$ (and $M \leq d$ for fermions) these states span the full Fock space \mathcal{F} associated with \mathcal{H} , satisfying

$$\langle 0|C_{\alpha}^{(M)}C_{\alpha'}^{(M')\dagger}|0\rangle = \delta^{MM'}\delta_{\alpha\alpha'}.$$
 (3)

The subspace \mathcal{F}_M of M-particle states has dimension $d_M = \binom{d+M-1}{M}$ for bosons and $d_M = \binom{d}{M}$ for fermions, and is generated by the d_M operators (2). These operators satisfy $C^{(M)}_{\boldsymbol{\alpha}} C^{(M)\dagger}_{\boldsymbol{\alpha}'} |0\rangle = \delta_{\boldsymbol{\alpha}\boldsymbol{\alpha}'} |0\rangle$ and (see App. A)

$$\sum_{\alpha} C_{\alpha}^{(M)\dagger} C_{\alpha}^{(M)} = \binom{N}{M}, \qquad (4)$$

for both fermions and bosons, where $\hat{N} = \sum_{i} c_{i}^{\dagger} c_{i}$ is the particle number operator and $\binom{\hat{N}}{M}$ the operator taking the value $\binom{N}{M} = \frac{N!}{M!(N-M)!}$ when applied to an N-particle state: $\binom{\hat{N}}{M}|\Psi\rangle = \binom{N}{M}|\Psi\rangle$ if $\hat{N}|\Psi\rangle = N|\Psi\rangle$. The sum over $\boldsymbol{\alpha}$ in (4) runs over all d_{M} operators (2), i.e. over all possible occupations (n_{1}, \ldots, n_{d}) with $\sum_{i} n_{i} = M$. For instance, $\sum_{\boldsymbol{\alpha}} C_{\boldsymbol{\alpha}}^{(2)\dagger} C_{\boldsymbol{\alpha}}^{(2)} = \sum_{i < j} c_{i}^{\dagger} c_{j}^{\dagger} c_{j} c_{i} + \sum_{i} \frac{c_{i}^{\dagger^{2}}}{\sqrt{2}} \frac{c_{i}^{2}}{\sqrt{2}} = \frac{1}{2} \sum_{i,j} c_{i}^{\dagger} c_{j}^{\dagger} c_{j} c_{i} = \frac{\hat{N}(\hat{N}-1)}{2} = \binom{\hat{N}}{2}$ for both bosons, and fermions (where i = j terms obviously vanish).

An arbitrary normalized pure state $|\Psi\rangle$ of N identical particles (bosons or fermions) can then be written as

$$|\Psi\rangle = \frac{1}{N!} \sum_{i_1,\dots,i_N} \Gamma_{i_1\dots i_N} c_{i_1}^{\dagger} \dots c_{i_N}^{\dagger} |0\rangle \qquad (5a)$$

$$= \sum_{\alpha} \Gamma_{\alpha}^{(N)} C_{\alpha}^{(N)\dagger} |0\rangle , \qquad (5b)$$

where $\Gamma_{i_1,...,i_N}$ is a fully symmetric (antisymmetric) tensor for bosons (fermions) and the sum over each i_j in (5a) runs over all d sp states, whereas that in (5b) over all distinct d_N operators (2), with (see App. A)

$$\Gamma_{\alpha}^{(N)} = \langle 0 | C_{\alpha}^{(N)} | \Psi \rangle = \frac{\Gamma_{i_1 \dots i_N}}{\sqrt{n_1! \dots \sqrt{n_d!}}}, \qquad (6)$$

for $c_{i_1}^{\dagger} \dots c_{i_N}^{\dagger} = c_1^{\dagger n_1} \dots c_d^{\dagger n_d}$ (and $i_1 < \dots < i_N$ for fermions). Here $|\Gamma_{\boldsymbol{\alpha}}^{(N)}|^2$ is the probability of finding the N particle state $C_{\boldsymbol{\alpha}}^{(N)\dagger}|0\rangle$ "occupied" in $|\Psi\rangle$, with $\langle \Psi|\Psi\rangle = \sum_{\boldsymbol{\alpha}} |\Gamma_{\boldsymbol{\alpha}}^{(N)}|^2 = \frac{1}{N!} \sum_{i_1,\dots,i_d} |\Gamma_{i_1\dots i_N}|^2 = 1.$

B. The (M, N - M) representation and Schmidt decomposition for bosons and fermions

We can rewrite the general N-particle state (5) in a bipartite-like form involving operators creating $M \leq N$ and N-M particles, such that side A refers to M particles (but not to any specific location in space or any other quantum number) and side B to N-M particles. Starting from Eq. (4) we obtain $\sum_{\alpha} C_{\alpha}^{(M)\dagger} C_{\alpha}^{(M)} |\Psi\rangle = {N \choose M} |\Psi\rangle$.

Then $|\Psi\rangle = \binom{N}{M}^{-1} \sum_{\alpha} C_{\alpha}^{(M)\dagger} C_{\alpha}^{(M)} |\Psi\rangle$ can be recast in the bipartite-like form

$$|\Psi\rangle = {\binom{N}{M}}^{-1} \sum_{\alpha,\beta} \Gamma^{(M)}_{\alpha\beta} C^{(M)\dagger}_{\alpha} C^{(N-M)\dagger}_{\beta} |0\rangle, \qquad (7)$$

for both bosons and fermions, where we have written

$$C_{\alpha}^{(M)}|\Psi\rangle = \sum_{\beta} \Gamma_{\alpha\beta}^{(M)} C_{\beta}^{(N-M)\dagger}|0\rangle, \qquad (8)$$

and sums over $\boldsymbol{\alpha}, \boldsymbol{\beta}$ run over all d_M and d_{N-M} operators $C^{(M)\dagger}_{\boldsymbol{\alpha}}, C^{(N-M)\dagger}_{\boldsymbol{\beta}}$ respectively. Here $\Gamma^{(M)}_{\boldsymbol{\alpha}\boldsymbol{\beta}} \equiv \Gamma^{(M,N-M)}_{\boldsymbol{\alpha}\boldsymbol{\beta}}$ is given by (see Eq. (3))

$$\Gamma_{\boldsymbol{\alpha}\boldsymbol{\beta}}^{(M)} = \langle 0|C_{\boldsymbol{\beta}}^{(N-M)}C_{\boldsymbol{\alpha}}^{(M)}|\Psi\rangle, \qquad (9)$$

and is directly related to $\Gamma_{i_1,...,i_n}$ in (5a) by Eq. (A5).

Eq. (7) is the (M, N-M) bipartite-like decomposition of $|\Psi\rangle$, expressing it as a linear combination of "products" of states in \mathcal{F}_M and \mathcal{F}_{N-M} . The coefficients $\Gamma_{\alpha\beta}^{(M)}$ determine the remnant N-M particle state (8) after annihilating in $|\Psi\rangle M$ particles in the state labeled by α , with $|\Gamma_{\alpha\beta}^{(M)}|^2$ proportional to the probability of having M particles in the state α and N-M in the state β . Eqs. (7) and (8) imply $\langle \Psi|\Psi\rangle = {N \choose M}^{-1} \sum_{\alpha,\beta} |\Gamma_{\alpha\beta}^{(M)}|^2$, such that for any normalized state,

$$\operatorname{Tr}\left[\Gamma^{(M)^{\dagger}}\Gamma^{(M)}\right] = \binom{N}{M},\qquad(10)$$

for both bosons and fermions.

As done for fermions [21], from the singular value decomposition (SVD) of the matrix $\Gamma^{(M)}$,

$$\Gamma^{(M)}_{\boldsymbol{\alpha\beta}} = \sum_{\nu} U^{(M)}_{\boldsymbol{\alpha\nu}} \sigma^{(M)}_{\nu} V^{(N-M)\dagger}_{\boldsymbol{\nu\beta}}, \qquad (11)$$

where $U^{(M)}, V^{(N-M)}$ are unitary $d_M \times d_M$ and $d_{N-M} \times d_{N-M}$ matrices and $\sigma_{\nu}^{(M)} > 0$ the singular values of $\Gamma^{(M)}$ (square roots of the nonzero eigenvalues of $\Gamma^{(M)\dagger}\Gamma^{(M)}$ or $\Gamma^{(M)}\Gamma^{(M)\dagger}$), we obtain from (7) the Schmidt-like diagonal (M, N-M) decomposition of a general bosonic or fermionic N-particle state,

$$|\Psi\rangle = {\binom{N}{M}}^{-1} \sum_{\nu=1}^{n_M} \sigma_{\nu}^{(M)} A_{\nu}^{(M)\dagger} B_{\nu}^{(N-M)\dagger} |0\rangle, \qquad (12)$$

where n_M is the rank of $\Gamma^{(M)}$ and

$$A_{\nu}^{(M)\dagger} = \sum_{\alpha} U_{\alpha\nu}^{(M)} C_{\alpha}^{(M)\dagger} ,$$

$$B_{\nu}^{(N-M)\dagger} = \sum_{\beta} V_{\beta\nu}^{(N-M)*} C_{\beta}^{(N-M)\dagger} ,$$
 (13)

are "collective" operators creating M and N-M particles. As $U^{(M)}$ and $V^{(N-M)}$ are unitary, they are again

orthogonal normalized operators satisfying

$$A_{\nu}^{(M)} A_{\nu'}^{(M)\dagger} |0\rangle = \delta_{\nu\nu'} |0\rangle = B_{\nu}^{(N-M)} B_{\nu'}^{(N-M)\dagger} |0\rangle, \quad (14a)$$
$$\sum_{\nu} A_{\nu}^{(M)\dagger} A_{\nu}^{(M)} = {\binom{\hat{N}}{M}} = \sum_{\nu} B_{\nu}^{(N-M)\dagger} B_{\nu}^{(N-M)}, \quad (14b)$$

for both bosons and fermions. Moreover, Eqs. (8)–(9) become diagonal in terms of these normal operators:

$$A_{\nu}^{(M)}|\Psi\rangle = \sigma_{\nu}^{(M)} B_{\nu}^{(N-M)\dagger}|0\rangle, \qquad (15a)$$

$$\langle 0|B_{\nu}^{(N-M)}A_{\nu'}^{(M)}|\Psi\rangle = \delta_{\nu\nu'}\,\sigma_{\nu}^{(M)}\,, \tag{15b}$$

such that $B_{\nu}^{(N-M)\dagger}|0\rangle$ is the state of remaining N-M particles after destroying M particles in the state labeled by ν . These states are orthogonal according to Eq. (14a), in analogy with the standard Schmidt decomposition and in contrast with the states (8) (see Eq. (16a)). On the other hand, the full terms $A_{\nu}^{(M)\dagger}B_{\nu}^{(N-M)\dagger}|0\rangle$ are not necessarily orthogonal for different ν .

The singular values $\sigma_{\nu}^{(M)}$ in (12) are characteristic of the state, i.e. independent of the choice of sp basis used to represent it (see (A6) in App. A). For N = 2 Eq. (12) becomes equivalent to the normal forms of refs. [10, 11].

C. The *M*-body density matrix and operator

The bipartite tensor $\Gamma_{\alpha\beta}^{(M)}$ is directly connected to the *M*-body density matrix $\rho^{(M)}$, of elements [50, 51]

$$\rho_{\alpha\alpha'}^{(M)} := \langle \Psi | C_{\alpha'}^{(M)\dagger} C_{\alpha}^{(M)} | \Psi \rangle$$
(16a)

$$= \sum_{\boldsymbol{\beta}} \Gamma^{(M)}_{\boldsymbol{\alpha}\boldsymbol{\beta}} \Gamma^{(M)*}_{\boldsymbol{\alpha}'\boldsymbol{\beta}} = (\Gamma^{(M)}\Gamma^{(M)\dagger})_{\boldsymbol{\alpha}\boldsymbol{\alpha}'}, \quad (16b)$$

i.e. $\rho^{(M)} = \Gamma^{(M)}\Gamma^{(M)\dagger}$ for both bosons and fermions, where in (16b) we used Eqs. (3)-(8). Here $\rho^{(M)}$ is a $d_M \times d_M$ positive semidefinite matrix, representing the hermitian covariance matrix of the linearly independent operators $C^{(M)}_{\alpha}$ in the state $|\Psi\rangle$. Eq. (10) implies

$$\operatorname{Tr} \rho^{(M)} = \begin{pmatrix} N \\ M \end{pmatrix}, \qquad (17)$$

for both bosons and fermions, in agreement with Eq. (4) [52]. The average of any bosonic or fermionic *M*-body operator can then be expressed as

$$\left\langle \sum_{\boldsymbol{\alpha},\boldsymbol{\alpha}'} O_{\boldsymbol{\alpha}\boldsymbol{\alpha}'}^{(M)} C_{\boldsymbol{\alpha}}^{(M)\dagger} C_{\boldsymbol{\alpha}'}^{(M)} \right\rangle = \operatorname{Tr}\left[\rho^{(M)} O^{(M)}\right].$$
(18)

Since Eq. (7) implies $\Gamma^{(N-M)} = (\pm 1)^{M(N-M)} \Gamma^{(M)t}$ for bosons (+) or fermions (-), the partner DM $\rho_{\beta\beta'}^{(N-M)} = \Gamma^{(N-M)} \Gamma^{(N-M)\dagger}$ is just $\Gamma^{(M)t} \Gamma^{(M)*}$.

From (11) we note that the squared singular values

$$\lambda_{\nu}^{(M)} = (\sigma_{\nu}^{(M)})^2 \,, \tag{19}$$

arising from the Schmidt decomposition (12), are precisely the nonzero eigenvalues of $\Gamma^{(M)}\Gamma^{(M)^{\dagger}} = \rho^{(M)}$ or equivalently $\Gamma^{(M)t}\Gamma^{(M)*} = \rho^{(N-M)}$, i.e., of the M and N-M-body DMs, which then have the same non-zero eigenvalues in any N-particle pure state $|\Psi\rangle$, for both bosons and fermions [53], with $U^{(M)}$ and $V^{(N-M)*}$ the corresponding eigenvector matrices. This result is analogous to that of the distinguishable bipartite case [2].

Moreover, the normal operators (13) are precisely those which diagonalize $\rho^{(M)}$ and $\rho^{(N-M)}$, constituting the M and (N-M)-body "natural orbitals":

$$\langle \Psi | A_{\nu'}^{(M)\dagger} A_{\nu}^{(M)} | \Psi \rangle = \lambda_{\nu}^{(M)} \delta_{\nu\nu'}$$
(20)
= $\langle \Psi | B_{\nu'}^{(N-M)\dagger} B_{\nu}^{(N-M)} | \Psi \rangle ,$

as follows from (14)–(15).

For M = N, $\rho^{(N)}$ has just a single eigenvalue $\lambda_1^{(N)} = 1$ corresponding to the operator $A_1^{(N)\dagger} = \sum_{\alpha} \Gamma_{\alpha}^{(N)} C_{\alpha}^{(N)\dagger}$ creating the state (5b), whereas for M = 1 we recover the one-body DM $\rho_{ij}^{(1)} = \langle \Psi | c_j^{\dagger} c_i | \Psi \rangle = (\Gamma^{(1)} \Gamma^{(1)^{\dagger}})_{ij}$, with $\operatorname{Tr} \rho^{(1)} = N$, isospectral with $\rho^{(N-1)}$ [20]. In this case $A_{\nu}^{\dagger} = c_{\nu}^{\dagger}$ are the standard sp "natural" creation operators diagonalizing $\rho^{(1)}$: $\langle c_{\nu'}^{\dagger} c_{\nu} \rangle = \delta_{\nu\nu'} \lambda_{\nu}^{(1)}$. We also mention that in an N-particle Fock state

We also mention that in an *N*-particle Fock state $|\Psi_{\beta}\rangle = C_{\beta}^{(N)\dagger}|0\rangle = \frac{c_1^{\dagger n_1}}{\sqrt{n_1!}} \dots \frac{c_d^{\dagger n_d}}{\sqrt{n_d}}|0\rangle \ (\beta = (n_1, \dots, n_d), \sum_i n_i = N)$, i.e. a permanent (bosons) or SD (fermions), $\rho^{(M)}$ is diagonal in the standard basis of operators $C_{\alpha}^{(M)\dagger} = \frac{c_1^{\dagger m_1}}{\sqrt{m_1!}} \dots \frac{c_d^{\dagger m_d}}{\sqrt{m_d!}} \ (\alpha = (m_1, \dots, m_d), \sum_i m_i = M)$, having just integer eigenvalues:

$$\langle C_{\boldsymbol{\alpha}}^{(M)\dagger}C_{\boldsymbol{\alpha}'}^{(M)}\rangle_{\boldsymbol{\beta}} = \delta_{\boldsymbol{\alpha}\boldsymbol{\alpha}'}\lambda_{\boldsymbol{\alpha}}^{(M)}, \quad \lambda_{\boldsymbol{\alpha}}^{(M)} = \prod_{i} \begin{pmatrix} n_{i} \\ m_{i} \end{pmatrix}, \quad (21)$$

which in the fermionic case reduce just to $\binom{N}{M}$ eigenvalues $\lambda_{\alpha}^{(M)} = 1$ [21]. In both cases they verify $\sum_{\alpha} \lambda_{\alpha}^{(M)} = {N \choose M}$. In the bosonic case the lowest rank obviously corresponds to a condensate (e.g. $n_i = N\delta_{i1}$) where there is a single nonzero eigenvalue $\lambda_1^{(M)} = {N \choose M}$. We can also define the *M*-body density operator (DO)

$$\hat{\rho}^{(M)} = \sum_{\beta} C_{\beta}^{(N-M)} |\Psi\rangle \langle \Psi | C_{\beta}^{(N-M)\dagger}$$
(22a)

$$= \sum_{\boldsymbol{\alpha},\boldsymbol{\alpha}'} \rho_{\boldsymbol{\alpha}\boldsymbol{\alpha}'}^{(M)} C_{\boldsymbol{\alpha}}^{(M)\dagger} |0\rangle \langle 0| C_{\boldsymbol{\alpha}'}^{(M)}, \qquad (22b)$$

where in (22b) we used Eq. (8) for $M \to N - M$ and (16b). It is the unique mixed *M*-particle state fulfilling

$$\operatorname{Tr}\left[\hat{\rho}^{(M)}C^{(M)\dagger}_{\boldsymbol{\alpha}'}C^{(M)}_{\boldsymbol{\alpha}}\right] = \rho^{(M)}_{\boldsymbol{\alpha}\boldsymbol{\alpha}'}$$
(23)

 $\forall \alpha, \alpha'$, for both bosons or fermions. Its diagonal form is

$$\hat{\rho}^{(M)} = \sum_{\nu} \lambda_{\nu}^{(M)} A_{\nu}^{(M)\dagger} |0\rangle \langle 0| A_{\nu}^{(M)} , \qquad (24)$$

in terms of the normal operators (13), such that $\hat{\rho}^{(M)}A^{(M)\dagger}_{\nu}|0\rangle = \lambda^{(M)}_{\nu}A^{(M)\dagger}_{\nu}|0\rangle$, having obviously the same eigenvalues as its matrix representation $\rho^{(M)}$. For M = N it reduces to $\hat{\rho}^{(N)} = |\Psi\rangle\langle\Psi|$.

Thus, the operation in (22a) can be seen as a partial trace over N - M particles, leading to the reduced state $\hat{\rho}^{(M)}$ of Eq. (22b) which determines the average of any Mbody (and hence L-body for L < M, see (29)) operator, in analogy with the standard distinguishable case.

Under unitary sp transformations of the state, $|\Psi\rangle \rightarrow \hat{U}|\Psi\rangle$ with $\hat{U} = e^{-i\sum_{ij}h_{ij}c_i^{\dagger}c_j}$, all $\rho^{(M)}$ and $\hat{\rho}^{(M)}$ will transform unitarily (see (A7)).

Measurements, M-body DM's and M-body D. entanglement

For both bosons and fermions, the operators

$$\mathcal{M}_{\boldsymbol{\beta}} := \binom{N}{M}^{-1/2} C_{\boldsymbol{\beta}}^{(N-M)} , \qquad (25)$$

can be considered as Kraus operators when acting on the subspace of states with definite particle number N, since by Eq. (4), they satisfy

$$\sum_{\beta} \mathcal{M}_{\beta}^{\dagger} \mathcal{M}_{\beta} = \mathbb{1}_{N}, \qquad (26)$$

i.e. $\sum_{\beta} \mathcal{M}_{\beta}^{\dagger} \mathcal{M}_{\beta} |\Psi\rangle = |\Psi\rangle \forall N$ -particle state $|\Psi\rangle$. Then they define a measurement on N-particle states in which N - M particles are annihilated (see also (B7) in App. B for a number conserving implementation): $|\Psi\rangle \rightarrow$ $\mathcal{M}_{\beta}|\Psi\rangle \propto C_{\beta}^{(N-M)}|\Psi\rangle$, with probabilities

$$p_{\boldsymbol{\beta}} = \langle \Psi | \mathcal{M}_{\boldsymbol{\beta}}^{\dagger} \mathcal{M}_{\boldsymbol{\beta}} | \Psi \rangle = \rho_{\boldsymbol{\beta}\boldsymbol{\beta}}^{(N-M)} / {N \choose M}, \qquad (27)$$

determined precisely by the (N - M)-body DM.

And according to Eqs. (22a)-(25), the ensuing postmeasurement state $\sum_{\beta} \mathcal{M}_{\beta} |\Psi\rangle \langle \Psi | \mathcal{M}_{\beta}^{\dagger}$ (without postselection) is just the normalized M-body DO $\hat{\rho}_n^{(M)}$:

$$\hat{\rho}_{n}^{(M)} := {\binom{N}{M}}^{-1} \hat{\rho}^{(M)} = \sum_{\beta} \mathcal{M}_{\beta} |\Psi\rangle \langle \Psi | \mathcal{M}_{\beta}^{\dagger}, \quad (28)$$

which satisfies $\operatorname{Tr} \hat{\rho}_n^{(M)} = 1$. The basic case is that of sp measurements $(N - M = 1, \mathcal{M}_i = c_i/\sqrt{N})$, with N - Mbody measurements based on the operators (25) being just compositions of sp measurements.

Regarding the L-body DM $\rho^{(L)}$ for $L \leq M$, we first note that using Eqs. (22a) and (4), it can be obtained from $\hat{\rho}^{(M)}$ as (see App. A)

$$\rho_{\boldsymbol{\gamma}\boldsymbol{\gamma}'}^{(L)} = {\binom{N-L}{N-M}}^{-1} \operatorname{Tr} \left[\hat{\rho}^{(M)} C_{\boldsymbol{\gamma}'}^{(L)\dagger} C_{\boldsymbol{\gamma}}^{(L)} \right] \quad (L \le M).$$
(29)

Then the expansions (22a) or (28) imply (see App. A)

$$\hat{\rho}_n^{(L)} = \sum_{\beta} p_{\beta} \, \hat{\rho}_{\beta n}^{(L)} \,, \tag{30}$$

where $\hat{\rho}_n^{(L)} = \hat{\rho}^{(L)}/{N \choose L}$ is the normalized *L*-body DO in the original state $|\Psi\rangle$ and $\hat{\rho}_{\beta n}^{(L)} = \hat{\rho}_{\beta}^{(L)}/{M \choose L}$ those in the postselected normalized *M*-particle states $|\Psi_{\beta}\rangle = \mathcal{M}_{\beta}|\Psi\rangle/\sqrt{p_{\beta}}$, with p_{β} the probabilities (27) of outcome β . Thus, for any $L \leq M$, the normalized DM $\rho_n^{(L)}$ is the average of the normalized post-measurement DMs $\hat{\rho}_{\beta n}^{(L)}$ in the post selected states.

This result is important since it implies the general majorization relation [21]

$$\boldsymbol{\lambda}(\rho_n^{(L)}) \prec \sum_{\boldsymbol{\beta}} p_{\boldsymbol{\beta}} \boldsymbol{\lambda}(\hat{\rho}_{\boldsymbol{\beta}n}^{(L)})$$
(31)

between the sorted (in decreasing order) eigenvalue spectrum $\boldsymbol{\lambda}$ of the original DM $\rho_n^{(L)}$ and those of $\hat{\rho}_{\beta n}^{(L)}$ (see e.g. [54, 55] for majorization properties). It entails the general entropic inequality

$$S(\hat{\rho}_{n}^{(L)}) \geq \sum_{\beta} p_{\beta} S(\hat{\rho}_{\beta n}^{(L)}), \qquad (32)$$

between the entropy of the original normalized *L*-body DM $\rho_n^{(L)}$ and the average entropy of the normalized *L*-body DMs in the post-measurement states. It is valid for any concave entropy $S(\rho)$, like the von Neumann entropy $S(\rho) = -\text{Tr} \rho \log_2 \rho$, or in general any trace-form entropy $S_f(\rho) = \text{Tr} f(\rho)$ with f concave and f(0) = f(1) = 0 [56], in both boson and fermion systems.

This result means that for $L \leq M$, the *L*-body entanglement, determined by the mixedness of the normalized *L*-body DM [21] and quantified by the associated entropy

$$E^{(L)}(|\Psi\rangle) = S(\hat{\rho}_n^{(L)}) = S(\hat{\rho}_n^{(N-L)}), \qquad (33)$$

cannot increase (and will typically decrease) on average under the N-M-body operation determined by the operators (25), for both bosons and fermions, and for any choice of entropy. This is in agreement with the fact that such measurement decreases the uncertainty about which L-body states are occupied.

Eqs. (26)–(33) remain valid for measurements based on the normal operators (13), $\tilde{\mathcal{M}}_{\nu} = {\binom{N}{M}}^{-\frac{1}{2}} B_{\nu}^{(N-M)}$, as they are unitarily related to the \mathcal{M}_{β} : just replace $\beta \rightarrow \nu$, with $p_{\beta} \rightarrow p_{\nu} = \lambda_{\nu}^{(N-M)} / {\binom{N}{M}}$ and $|\Psi_{\beta}\rangle \rightarrow$ $|\Psi_{\nu}\rangle = A_{\nu}^{(M)\dagger}|0\rangle (\propto B_{\nu}^{(N-M)}|\Psi\rangle)$ in (30)–(32).

Hence, a resource theory based on the previous quantum operations (plus free operations like unitary sp transformations) is in principle feasible, with the *L*-body entanglement, determined by the mixedness of the reduced densities $\hat{\rho}_n^{(L)}$ (which is fully independent of the choice of sp modes) as a basic resource which cannot increase on average under these operations (see also App. B).

III. REDUCED EXACT DECOMPOSITIONS

The presence of symmetries in $|\Psi\rangle$ can simplify the DMs $\rho^{(M)}$ into a blocked structure in an obvious basis,

reducing the effective number of nonzero elements. A common example is the conservation of the number of particles in a certain subspace $S \subset \mathcal{H}$ of sp modes,

$$\hat{N}_{\mathcal{S}} = \sum_{i \in \mathcal{S}} c_i^{\dagger} c_i \,, \tag{34}$$

as in the case of eigenstates $|\Psi\rangle$ of Hamiltonians satisfying $[H, \hat{N}_{S}] = 0$, such that

$$\hat{N}_{\mathcal{S}}|\Psi\rangle = N_{\mathcal{S}}|\Psi\rangle. \tag{35}$$

If $|\Psi\rangle$ has definite particle number N, (35) also implies $\hat{N}_{\bar{S}}|\Psi\rangle = N_{\bar{S}}|\Psi\rangle$ for $N_{\bar{S}} = N - N_{\bar{S}}$ the number of particles in the orthogonal complement \bar{S} , such that $\mathcal{H} = S \oplus \bar{S}$.

Common well-known cases are e.g. systems with pairing-type two-body couplings, where the number of particles in positive and negative quasimomentum states N_S , $N_{\bar{S}}$, are conserved (see sec. IV), Hubbard-type Hamiltonians in solid state physics [46, 57, 58], which conserve the number of particles $N_{\sigma} = \sum_i c_{i\sigma}^{\dagger} c_{i\sigma}$ with definite spin component σ , and the strong nuclear force in an atomic nucleus within the isospin formalism [50], which conserves $T_z = \frac{N-Z}{2}$, i.e., the number of neutrons $N = \sum_k c_{k+}^{\dagger} c_{k+}$ and protons $Z = \sum_k c_{k-}^{\dagger} c_{k-}$ for N + Zfixed and k denoting remaining quantum numbers. It is also the case of any system with fixed number of particles at distinct sites (corresponding to orthogonal sp subspaces S_i), like entangled states of spatially separated Mand N - M particles, where the standard distinguishable scenario emerges naturally as a special case (sec. III D).

Eq. (35) implies that elements of $\rho^{(M)}$ which do not conserve the number of particles in \mathcal{S} will vanish: $\langle C^{(M)\dagger}_{\boldsymbol{\alpha}} C^{(M)}_{\boldsymbol{\alpha}'} \rangle = 0$ if $[C^{(M)\dagger}_{\boldsymbol{\alpha}} C^{(M)}_{\boldsymbol{\alpha}'}, \hat{N}_{\mathcal{S}}] \neq 0$, leading to a blocked $\rho^{(M)}$ where each block corresponds to a fixed number m of operators $c^{\dagger}_{i\in\mathcal{S}}, c_{i'\in\mathcal{S}}$ in $C^{(M)\dagger}_{\boldsymbol{\alpha}}, C^{(M)}_{\boldsymbol{\alpha}'}$. We show here that reduced exact (M, N-M) expansions of $|\Psi\rangle$ associated with each of these blocks are also feasible.

A. One-body case

We start with the simplest case M = 1. It is easily seen that Eq. (35) implies the following blocked form of the one-body DM $\rho^{(1)}$,

$$\rho^{(1)} = \begin{pmatrix} \rho_{\mathcal{S}}^{(1)} & 0\\ 0 & \rho_{\bar{\mathcal{S}}}^{(1)} \end{pmatrix}, \qquad (36)$$

where $(\rho_{\mathcal{S}}^{(1)})_{ij} = \langle c_j^{\dagger} c_i \rangle$, $(\rho_{\bar{\mathcal{S}}}^{(1)})_{ij} = \langle c_{\bar{j}}^{\dagger} c_{\bar{i}} \rangle$ are the onebody DM's in each subspace, since remaining contractions $\langle c_i^{\dagger} c_{\bar{j}} \rangle$ vanish due to the conservation of $\hat{N}_{\mathcal{S}}$. These blocks have a fixed trace $\operatorname{Tr} \rho_{\mathcal{S}}^{(1)} = N_{\mathcal{S}}$, $\operatorname{Tr} \rho_{\bar{\mathcal{S}}}^{(1)} = N_{\bar{\mathcal{S}}}$. Moreover, if $N_{\mathcal{S}} \geq 1$, $N_{\bar{\mathcal{S}}} \geq 1$, each block can be associ-

Moreover, if $N_{\mathcal{S}} \geq 1$, $N_{\bar{\mathcal{S}}} \geq 1$, each block can be associated to an own (1, N-1) expansion and Schmidt decomposition of $|\Psi\rangle$: Starting from (35), $|\Psi\rangle = \frac{1}{N_{\mathcal{S}}}\hat{N}_{\mathcal{S}}|\Psi\rangle$,

and writing $c_i |\Psi\rangle = \sum_{\beta} \Gamma_{i\beta}^{(1)} C_{\beta}^{(N-1)\dagger} |0\rangle$ with $\Gamma_{i\beta}^{(1)} =$ $\langle 0|C_{\boldsymbol{\beta}}^{(N-1)}c_i|\Psi\rangle$, we obtain the reduced exact expansion

$$|\Psi\rangle = N_{\mathcal{S}}^{-1} \sum_{i \in \mathcal{S}, \beta} \Gamma_{i\beta}^{(1)} c_i^{\dagger} C_{\beta}^{(N-1)\dagger} |0\rangle \qquad (37a)$$

$$= N_{S}^{-1} \sum_{\nu \in S} \sigma_{\nu}^{(1)} a_{\nu}^{\dagger} B_{\nu}^{(N-1)\dagger} |0\rangle , \qquad (37b)$$

which just involves the block $\Gamma_{\mathcal{S}}^{(1)}$ of elements $\Gamma_{i\in\mathcal{S},\mathcal{G}}^{(1)}$ of the full $\Gamma^{(1)}$ in (7), where β spans (N-1)-particle states with $N_{\mathcal{S}} - 1$ particles in \mathcal{S} and $N_{\bar{\mathcal{S}}}$ in $\bar{\mathcal{S}}$. In (37b), $\sigma_{\nu}^{(1)}$ are the singular values of $\Gamma_{\mathcal{S}}^{(1)}$ and $a_{\nu} = \sum_{i \in \mathcal{S}} U_{i\nu}^{(1)} c_i^{\dagger}$, $B_{\nu}^{\dagger} = \sum_{\beta} V_{\beta\nu}^{(N-1)*} C_{\beta}^{(N-1)\dagger}$ the associated normal operators (13) (M = 1). It leads to $\langle c_i^{\dagger} c_i \rangle = (\Gamma_{\mathcal{S}}^{(1)} \Gamma_{\mathcal{S}}^{(1)\dagger})_{ij}$ for $i, j \in \mathcal{S}$ and hence to the upper block of $\rho^{(1)}$,

$$\rho_{\mathcal{S}}^{(1)} = \Gamma_{\mathcal{S}}^{(1)} \Gamma_{\mathcal{S}}^{(1)\dagger} , \qquad (38)$$

with eigenvalues $\lambda_{\nu}^{(1)} = (\sigma_{\nu}^{(1)})^2$ for $\nu \in \mathcal{S}$.

Similar expressions with $S \to \bar{S}$ in (37)-(38) obviously hold for the expansion of $|\Psi\rangle$ associated to the second block $\rho_{\bar{S}}^{(1)}$ in (36), determined by $\Gamma_{\bar{S}}^{(1)}$ of elements $\Gamma_{\bar{i}\in\bar{S},\beta}^{(1)}$. Both expansions, Eq. (37) and the analogous one based on \overline{S} , are exact but run in general over distinct singular values $\sigma_{\nu}^{(1)}$, $\sigma_{\bar{\nu}}^{(1)}$ and operators a_{ν} , $a_{\bar{\nu}}$. For composite systems with $N_{S_i} \geq 1$ particles in *n* orthogonal subspaces \mathcal{S}_i , analogous expansions hold for each subspace.

Eq. (35) also leads to a similar blocked structure of the partner isospectral (N-1)-body DM,

$$\rho^{(N-1)} = \begin{pmatrix} \rho_{\mathcal{S}}^{(N-1)} & 0\\ 0 & \rho_{\bar{\mathcal{S}}}^{(N-1)} \end{pmatrix}$$
(39)

where $\rho_{\mathcal{S}}^{(N-1)} = \Gamma_{\mathcal{S}}^{(1)T} \Gamma_{\mathcal{S}}^{(1)*}$ contains just those elements $\rho_{\beta'\beta}^{(N-1)} \text{ with } \beta, \beta' \text{ involving } N_{\mathcal{S}} - 1 \text{ particles in } \mathcal{S}, \text{ and} \\ \rho_{\bar{\mathcal{S}}}^{(N-1)} = \Gamma_{\bar{\mathcal{S}}}^{(1)T} \Gamma_{\bar{\mathcal{S}}}^{(1)*} \text{ those with } N_{\bar{\mathcal{S}}} - 1 \text{ particles in } \bar{\mathcal{S}}$ (and hence $N_{\mathcal{S}}$ in \mathcal{S}). Notice, however, that $\rho_{\mathcal{S}}^{(1)}$ and $\rho_{\bar{\mathcal{S}}}^{(1)}$ are not isospectral in general.

В. Two-body case

The implications of (35) become even more important for the two-body DM. For $N_{\mathcal{S}} \geq 2$, $N_{\bar{\mathcal{S}}} \geq 2$, Eq. (35) entails that the full $\rho^{(2)}$ will have three principal blocks:

$$\rho^{(2)} = \begin{pmatrix} \rho_{\mathcal{S}}^{(2)} & 0 & 0\\ 0 & \rho_{\mathcal{S}\bar{\mathcal{S}}}^{(2)} & 0\\ 0 & 0 & \rho_{\bar{\mathcal{S}}}^{(2)} \end{pmatrix}, \qquad (40)$$

where

$$(\rho_{\mathcal{S}}^{(2)})_{ij,kl} = \langle c_k^{\dagger} c_l^{\dagger} c_j c_i \rangle, \quad (\rho_{\bar{\mathcal{S}}}^{(2)})_{\bar{i}\bar{j},\bar{k}\bar{l}} = \langle c_{\bar{k}}^{\dagger} c_{\bar{l}}^{\dagger} c_{\bar{j}} c_{\bar{i}} \rangle, \quad (41)$$

contain the contractions within S and \overline{S} respectively (including diagonal elements $(\rho_{S}^{(2)})_{ii,kl} = \langle c_{k}^{\dagger} c_{l}^{\dagger} c_{i}^{2} \rangle / \sqrt{2}$, $(\rho_{\mathcal{S}}^{(2)})_{ii,kk} = \langle c_k^{\dagger 2} c_i^2 \rangle/2$, etc. in the bosonic case) and

$$(\rho_{\mathcal{S}\bar{\mathcal{S}}}^{(2)})_{i\bar{j},k\bar{l}} = \langle c_k^{\dagger} c_{\bar{l}}^{\dagger} c_{\bar{j}} c_i \rangle, \qquad (42)$$

those involving one particle in \mathcal{S} and one in $\overline{\mathcal{S}}$. All remaining contractions vanish due to the conserved $N_{\mathcal{S}}$. These blocks have fixed traces

$$\operatorname{Tr} \rho_{\mathcal{S}}^{(2)} = {N_{\mathcal{S}} \choose 2}, \ \operatorname{Tr} \rho_{\bar{\mathcal{S}}}^{(2)} = {N_{\bar{\mathcal{S}}} \choose 2}, \ \operatorname{Tr} \rho_{\mathcal{S}\bar{\mathcal{S}}}^{(2)} = N_{\mathcal{S}} N_{\bar{\mathcal{S}}},$$
(43)

verifying $\binom{N_{\mathcal{S}}}{2} + \binom{N_{\mathcal{S}}}{2} + N_{\mathcal{S}}N_{\mathcal{S}} = \binom{N_{\mathcal{S}}+N_{\mathcal{S}}}{2}$. Moreover, Eq. (35) implies

$$\sum_{\boldsymbol{\alpha}\in\mathcal{S}} C_{\boldsymbol{\alpha}}^{(2)\dagger} C_{\boldsymbol{\alpha}}^{(2)} |\Psi\rangle = \binom{N_{\mathcal{S}}}{2} |\Psi\rangle, \qquad (44)$$

for $C^{(2)\dagger}_{\alpha\in\mathcal{S}} \equiv c_i^{\dagger}c_j^{\dagger}$ (i < j) or $(c_i^{\dagger})^2/2$, with a similar expression for $\mathcal{S} \to \overline{\mathcal{S}}$, and also

$$\sum_{\boldsymbol{\gamma}\in\mathcal{S}\bar{\mathcal{S}}} C_{\boldsymbol{\gamma}}^{(2)\dagger} C_{\boldsymbol{\gamma}}^{(2)} |\Psi\rangle = N_{\mathcal{S}} N_{\bar{\mathcal{S}}} |\Psi\rangle \,. \tag{45}$$

for $C^{(2)\dagger}_{\gamma \in S\bar{S}} \equiv c_i^{\dagger} c_{\bar{i}}^{\dagger}$. Then the tensor $\Gamma^{(2)}$ will also have three corresponding blocks $\Gamma_{\mathcal{S}}^{(2)}$, $\Gamma_{\mathcal{S}\bar{\mathcal{S}}}^{(2)}$ and $\Gamma_{\bar{\mathcal{S}}}^{(2)}$, each of them generating an *exact* reduced expansion and Schmidt decomposition of $|\Psi\rangle$: The first one, emerging from (44),

$$|\Psi\rangle = {\binom{N_{\mathcal{S}}}{2}}^{-1} \sum_{\boldsymbol{\alpha} \in S, \boldsymbol{\beta}} (\Gamma_{\mathcal{S}}^{(2)})_{\boldsymbol{\alpha}\boldsymbol{\beta}} C_{\boldsymbol{\alpha}}^{(2)\dagger} C_{\boldsymbol{\beta}}^{(N-2)\dagger} |0\rangle$$
(46a)

$$= \binom{N_{\mathcal{S}}}{2}^{-1} \sum_{\nu \in \mathcal{S}} \sigma_{\nu}^{(2)} A_{\nu}^{(2)\dagger} B_{\nu}^{(N-2)\dagger} |0\rangle, \qquad (46b)$$

where $(\Gamma_{\mathcal{S}}^{(2)})_{\alpha\beta} = \langle 0|C_{\beta}^{(N-2)}C_{\alpha\in\mathcal{S}}^{(2)}|\Psi\rangle$ and $\sigma_{\nu}^{(2)}$ are the singular values of $\Gamma_{\mathcal{S}}^{(2)}$, is related to $\rho_{\mathcal{S}}^{(2)} = \Gamma_{\mathcal{S}}^{(2)}\Gamma_{\mathcal{S}}^{(2)\dagger}$ and its eigenvalues $\lambda_{\nu}^{(2)} = (\sigma_{\nu}^{(2)})^2$ for $\nu \in \mathcal{S}$. The third one is analogous for $S \to \bar{S}$ and is related to $\rho_{\bar{S}}^{(2)} = \Gamma_{\bar{S}}^{(2)} \Gamma_{\bar{S}}^{(2)\dagger}$. Finally, the second one, emerging from (45),

$$|\Psi\rangle = \frac{1}{N_{\mathcal{S}}N_{\mathcal{S}}} \sum_{\gamma \in \mathcal{S}\bar{\mathcal{S}}, \beta} (\Gamma_{\mathcal{S}\bar{\mathcal{S}}}^{(2)})_{\gamma\beta} C_{\gamma}^{(2)\dagger} C_{\beta}^{(N-2)\dagger} |0\rangle \quad (47a)$$

$$= \frac{1}{N_{\mathcal{S}}N_{\bar{\mathcal{S}}}} \sum_{\tilde{\nu}} \sigma_{\tilde{\nu}}^{(2)} A_{\tilde{\nu}}^{(2)\dagger} B_{\tilde{\nu}}^{(N-2)\dagger} |0\rangle, \qquad (47b)$$

where $(\Gamma^{(2)}_{S\bar{S}})_{\gamma\beta} = \langle 0|C^{(N-2)}_{\beta}C^{(2)}_{\gamma\in S\bar{S}}|\Psi\rangle$ and $\sigma^{(2)}_{\tilde{\nu}}$ are the singular values of $\Gamma_{S\bar{S}}^{(2)}$, determines the central block $\rho_{S\bar{S}}^{(2)} = \Gamma_{S\bar{S}}^{(2)}\Gamma_{S\bar{S}}^{(2)\dagger}$ and its eigenvalues $\lambda_{\tilde{\nu}}^{(2)} = (\sigma_{\tilde{\nu}}^{(2)})^2$. It exposes the two-body correlations between the particles in \mathcal{S} and those in $\overline{\mathcal{S}}$. Summing the three previous expansions with their relative weights $p_{\mathcal{S}} = \binom{N_{\mathcal{S}}}{2} / \binom{N}{2}$, $p_{\bar{S}} = {N_{\bar{S}} \choose 2} / {N \choose 2}$ and $p_{S\bar{S}} = N_S N_{\bar{S}} / {N \choose 2}$ leads to the original expansion (7) for M = 2.

Previous blocked structure and expansions also hold for the partner (N-2)-body DM $\rho^{(N-2)}$, with $\rho_{S}^{(N-2)} = \Gamma_{S\bar{S}}^{(2)T}\Gamma_{S\bar{S}}^{(2)*}$, $\rho_{S\bar{S}}^{(N-2)} = \Gamma_{S\bar{S}}^{(2)T}\Gamma_{S\bar{S}}^{(2)*}$ and $\rho_{\bar{S}}^{(N-2)} = \Gamma_{\bar{S}}^{(2)T}\Gamma_{\bar{S}}^{(2)*}$ containing elements involving $N_{S} - 2$, $N_{S} - 1$ and N_{S} particles in S respectively.

The expansion (47) is convenient when $\rho_{S\bar{S}}^{(2)}$ possesses one or a few large dominant eigenvalues $\lambda_{\bar{\nu}}^{(2)} > 1$ which absorb most of the sum (see sec. IV), and which reflect pairing-like correlations between particles in S and those in \bar{S} [48, 59, 60]. For any "product" state

$$|\Psi\rangle = A_{\mathcal{S}}^{(N_{\mathcal{S}})\dagger} B_{\bar{\mathcal{S}}}^{(N_{\bar{\mathcal{S}}})\dagger} |0\rangle, \qquad (48a)$$

where $A_{S}^{(N_{S})\dagger}(B_{\bar{S}}^{(N_{\bar{S}})\dagger})$ creates an arbitrary state of N_{S} $(N_{\bar{S}})$ particles in $S(\bar{S})$, $\rho_{S\bar{S}}^{(2)}$ becomes a direct product of one-body densities in both fermion and boson systems,

$$(\rho_{S\bar{S}}^{(2)})_{i\bar{j},k\bar{l}} = (\rho_{S}^{(1)})_{ik} (\rho_{\bar{S}}^{(1)})_{\bar{j}\bar{l}}, \qquad (48b)$$

becoming diagonal in the natural sp bases which diagonalize $\rho_{\mathcal{S}}^{(1)}$ and $\rho_{\mathcal{S}}^{(1)}$: $(\rho_{\mathcal{S}\overline{\mathcal{S}}}^{(2)})_{i\overline{j},k\overline{l}} = \delta_{ik}\delta_{\overline{j}\overline{l}}\lambda_i^{(1)}\lambda_{\overline{j}}^{(1)}$, with $\lambda_i^{(1)}\lambda_{\overline{j}}^{(1)} \leq 1$ for fermions.

Hence an immediate consequence for fermions is the following: If in a state with definite fermion numbers $N_{\mathcal{S}}$, $N_{\bar{\mathcal{S}}}$ in orthogonal sp subspaces \mathcal{S} , $\bar{\mathcal{S}}$, the joint twobody DM $\rho_{S\bar{\mathcal{S}}}^{(2)}$ has an eigenvalue $\lambda_{\nu}^{(2)} > 1$, there is *bipartite entanglement between the* $N_{\mathcal{S}}$ and $N_{\bar{\mathcal{S}}}$ fermions, in the sense of not being a product state (48a) (see III D).

Such dominant eigenvalue also indicates that $|\Psi\rangle$ cannot be an independent fermion state (SD) either, since in these states all nonzero eigenvalues of $\rho^{(2)}$ have the value 1. Moreover, if $|\Psi\rangle$ is a SD $((\rho^{(1)})^2 = \rho^{(1)})$ and $N_S, N_{\bar{S}}$ are conserved, then (48a) necessarily holds, with $A_S^{(N_S)\dagger} = C_{\alpha\in S}^{(N_{\bar{S}})\dagger}, B_{\bar{S}}^{(N_{\bar{S}})\dagger} = C_{\beta\in \bar{S}}^{(N_{\bar{S}})\dagger}$ simple product operators of the form (2), since the blocked structure (36) then implies $(\rho_S^{(1)})^2 = \rho_S^{(1)}, (\rho_{\bar{S}}^{(1)})^2 = \rho_{\bar{S}}^{(1)}$, entailing that both $A_S^{(N_S)\dagger}|0\rangle, B_S^{(N_S)\dagger}|0\rangle$ are SDs.

C. General M-body case

For a general *M*-body DM in a system with $N_{\mathcal{S}} \geq M$ particles in \mathcal{S} and $N_{\bar{\mathcal{S}}} \geq M$ in $\bar{\mathcal{S}}$, Eq. (35) implies that $\rho^{(M)}$ will be blocked into M + 1 subdensities $\rho^{(m,l)}_{\mathcal{S}\bar{\mathcal{S}}}$ involving *m* particles in \mathcal{S} and l = M - m particles in $\bar{\mathcal{S}}$, with $m = 0, 1, \ldots, M$:

$$\rho^{(M)} = \begin{pmatrix}
\rho_{\mathcal{S}}^{(M,0)} & 0 & \dots & 0 \\
0 & \rho_{\mathcal{S}\bar{\mathcal{S}}}^{(1,M-1)} & \dots & \vdots \\
\vdots & \dots & \ddots & \vdots \\
0 & \dots & 0 & \rho_{\bar{\mathcal{S}}}^{(0,M)}
\end{pmatrix}, \quad (49)$$

where $(\rho_{S\bar{S}}^{(m,l)})_{\alpha\alpha'} = \langle C_{\alpha'}^{(m,l)\dagger} C_{\alpha}^{(m,l)} \rangle$, with $C_{\alpha}^{(m,l)\dagger} = C_{\alpha\in S}^{(m)\dagger} C_{\bar{\alpha}\in\bar{S}}^{(l)\dagger}$ creating *m* particles in *S* and *l* in \bar{S} . The

blocked form (49) is equivalent to the condition

$$[\hat{\rho}^{(M)}, \hat{N}_{\mathcal{S}}] = 0 \tag{50}$$

on the *m*-body density operator (22), implied by a state fulfilling Eq. (35). The operators $C_{\alpha}^{(m,l)\dagger}$ satisfy

$$\sum_{\alpha} C_{\alpha}^{(m,l)\dagger} C_{\alpha}^{(m,l)} = {\binom{\hat{N}_{\mathcal{S}}}{m}} {\binom{\hat{N}_{\mathcal{S}}}{l}}.$$
 (51)

Therefore, each block in (49) has a definite trace $\operatorname{Tr} \rho^{(l,m)} = {N_{\mathcal{S}} \choose m} {N_{\mathcal{S}} \choose l}$. For M = 1 and 2 we recover the blocks of Eqs. (36) and (40).

The tensor $\Gamma^{(M)}$ will also be decomposed into M + 1blocks $\Gamma^{(m,l)}$, each one generating an exact expansion and Schmidt-like decomposition of $|\Psi\rangle$, since for each m, lEq. (51) implies $\sum_{\alpha} C_{\alpha}^{(m,l)\dagger} C_{\alpha}^{(m,l)} |\Psi\rangle = \binom{N_{\mathcal{S}}}{m} \binom{N_{\mathcal{S}}}{l} |\Psi\rangle$ for states satisfying (35). Thus, we obtain

$$\Psi \rangle = \frac{1}{\binom{N_{\mathcal{S}}}{m}\binom{N_{\mathcal{S}}}{l}} \sum_{\boldsymbol{\alpha},\boldsymbol{\beta}} \Gamma_{\boldsymbol{\alpha}\boldsymbol{\beta}}^{(m,l)} C_{\boldsymbol{\alpha}}^{(m,l)\dagger} C_{\boldsymbol{\beta}}^{(N_{\mathcal{S}}-m,N_{\mathcal{S}}-l)\dagger} |0\rangle, \quad (52a)$$
$$= \frac{1}{\binom{N_{\mathcal{S}}}{m}\binom{N_{\mathcal{S}}}{l}} \sum_{\nu} \sigma_{\nu}^{(m,l)} A_{\nu}^{(m,l)\dagger} B_{\nu}^{(N_{\mathcal{S}}-m,N_{\mathcal{S}}-l)\dagger} |0\rangle, \quad (52b)$$

with $\Gamma_{\alpha\beta}^{(m,l)} = \langle 0|C_{\beta}^{(N_{S}-m,N_{S}-l)}C_{\alpha}^{(m,l)}|\Psi\rangle$, such that $C_{\alpha}^{(m,l)}|\Psi\rangle = \sum_{\beta}\Gamma_{\alpha\beta}^{(m,l)}C^{(N_{S}-m,N_{S}-l)\dagger}|0\rangle$ and $\rho^{(m,l)} = \Gamma^{(m,l)}\Gamma^{(m,l)\dagger}$. In (52b) $\sigma_{\nu}^{(m,l)}$ are the singular values of $\Gamma^{(m,l)}$ and $A_{\nu}^{(m,l)\dagger}$, $B_{\nu}^{(N_{S}-m,N_{S}-l)\dagger}$ the normal operators obtained from its SVD. These block-DMs and its eigenvalues can be used to characterize the *M*-body correlations between particles at S and \bar{S} .

The (m, l)-body density operator corresponding to such block can be likewise obtained as

$$\hat{\rho}^{(m,l)} = \sum_{\beta} C^{(N_{\mathcal{S}}-m,N_{\bar{\mathcal{S}}}-l)}_{\beta} |\Psi\rangle \langle\Psi| C^{(N_{\mathcal{S}}-m,N_{\bar{\mathcal{S}}}-l)\dagger}_{\beta}$$
$$= \sum_{\alpha,\alpha'} \rho^{(m,l)}_{\alpha\alpha'} C^{(m,l)\dagger}_{\alpha} |0\rangle \langle 0| C^{(m,l)}_{\alpha'} .$$
(53)

Similarly, reduced measurements based on the operators

$$\mathcal{M}_{\boldsymbol{\alpha}}^{(m,l)} = \left[\binom{N_{\mathcal{S}}}{m} \binom{N_{\bar{\mathcal{S}}}}{l} \right]^{-1/2} C_{\boldsymbol{\alpha}}^{(m,l)} , \qquad (54)$$

which satisfy $\sum_{\alpha} \mathcal{M}_{\alpha}^{(m,l)\dagger} \mathcal{M}_{\alpha}^{(m,l)} = \mathbb{1}_{N_S N_S}$ on the subspace of states with definite particle number N and subsystem particle number N_S , become feasible, leading to a direct extension of previous Eqs. (27)–(33). The associated entanglement, i.e. the mixedness of each of these blocks, will not increase on average under these measurements, and can then be also considered as a resource in this scenario.

D. Connection with entanglement between distinguishable systems

In the special case $M = N_S$, the expansion (52) associated to the first block (m = M, l = 0) in (49) becomes

$$|\Psi\rangle = \sum_{\boldsymbol{\alpha}\in\mathcal{S},\boldsymbol{\beta}\in\bar{\mathcal{S}}} \Gamma_{\boldsymbol{\alpha}\boldsymbol{\beta}}^{(N_{\mathcal{S}})} C_{\boldsymbol{\alpha}}^{(N_{\mathcal{S}})\dagger} C_{\boldsymbol{\beta}}^{(N_{\mathcal{S}})\dagger} |0\rangle \qquad (55a)$$

$$= \sum_{\nu} \sigma_{\nu}^{(N_{\mathcal{S}})} A_{\nu}^{(N_{\mathcal{S}})\dagger} B_{\nu}^{(N_{\mathcal{S}})\dagger} |0\rangle , \qquad (55b)$$

with $\Gamma_{\alpha\beta}^{(N_S)} \equiv \Gamma_{\alpha\beta}^{(N_S,N_{\bar{S}})} = \langle 0|C_{\beta}^{(N_S)}C_{\alpha}^{(N_S)}|\Psi\rangle$ for $\alpha \in S$, $\beta \in \bar{S}$. Eq. (55a) is just the standard decomposition of a bipartite state of two distinguishable systems (the N_S particles at S and the $N_{\bar{S}}$ at \bar{S}) in terms of local states $(C_{\alpha\in S}^{(N_S)\dagger}|0\rangle$ and $C_{\beta\in \bar{S}}^{(N_S)\dagger}|0\rangle$) expressed in second quantized form. The N_S particles at S can be distinguished from the $N_{\bar{S}}$ at \bar{S} since they occupy orthogonal subspaces and have then a distinct quantum number. The diagonal representation (55b) is the standard Schmidt decomposition, with Eq. (48a) corresponding to the separable case.

Accordingly, all terms in the sums (55a)-(55b) are now mutually orthogonal. The associated isospectral N_{S} - and $N_{\bar{S}}$ -body densities,

$$\rho_{\mathcal{S}}^{(N_{\mathcal{S}})} = \Gamma^{(N_{\mathcal{S}})} \Gamma^{(N_{\mathcal{S}})\dagger}, \ \rho_{\bar{\mathcal{S}}}^{(N_{\bar{\mathcal{S}}})} = \Gamma^{(N_{\mathcal{S}})t} \Gamma^{(N_{\mathcal{S}})*}, \quad (56)$$

where $\rho_{\mathcal{S}}^{(N_{\mathcal{S}})} = \rho_{\mathcal{S}}^{(m,0)}$ is the first block in (49), represent the standard local isospectral density matrices of the $N_{\mathcal{S}}$ particles at \mathcal{S} and those at $\bar{\mathcal{S}}$, having the same eigenvalues $\lambda_{\nu}^{(N_{\mathcal{S}})} = (\sigma_{\nu}^{(N_{\mathcal{S}})})^2$ and satisfying $\operatorname{Tr} \rho_{\mathcal{S}}^{(N_{\mathcal{S}})} = {N_{\mathcal{S}} \choose N_{\mathcal{S}}} = 1$, $\operatorname{Tr} \rho_{\bar{\mathcal{S}}}^{(N_{\bar{\mathcal{S}}})} = {N_{\bar{\mathcal{S}}} \choose N_{\bar{\mathcal{S}}}} = 1$, with

$$\hat{\rho}_{\mathcal{S}}^{(N_{\mathcal{S}})} = \sum_{\boldsymbol{\beta}\in\bar{\mathcal{S}}} C_{\boldsymbol{\beta}}^{(N_{\bar{\mathcal{S}}})} |\Psi\rangle \langle\Psi| C_{\boldsymbol{\beta}}^{(N_{\bar{\mathcal{S}}})\dagger}, \qquad (57a)$$

$$\hat{\rho}_{\bar{\mathcal{S}}}^{(N_{\bar{\mathcal{S}}})} = \sum_{\boldsymbol{\alpha}\in\mathcal{S}} C_{\boldsymbol{\alpha}}^{(N_{\mathcal{S}})} |\Psi\rangle \langle \Psi| C_{\boldsymbol{\alpha}}^{(N_{\mathcal{S}})\dagger}, \qquad (57b)$$

the corresponding local reduced states. Their common entropy is the standard bipartite entanglement entropy of the N_S and $N_{\bar{S}}$ particles:

$$E(\mathcal{S},\bar{\mathcal{S}}) = S(\hat{\rho}^{(N_{\mathcal{S}})}) = S(\hat{\rho}^{(N_{\bar{\mathcal{S}}})}).$$
(58)

We finally notice that for a quantum operation transforming an N particle state $|\Psi_0\rangle$ with support on a sp subspace $S_0 \equiv \bar{S}$, to a state $|\Psi\rangle$ with $M = N_S$ particles in a subspace S orthogonal to \bar{S} , and $N_{\bar{S}} = N - M$ particles in \bar{S} , satisfying then Eq. (35), the result derived in [21] for fermions also holds for bosons: The entropy of the original normalized M-body DM $\rho_{0n}^{(M)}$ in $|\Psi_0\rangle$ is an upper bound to the average bipartite entanglement (58) in the final states (see App. B).

IV. EXAMPLES

We now discuss some examples illustrating previous considerations in both boson and fermion systems. We focus on N-particle paired states, which arise as ground states (GS) of systems with attractive pairing interactions. The latter are well-known to be most relevant in several distinct contexts, from the standard BCS theory of superconductivity [61] and its extension for describing He³ superfluidity [62], to the description of pairing effects in nuclear systems and neutron stars [50, 63], including also ultracold quantum gases [64]. BCS-like pairing models for bosons have also been considered [65–67]. Such paired states are strongly correlated, requiring, as is well known, at least a particle number violating BCS [61] or Bogoliubov approach [50] for an approximate treatment at the mean field level. We will here consider exact results in finite N-particle systems, focusing on the eigenvalues of the one- and, specially, two-body DM and on the associated entropies and bipartite expansions, of some typical paired states, including the GS of a finite pairing model.

A. Maximally paired states in fermionic and bosonic systems

We start from the uniform pair creation operator

$$A^{\dagger} = \frac{1}{\sqrt{n}} \sum_{k=1}^{n} c_{k}^{\dagger} c_{\bar{k}}^{\dagger} , \qquad (59)$$

where k, \bar{k} label n orthogonal sp states belonging to orthogonal subspaces S and \bar{S} respectively (e.g. k, \bar{k} may label opposite quasimomentum states) and $c_k^{\dagger}, c_{\bar{k}}^{\dagger}$ can be either bosonic or fermionic creation operators. It creates a maximally entangled pair state $|\Psi_1\rangle = A^{\dagger}|0\rangle = \frac{1}{\sqrt{n}} \sum_k c_k^{\dagger} c_{\bar{k}}^{\dagger} |0\rangle$, both in the sense of leading to a maximally mixed one-body DM $\rho^{(1)} = 1/n$ for 2 particles in 2n levels, i.e. maximal one-body entanglement $E^{(1)}$ in Eq. (33) for L = 1, as well as maximum bipartite entanglement between the two particles (which occupy orthogonal subspaces), i.e. maximally mixed $\hat{\rho}^{(N_S)}$, $\hat{\rho}^{(N_S)}$ in (57) and hence maximal $E(S, \bar{S})$ in (58), for $N_S = N_{\bar{S}} = 1$.

The operator (59) fulfills the commutation relation (in what follows + corresponds to bosons, - to fermions)

$$[A, A^{\dagger}] = 1 \pm \hat{N}/n, \qquad (60)$$

where $\hat{N} = \sum_{k} c_{k}^{\dagger} c_{k} + c_{\bar{k}}^{\dagger} c_{\bar{k}} = \hat{N}_{S} + \hat{N}_{\bar{S}}$ is the total number operator. Using (60) it is straightforward to show that the ensuing normalized *m*-pair state created by $(A^{\dagger})^{m}$ is

$$|\Psi_m\rangle := \frac{1}{m!} \sqrt{\frac{n^m}{\mathcal{N}_m}} (A^{\dagger})^m |0\rangle$$
 (61a)

$$= \frac{1}{\sqrt{\mathcal{N}_m}} \sum_{\substack{m_1,\dots,m_n\\\sum_k m_k = m}} |m_1,\dots,m_n\rangle, \quad (61b)$$

where m_k is the number of pairs in states (k, \bar{k}) , with $m_k = 0, 1, 2, ...$ for bosons and $m_k = 0, 1$ for fermions. In (61b), $\mathcal{N}_m = \binom{n+m-1}{m}$ (bosons) or $\mathcal{N}_m = \binom{n}{m}$ (fermions) is the number of ways of distributing m indistinguishable pairs in n pair states (with single occupancy in the

fermion case and $m \ge 0$ for bosons, $0 \le m \le n$ for fermions), and

$$|m_1,\ldots,m_n\rangle = \prod_{k=1}^n \frac{(c_k^{\dagger} c_{\bar{k}}^{\dagger})^{m_k}}{m_k!} |0\rangle = s_{\alpha} C_{\alpha}^{(m)\dagger} C_{\bar{\alpha}}^{(m)\dagger} |0\rangle, (62)$$

are basic normalized *m*-pair states, with $C_{\alpha}^{(m)\dagger} = \prod_{k} \frac{(c_{k}^{\dagger})^{m_{k}}}{\sqrt{m_{k}!}}, C_{\overline{\alpha}}^{(m)\dagger} = \prod_{k} \frac{(c_{k}^{\dagger})^{m_{k}}}{\sqrt{m_{k}!}}$ for $\alpha = (m_{1}, \ldots, m_{n})$ and $s_{\alpha} = \pm$ a phase factor for the fermionic case.

Hence, the states (61) are just a uniform superposition of these \mathcal{N}_m basic *m*-pair states, satisfying Eq. (35),

$$\hat{N}_{\mathcal{S}}|\Psi_m\rangle = \hat{N}_{\bar{\mathcal{S}}}|\Psi_m\rangle = m|\Psi_m\rangle.$$
(63)

They arise as exact GS of simple pairing Hamiltonians in the strong coupling limit (see section IV C). From (60)-(61) it can be shown that

$$A^{\dagger}|\Psi_{m-1}\rangle = \sqrt{m(1\pm\frac{m-1}{n})}|\Psi_{m}\rangle, \qquad (64a)$$

$$A^{\dagger}A|\Psi_m\rangle = m(1\pm\frac{m-1}{n})|\Psi_m\rangle,$$
 (64b)

with $A|\Psi_m\rangle = \sqrt{m(1\pm\frac{m-1}{n})}|\Psi_{m-1}\rangle$, such that $A^{\dagger}A$ counts essentially the number *m* of pairs for $n \gg m$.

The states (61) have again maximum one-body entanglement (for fixed N = 2m particles) for both fermions and bosons: Eq. (63) implies $\rho^{(1)}$ has the blocked form (36), with $\rho_{\mathcal{S}}^{(1)}$, $\rho_{\mathcal{S}}^{(1)}$ diagonal and maximally mixed, as all sp states k, \bar{k} have the same occupation:

$$\rho_{\mathcal{S}}^{(1)} = \rho_{\bar{\mathcal{S}}}^{(1)} = \lambda^{(1)} \,\mathbb{1} \,, \quad \lambda^{(1)} = m/n \,, \tag{65}$$

i.e. $\langle c_k^{\dagger} c_{k'} \rangle = \langle c_{\bar{k}}^{\dagger} c_{\bar{k}'} \rangle = \delta_{kk'} \lambda^{(1)}$, verifying Tr $\rho_{\mathcal{S}}^{(1)} =$ Tr $\rho_{\mathcal{S}}^{(1)} = \frac{1}{2}$ Tr $\rho^{(1)} = m$ and leading to maximum $E^{(1)}$ in (33), i.e. $E^{(1)} = \log_2(2n)$ for the von Neumann entropy.

Similarly, the states (61) have also maximum bipartite entanglement $E(S, \bar{S})$ in Eq. (58), between the $N_S = m$ particles in S and the m ones in \bar{S} , for both fermions and bosons: Eqs. (61b)-(62) are already the Schmidt decomposition for such partition, $|\Psi_m\rangle = \frac{1}{\sqrt{N_m}} \sum_{\alpha} s_{\alpha} C_{\alpha}^{(m)\dagger} C_{\bar{\alpha}}^{(m)\dagger} |0\rangle$ with $|s_{\alpha}| = 1$, hence leading to maximally mixed reduced densities $\rho_S^{(m)} = \rho_{\bar{S}}^{(m)} = \mathcal{N}_m^{-1} \mathbb{1}$ and maximum entanglement entropy

$$E(\mathcal{S}, \bar{\mathcal{S}}) = \log_2 \mathcal{N}_m. \tag{66}$$

On the other hand, the two-body DM $\rho^{(2)}$ determined by the state (61) is not maximally mixed, as can be seen from its eigenvalues $\lambda_i^{(2)}$: Eq. (63) ensures it will have the blocked structure of Eq. (40), with still maximally mixed diagonal blocks $\rho_{\mathcal{S}}^{(2)}$, $\rho_{\bar{\mathcal{S}}}^{(2)}$ (of length $n(n \pm 1)/2$),

$$\rho_{\mathcal{S}}^{(2)} = \rho_{\mathcal{S}}^{(2)} = \lambda_2^{(2)} \mathbb{1} , \quad \lambda_2^{(2)} = \frac{m(m-1)}{n(n\pm 1)} , \qquad (67)$$

since $\langle c_k^{\dagger} c_{k''}^{\dagger} c_{k'''} c_{k'''} \rangle = \delta_{kk''} \delta_{k'k'''} \lambda_2^{(2)}$ for k < k', k'' < k''', and additionally $\langle \frac{c_k^{\dagger}^2}{\sqrt{2}} \frac{c_{k'}^2}{\sqrt{2}} \rangle = \delta_{kk'} \lambda_2^{(2)}$ for bosons (see App. A), with identical expressions in $\bar{\mathcal{S}}$, verifying $\operatorname{Tr} \rho_{\mathcal{S}}^{(2)} = \operatorname{Tr} \rho_{\bar{\mathcal{S}}}^{(2)} = {m \choose 2}$ for bosons and fermions.

However, the remaining block $\rho_{S\bar{S}}^{(2)}$ in (40), of length n^2 for bosons and fermions, becomes itself blocked in two submatrices (see Eq. (A14) in App. A),

$$\langle c_{k}^{\dagger} c_{\bar{k}'}^{\dagger} c_{\bar{k}'''} c_{k'''} \rangle = (1 - \delta_{kk'}) \delta_{kk''} \delta_{k'k'''} \lambda_{2}^{(2)}$$
(68a)
+ $\delta_{kk'} \delta_{k''k'''} [\frac{m(n \pm m)}{n(n \pm 1)} + \delta_{kk''} \lambda_{2}^{(2)}],$ (68b)

where the first block (68a) is diagonal and similar to (67), while the second block (68b), of length n, is nondiagonal and exposes the two-body pairing correlations between particles in S and \bar{S} . It has two distinct eigenvalues: one given again by $\lambda_2^{(2)}$, Eq. (67), n-1 degenerate in this sub-block, while the remaining one (see App. A),

$$\lambda_1^{(2)} = m \left(1 \pm \frac{m-1}{n} \right) ,$$
 (69)

is the single dominant nondegenerate eigenvalue of $\rho^{(2)}$, satisfying $\lambda_1^{(2)} \ge m$ for bosons and $\lambda_1^{(2)} \ge 1$ for fermions (with $\lambda_1^{(2)} > m$ for bosons if m > 1 and $\lambda_1^{(2)} > 1$ for fermions if 1 < m < n). It corresponds to the flat normal operator $A_1^{(2)\dagger} = A^{\dagger}$ of $\rho^{(2)}$, as $A^{\dagger}A|\Psi_m\rangle = \lambda_1^{(2)}|\Psi_m\rangle$ (Eq. (64b)) and hence $\langle A^{\dagger}A \rangle = \lambda_1^{(2)}$.

Thus, for $n \gg m$, $\lambda_1^{(2)} \approx m$ is essentially the number mof pairs while $\lambda_2^{(2)} \approx (m/n)^2$ becomes small, in agreement with the approximate bosonic interpretation of A^{\dagger} for $n \gg N$ (Eq. (60)), in which case the state (61a) can be seen as an m-boson condensate. Nonetheless, their exact values are required for fulfilling Eqs. (17)–(43): Tr $\rho^{(2)} =$ $\lambda_1^{(2)} + (n(2n \pm 1) - 1)\lambda_2^{(2)} = \binom{2m}{2}$, Tr $\rho_{S\bar{S}}^{(2)} = \lambda_1^{(2)} + (n^2 1)\lambda_2^{(2)} = m^2$, and are important when $m \sim n$ or m > n(bosonic case). In the fermionic case Eq. (69) is also the *largest* value the maximum eigenvalue of $\rho^{(2)}$ can reach among any state of N = 2m particles in a 2n-dimensional sp space [48, 59].

We can now verify expansions (37) and (46)–(47). Blocks $\rho_{S}^{(1)}$ and $\rho_{\bar{S}}^{(1)}$ generate similar (1, N - 1) uniform expansions (37b) of $|\Psi_{m}\rangle$: From (61a), $c_{k}(A^{\dagger})^{m}|0\rangle = \frac{m}{\sqrt{n}}c_{\bar{k}}^{\dagger}(A^{\dagger})^{(m-1)}|0\rangle$, such that (15a) is verified for M = 1, $A_{\nu}^{(1)} = c_{k}$: $c_{k}|\Psi_{m}\rangle = \sqrt{\lambda^{(1)}}B_{k}^{(2m-1)\dagger}|0\rangle$, with $B_{k}^{(2m-1)\dagger}|0\rangle = \frac{c_{\bar{k}}^{\dagger}}{\sqrt{1\pm\frac{m-1}{n}}}|\Psi_{m-1}\rangle$ the normalized state of remaining particles after one in sp state k is annihilated. Eq. (37b) is then fulfilled: $\frac{1}{m}\sqrt{\frac{m}{n}}\sum_{k}c_{k}^{\dagger}B_{k}^{(2m-1)\dagger}|0\rangle = \frac{A^{\dagger}|\Psi_{m-1}\rangle}{\sqrt{m(1\pm\frac{m-1}{n})}} = |\Psi_{m}\rangle$, according to (64a).

Likewise, blocks $\rho_{\mathcal{S}}^{(2)}$, $\rho_{\bar{\mathcal{S}}}^{(2)}$ generate similar uniform expansions (46) of $|\Psi_m\rangle$: for $k \neq k'$, $c_{k'}c_k(A^{\dagger})^m|0\rangle$ $= \frac{m(m-1)}{n}c_{\bar{k}'}^{\dagger}c_{\bar{k}}^{\dagger}(A^{\dagger})^{(m-2)}|0\rangle$ and hence $c_{k'}c_k|\Psi_m\rangle =$
$$\begin{split} &\sqrt{\lambda_2^{(2)}}B_{k'k}^{(2m-2)\dagger}|0\rangle, \text{ where } B_{k'k}^{(2m-2)\dagger}|0\rangle = \frac{c_{k'\ell}^{\dagger}c_{k}^{\dagger}|\Psi_{2m-2}\rangle}{\langle c_{k}c_{k'}c_{k}^{\dagger}c_{k}^{\dagger}\rangle^{1/2}} \text{ is the normalized state of remaining particles after annihilating two particles in states } k,k'. Eq. (46) is then verified: <math display="block">\frac{\sqrt{\lambda_2^{(2)}}}{m(m-1)}\sum_{k,k'}c_{k}^{\dagger}c_{k'}^{\dagger}B_{k'k}^{(2m-2)\dagger}|0\rangle = |\Psi_m\rangle. \end{split}$$

On the other hand the expansion (47) based on $\rho_{S\bar{S}}^{(2)}$ has here a single dominant term: Eq. (64a) is just (15a) for the main eigenvalue $\lambda_1^{(2)}$ and eigenvector $A_1^{(2)} = A$: $A|\Psi_m\rangle = \sqrt{\lambda_1^{(2)}}|\Psi_{m-1}\rangle$, with $B_1^{(N-2)\dagger} \propto (A^{\dagger})^{m-1}$. Moreover, the first term alone in (47b) is already proportional to the exact state, as $A^{\dagger}A|\Psi_m\rangle \propto |\Psi_m\rangle$ according to (64b). The sum of all remaining terms in (47b) is in this case proportional to this first term.

B. General paired states

Let us now consider the general m-pair state

$$|\Psi_m\rangle = \sum_{\alpha} \Gamma_{\alpha} C_{\alpha}^{(m)\dagger} C_{\bar{\alpha}}^{(m)\dagger} |0\rangle$$
 (70a)

$$= \sum_{\substack{m_1,\ldots,m_n\\\sum_k m_k=m}} \Gamma_{m_1\ldots m_n} | m_1,\ldots,m_n \rangle \quad (70b)$$

where $|m_1, \ldots, m_n\rangle$ are the previous states (62) and $\Gamma_{\alpha} = \Gamma_{m_1...m_n}$ arbitrary coefficients satisfying $\sum_{\alpha} |\Gamma_{\alpha}|^2 = 1$, with $m_k = 0, 1, 2, \ldots (0, 1)$ for bosons (fermions). Like (61), these states contain all N = 2m particles in mpairs (k, \bar{k}) and arise as GS of pairing Hamiltonians at finite couplings strengths (see sec. IV C). They satisfy Eqs. (35)-(63), then leading to the same blocked structure (36)-(40) of $\rho^{(1)}$ and $\rho^{(2)}$ for fermions and bosons, with $\rho^{(1)}$ and $\rho_{S}^{(2)}$, $\rho_{S}^{(2)}$ again diagonal in the standard basis:

$$\langle c_{k}^{\dagger}c_{k'}\rangle = \langle c_{\bar{k}}^{\dagger}c_{\bar{k}'}\rangle = \delta_{kk'}\lambda_{k}^{(1)}, \qquad (71a)$$

$$\langle c_k^{\dagger} c_{k'}^{\dagger} c_{k'''} c_{k'''} \rangle = \delta_{kk''} \delta_{k'k'''} \lambda_{kk'}^{(2)} , \qquad (71b)$$

for k < k', k'' < k''', and similarly for $k \to \bar{k}$, with $\lambda_{\bar{k}\bar{k}'}^{(2)} = \lambda_{kk'}^{(2)}$ and $\lambda_{kk}^{(2)} = \frac{1}{2} \langle c_k^{\dagger 2} c_k^2 \rangle = \lambda_{\bar{k}\bar{k}}^{(2)}$ for bosons. And for $\rho_{S\bar{S}}^{(2)}$, Eq. (68) is replaced by

$$\langle c_k^{\dagger} c_{\bar{k}^{\prime\prime}}^{\dagger} c_{\bar{k}^{\prime\prime\prime\prime}} c_{k^{\prime\prime\prime}} \rangle = (1 - \delta_{kk^{\prime\prime}}) \delta_{kk^{\prime\prime}} \delta_{k^{\prime}k^{\prime\prime\prime}} \lambda_{kk^{\prime}}^{(2)}$$
(72a)

$$+\delta_{kk'}\delta_{k''k'''}\rho_{c\,kk''}^{(2)} \tag{72b}$$

such that $\rho_{S\bar{S}}^{(2)} = \begin{pmatrix} \rho_d^{(2)} & 0 \\ 0 & \rho_c^{(2)} \end{pmatrix}$, with $\rho_d^{(2)}$ the diagonal subblock (72a) having the same elements (71b), and $\rho_c^{(2)}$ the nondiagonal $n \times n$ "collective" sub-block (72b) containing the two-body pairing contractions $\langle c_k^{\dagger} c_{\bar{k}}^{\dagger} c_{\bar{k}'} c_{k'} \rangle$.

In the *fermionic* case, this sub-block yields itself to an exact (2, N-2) reduced expansion of $|\Psi_m\rangle$ containing at most *n* terms, since $\hat{N}_p \equiv \sum_k c_k^{\dagger} c_k^{\dagger} c_k c_k$ just counts

for fermions the number of pairs, satisfying $\hat{N}_p |\Psi_m\rangle = m |\Psi_m\rangle$. Thus, $|\Psi_m\rangle = \frac{1}{m} \hat{N}_p |\Psi_m\rangle$ can be expanded as

$$|\Psi_m\rangle = \frac{1}{m} \sum_{k,\beta} \Gamma_{k\beta}^{(2)} c_k^{\dagger} c_{\bar{k}}^{\dagger} B_{\beta}^{(N-2)} |0\rangle$$
(73a)

$$= \frac{1}{m} \sum_{\tilde{\nu}} \sigma_{\tilde{\nu}}^{(2)} A_{\tilde{\nu}}^{(2)\dagger} B_{\tilde{\nu}}^{(N-2)\dagger} |0\rangle, \qquad (73b)$$

where we have written $c_{\bar{k}}c_k|\Psi_m\rangle = \sum_{\beta} \Gamma_{k\beta}^{(2)} B_{\beta}^{(N-2)\dagger}|0\rangle$ with $\Gamma_{k\beta}^{(2)} = \langle 0|B_{\beta}^{(N-2)}c_{\bar{k}}c_k|\Psi_m\rangle$ the elements of the "collective" sub-block $\Gamma_c^{(2)}$ of the full $\Gamma_{S\bar{S}}^{(2)}$ in (47), such that $\Gamma_c^{(2)}\Gamma_c^{(2)\dagger} = \rho_c^{(2)}$ in (72). In (73b) $\sigma_{\nu}^{(2)}$ are the singular values of this sub-block, with $\lambda_{\nu}^{(2)} = (\sigma_{\nu}^{(2)})^2$ the eigenvalues of $\rho_c^{(2)}$ and $A_{\nu}^{(2)\dagger}, B_{\nu}^{(N-2)\dagger}$ the associated normal operators determined by its SVD. In the presence of a dominant eigenvalue, a good approximation to $|\Psi_m\rangle$ can be obtained with just a few terms in (73) (see IV C).

In particular, for a pair creation operator of the form

$$A^{\dagger} = \sum_{k=1}^{m} \sigma_k c_k^{\dagger} c_{\bar{k}}^{\dagger} , \qquad (74)$$

where $\sum_{k} |\sigma_{k}|^{2} = 1$ and we can set σ_{k} real ≥ 0 by adjusting the phases of the c_{k}^{\dagger} , an example of (70) is

$$\begin{aligned}
|\Psi_m\rangle &= \frac{1}{m!\sqrt{\mathcal{N}'_m}} (A^{\dagger})^m |0\rangle \\
&= \frac{1}{\sqrt{\mathcal{N}'_m}} \sum_{\substack{m_1,\dots,m_n\\ \sum_k m_k = m}} \sigma_1^{m_1} \dots \sigma_n^{m_n} |m_1,\dots,m_n\rangle,
\end{aligned} \tag{75}$$

where $\mathcal{N}'_m = \sum_{m_1,...,m_n} \sigma_1^{2m_1} \dots, \sigma_n^{2m_n}$ (with the same previous restrictions on the m_k for bosons or fermions). These states are just particle number projected BCS-like states: $|\Psi_m\rangle \propto P_m |\text{BCS}\rangle$, where $|\text{BCS}\rangle \propto \exp[A^{\dagger}]|0\rangle = \prod_k \exp(\sigma_k c_k^{\dagger} c_{\bar{k}}^{\dagger})|0\rangle$ [38, 50] (= $\prod_k (1 + \sigma_k c_k^{\dagger} c_{\bar{k}}^{\dagger})|0\rangle$ for fermions) and P_m is the projector onto *m*-pair (2*m*-particle) states.

We now prove that in *all* states (75) (but not (70)) the largest eigenvalue $\lambda_1^{(2)}$ of $\rho_{S\bar{S}}^{(2)}$ (stemming from $\rho_c^{(2)}$ in (72)) satisfies

$$\lambda_1^{(2)} \ge \begin{cases} m \text{ (bosons)} \\ 1 \text{ (fermions)} \end{cases}, \tag{76}$$

for arbitrary $\{\sigma_k\}$: A straightforward evaluation of the average in the state (75) yields (see (A.15)-(A.16)),

$$\langle A^{\dagger}A \rangle = m \pm (m-1)\sum_{k} \sigma_{k}^{2} \langle c_{k}^{\dagger}c_{k} \rangle$$
 (77a)

$$= 1 + (m-1)(1 \pm \sum_{k} \sigma_{k}^{2} \langle c_{k}^{\dagger} c_{k} \rangle), \qquad (77b)$$

for bosons (+) or fermions (-), such that (77a) implies $\langle A^{\dagger}A \rangle \geq m$ for bosons and (77b) $\langle A^{\dagger}A \rangle \geq 1$ for fermions (where $\sum_{k} \sigma_{k}^{2} \langle c_{k}^{\dagger}c_{k} \rangle \leq \sum_{k} \sigma_{k}^{2} = 1$). This proves Eq.

(76) since the largest eigenvalue $\lambda_1^{(2)}$ of $\rho_{S\bar{S}}^{(2)}$ should satisfy $\lambda_1^{(2)} \ge \langle A^{\dagger}A \rangle$.

And to see it does not hold for all states (70), just take a superposition of states with orthogonal sp support, e.g. $\frac{1}{\sqrt{2}}(\prod_{k=1}^{m} c_{k}^{\dagger} c_{\bar{k}}^{\dagger} + \prod_{k=m+1}^{2m} c_{k}^{\dagger} c_{\bar{k}}^{\dagger})|0\rangle$, for which the nonzero eigenvalues of $\rho^{(2)}$ are just all 1/2 for both bosons and fermions $\forall m \geq 2$.

Eq. (77) also shows that $A^{\dagger}A$ has itself a largest eigenvalue $\lambda_{\max}^N \geq m$ (1) for bosons (fermions) amongst N = 2m-particle states, since again $\lambda_{\max}^N \geq \langle A^{\dagger}A \rangle$ for the average taken in any N particle state. Notice, however, that for general σ_k the state (75) is no longer an exact eigenstate of $A^{\dagger}A$, nor is A^{\dagger} a normal mode of the associated $\hat{\rho}^{(2)}$.

In the uniform case $\sigma_k = 1/\sqrt{n} \forall k$, $\langle c_k^{\dagger} c_k \rangle = m/n \forall k$ and we recover from (77) the result (69). Moreover, while (as previously mentioned) in fermion systems this is the maximum value $\lambda_1^{(2)}$ can reach among all 2*m*-particle states in a 2*n*-dimensional sp space, in boson systems Eq. (69) represents actually the minimum value reached by the maximum eigenvalue $\lambda_1^{(2)}$ among the states (75): Since for nonuniform $\sigma_k > 0$, $\langle c_k^{\dagger} c_k \rangle > \langle c_{k'}^{\dagger} c_{k'} \rangle$ if $\sigma_k > \sigma_{k'}$ while $\sum_k \langle c_k^{\dagger} c_k \rangle = m$ and $\sum_k \sigma_k^2 = 1$ are fixed, we obtain, writing $\langle c_k^{\dagger} c_k \rangle = m/n + \delta_k$, with $\sum_k \delta_k = 0$, the bound $\sum_k \sigma_k^2 \langle c_k^{\dagger} c_k \rangle \geq m/n$, such that (77a) and previous fermionic result imply, for all states (75),

$$\lambda_1^{(2)} \ge m(1 + \frac{m-1}{n}) \quad \text{(bosons)} \\ \lambda_1^{(2)} \le m(1 - \frac{m-1}{n}) \quad \text{(fermions)} \quad , \tag{78}$$

which for bosons is stronger than (76). Maximum $\lambda_1^{(2)}$ in the bosonic case amongst the states (75) is obtained when all pairs are in a single state k ($\lambda_1^{(2)} = m^2$).

Finally, we recall that in the fermion case any pair creation operator $A^{\dagger} = \frac{1}{2} \sum_{i,j} \Gamma_{ij} c_i^{\dagger} c_j^{\dagger}$ (with $\Gamma_{ji} = -\Gamma_{ij}$) can be written in the previous normal form (74) [11] (directly related to the normal form (12) for N = 2, M = 1of the state $A^{\dagger}|0\rangle$), with σ_k the singular values of Γ and the $c_k^{\dagger}, c_{L}^{\dagger}$, unitarily related to the c_i^{\dagger} .

C. Finite pairing system

We finally consider the exact ground state (GS) of a finite discrete pairing model. Such model describes finite superconducting systems in the fermionic case (see e.g. [50, 68, 69]), while its bosonic version has also been considered [65–67]. Studies of entanglement in such systems have mainly focused on mode-type entanglement in the approximate BCS GS [70–73] or on the fermionic one-body entanglement and concurrence [38]. Here we will concentrate on the two-body entanglement determined by $\rho^{(2)}$ and the associated state expansions, in both the fermionic and bosonic version of the model.

As in previous examples, we will work within an effective single particle subspace of dimension 2n, spanned

by *n* states *k* and *n* states \bar{k} , with sp levels of energies $\varepsilon_k = \varepsilon_{\bar{k}}$. The Hamiltonian is

$$H = \sum_{k} \varepsilon_k (c_k^{\dagger} c_k + c_{\bar{k}}^{\dagger} c_{\bar{k}}) - \sum_{k,k'} G_{kk'} c_{k'}^{\dagger} c_{\bar{k}'}^{\dagger} c_{\bar{k}} c_k , \quad (79)$$

where the second term is the pairing interaction. It conserves the number of particles in states k (subspace S) and \bar{k} (subspace \bar{S}), satisfying

$$[H, N_{\mathcal{S}}] = [H, N_{\bar{\mathcal{S}}}] = 0.$$
(80)

Hence its exact eigenstates, and in particular its GS, will satisfy Eqs. (35)-(63). For even N = 2m and $G_{kk'} > 0$ $\forall k, k'$, the exact GS will be of the form (70) for both bosons and fermions, since in order to minimize its energy it will have all N particles in m pairs (k, \bar{k}) , without broken pairs.

In what follows we consider a constant sp spacing $\varepsilon_{k+1} - \varepsilon_k = \varepsilon \ \forall k$ and uniform coupling strength $G_{kk'} = G \ge 0 \ \forall k, k'$, such that the interaction in (79) becomes $nGA^{\dagger}A$ with A^{\dagger} the uniform pair creation operator (59).

Thus, for $g \equiv G/\varepsilon \to \infty$, the GS of H will approach that of $-nGA^{\dagger}A$, which is the maximally paired state $|\Psi_m\rangle \propto (A^{\dagger})^m |0\rangle$, Eq. (61), as it maximizes $\langle A^{\dagger}A \rangle$ for any fixed N = 2m. For a uniform spectrum centered at 0, $\varepsilon_k = \varepsilon(k - \frac{n+1}{2})$, $k = 1, \ldots, n$, the energy $E_m =$ $\langle \Psi_m | H | \Psi_m \rangle$ of such state is (Eqs. (64b)-(69))

$$E_m = -nG\lambda_1^{(2)} = -mG[n \pm (m-1)], \qquad (81)$$

where +(-) is for bosons (fermions).

On the other hand, for $g \to 0^+$ the GS will approach $|\Psi_m^0\rangle = |m, 0, \ldots, 0\rangle$ for bosons, $|\Psi_m^0\rangle = |1_1 \ldots 1_m, 0, \ldots, 0\rangle$ for fermions (in terms of the paired states (62)), with $E_m^0 = \langle \Psi_m^0 | H | \Psi_m^0 \rangle = -m[\varepsilon(n-a)+bG]$ and a = 1 (m), b = m (1) for bosons (fermions). Therefore, $E_m - E_m^0 = m(\varepsilon - G)(n-a) < 0$ already for $G > \varepsilon$. The exact GS for finite n, m will then evolve continuously from $|\Psi_m^0\rangle$ to the state (61) as g increases from 0 to ∞ , through states of the form (70).

1. Fermionic system

We have analyzed in [38] the one-body entanglement determined by $\rho^{(1)}$ in the fermionic version of this system, together with the fermionic concurrence of the reduced state of 4 modes and other related aspects. We will here focus on the two-body DM and the associated entanglement entropy and exact expansions of the GS.

We first depict in Fig. 1 the exact eigenvalues (i.e. the entanglement spectrum) of the one- and two-body DMs in the fermionic case as a function of $g = G/\varepsilon$. We have considered a half-filled system N = 2m = n, with n = 10. At g = 0 the GS $|\Psi_0^m\rangle$ has just the bottom half levels occupied, such that all eigenvalues of $\rho^{(1)}$ and $\rho^{(2)}$ start from 1 or 0 at g = 0. Both the eigenvalues $\lambda_k^{(1)}$ of $\rho_S^{(1)}$ (top panel) and $\lambda_{kk'}^{(2)}$ of $\rho_S^{(2)}$ (bottom panel), Eq. (71),



FIG. 1. Eigenvalues of the one-body (top) and two-body (center and bottom) density matrices $\rho^{(1)}$ and $\rho^{(2)}$ as a function of the scaled coupling strength $g = G/\varepsilon$ (dimensionless) in the GS of the Hamiltonian (79) for a finite half-filled fermionic case (n = 10). The central panel depicts those of the central block $\rho_{S\bar{S}}^{(2)}$ in (40), containing the dominant eigenvalue $\lambda_1^{(2)} \geq 1$, while the bottom panel those of $\rho_S^{(2)} = \rho_{\bar{S}}^{(2)}$.

identical to those of $\rho_{\mathcal{S}}^{(1)}$, $\rho_{\mathcal{S}}^{(2)}$, become "more mixed" and < 1 as the coupling g increases, reflecting the departure of the GS from a SD as all levels above the Fermi level start to be occupied. They exhibit maximum variation around the transition region $g \approx 1$ and approach the maximally mixed limit (for such N) for $g \to \infty$, where they all coalesce with the values (65)-(67), i.e., $\lambda_k^{(1)} \to 1/2$, $\lambda_{kk'}^{(2)} \to \frac{1}{4}\frac{1-2/n}{1-1/n} \approx \frac{1}{4}(1-\frac{1}{n})$ in the half-filled case. These results imply a monotonously increasing one- and two-body entanglement within \mathcal{S} for increasing pairing strength, saturating for $g \to \infty$.

In contrast, the two-body DM block $\rho_{S\bar{S}}^{(2)}$ (central panel) exhibits instead a single *dominant eigenvalue* $\lambda_1^{(2)} > 1 \ \forall g > 1$ which departs from the rest (that behave as those of $\rho_S^{(2)}$) and increases for increasing g, approaching the limit (69) (= $\frac{n}{4} + \frac{1}{2}$ in the half-filled case) for $g \to \infty$. It is characteristic of pairing correlations and stems from the collective sub-block $\rho_c^{(2)}$, Eq. (72b), re-



FIG. 2. The entropy increment (82) of the normalized oneand two-body DMs as a function of the scaled strength $g = G/\varepsilon$ in the fermionic case of Fig. 1. The top panel depicts results for that of $\rho_{S}^{(1)}$ ($\Delta S_{S}^{(1)}$) and of the two-body block $\rho_{S}^{(2)}$ ($\Delta S_{S}^{(2)}$) in (40), whereas the bottom panel those of the central block $\rho_{S\bar{S}}^{(2)}$ and its collective sub-block $\rho_{c}^{(2)}$ ($\Delta_{S\bar{S}}^{(2)'}$), Eq. (72b). The dashed lines indicate their maximum values (reached for a maximally mixed DM).

flecting for large g the "multiple occupation" of the collective pair state created by A^{\dagger} . It prevents this block from becoming more mixed as g increases, implying a two-body $S\bar{S}$ entanglement below maximum for $g \to \infty$.

In Fig. 2 we depict the associated one- and two-body entropy increments ($\alpha = S$ or $S\bar{S}$, i = 1, 2)

$$\Delta S_{\alpha}^{(i)} = S[\rho_{n\alpha}^{(i)}(g)] - S[\rho_{n\alpha}^{(i)}(0)], \qquad (82)$$

which quantify the entanglement generated by the coupling, where $S(\rho) = -\text{Tr} \rho \log_2 \rho$ and $\rho_{n\alpha}^{(i)}(g)$ denotes the normalized *i*-body DM of block α at coupling *g*. Accordingly, for $\rho_S^{(1)}$ and $\rho_S^{(2)}$ (upper panel) this difference increases monotonously from 0 as *g* increases, reaching its saturation value for $g \to \infty$, where $\Delta_S^{(1)} \to \log_2 \frac{2n}{n} = 1$, $\Delta_S^{(2)} \to \log_2[\binom{n}{2}/\binom{n/2}{2}] \approx 2 + \frac{1}{n \ln 2}$ in the half-filled case. In contrast, for $\rho_{SS}^{(2)}$ (bottom), though increasing with g, $\Delta_{SS}^{(2)}$ stays below the saturation value $\log_2 \frac{n^2}{(n/2)^2} = 2$ due to the dominant eigenvalue $\lambda_1^{(2)}$, reaching for $g \to \infty$ the lower limit $-p \log_2 p - (1-p) \log_2 \frac{1-p}{n^2-1} - 2 \log_2 \frac{n}{2} \approx 2 - \frac{\log_2(n/e)}{n}$, where $p = \frac{\lambda_1^{(2)}}{(n/2)^2}$ with $\lambda_1^{(2)}$ given by (69). We also depict in the lower panel the entropy increment $\Delta_{SS}^{(2)}$ of the collective $n \times n$ sub-block $\rho_{Cn}^{(2)}$ of $\rho_{SS}^{(2)}$.

We also depict in the lower panel the entropy increment $\Delta_{SS}^{(2)'}$ of the collective $n \times n$ sub-block $\rho_{cn}^{(2)}$ of $\rho_{S\bar{S}}^{(2)}$, containing just the contractions $\langle c_k^{\dagger} c_{\bar{k}}^{\dagger} c_{\bar{k}'} c_{k'} \rangle$ (Eq. (72b)) and hence the dominant eigenvalue, which best reflects its effect. This increment actually becomes *negative* for



FIG. 3. Eigenvalues of the one-body (top) and two-body (center and bottom) density matrices $\rho_S^{(1)}$ and $\rho_{S\bar{S}}^{(2)}$, $\rho_S^{(2)}$ as a function of the scaled strength G/ε in the GS of Hamiltonian (79) for the N = n bosonic case. Details are similar to those of Fig. 1). The dominant eigenvalue of each block is indicated.

large g, approaching $\approx -\frac{1}{2}\log_2 \frac{n}{16} + O(\frac{\log_2 n}{n})$ for large n, well below its saturation value $\log_2 \frac{n}{n/2} = 1$. Thus, in this limit the entropy of this sub-block becomes *lower* than in the noninteracting case, reflecting the "separable-like" (2, N-2) form of the limit state (61a).

2. Bosonic system

Figs. 3-4 depict previous quantities in the bosonic case. The main difference is the behavior for weak coupling, since for $g \to 0^+$ all m pairs now fall to the lowest sp level. This implies a dominant eigenvalue in all blocks $\rho_S^{(1)}$, $\rho_S^{(2)}$ and $\rho_{SS}^{(2)}$ at low g, with $\lambda_k^{(1)} \to \frac{n}{2}\delta_{k1}$, $\lambda_{kk'}^{(2)} \to \delta_{kk'}\delta_{k1}\binom{n/2}{2}$ and $\rho_{ckk''}^{(2)} = \delta_{kk''}\delta_{k1}(\frac{n}{2})^2$ for $g \to 0$ in (71)-(72). As g increases all levels become occupied and all eigenvalues of $\rho_S^{(1)}$, $\rho_S^{(2)}$ become < 1 for large g, approaching for $g \to \infty$ the maximally mixed limits (65)-(67) ($\lambda_k^{(1)} \to \frac{1}{2}$, $\lambda_{kk'}^{(2)} \to \frac{1}{4}\frac{1-2/n}{1+1/n} \approx \frac{1}{4}(1-\frac{3}{n})$). However, in $\rho_{S\bar{S}}^{(2)}$ the dominant eigenvalue $\lambda_1^{(2)}$, though also decreasing for increasing g, stays well above m = n/2, approaching (69)



FIG. 4. The entropies $S^{(i)} = S(\rho_{Sn}^{(i)})$ (top) and $S(\rho_{S\bar{S}n}^{(2)})$ (bottom) of the normalized one- and two-body DM blocks in the bosonic case of Fig. 3. The bottom panel also depicts that of the normalized collective sub-block (72b), $S_{S\bar{S}}^{(2)'} = S(\rho_{cn}^{(2)})$. Dashed lines indicate again their maximum values.

 $(=\frac{3n}{4}-\frac{1}{2}$ for N=n) for $g \to \infty$. This reflects the strong deviation of the GS from a permanent as g increases, becoming approximately a bosonic coboson condensate, where a prominent eigenvalue remains in $\rho^{(2)}$ but not in $\rho^{(1)}$, in contrast with a standard condensate. The paired structure of the bosonic GS for large g can thus be also clearly identified through the spectra of $\rho^{(2)}$ and $\rho^{(1)}$.

The associated entropies (using the normalized DM blocks) are depicted in Fig. 4. Now they all vanish for $g \to 0$, while for $g \to \infty$ behave as in the fermionic case: those of $\rho_S^{(1)}$ and $\rho_S^{(2)}$ approach their saturation values $(S(\rho_{Sn}^{(1)}) \to \log_2 n, S(\rho_{Sn}^{(2)}) \to \log_2 \frac{n(n+1)}{2})$, while those of $\rho_{S\bar{S}n}^{(2)}$ and the collective sub-block $\rho_{cn}^{(2)}$ stabilize well below their maximum values $(\log_2 n^2 \text{ and } \log_2 n, \text{ dashed lines})$, reaching the lower limits $\approx \log_2 n^2 - \frac{3}{n} \log_2 \frac{3n}{2}$ and $\approx \frac{1}{4} \log_2 9.5n$ (plus $O(n^{-1})$ terms) respectively, reflecting the effect of the remnant dominant eigenvalue.

3. Approximate expansions

We finally depict in Fig. 5 the overlap between the exact GS $|\Psi\rangle$ and the approximate normalized GS $|\Psi_k\rangle$ obtained by taking just the first k terms in the Schmidt-like (2, N-2) expansion of Eq. (47b), $|\Psi_k\rangle \propto \sum_{\tilde{\nu}=1}^k \sigma_{\tilde{\nu}}^{(2)} A_{\tilde{\nu}}^{(2)\dagger} B_{\tilde{\nu}}^{(N-2)\dagger} |0\rangle$, based on the two-body DM block $\rho_{S\bar{S}}^{(2)}$. Terms are sorted in decreasing order of $\sigma_{\tilde{\nu}}^{(2)}$, i.e. of the eigenvalues $\lambda_{\nu}^{(2)} = (\sigma_{\nu}^{(2)})^2$ of this block. For fermions the expansion can be reduced to the sum (73b) based on the collective sub-block $\rho_c^{(2)}$, involving just n terms. In particular, for k = 1 the approximation corre-



FIG. 5. The overlap $\langle \Psi | \Psi_k \rangle$ between the exact GS $| \Psi \rangle$ and the approximate GS $| \Psi_k \rangle$ obtained by conserving just the first k terms in the 2 – (N – 2) Schmidt-like expansion (47b) associated to the central block $\rho_{S\bar{S}}^{(2)}$ in (40) of the two-body DM, as a function of the scaled strength G/ε , in the fermionic (top) and bosonic (bottom) systems of previous figures. Saturation $\langle \Psi | \Psi_k \rangle = 1$ (thin line) is always reached for the full expansion associated to this block. In the fermionic case the reduced expansion (73b) can be used instead of (47b).

sponds to the dominant eigenvalue $\lambda_1^{(2)}$ and is determined by its normal eigenvector $A_1^{(2)\dagger}$:

$$|\Psi_1\rangle \propto \sigma_1^{(2)} A_1^{(2)\dagger} B_1^{(N-2)\dagger} |0\rangle,$$
 (83)

where $\sigma_1^{(2)} B_1^{(N-2)\dagger} |0\rangle = A_1^{(2)} |\Psi\rangle$ (Eq. (15a)). It is first verified that the full sum always yields the

exact GS in all expansions. Nonetheless, for those based on $\rho_{S\bar{S}}^{(2)}$ (or $\rho_c^{(2)}$ for fermions), already the k = 1 approximation (83) is seen to provide a very good overlap $|\langle \Psi | \Psi_k \rangle| \gtrsim 0.9$ for all values of g, minimum just at the transition region around $q \approx 1$. Moreover, this approximation becomes exact for both $q \to \infty$ and $q \to 0$ in both the fermionic and bosonic systems, since in these limits the exact GS becomes of the form (75), i.e. Eq. (83) with $B_1^{(N-2)\dagger} = (A_1^{(2)\dagger})^{m-1}$ and $A_1^{(2)\dagger}$ eigenvector of $\rho^{(2)}$: For $g \to \infty$, $A_1^{(2)\dagger}$ becomes the uniform pair creation operator (59), as the GS approaches the state (61a), whereas for $g \to 0$, $A_1^{(2)\dagger} \to c_1^{\dagger} c_1^{\dagger}$ for bosons while $A_1^{(2)\dagger} \rightarrow \frac{1}{\sqrt{m}} \sum_{k=1}^m c_k^{\dagger} c_{\bar{k}}^{\dagger}$ for fermions (as the SD $|\Psi_m^0\rangle$ can be written as $\frac{m^{m/2}}{m!} (A_1^{(2)\dagger})^m |0\rangle$). In the fermionic system the k = 2 approximation in (73b) further improves the k = 1 result in the transition region, staying also reliable for large q, while for bosons more terms are required for obtaining a better approximation for all q. As the terms in these expansions are not necessarily linearly independent, the convergence to the exact $|\Psi\rangle$ is not necessarily monotonous as k increases for all values of q.

V. CONCLUSIONS

We have presented a unified formalism for analyzing general pure states of systems of N indistinguishable particles in terms of exact bipartite-like (M, N - M) decompositions, valid for both fermions and bosons. It is directly connected with the isospectral M and N - M-body DMs, whose eigenvalues acquire here meaning also as coefficients of the associated diagonal Schmidt decomposition. The ensuing M-body entanglement, quantified through the entropy of the normalized M-body DM, was shown to fulfill general monotonicity relations under certain quantum operations.

We have analyzed in addition the exact reduced expansions emerging when the number of particles in a certain subspace S of the full sp space is fixed, which are associated to the ensuing blocks of the DMs. Then both local (in S) and "mixed" (in S and its complementary subspace \bar{S}) DMs and corresponding exact (M, N-M)expansions arise in connection with these blocks, whose eigenvalues and Schmidt decomposition characterize the system correlations. The standard reduced DMs and Schmidt decomposition of distinguishable bipartite systems emerge as a particular case in this scenario.

As example, we have analyzed in detail the behavior of the one- and two-body DMs in the GS a finite pairing system. These systems are characterized by a large dominant eigenvalue of $\rho^{(2)}$ in the superfluid phase, which in the present formalism enables a reliable description of the exact GS with just a few terms (in fact just one term) of the associated (2, N - 2) expansion, in both fermion and boson systems. We have also provided exact results for the eigenvalues of $\rho^{(2)}$ in the maximally paired states for bosons and fermions, as well as bounds for its dominant eigenvalue in some general paired states, again for both bosons and fermions. Applications to more complex systems and further analysis of the role of present modeindependent entanglement measures in quantum information are under investigation.

ACKNOWLEDGMENTS

Authors acknowledge support from CONICET and CIC (R.R.) of Argentina. Work supported by CONICET PIP Grant No. 11220200101877CO.

Appendix A: Proofs of main expressions

In order to prove Eq. (4), we first show that

$$\frac{1}{M!} \sum_{i_1,\dots,i_M} c_{i_1}^{\dagger} \dots c_{i_M}^{\dagger} c_{i_M} \dots c_{i_1} = \binom{N}{M}$$
(A1)

for both fermions and bosons, where the sum over each i_j (j = 1, ..., M) runs over all d sp states and $\hat{N} = \sum_i c_i^{\dagger} c_i$ is the particle number operator.

Proof: When applying the sum in (A1) to an arbitrary N particle state $|\Psi\rangle$ with $N \ge M$ (for N < M we set $\binom{N}{M} = 0$), the innermost sum $\sum_{i_m} c_{i_m}^{\dagger} c_{i_m} = \hat{N}$ takes the value N - M + 1, and then, successively, the sums $\sum_{i_j} c_{i_j}^{\dagger} c_{i_j}$ take the values N - j + 1 for $j = M, \ldots, 1$. This leads to Eq. (A1). Then, using relations (1),

$$\sum_{\substack{i_1,\dots,i_M \\ n_1+\dots+n_d=M}} \frac{c_{i_1}^{\dagger}\dots c_{i_M}^{\dagger}c_{i_M}\dots c_{i_1}}{M!} = \sum_{\substack{i_1 \le i_2 \dots \le i_M \\ i_1 \le \dots \le i_M}} \frac{c_{i_1}^{\dagger}\dots c_{d_1}^{\dagger}c_{d_1}^{n}\dots c_{i_1}^{n_1}}{n_1!\dots n_d!} = \sum_{\boldsymbol{\alpha}} C_{\boldsymbol{\alpha}}^{(M)\dagger} C_{\boldsymbol{\alpha}}^{(M)}$$
(A2)

where the sum in (A2) runs over all d_M operators (2), which leads to Eq. (4). Here n_j (j = 1, ..., d) is the number of times sp state j appears in the string $(i_1, ..., i_M)$, such that $c_{i_1}^{\dagger} \ldots c_{i_M}^{\dagger} = c_1^{\dagger n_1} \ldots c_d^{\dagger n_d}$, with $n_1 + \ldots + n_d = M$. The sum over the ordered i_j 's is equivalent to that over these occupations with previous constraint, with each configuration (n_1, \ldots, n_d) appearing $\frac{M!}{n_1 \ldots n_d!}$ times in the first unrestricted sum. These arguments hold for both bosons and fermions, but for the latter are trivial as $n_i = 0, 1$.

We now prove Eqs. (5b)–(6). Using the same previous reasoning, we obtain, for a completely symmetric (bosons) or antisymmetric (fermions) tensor $\Gamma_{i_1,...,i_N}$,

$$\frac{1}{N!} \sum_{\substack{i_1,\dots,i_N\\i_1,\dots,i_N}} \Gamma_{i_1\dots i_N} c_{i_1}^{\dagger} \dots c_{i_N}^{\dagger} |0\rangle = \sum_{\substack{i_1 \leq \dots \leq i_N\\i_1 \leq \dots \leq i_N}} \Gamma_{i_1\dots i_N} \frac{c_{i_1}^{\dagger} \dots c_{i_N}^{\dagger}}{n_1! \dots n_d!} |0\rangle = \sum_{\boldsymbol{\alpha}} \Gamma_{\boldsymbol{\alpha}}^{(N)} C_{\boldsymbol{\alpha}}^{(N)\dagger} |0\rangle, \quad (A3)$$

where $\Gamma_{n_1...n_d} = \Gamma_{i_1...i_N}$ if $c_{i_1}^{\dagger} \dots c_{i_N}^{\dagger} = c_1^{\dagger n_1} \dots c_d^{\dagger n_d}$ $(i_1 \leq \dots \leq i_N \text{ for fermions})$ and $\Gamma_{\boldsymbol{\alpha}}^{(N)} = \frac{\Gamma_{n_1...n_d}}{\sqrt{n_1!...\sqrt{n_d!}}}$. \square Relation between $\Gamma_{\boldsymbol{\alpha}\boldsymbol{\beta}}^{(M)}$ and $\Gamma_{i_1...i_N}$. Rewriting (A3) as

$$\begin{split} |\Psi\rangle &= \frac{M!(N-M)!}{N!} \sum_{\substack{i_{1} \leq \dots \leq i_{M}, \\ i_{M+1} \leq \dots \leq i_{N}}} \Gamma_{i_{1}\dots i_{M} i_{M+1}\dots i_{N}} \frac{c_{i_{1}}^{\dagger} \dots c_{i_{M}}^{\dagger} c_{i_{M+1}}^{\dagger} \dots c_{i_{N}}^{\dagger}}{n_{1} \dots n_{d}! n_{1}' \dots n_{d}'!} |0\rangle \\ &= \binom{N}{M}^{-1} \sum_{\substack{n_{1},\dots,n_{d},n_{1}',\dots,n_{d}' \\ \sum_{j} n_{j} = M, \sum_{j} n_{j}' = N-M}} \Gamma_{n_{1}\dots n_{d},n_{1}',\dots,n_{d}'}^{(M)} \frac{c_{1}^{\dagger n_{1}} \dots c_{d}^{\dagger n_{d}} c_{1}^{\dagger n_{1}'} \dots c_{d}^{\dagger n_{d}'}}{n_{1} \dots n_{d}! n_{1}' \dots n_{d}'!} |0\rangle \\ &= \binom{N}{M}^{-1} \sum_{\boldsymbol{\alpha},\boldsymbol{\beta}} \Gamma_{\boldsymbol{\alpha}\boldsymbol{\beta}}^{(M)} C_{\boldsymbol{\alpha}}^{(M)\dagger} C_{\boldsymbol{\beta}}^{(N-M)\dagger} |0\rangle , \end{split}$$
(A4)

where $\Gamma_{n_1...n_d,n'_1...n'_d}^{(M)} = \Gamma_{i_1,...,i_M,i_{M+1}...i_N}$ for n_j , n'_j the number of times sp state j appears in $(i_1,...,i_M)$ and $(i_{M+1},...,i_N)$ $(i_1 < ... < i_M, i_{M+1} < ... < i_N$ for fermions) such that $\sum_j n_j = M$, $\sum_j n'_j = N - M$, it is seen that

$$\Gamma_{\alpha\beta}^{(M)} = \frac{\Gamma_{n_1...n_d,n'_1...n'_d}^{(M)}}{\sqrt{n_1!...n_d!}\sqrt{n'_1!...n'_d!}}.$$
 (A5)

Behavior under unitary sp transformations. If

$$c_i \to \hat{U}^{\dagger} c_i \hat{U} = \sum_k U_{ki} c_k , \quad \hat{U} = e^{-i \sum_{i,j} h_{ij} c_i^{\dagger} c_j} , \quad (A6)$$

where $h^{\dagger} = h$, U is the matrix $U = \exp[-ih]$ and (A6) holds for both bosons and fermions, the operators $C_{\alpha}^{(M)}$ transform unitarily: $C_{\alpha}^{(M)} \rightarrow \hat{U}^{\dagger}C_{\alpha}^{(M)}\hat{U} = \sum_{\alpha'} U_{\alpha'\alpha}^{(M)} C_{\alpha'}^{(M)}$ for $U^{(M)}$ a unitary symmetrized or antisymmetrized tensor product of M matrices U. This implies $\Gamma^{(M)} \rightarrow U^{(M)*}\Gamma^{(M)}U^{(N-M)\dagger}$, then leaving its singular values $\sigma_{\nu}^{(M)}$ unchanged $\forall M$. And under similar sp transformations of the state,

$$|\Psi\rangle \to \hat{U}|\Psi\rangle , \quad \hat{U} = e^{-i\sum_{i,j}h_{ij}c_i^{\dagger}c_j} , \quad (A7)$$

we have $\langle \hat{O} \rangle \rightarrow \langle \hat{U}^{\dagger} \hat{O} \hat{U} \rangle$ and hence $\rho^{(M)} \rightarrow U^{(M)t} \rho^{(M)} U^{(M)t\dagger}$, with $\hat{\rho}^{(M)} \rightarrow \hat{U} \hat{\rho}^{(M)} \hat{U}^{\dagger}$ as is apparent from (22). Its eigenvalues $\lambda_{\nu}^{(M)}$ remain then unchanged, in agreement with their direct relation with $\sigma_{\nu}^{(M)}$.

Proof of Eqs. (29)–(30): From Eq. (22a) we obtain $\operatorname{Tr} \hat{\rho}^{(M)} C_{\gamma'}^{(L)\dagger} C_{\gamma}^{(L)} = \sum_{\beta} \langle \Psi | C_{\beta}^{(N-M)\dagger} C_{\gamma'}^{(L)\dagger} C_{\gamma}^{(L)} C_{\beta}^{(N-M)} | \Psi \rangle$ $= \binom{N-L}{N-M} \rho_{\gamma\gamma'}^{(L)} \text{ for } L \leq M \leq N, \text{ by using Eq. (4) for } \sum_{\beta} C_{\beta}^{(N-M)\dagger} C_{\beta}^{(N-M)} \text{ applied on the } N-L\text{-particle state } C_{\gamma}^{(L)} | \Psi \rangle, \text{ with } \rho_{\gamma\gamma'}^{(L)} = \langle \Psi | C_{\gamma'}^{(L)\dagger} C_{\gamma}^{(L)} | \Psi \rangle. \text{ This leads to Eq. (29). Then, by replacing expression (22a) or (28) in (29), we obtain, for the normalized DM <math>\rho_n^{(L)} = \rho^{(L)} / \binom{N}{L},$

$$\rho_n^{(L)} = \frac{\binom{N}{M}\binom{L}{L}}{\binom{N}{L}\binom{N-L}{N-M}} \sum_{\boldsymbol{\beta}} p_{\boldsymbol{\beta}} \rho_{\boldsymbol{\beta}n}^{(L)}, \qquad (A8)$$

where $\rho_{\beta n}^{(L)} = \rho_{\beta}^{(L)} / {M \choose L}$ are the normalized *L*-body DMs in the normalized *M*-particle states $|\Psi_{\beta}\rangle = \frac{1}{\sqrt{p_{\beta}}}\mathcal{M}_{\beta}|\Psi\rangle$, with p_{β} the probabilities (27). This leads to Eq. (30) since ${N \choose L} / [{N \choose L} {N-L \choose N-M}] = 1$, in agreement with normalization: $\operatorname{Tr} \rho_n^{(L)} = \operatorname{Tr} \rho_{\beta n}^{(L)} = 1 = \sum_{\beta} p_{\beta}$.

We now prove the contractions and results (65),(67) and (68) in the maximally paired state (61) for bosons and fermions. The number of states with m pairs (k, \bar{k}) in a sp space of n sp states k and n sp states \bar{k} is

$$\mathcal{N}_{n,m} = \begin{cases} \binom{n+m-1}{m} & \text{(bosons)} \\ \binom{n}{m} & \text{(fermions)} \end{cases} .$$
(A9)

where $m \ge 0$ for bosons and $0 \le m \le n$ for fermions. Then, for averages $\langle O \rangle = \langle \Psi_m | O | \Psi_m \rangle$ in the state (61), we obtain in the first place the expected obvious result

$$\langle c_k^{\dagger} c_{k'} \rangle = \delta_{kk'} \sum_l \frac{l \mathcal{N}_{n-1,m-l}}{\mathcal{N}_{n,m}} = \frac{m}{n}$$
 (A10)

for both bosons and fermions, where the sum runs over l = 0, 1, 2, ..., m for bosons and l = 0, 1 for fermions. The same holds for $k \to \bar{k}$. For two body contractions, assuming in what follows $k \neq k'$, we obtain

$$\langle c_k^{\dagger} c_{k'}^{\dagger} c_{k'} c_k \rangle = \sum_{l,l'} \frac{ll' \mathcal{N}_{n-2,m-l-l'}}{\mathcal{N}_{n,m}} = \frac{m(m-1)}{n(n\pm 1)} = \lambda_2^{(2)}$$
 (A11)

where the sum runs over $l + l' \leq m$ for bosons (+) and l = l' = 1 for fermions (-). This same result is obtained for $\langle c_k^{\dagger} c_{\bar{k}'}^{\dagger} c_{\bar{k}'} c_k \rangle$, $\langle \frac{c_k^{\dagger 2} c_k^2}{2!} \rangle$ (boson case) and $k, k' \to \bar{k}, \bar{k'}$. On the other hand,

$$\langle c_k^{\dagger} c_{\bar{k}}^{\dagger} c_{\bar{k}} c_k \rangle = \sum_l \frac{l^2 \mathcal{N}_{n-1,m-l}}{\mathcal{N}_{n,m}} = \frac{m(n+m-1\pm m)}{n(n\pm 1)}$$
(A12)

$$\langle c_k^{\dagger} c_{\bar{k}}^{\dagger} c_{\bar{k}'} c_{k'} \rangle = \sum_{l,l'} \frac{l(l'+1) \mathcal{N}_{n-2,m-l-l'}}{\mathcal{N}_{n,m}} = \frac{m(n\pm m)}{n(n\pm 1)}.$$
 (A13)

These exact results lead to Eqs. (65)-(68).

Then blocks $\rho_{S}^{(2)}$ and $\rho_{\bar{S}}^{(2)}$ of $\rho^{(2)}$ become here identical and proportional to $\lambda_{2}^{(2)}\mathbb{1}$, whereas the mixed block $\rho_{S\bar{S}}^{(2)}$ becomes itself blocked in two submatrices:

$$\rho_{\mathcal{S}\bar{\mathcal{S}}}^{(2)} = \begin{pmatrix} \lambda_2^{(2)} \mathbb{1} & 0\\ 0 & \rho_c^{(2)} \end{pmatrix}, \qquad (A14)$$

where $\rho_{c_{kk'}}^{(2)} = \langle c_k^{\dagger} c_k^{\dagger} c_{k'} c_{k'} \rangle = a \delta_{kk'} + b(1 - \delta_{kk'})$ is the $n \times n$ block containing the two-body pairing correlations, with a, b given by (A12)–(A13). Its eigenvalues are then $\lambda_1^{(2)} = (n-1)b + a$, Eq. (69), non-degenerate and dominant, and $\lambda_2^{(2)} = a - b$, Eqs. (67)–(A11), n - 1-fold degenerate. \Box Proof of Eq. (77): For bosons, a direct evaluation of

 $\langle A^{\dagger}A \rangle$ in the state (75), with A given by (74), yields

$$\langle A^{\dagger}A \rangle = \sum_{k} \sigma_{k}^{2} \langle m_{k}^{2} \rangle + \sum_{k \neq k'} \sigma_{k}^{2} \langle (m_{k}+1)m_{k'} \rangle$$

$$= m + (m-1) \sum_{k} \sigma_{k}^{2} \langle c_{k}^{\dagger}c_{k} \rangle$$
(A15)

where we used $\sum_k \sigma_k^2 = 1$, $\sum_k m_k = m$ and $\langle m_k \rangle = \langle c_k^{\dagger} c_k \rangle = \langle c_k^{\dagger} c_k \rangle$. Similarly, for fermions we obtain

$$\langle A^{\dagger}A \rangle = \sum_{k} \sigma_{k}^{2} \langle m_{k} \rangle + \sum_{k \neq k'} \sigma_{k}^{2} \langle (1 - m_{k})m_{k'} \rangle$$

$$= m - (m - 1) \sum_{k} \sigma_{k}^{2} \langle c_{k}^{\dagger}c_{k} \rangle .$$
(A16)

Eqs. (A15)-(A16) then lead to Eq. (77).

Appendix B: *M*-body entanglement and bounds to bipartite entanglement after particle transfer

Let us consider a general initial state (Eq. (7))

$$|\Psi_0\rangle = {\binom{N}{M}}^{-1} \sum_{\boldsymbol{\alpha}\boldsymbol{\beta}\in\mathcal{S}_0} \Gamma^{(M)}_{\boldsymbol{\alpha},\boldsymbol{\beta}} C^{(M)\dagger}_{\boldsymbol{\alpha}} C^{(N-M)\dagger}_{\boldsymbol{\beta}} |0\rangle \qquad (B1)$$

of N indistinguishable particles (fermions or bosons), occupying sp states just within a subspace $S_0 \subset \mathcal{H}$ of the full sp space \mathcal{H} . We then consider a completely positive trace preserving (CPTP) operation \mathcal{T} which transfers M < N particles from S_0 to an initially empty subspace S orthogonal to S_0 (e.g., M particles from a group of N localized within some bounded region of space to a distinct non overlapping region, or from low lying energy levels to higher orthogonal levels). For this purpose we define the *M*-body operators \hat{T}_r such that

$$\hat{T}_{r}|\Psi_{0}\rangle = \binom{N}{M}^{-1/2} \sum_{\boldsymbol{\gamma}\in\mathcal{S},\boldsymbol{\alpha}\in\mathcal{S}_{0}} T_{r_{\boldsymbol{\gamma}\boldsymbol{\alpha}}} C^{(M)\dagger}_{\boldsymbol{\gamma}} C^{(M)}_{\boldsymbol{\alpha}} |\Psi_{0}\rangle (B2a)$$
$$= \sum_{\boldsymbol{\gamma}\in\mathcal{S},\boldsymbol{\beta}\in\mathcal{S}_{0}} \Gamma^{(M)}_{r_{\boldsymbol{\gamma}\boldsymbol{\beta}}} C^{(M)\dagger}_{\boldsymbol{\gamma}} C^{(N-M)\dagger}_{\boldsymbol{\beta}} |0\rangle, \quad (B2b)$$

is a state satisfying (35) for $N_{\mathcal{S}} = M$, where $\Gamma_r^{(M)} = \binom{N}{M}^{-1/2} T_r \Gamma^{(M)}$ (Eq. (8)). If we now assume $\sum_r T_r^{\dagger} T_r = \mathbb{1}$ and use Eq. (4) for $N \to M$, then for any such $|\Psi_0\rangle$,

$$\sum_{r} \hat{T}_{r}^{\dagger} \hat{T}_{r} |\Psi_{0}\rangle = |\Psi_{0}\rangle, \qquad (B3)$$

since $C_{\gamma'}^{(M)}C_{\gamma}^{(M)\dagger}C_{\alpha}^{(M)}|\Psi_{0}\rangle = \delta_{\gamma'\gamma}C_{\alpha}^{(M)}|\Psi_{0}\rangle$ for $\gamma', \gamma \in \mathcal{S}, \alpha \in \mathcal{S}_{0}$, for both fermions and bosons. Hence we can consider the set of operators \hat{T}_{r} as a quantum operation mapping $|\Psi_{0}\rangle$ to states $|\Psi_{r}\rangle \propto \hat{T}_{r}|\Psi\rangle$ with probabilities

$$p_r = \langle \Psi_0 | \hat{T}_r^{\dagger} \hat{T}_r | \Psi_0 \rangle = \operatorname{Tr}[\Gamma_r^{(M)\dagger} \Gamma_r^{(M)}], \qquad (B4)$$

satisfying $\sum_{r} p_{r} = \text{Tr} \left[\Gamma^{(M)\dagger} \Gamma^{(M)} \right] / {N \choose M} = 1.$ Using result (56), the reduced state of the $N_{\mathcal{S}} =$

Using result (56), the reduced state of the $N_{\mathcal{S}} = M$ particles at \mathcal{S} in the normalized state $|\Psi_r\rangle$ is $\rho_{\mathcal{S}r}^{(M)} = \Gamma_r^{(M)} \Gamma_r^{(M)\dagger} / p_r$. Since the nonzero eigenvalues of $\Gamma_r^{(M)} \Gamma_r^{(M)\dagger}$ are the same as those of $\Gamma_r^{(M)\dagger} \Gamma_r^{(M)}$ and since $\sum_r \Gamma_r^{(M)\dagger} \Gamma_r^{(M)} = \Gamma^{(M)\dagger} \Gamma_r^{(M)} / {N \choose M}$ has then the same nonzero eigenvalues as the normalized DM $\rho_{0n}^{(M)} = \Gamma^{(M)} \Gamma^{(M)\dagger} / {N \choose M}$ in the original state $|\Psi_0\rangle$, we obtain the following majorization relation

$$\boldsymbol{\lambda}(\hat{\rho}_{0n}^{(M)}) \prec \sum_{r} p_r \boldsymbol{\lambda}(\hat{\rho}_{\mathcal{S}r}^{(M)})$$
(B5)

between the sorted eigenvalues of $\rho_{0n}^{(M)}$ and those of $\rho_{Sr}^{(M)}$, the latter determining the entanglement between the $M = N_S$ particles at S and the remaining N - M particles at the orthogonal subspace S_0 . It implies the following entropic inequality

$$S(\hat{\rho}_{0n}^{(M)}) \ge \sum_{r} p_r S(\hat{\rho}_{\mathcal{S}r}^{(M)}), \qquad (B6)$$

between the entropy of the normalized DM $\rho_{0n}^{(M)}$ in $|\Psi_0\rangle$, which measures its *M*-body entanglement, and the average entanglement entropy between the (now distinguishable) *M* particles at *S* and the remaining N-M particles at S_0 in the post-selected states $|\Psi_r\rangle$, which then cannot surpass the original *M*-body entropy. It is valid again for any concave concave entropy *S*. All other results derived in [21] for the conversion $|\Psi\rangle \rightarrow \{|\Psi_r\rangle\}$ in fermion systems remain then valid for bosons.

Finally, note that if $T_{r_{\gamma\alpha}} = T_{\gamma\alpha}\delta_{r\alpha} \ (\alpha \in S_0)$, then

$$\hat{T}_{\alpha}|\Psi_{0}\rangle = {\binom{N}{M}}^{-1/2} \sum_{\gamma \in \mathcal{S}} T_{\gamma \alpha} C_{\gamma}^{(M)\dagger} C_{\alpha}^{(M)} |\Psi_{0}\rangle, \quad (B7)$$

with $\sum_{\gamma} |T_{\gamma\alpha}|^2 = 1$ and hence $\sum_{\alpha} \hat{T}^{\dagger}_{\alpha} \hat{T}_{\alpha} |\Psi_0\rangle = |\Psi_0\rangle$. Thus, this map implements the measurement based on

the operators (25) (for $N - M \to M$) on S_0 through a particle number conserving map in the full system $S_0 \oplus S$, transferring M particles from S_0 to S.

- F. Benatti, R. Floreanini, F. Franchini, and U. Marzolino, "Entanglement in indistinguishable particle systems," Phys. Rep. 878, 1 (2020).
- [2] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Camb. Univ. Press, Cambridge UK, 2000).
- [3] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, "Quantum entanglement," Rev. Mod. Phys. 81, 865 (2009).
- [4] P. Zanardi, "Quantum entanglement in fermionic lattices," Phys. Rev. A 65, 042101 (2002).
- [5] Y. Shi, "Quantum entanglement of identical particles," Phys. Rev. A 67, 024301 (2003).
- [6] S. Ashhab, K. Maruyama, and F. Nori, "Observing quantum nonlocality in the entanglement between modes of massive particles," Phys. Rev. A 75, 022108 (2007).
- [7] L. Heaney and V. Vedral, "Natural mode entanglement as a resource for quantum communication," Phys. Rev. Lett. 103, 200502 (2009), L. Heaney and J. Anders, "Bell-inequality test for spatial- mode entanglement of a single massive particle", Phys. Rev. A 80, 032104 (2009).
- [8] N. Friis, A.R. Lee, and D.E. Bruschi, "Fermionic-mode entanglement in quantum information," Phys. Rev. A 87, 022338 (2013).
- [9] C. Spee, K. Schwaiger, G. Giedke, and B. Kraus, "Mode entanglement of gaussian fermionic states," Phys. Rev. A 97, 042325 (2018).
- [10] J. Schliemann, J. I. Cirac, M. Kuś, M. Lewenstein, and D. Loss, "Quantum correlations in two-fermion systems," Phys. Rev. A 64, 022303 (2001).
- [11] K Eckert, J. Schliemann, D. Bruß, and M. Lewenstein, "Quantum correlations in systems of indistinguishable particles," Ann. Phys. 299, 88 (2002).
- [12] R. Paškauskas and L. You, "Quantum correlations in twoboson wave functions," Phys. Rev. A 64, 042310 (2001).
- [13] Y.S. Li, B. Zeng, X.S. Liu, and G.L. Long, "Entanglement in a two-identical-particle system," Phys. Rev. A 64, 054302 (2001).
- [14] G. Ghirardi, L. Marinatto, and T. Weber, "Entanglement and properties of composite quantum systems: a conceptual and mathematical analysis," J. Stat. Phys. 108, 49–122 (2002), G.C. Ghirardi and L. Marinatto, "General criterion for the entanglement of two indistinguishable particles", Phys. Rev. A 70, 012109 (2004).
- [15] H. M. Wiseman and J. A. Vaccaro, "Entanglement of indistinguishable particles shared between two parties," Phys. Rev. Lett. **91**, 097902 (2003).
- [16] F. Iemini and R.O. Vianna, "Computable measures for the entanglement of indistinguishable particles," Phys. Rev. A 87, 022327 (2013).
- [17] N. Gigena and R. Rossignoli, "Entanglement in fermion systems," Phys. Rev. A 92, 042326 (2015).
- [18] A.P. Majtey, P.A. Bouvrie, A. Valdés-Hernández, and A.R. Plastino, "Multipartite concurrence for identicalfermion systems," Phys. Rev. A 93, 032335 (2016).
- [19] L.S. Souza, T. Debarba, D. L. Braga Ferreira, F. Iemini,

and R. O. Vianna, "Completely positive maps for reduced states of indistinguishable particles," Phys. Rev. A **98**, 052135 (2018).

- [20] N. Gigena, M. Di Tullio, and R. Rossignoli, "One-body entanglement as a quantum resource in fermionic systems," Phys. Rev. A 102, 042410 (2020).
- [21] N. Gigena, M. Di Tullio, and R. Rossignoli, "Many-body entanglement in fermion systems," Physical Review A 103, 052424 (2021).
- [22] H. Barnum, E. Knill, G. Ortiz, R. Somma, and L. Viola, "A subsystem-independent generalization of entanglement," Phys. Rev. Lett. **92**, 107902 (2004).
- [23] P. Zanardi, D.A. Lidar, and S. Lloyd, "Quantum tensor product structures are observable induced," Phys. Rev. Lett. 92, 060402 (2004).
- [24] T. Sasaki, T.Ichikawa, I.Tsutsui, "Entanglement of indistinguishable particles," Phys. Rev. A 83, 012113 (2011).
- [25] A.P. Balachandran, T.R. Govindarajan, A.R. de Queiroz, A.F. Reyes-Lega, "Entanglement and particle identity: a unifying approach," Phys. Rev. Lett. **110**, 080503 (2013); "Algebraic approach to entanglement and entropy", Phys. Rev. A **88**, 022301 (2013).
- [26] F. Benatti, R. Floreanini, and U. Marzolino, "Entanglement in fermion systems and quantum metrology," Phys. Rev. A 89, 032326 (2014).
- [27] D. Cavalcanti, L.M. Malard, F.M. Matinaga, M.O. Terra Cunha, M. F. Santos, "Useful entanglement from the Pauli principle," Phys. Rev. B 76, 113304 (2007).
- [28] N. Killoran, M. Cramer, and M. B. Plenio, "Extracting entanglement from identical particles," Phys. Rev. Lett. 112, 150501 (2014).
- [29] G. Compagno, A. Castellini, and R. Lo Franco, "Dealing with indistinguishable particles and their entanglement," Phil. Trans. R. Soc. 376, 20170317 (2018).
- [30] R. Lo Franco and G. Compagno, "Indistinguishability of elementary systems as a resource for quantum information processing," Phys. Rev. Lett. **120**, 240403 (2018).
- [31] B. Morris, B. Yadin, M. Fadel, T. Zibold, P. Treutlein, and G. Adesso, "Entanglement between identical particles is a useful and consistent resource," Phys. Rev. X 10, 041012 (2020).
- [32] F. Iemini, T.O. Maciel, and R.O. Vianna, "Entanglement of indistinguishable particles as a probe for quantum phase transitions in the extended Hubbard model," Phys. Rev. B 92, 075423 (2015).
- [33] D. Dasenbrook, J. Bowles, J. Bohr Brask, P.P. Hofer, C. Flindt, and N. Brunner, "Single-electron entanglement and nonlocality," New J. Phys. 18, 043036 (2016).
- [34] F. Benatti, R. Floreanini, F. Franchini, and U. Marzolino, "Remarks on entanglement and identical particles," Open Syst. Inf. Dyn. 24, 17400042 (2017).
- [35] N. Gigena and R. Rossignoli, "Bipartite entanglement in fermion systems," Phys. Rev. A 95, 062320 (2017).
- [36] T. Debarba, F. Iemini, G. Giedke, and N. Friis, "Teleporting quantum information encoded in fermionic modes," Phys. Rev. A 101, 052326 (2020).

- [37] L. Ding and C. Schilling, "Correlation paradox of the dissociation limit: A quantum information perspective," J. Chem. Theory Comput. 16, 4159 (2020), L. Ding, S. Mardazad, S. Das, S. Szalay, U. Schollwočk, A. Zimborás, C. Schilling, "Concept of orbital entanglement and correlation in quantum chemistry", J. Chem. Theory Comput. **17**, 79 (2021).
- [38] M. Di Tullio, N. Gigena, and R. Rossignoli, "Fermionic entanglement in superconducting systems," Phys. Rev. A 97, 062109 (2018).
- [39] M. Di Tullio, R. Rossignoli, M. Cerezo, and N. Gigena, "Fermionic entanglement in the Lipkin model," Phys. Rev. A 100, 062104 (2019).
- [40] J. Faba, V. Martín, and L. Robledo, "Correlation energy and quantum correlations in a solvable model," Phys. Rev. A 104, 032428 (2021), J. Faba, V. Martín, L. Robledo, "Analysis of quantum correlations within the ground state of a three-level Lipkin model", Phys. Rev. A 105, 062449 (2022).
- [41] A. T. Kruppa, J. Kovács, P. Salamon, Ö. Legeza, and G. Zaránd, "Entanglement and seniority," Phys. Rev. C 106, 024303 (2022).
- [42] Catren G., "Identity, individuality and indistinguishability in physics and mathematics," Philos. Trans. R. Soc. A. 381, 20220109 (2023).
- [43] S.M. Hengstenberg, C.E.P. Robin, M.J. Savage, "Multibody entanglement and information rearrangement in nuclear many-body systems: a study of the Lipkin -Meshkov-Glick model," Eur. Phys. J. A 59, 231 (2023).
- [44] E. Chitambar and G. Gour, "Quantum resource theories," Rev. Mod. Phys. 91, 025001 (2019).
- [45] D. Egloff, J.M. Matera, T. Theurer, and M.B. Plenio, "Of local operations and physical wires," Phys. Rev. X 8, 031005 (2018).
- [46] D.L.B. Ferreira, T.O. Maciel, R.O. Vianna, F. Iemini, "Quantum correlations, entanglement spectrum and coherence of two-particle reduced density matrix in the extended Hubbard model" Phys.Rev. B 105 115145 (2022).
- [47] G.G. Amosov and S.N. Filippov, "Spectral properties of reduced fermionic density operators and parity superselection rule," Quantum Inf. Process. 16, 2 (2017).
- [48] M. Sambataro and N. Sandulescu, "Pairing correlations and eigenvalues of two-body density matrix in atomic nuclei," Ann. Phys. 413, 168061 (2020).
- [49] N.L. Diaz, D. García-Martín, S. Kazi, M. Larocca and M. Cerezo, "Showcasing a barren plateau theory beyond the dynamical lie algebra," arXiv:2310.11505 (2023).
- [50] P. Ring and P. Schuck, The Nuclear Many-Body Problem (Springer, Berlin, Heidelberg, 1980).
- [51] T. Ando, "Properties of fermion density matrices," Rev.
- Mod. Phys. **35**, 690 (1963). [52] Here $\rho_{\alpha\alpha'}^{(M)} = \langle C_{\alpha'}^{(M)\dagger} C_{\alpha}^{(M)} \rangle$ is defined over linearly independent operators $C_{\alpha}^{(M)\dagger}$ creating normalized orthogonal M-particle states, such that its trace and eigenvalues differ by a factor 1/M! from those obtained with definitions covering all averages $\langle c_{i_1}^{\dagger} \dots c_{i_M}^{\dagger} c_{i'_1} \dots c_{i'_M} \rangle$ for arbitrary labels, where elements are then repeated M! times. The rank of $\rho^{(M)}$ remains nonetheless unchanged $\forall M$.
- [53] B. C. Carlson and J. M. Keller, "Eigenvalues of density

matrices," Phys. Rev. 121, 659 (1961).

- [54] R. Bhatia, *Matrix Analysis* (Springer, New York, 1997).
- [55] A. W. Marshall, I. Olkin, and B. C. Arnold, Inequalities: Theory of Majorization and its Applications (Springer, New York, 2011).
- [56] R. Rossignoli, N. Canosa, "Generalized entropic criterion for separability," Phys. Rev. A 66, 042306 (2002); "Violation of majorization relations in entangled states and its detection by means of generalized entropic forms". Phys. Rev. A 67, 042302 (2003); N. Canosa, R. Rossignoli, "Generalized nonadditive entropies and quantum entanglement", Phys. Rev. Lett. 88, 170401 (2002).
- [57] J. Hubbard, "Electron correlations in narrow energy bands," Proc. R. Soc. A 276, 238 (1963).
- [58] J.P.F. LeBlanc et al "Solutions of the two-dimensional Hubbard model: benchmarks and results from a wide range of numerical algorithms," Phys. Rev. X 5, 041041 (2015).
- [59] C.N. Yang, "Pairing correlations and eigenvalues of twobody density matrix in atomic nuclei," Rev. Mod. Phys. **34**, 694 (1962).
- [60] C.V. Kraus, M.M. Wolf, J.I. Cirac, and G. Giedke, "Pairing in fermionic systems: A quantum-information perspective," Phys. Rev. A 79, 012306 (2009).
- [61] J. Bardeen, L.N. Cooper, and J.R. Schrieffer, "Theory of superconductivity," Phys. Rev. 108, 1175 (1957).
- [62] P.W. Anderson and P. Morel, "Generalized Bardeen-Cooper-Schrieffer states and the proposed lowtemperature phase of liquid He³," Phys. Rev. 123, 1911 (1961).
- [63] D.J. Dean and M. Hjorth-Jensen, "Pairing in nuclear systems: from neutron stars to finite nuclei," Rev. Mod. Phys. 75, 607 (2003).
- [64] M.W. Zwierlein, J.R. Abo-Shaeer, A. Schirotzek, C.H. Schunck. W. Ketterle, "Vortices and superfluidity in a strongly interacting fermi gas," Nature 435, 1047 (2005).
- [65] A. A. Ovchinnikov, "On the exactly solvable pairing models for bosons," J. Stat. Mech. 2004, P07004 (2004).
- [66] J. Dukelsky and P. Schuck, "Condensate fragmentation in a new exactly solvable model for confined bosons," Phys. Rev. Lett. 86, 4207 (2001).
- [67] R.W. Richardson, "Exactly solvable many-boson model," J. Math. Phys. 9, 1327 (1968).
- [68] J. Von Delft and D.C. Ralph, "Spectroscopy of discrete energy levels in ultrasmall metallic grains," Phys. Rep. **345**, 61 (2001).
- [69] R. Rossignoli, N. Canosa, and P. Ring, "Thermal and quantal fluctuations for fixed particle number in finite superfluid systems," Phys. Rev. Lett. 80, 1853 (1998).
- [70] X.M. Puspus, K.H. Villegas, and F.N.C. Paraan, "Entanglement spectrum and number fluctuations in the spin-partitioned BCS ground state," Phys. Rev. B 90, 155123 (2014).
- [71] C. Dunning, J. Links, and H-Q. Zhou, "Ground-state entanglement of the BCS model," Phys. Rev. Lett. 94, 227002 (2005).
- [72] C.K. Chung and C.K. Law, "Negativity and momentum entanglement in Bardeen-Cooper-Schrieffer states at finite temperature," Phys. Rev. A 78, 034302 (2008).
- [73]S. Oh and J. Kim, "Entanglement of electron spins in superconductors," Phys. Rev. B 71, 144523 (2005).