



# On a non-linear moving boundary problem for a diffusion–convection equation

A.C. Briozzo<sup>a,b,\*</sup>, M.F. Natale<sup>a</sup>

<sup>a</sup> Depto. de Matemática, F.C.E., Universidad Austral, Paraguay 1950, S2000FZF Rosario, Argentina

<sup>b</sup> CONICET, Argentina

## ARTICLE INFO

### Article history:

Received 30 September 2011

Received in revised form

30 November 2011

Accepted 30 November 2011

Available online 13 December 2011

### Keywords:

Free boundary problem

Nonlinear diffusion–convection equation

Volterra integral equation

## ABSTRACT

We study a one-dimensional free boundary problem for a non-linear diffusion–convection equation whose diffusivity is heterogeneous in space as well as being non-linear. Under the Bäcklund transformation the problem is reduced to an associated free boundary problem. We prove the existence and uniqueness, local in time, of the solution by using the Friedman Rubinstein integral representation method and the Banach contraction theorem.

© 2011 Elsevier Ltd. All rights reserved.

## 1. Introduction

We study a non-linear free boundary problem for a semi-infinite region  $x > 0$  with a Dirichlet boundary condition at the fixed  $x=0$  given by condition (2). It is required to determine the evolution of the moving phase separation  $x=s(t)$  and the distribution  $\theta(x,t)$ . The modeling of this kind of systems is a problem with a great mathematical and industrial significance. Phase-change problems appear frequently in industrial processes and other problems of technological interest [1–10].

Owing to [13] we consider the following a free boundary problem with non-linear diffusion equation and a convective term

$$\frac{\partial \theta}{\partial t} = \frac{\partial}{\partial x} \left( D(\theta, x) \frac{\partial \theta}{\partial x} \right) - v(\theta) \frac{\partial \theta}{\partial x}, \quad 0 < x < s(t), \quad t > 0 \quad (1)$$

$$\theta(0, t) = g(t) \geq 0, \quad t > 0 \quad (2)$$

$$D(\theta(s(t), t), s(t)) \frac{\partial \theta}{\partial x}(s(t), t) = -\alpha s(t), \quad t > 0 \quad (3)$$

$$\theta(s(t), t) = 0, \quad t > 0, \quad s(0) = B \quad (4)$$

$$\theta(x, 0) = f(x) \geq 0, \quad 0 \leq x \leq B \quad (5)$$

where the velocity  $v(\theta)$  and the medium diffusivity  $D(\theta, x)$  are given by

$$v(\theta) = \frac{d}{2(a+b\theta)^2}, \quad D(\theta, x) = \frac{1+dx}{(a+b\theta)^2} \quad (6)$$

with positive parameters  $a, b, d$  and  $\alpha$ . This kind of non-linear conductivity or diffusion coefficients was considered in numerous papers, e.g. [11–21]. In [18] it was founded that the non-linear diffusion equation can be transformed to the linear form that possesses a well-known analytical solution when the soil water diffusivity is given by  $D(\theta) = a(b-\theta)^{-2}$ , where  $a$  and  $b$  are constant. Moreover, they argued that the diffusivity functional form is plausible for soil water diffusivities. The non-linear transport equation (1) arises in connection with unsaturated flow in heterogeneous porous media. If we set  $d=0$  and  $b=0$  in the free boundary problem (1)–(6) then we retrieve the classical one-phase Lamé–Clapeyron–Stefan problem.

We follow [13] where it was studied a analogous problem but the boundary conditions allowed to use the similarity method.

The goal of this paper is to prove the local existence and uniqueness in time of the solution to the problem given by (1)–(6). First, under the Bäcklund transformation, we reduce the problem to an associated free boundary problem and we prove that the problem is equivalent to solve a system of Volterra integral equations (38) and (39) [22,23] following Friedman–Rubinstein's method given in [24,25]. Then we prove that the system of equations (38)–(39) has a unique local solution by using the Banach contraction theorem.

\* Corresponding author at: Depto. de Matemática, F.C.E., Universidad Austral, Paraguay 1950, S2000FZF Rosario, Argentina.

E-mail addresses: [abriozzo@austral.edu.ar](mailto:abriozzo@austral.edu.ar) (A.C. Briozzo), [fnatale@austral.edu.ar](mailto:fnatale@austral.edu.ar) (M.F. Natale).

## 2. Statement of the problem

We consider the free boundary problem (1)–(5). We will make the following assumptions on the initial and boundary data:

- (i) Let  $a, b, d, \alpha \in \mathbb{R}^+$  with  $\alpha < a/b$ .
- (ii) Let  $f \in C^1[0, B]$  be a non-negative function with  $f'(0) > 0$ .
- (iii) Let  $g \in C^1[0, 1]$  be a non-negative function and  $D = \max_{t \in [0, 1]} g(t)$ .
- (iv) Compatibility conditions:  $f(0) = g(0)$  and  $f(B) = 0$ .

Taking into account (6) we can put our problem as

$$\frac{\partial \theta}{\partial t} = \frac{\partial}{\partial x} \left( \frac{1+dx}{(a+b\theta)^2} \frac{\partial \theta}{\partial x} + \frac{d}{2b(a+b\theta)} \right), \quad 0 < x < s(t), \quad t > 0 \quad (7)$$

$$\theta(0, t) = g(t), \quad t > 0 \quad (8)$$

$$\frac{1+ds(t)}{a^2} \frac{\partial \theta}{\partial x}(s(t), t) = -\alpha s(t), \quad t > 0 \quad (9)$$

$$\theta(s(t), t) = 0, \quad t > 0, \quad s(0) = B \quad (10)$$

$$\theta(x, 0) = f(x), \quad 0 \leq x \leq B \quad (11)$$

If we define the transformations in the same way as in [11,13,21]

$$\begin{cases} y = \frac{2}{d}[(1+dx)^{1/2}-1] \\ \bar{\theta}(y, t) = \theta(x, t) \end{cases} \quad (12)$$

we obtain the following associated free boundary problem:

$$\frac{\partial \bar{\theta}}{\partial t} = \frac{\partial}{\partial y} \left( \frac{1}{(a+b\bar{\theta})^2} \frac{\partial \bar{\theta}}{\partial y} \right), \quad 0 < y < \bar{s}(t), \quad t > 0 \quad (13)$$

$$\bar{\theta}(0, t) = g(t), \quad t > 0 \quad (14)$$

$$\frac{1}{a^2} \frac{\partial \bar{\theta}}{\partial y}(\bar{s}(t), t) = -\alpha \bar{s}(t), \quad t > 0 \quad (15)$$

$$\bar{\theta}(\bar{s}(t), t) = 0, \quad t > 0, \quad \bar{s}(0) = \bar{B} \quad (16)$$

$$\bar{\theta}(y, 0) = \bar{f}(y), \quad 0 \leq y \leq \bar{B} \quad (17)$$

where

$$\bar{B} = \frac{2}{d}[(1+dB)^{1/2}-1], \quad \bar{f}(y) = f\left(\frac{1}{d}\left[\left(\frac{d}{2}y+1\right)^2-1\right]\right) \quad (18)$$

and

$$\bar{s}(t) = \frac{2}{d}[(1+ds(t))^{1/2}-1] \quad (19)$$

is the free boundary.

By using the Bäcklund transformation we consider the new transformation

$$\begin{cases} y^* = y^*(y, t) = \int_{\bar{s}(t)}^y (a+b\bar{\theta}(\sigma, t)) d\sigma + (-\alpha b + a)\bar{s}(t) \\ \theta^*(y^*, t) = \frac{1}{a+b\bar{\theta}(y, t)} \end{cases} \quad (20)$$

In order to obtain an alternative expression for  $y^*$  we compute

$$\begin{aligned} \frac{\partial y^*}{\partial t} &= -(a+b\bar{\theta}(\bar{s}(t), t))\bar{s}'(t) + \int_{\bar{s}(t)}^y b \frac{\partial \bar{\theta}}{\partial t}(\sigma, t) d\sigma + (-\alpha b + a)\bar{s}'(t) \\ &= -\alpha b \bar{s}'(t) + \int_{\bar{s}(t)}^y b \frac{\partial}{\partial \sigma} \left( \frac{1}{(a+b\bar{\theta}(\sigma, t))^2} \frac{\partial \bar{\theta}}{\partial \sigma} \right) d\sigma \end{aligned}$$

$$\begin{aligned} &= -\alpha b \bar{s}'(t) + b \left( \frac{1}{(a+b\bar{\theta}(y, t))^2} \frac{\partial \bar{\theta}}{\partial y}(y, t) - \frac{1}{a^2} \frac{\partial \bar{\theta}}{\partial y}(\bar{s}(t), t) \right) \\ &= \frac{b}{(a+b\bar{\theta}(y, t))^2} \frac{\partial \bar{\theta}}{\partial y}(y, t) \\ &= \int_0^y \frac{\partial}{\partial \sigma} \left( \frac{b}{(a+b\bar{\theta}(\sigma, t))^2} \frac{\partial \bar{\theta}}{\partial \sigma}(\sigma, t) \right) d\sigma + \frac{b}{(a+b\bar{\theta}(0, t))^2} \frac{\partial \bar{\theta}}{\partial y}(0, t) \end{aligned} \quad (20)$$

$$\frac{\partial y^*}{\partial t} = b \int_0^y \frac{\partial \bar{\theta}}{\partial \sigma}(\sigma, t) d\sigma + \frac{b}{(a+b\bar{\theta}(0, t))^2} \frac{\partial \bar{\theta}}{\partial y}(0, t) \quad (21)$$

and therefore the new expression for  $y^*$  is given by

$$\begin{aligned} y^*(y, t) &= \int_0^t \left( \int_0^y \frac{\partial}{\partial \sigma} (a+b\bar{\theta}(\sigma, \tau)) d\sigma + \frac{b}{(a+b\bar{\theta}(0, \tau))^2} \frac{\partial \bar{\theta}}{\partial y}(0, \tau) \right) d\tau \\ &\quad + \int_0^y (a+b\bar{\theta}(\sigma, 0)) d\sigma + \frac{b}{(a+b\bar{\theta}(0, 0))^2} \frac{\partial \bar{\theta}}{\partial y}(0, 0) \\ &= \int_0^y (a+b\bar{\theta}(\sigma, t)) d\sigma + \int_0^t \frac{b}{(a+b\bar{\theta}(0, \tau))^2} \frac{\partial \bar{\theta}}{\partial y}(0, \tau) d\tau + M^* \end{aligned} \quad (22)$$

where

$$M^* = \frac{b}{(a+b\bar{\theta}(0, 0))^2} f'(0)$$

Now, applying (12) and (19) the problem (7)–(11) is transformed in a free boundary problem with a Dirichlet boundary condition given by

$$\frac{\partial \theta^*}{\partial t} = \frac{\partial^2 \theta^*}{\partial y^{*2}}, \quad y_0^*(t) < y^* < s^*(t), \quad t > 0 \quad (23)$$

$$\theta^*(y_0^*(t), t) = g^*(t), \quad t > 0, \quad y_0^*(0) = M^* \quad (24)$$

$$\frac{\partial \theta^*}{\partial y^*}(y_0^*(t), t) = -g^{*'}(t) y_0^{*'}(t), \quad t > 0 \quad (25)$$

$$\frac{\partial \theta^*}{\partial y^*}(s^*(t), t) = \alpha^* s^{*'}(t), \quad t > 0 \quad (26)$$

$$\theta^*(s^*(t), t) = \theta_f^*, \quad t > 0 \quad (27)$$

$$\theta^*(y^*, 0) = f^*(y^*), \quad M^* \leq y^* \leq B^* \quad (28)$$

$$s^*(0) = B^* \quad (29)$$

where

$$\alpha^* = \frac{\alpha b}{a(-\alpha b + a)}, \quad \theta_f^* = \frac{1}{a}, \quad B^* = (-\alpha b + a)\bar{B} \quad (30)$$

$$g^*(t) = \frac{1}{a+b\bar{g}(t)} \geq \frac{1}{a+bD}, \quad f^*(y^*) = \frac{1}{a+b\bar{f}(y)} \quad (31)$$

and

$$y_0^*(t) = y^*|_{y=0} = \int_0^t \frac{b}{(a+b\bar{g}(\tau))^2} \frac{\partial \bar{\theta}}{\partial y}(0, \tau) d\tau + M^* \quad (32)$$

$$s^*(t) = y^*|_{y=\bar{s}(t)} = (-\alpha b + a)\bar{s}(t) \quad (33)$$

are the free boundaries.

## 3. Existence and uniqueness of solutions

We have the following equivalence for the existence of solutions to the free boundary problem (23)–(29).

**Theorem 1.** The solution to the free boundary problem (23)–(29) is given by the following expression:

$$\begin{aligned} \theta^*(y^*, t) = & \int_{M^*}^{B^*} G(y^*, t; \xi, 0) f^*(\xi) d\xi + (\theta_f^* + \alpha^*) \int_0^t G(y^*, t; S^*(\tau), \tau) W(\tau) d\tau \\ & - \theta_f^* \int_0^t G_\xi(y^*, t; S^*(\tau), \tau) d\tau + \int_0^t G_\xi(y^*, t; y_0^*(\tau), \tau) g^*(\tau) d\tau \end{aligned} \quad (34)$$

and

$$y_0^*(t) = M^* - \int_0^t w(\tau) d\tau \quad (35)$$

$$S^*(t) = B^* + \int_0^t W(\tau) d\tau \quad (36)$$

where the functions  $w, W$  are defined by

$$w(t) = \frac{-1}{g^*(t)} \frac{\partial \theta^*}{\partial y^*}(y_0^*(t), t), \quad W(t) = \frac{1}{\alpha^*} \frac{\partial \theta^*}{\partial y^*}(S^*(t), t) \quad (37)$$

and they must satisfy the following system of two Volterra integral equations:

$$\begin{aligned} w(t) = & \frac{1}{g^*(t)} \left\{ \int_{M^*}^{B^*} N(y_0^*(t), t; \xi, 0) f^*(\xi) d\xi \right. \\ & + (\theta_f^* + \alpha^*) \int_0^t G_{y^*}(y_0^*(t), t; S^*(\tau), \tau) W(\tau) d\tau \\ & \left. - \int_0^t N(y_0^*(t), t; y_0^*(\tau), \tau) \dot{g}^*(\tau) d\tau \right\} \end{aligned} \quad (38)$$

$$\begin{aligned} W(t) = & \frac{2}{\alpha^* - \theta_f^*} \left\{ \int_{M^*}^{B^*} N(S^*(t), t; \xi, 0) f^*(\xi) d\xi \right. \\ & + (\theta_f^* + \alpha^*) \int_0^t G_{y^*}(S^*(t), t; S^*(\tau), \tau) W(\tau) d\tau \\ & \left. - \int_0^t N(S^*(t), t; y_0^*(\tau), \tau) \dot{g}^*(\tau) d\tau \right\} \end{aligned} \quad (39)$$

where  $G, N$  are the Green and Neumann functions and  $K$  is the fundamental solution to the heat equation, defined respectively by

$$G(x, t, \xi, \tau) = K(x, t, \xi, \tau) - K(-x, t, \xi, \tau) \quad (40)$$

$$N(x, t, \xi, \tau) = K(x, t, \xi, \tau) + K(-x, t, \xi, \tau) \quad (41)$$

$$K(x, t, \xi, \tau) = \begin{cases} \frac{1}{2\sqrt{\pi(t-\tau)}} \exp\left(-\frac{(x-\xi)^2}{4(t-\tau)}\right), & t > \tau \\ 0, & t \leq \tau \end{cases} \quad (42)$$

and  $y_0^*$  and  $S^*$  are given by (35) and (36) respectively.

**Proof.** Let  $\theta^*(y^*, t)$  be the solution to the problem (23)–(29). We integrate on the domain

$$D_{t,\varepsilon} = \{(\xi, \tau) / y_0^*(\tau) < \xi < S^*(\tau), \varepsilon < \tau < t - \varepsilon\} \quad (\varepsilon > 0)$$

the Green identity

$$(G\theta_\xi^* - \theta^* G_\xi)_\xi - (G\theta^*)_t = 0 \quad (43)$$

and we let  $\varepsilon \rightarrow 0$ , to obtain the integral representation for  $\theta^*(y^*, t)$  [24,25]

$$\begin{aligned} \theta^*(y^*, t) = & \int_{M^*}^{B^*} G(y^*, t; \xi, 0) f^*(\xi) d\xi + (\theta_f^* + \alpha^*) \int_0^t G(y^*, t; S^*(\tau), \tau) W(\tau) d\tau \\ & - \theta_f^* \int_0^t G_\xi(y^*, t; S^*(\tau), \tau) d\tau + \int_0^t G_\xi(y^*, t; y_0^*(\tau), \tau) g^*(\tau) d\tau \end{aligned} \quad (44)$$

by using the definitions of  $w(t)$  and  $W(t)$  given by (37). Moreover, if we differentiate (44) in variable  $y^*$  and we let  $y^* \rightarrow y_0^{*+}(t)$  and  $y^* \rightarrow S^{*-}(t)$ , by the jump relations [24], we obtain the system of integral equations (38) and (39) for  $w$  and  $W$ .

Conversely the function  $\theta^*(y^*, t)$  defined by (34), where  $w$  and  $W$  are the solutions of (38) and (39), satisfies the conditions (25), (26), (28) and (29). In order to prove the conditions (24) and (27) we define

$$\varphi_1(t) = \theta^*(S^*(t), t) - \theta_f^* \quad \text{and} \quad \varphi_2(t) = \theta^*(y_0^*(t), t) - g^*(t)$$

If we integrate the Green identity (43) over the domain  $D_{t,\varepsilon}$  ( $\varepsilon > 0$ ) and we let  $\varepsilon \rightarrow 0$ , we obtain that

$$\begin{aligned} \theta^*(y^*, t) = & \int_{M^*}^{B^*} G(y^*, t; \xi, 0) f^*(\xi) d\xi + \int_0^t G(y^*, t; S^*(\tau), \tau) \theta^*(S^*(\tau), \tau) W(\tau) d\tau \\ & + \int_0^t [G(y^*, t; S^*(\tau), \tau) \alpha^* W(\tau) - \theta^*(S^*(\tau), \tau) G_\xi(y^*, t; S^*(\tau), \tau)] d\tau \\ & + \int_0^t G(y^*, t; y_0^*(\tau), \tau) \theta^*(y_0^*(\tau), \tau) w(\tau) d\tau \\ & - \int_0^t [G(y^*, t; y_0^*(\tau), \tau) g^*(\tau) w(\tau) - \theta^*(y_0^*(\tau), \tau) G_\xi(y^*, t; y_0^*(\tau), \tau)] d\tau \end{aligned} \quad (45)$$

Then, if we compare this last expression (45) with (34) we deduce that

$$\begin{aligned} \int_0^t [G(y^*, t; S^*(\tau), \tau) W(\tau) - G_\xi(y^*, t; S^*(\tau), \tau)] \varphi_1(\tau) d\tau \\ + \int_0^t [G_\xi(y^*, t; y_0^*(\tau), \tau) - G(y^*, t; y_0^*(\tau), \tau) w(\tau)] \varphi_2(\tau) d\tau \equiv 0 \end{aligned} \quad (46)$$

We let  $y^* \rightarrow S^{*-}(t)$  and  $y^* \rightarrow y_0^{*+}(t)$  in (46) and we use the jump relations to obtain that  $\varphi_1$  and  $\varphi_2$  must satisfy the following system of Volterra integral equations:

$$\begin{aligned} \varphi_1(t) = & -2 \int_0^t \{ [G(S^*(t), t; S^*(\tau), \tau) W(\tau) + G_\xi(S^*(t), t; S^*(\tau), \tau)] \varphi_1(\tau) \\ & + [G_\xi(S^*(t), t; y_0^*(\tau), \tau) + G(S^*(t), t; y_0^*(\tau), \tau) w(\tau)] \varphi_2(\tau) \} d\tau \end{aligned} \quad (47)$$

$$\begin{aligned} \varphi_2(t) = & 2 \int_0^t \{ [G(y_0^*(t), t; S^*(\tau), \tau) W(\tau) - G_\xi(y_0^*(t), t; S^*(\tau), \tau)] \varphi_1(\tau) \\ & + [G_\xi(y_0^*(t), t; y_0^*(\tau), \tau) + G(y_0^*(t), t; y_0^*(\tau), \tau) w(\tau)] \varphi_2(\tau) \} d\tau \end{aligned} \quad (48)$$

Following [23] it is easy to see that there exists a unique solution  $\varphi_1 \equiv \varphi_2 \equiv 0$  to the system of Volterra integral equations (47)–(48). Then (24) and (27) holds.  $\square$

Next, we use the Banach fixed point theorem in order to prove the local existence and uniqueness of solution  $w, W \in C^0[0, \sigma]$  to the system of two Volterra integral equations (38) and (39) where  $\sigma$  is a positive small number to be determinate. We consider the Banach Space:

$$C_{R,\sigma} = \left\{ \vec{w}^* = \begin{pmatrix} w \\ W \end{pmatrix} / w, W : [0, \sigma] \rightarrow \mathbb{R}, \text{ continuous, with } \|\vec{w}^*\|_\sigma \leq R \right\}$$

where

$$\|\vec{w}^*\|_\sigma := \max_{t \in [0, \sigma]} |w(t)| + \max_{t \in [0, \sigma]} |W(t)|$$

We define the map  $F : C_{R,\sigma} \rightarrow C_{R,\sigma}$ , such that

$$\vec{w}^*(t) = F(\vec{w}^*(t)) = \begin{pmatrix} F_1(w(t), W(t)) \\ F_2(w(t), W(t)) \end{pmatrix}$$

where

$$F_1(w(t), W(t)) = \frac{1}{g^*(t)} \left\{ \int_{M^*}^{B^*} N(y_0^*(t), t; \zeta, 0) f^*(\zeta) d\zeta \right. \\ \left. + (\theta_f^* + \alpha^*) \int_0^t G_{y^*}(y_0^*(t), t; S^*(\tau), \tau) W(\tau) d\tau \right. \\ \left. - \int_0^t N(y_0^*(t), t; y_0^*(\tau), \tau) \dot{g}^*(\tau) d\tau \right\} \quad (49)$$

and

$$F_2(w(t), W(t)) = \frac{2}{\alpha^* - \theta_f^*} \left\{ \int_{M^*}^{B^*} N(S^*(t), t^*; \zeta, 0) f^*(\zeta) d\zeta \right. \\ \left. + (\theta_f^* + \alpha^*) \int_0^t G_{y^*}(S^*(t), t; S^*(\tau), \tau) W(\tau) d\tau \right. \\ \left. - \int_0^t N(S^*(t), t^*; y_0^*(\tau), \tau) \dot{g}^*(\tau) d\tau \right\} \quad (50)$$

**Lemma 2.** Let  $w, W \in C^0[0, \sigma]$ ,  $\max_{t \in [0, \sigma]} |w(t)| \leq R$ ,  $\max_{t \in [0, \sigma]} |W(t)| \leq R$  and  $2R\sigma \leq M^* < B^*/3$  then  $y_0^*$  and  $S^*$  defined by (35) and (36) satisfies

$$|y_0^*(t) - y_0^*(\tau)| \leq R|t - \tau|, \quad \forall \tau, t \in [0, \sigma]$$

$$\frac{M^*}{2} \leq y_0^*(t) \leq \frac{3M^*}{2}, \quad \forall t \in [0, \sigma]$$

$$|S^*(t) - S^*(\tau)| \leq R|t - \tau|, \quad \forall \tau, t \in [0, \sigma]$$

$$\frac{B^*}{2} \leq S^*(t) \leq \frac{3B^*}{2}, \quad \forall t \in [0, \sigma]. \quad \square \quad (51)$$

To prove the following lemmas we need the classical inequality:

$$\frac{\exp\left(\frac{-x^2}{\alpha(t-\tau)}\right)}{(t-\tau)^{n/2}} \leq \left(\frac{n\alpha}{2ex^2}\right)^{n/2}, \quad \alpha, x > 0, t > \tau, n \in \mathbb{N}. \quad (52)$$

**Lemma 3.** Let  $\sigma \leq 1$ . Under the hypothesis of Lemma 2 we have the following properties:

$$\int_{M^*}^B |f^*(\zeta)| |N(y_0^*(t), t; \zeta, 0)| d\zeta \leq \|f^*\| \quad (53)$$

$$\int_0^t |G_{y^*}(y_0^*(t), t; S^*(\tau), \tau) W(\tau)| d\tau \leq C_1(R, B^*, M^*)t \quad (54)$$

$$\int_0^t |\dot{g}^*(\tau)| |N(y_0^*(t), t; y_0^*(\tau), \tau)| d\tau \leq 2\sqrt{\frac{t}{\pi}} \|\dot{g}^*\| \quad (55)$$

$$\int_{M^*}^B |f^*(\zeta)| |N(S^*(t), t; \zeta, 0)| d\zeta \leq \|f^*\| \quad (56)$$

$$\int_0^t |G_{y^*}(S^*(t), t; S^*(\tau), \tau) W(\tau)| d\tau \leq C_2(R, B^*)\sqrt{t} \quad (57)$$

$$\int_0^t |\dot{g}^*(\tau)| |N(S^*(t), t^*; y_0^*(\tau), \tau)| d\tau \leq 2\sqrt{\frac{t}{\pi}} \|\dot{g}^*\| \quad (58)$$

where

$$C_1(R, B^*, M^*) = \frac{\sqrt{6}}{\sqrt{e\pi}} \left[ \frac{1}{(B^* - 3M^*)^2} + \frac{1}{(B^* + M^*)^2} \right] R$$

and

$$C_2(R, B^*) = \frac{R}{4\sqrt{\pi}} \left[ 2R + \frac{3}{B^{*2}} \left( \frac{2}{3e} \right)^{3/2} \right]$$

**Proof.** To prove (53) we consider

$$\int_{M^*}^B |f^*(\zeta)| |N(y_0^*(t), t^*; \zeta, 0)| d\zeta \leq \|f^*\| \int_0^\infty |N(y_0^*(t), t^*; \zeta, 0)| d\zeta \leq \|f^*\|$$

We have

$$|G_{y^*}(y_0^*(t), t^*; S^*(\tau), \tau)| \leq \frac{1}{4\sqrt{\pi}} \left\{ |y_0^*(t) - S^*(\tau)| \frac{\exp\left(\frac{-(y_0^*(t) - S^*(\tau))^2}{4(t-\tau)}\right)}{(t-\tau)^{3/2}} \right. \\ \left. + |y_0^*(t) + S^*(\tau)| \frac{\exp\left(\frac{-(y_0^*(t) + S^*(\tau))^2}{4(t-\tau)}\right)}{(t-\tau)^{3/2}} \right\}$$

and by using Lemma 2 and (52) we obtain

$$|G_{y^*}(y_0^*(t), t^*; S^*(\tau), \tau)| \leq \frac{\sqrt{6}}{\sqrt{e\pi}} \left[ \frac{1}{(B^* - 3M^*)^2} + \frac{1}{(B^* + M^*)^2} \right] \quad (59)$$

Then (54) holds.

To prove (55) by taking into account that

$$|N(y_0^*(t), t^*; y_0^*(\tau), \tau)| \leq \frac{1}{\sqrt{\pi(t-\tau)}}$$

so, we obtain

$$\int_0^t |\dot{g}^*(\tau)| |N(y_0^*(t), t^*; y_0^*(\tau), \tau)| d\tau \leq 2\sqrt{\frac{t}{\pi}} \|\dot{g}^*\|$$

The inequality (56) is proved in the same way as (53).

In [26] it was proved

$$|G_{y^*}(S^*(t), t; S(\tau), \tau)| \leq \frac{1}{4\sqrt{\pi}} \left[ \frac{R}{\sqrt{t-\tau}} + \frac{3}{B^{*2}} \left( \frac{2}{3e} \right)^{3/2} \right]$$

then we have (57).

The inequality (58) is proved in the same way as (55).  $\square$

**Lemma 4.** Let  $S_1^*$  and  $S_2^*$  be the functions corresponding to  $W_1$  and  $W_2$  in  $C^0[0, \sigma]$  respectively and  $y_{01}^*$  and  $y_{02}^*$  be the functions corresponding to  $w_1$  and  $w_2$  in  $C^0[0, \sigma]$  respectively with

$$\max_{t \in [0, \sigma]} |w_i(t)| \leq R, \quad \max_{t \in [0, \sigma]} |W_i(t)| \leq R,$$

then we have

$$\begin{cases} |S_2^*(t) - S_1^*(t)| \leq t \|W_2 - W_1\|_\sigma \\ |S_i^*(t) - S_i^*(\tau)| \leq R|t - \tau|, \quad i = 1, 2 \\ \frac{B^*}{2} \leq S_i^*(t) \leq \frac{3B^*}{2}, \quad \forall t \in [0, \sigma], i = 1, 2 \end{cases} \quad (60)$$

and

$$\begin{cases} |y_{01}^*(t) - y_{02}^*(t)| \leq t \|w_2 - w_1\|_\sigma \\ |y_{0i}^*(t) - y_{0i}^*(\tau)| \leq R|t - \tau|, \quad i = 1, 2 \\ \frac{M^*}{2} \leq y_{0i}^*(t) \leq \frac{3M^*}{2}, \quad \forall t \in [0, \sigma], i = 1, 2. \end{cases} \quad \square \quad (61)$$

**Lemma 5.** If we take  $R^2\sigma \leq 1$ ,  $\sigma \leq 1$  and  $2R\sigma \leq M^* < B^*/3$  then we have

$$\int_0^t |W_2(\tau)G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - W_1(\tau)G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))| d\tau \leq [P_3(M^*, B^*)\|W_1 - W_2\| + RP_1(M^*, B^*)\|w_1 - w_2\| + RP_2(M^*, B^*)\|W_1 - W_2\|]\sqrt{t} \quad (62)$$

$$\int_{M^*}^{B^*} |f^{*'}(\xi)| |N(y_{0_2}^*(t, t; \xi, 0) - N(y_{0_1}^*(t, t; \xi, 0))| d\xi \leq \frac{2\|f^{*'}\|}{\sqrt{\pi}} \|w_1 - w_2\| \sqrt{\sigma} \quad (63)$$

$$\int_0^t |g^*(\tau)| |N(y_{0_2}^*(t, t; y_{0_2}^*(\tau), \tau) - N(y_{0_1}^*(t, t; y_{0_1}^*(\tau), \tau))| d\tau \leq P_4(M^*, R)\|g^*\| \|w_1 - w_2\| \sqrt{t} \quad (64)$$

$$\int_{M^*}^{B^*} |f^{*'}(\xi)| |N(S_2^*(t, t; \xi, 0) - N(S_1^*(t, t; \xi, 0))| d\xi \leq \|f^{*'}\| \frac{2}{\sqrt{\pi}} \|W_1 - W_2\| \sqrt{t} \quad (65)$$

$$\int_0^t |W_2(\tau)G_{y^*}(S_2^*(t, t; S_2^*(\tau), \tau) - W_1(\tau)G_{y^*}(S_1^*(t, t; S_1^*(\tau), \tau))| d\tau \leq P_5(R, B^*)\|W_1 - W_2\| t \quad (66)$$

$$\int_0^t |g^*(\tau)| |N(S_2^*(t, t; y_{0_2}^*(\tau), \tau) - N(S_1^*(t, t; y_{0_1}^*(\tau), \tau))| d\tau \leq \|g^*\| P_6(M^*, B^*)(\|w_1 - w_2\| + \|W_1 - W_2\|)\sqrt{t} \quad (67)$$

where

$$P_1(M^*, B^*) = \frac{1}{\sqrt{\pi}e^{3/2}} \left[ \frac{\sqrt{6}(3B^* - M^*)^2}{16(B^* - 3M^*)^3} + \frac{27\sqrt{3}}{4} + \frac{12\sqrt{6}}{(B^* - 3M^*)^3} + \frac{6\sqrt{3}}{(B^* + M^*)^3} \right] \quad (68)$$

$$P_2(M^*, B^*) = \frac{12\sqrt{6}}{\sqrt{\pi}e^{3/2}} \left[ \frac{1}{(B^* - 3M^*)^3} + \frac{9}{8} + \frac{(3B^* - M^*)^2}{8(B^* - 3M^*)^3} + \frac{1}{(B^* + M^*)^2} \right] \quad (69)$$

$$P_3(M^*, B^*) = \frac{\sqrt{6}}{\sqrt{\pi}e} \left[ \frac{1}{(B^* - 3M^*)^2} + \frac{1}{(B^* + M^*)^2} \right] \quad (70)$$

$$P_4(M^*, B^*) = \frac{R^2}{\sqrt{\pi}} + \left(\frac{6}{e}\right)^{3/2} \frac{3R}{M^{*2}\sqrt{\pi}}$$

$$P_5(R, B^*) = \frac{1}{4\sqrt{\pi}} \left[ 6R + \frac{3}{B^{*2}} \left(\frac{2}{3e}\right)^{3/2} + \frac{6R}{B^{*2}} \left(\frac{6}{e}\right)^{3/2} \right] \quad (71)$$

$$P_6(M^*, B^*) = \frac{6^{3/2}}{\sqrt{\pi}e^{3/2}} \left[ \frac{3B^* - M^*}{(B^* - 3M^*)^3} + \frac{3}{(B^* + M^*)^2} \right] \quad (72)$$

**Proof.** To prove (62) we have

$$|W_2(\tau)G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - W_1(\tau)G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))| \leq |W_1(\tau) - W_2(\tau)| |G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))| + |W_1(\tau)| |G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))|$$

Taking into account that

$$|G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))|$$

$$\leq |G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_2^*(\tau), \tau))| + |G_{y^*}(y_{0_1}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))| \quad (73)$$

and by using the mean value theorem we have that there exists  $m = m(t)$  between  $y_{0_1}^*(t)$  and  $y_{0_2}^*(t)$  such that

$$|G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_2^*(\tau), \tau))| = |G_{y^*y^*}(m(t), t; S_2^*(\tau), \tau)| |y_{0_1}^*(t) - y_{0_2}^*(t)| = |G_t(m(t), t; S_2^*(\tau), \tau)| |y_{0_1}^*(t) - y_{0_2}^*(t)|$$

and there exists  $n = n(\tau)$  between  $S_1^*(\tau)$  and  $S_2^*(\tau)$  such that

$$|G_{y^*}(y_{0_1}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))| = |G_{y^*S_2^*}(y_{0_1}^*(t, t; n(\tau), \tau)| |S_1^*(\tau) - S_2^*(\tau)| = |N_\tau(y_{0_1}^*, t; n(\tau), \tau)| |S_1^*(\tau) - S_2^*(\tau)|.$$

Taking into account that

$$|G_t(m(t), t; S_2^*(\tau), \tau)| \leq \frac{1}{8\sqrt{\pi}(t-\tau)^{3/2}} \left\{ (m(t) - S_2^*(\tau))^2 \exp\left(\frac{-(m(t) - S_2^*(\tau))^2}{4(t-\tau)}\right) + (m(t) + S_2^*(\tau))^2 \exp\left(\frac{-(m(t) + S_2^*(\tau))^2}{4(t-\tau)}\right) \right\} + \frac{1}{4\sqrt{\pi}(t-\tau)^{3/2}} \left\{ \exp\left(\frac{-(m(t) - S_2^*(\tau))^2}{4(t-\tau)}\right) + \exp\left(\frac{-(m(t) + S_2^*(\tau))^2}{4(t-\tau)}\right) \right\}$$

and

$$|N_\tau(y_{0_1}^*, t; n(\tau), \tau)| \leq \frac{1}{8\sqrt{\pi}(t-\tau)^{3/2}} \left\{ (y_{0_1}^* - n(\tau))^2 \exp\left(\frac{-(y_{0_1}^* - n(\tau))^2}{4(t-\tau)}\right) + (y_{0_1}^* + n(\tau))^2 \exp\left(\frac{-(y_{0_1}^* + n(\tau))^2}{4(t-\tau)}\right) \right\} + \frac{1}{4\sqrt{\pi}(t-\tau)^{3/2}} \left\{ \exp\left(\frac{-(y_{0_1}^* - n(\tau))^2}{4(t-\tau)}\right) + \exp\left(\frac{-(y_{0_1}^* + n(\tau))^2}{4(t-\tau)}\right) \right\}$$

from (52) and Lemma 2 we have

$$|G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))| \leq P_1(M^*, B^*)t\|w_1 - w_2\|_t + P_2(M^*, B^*)\tau\|W_1 - W_2\|_t$$

Moreover, by (59)

$$|G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau)| \leq \frac{\sqrt{6}}{\sqrt{e}\pi} \left[ \frac{1}{(B^* - 3M^*)^2} + \frac{1}{(B^* + M^*)^2} \right] = P_3(M^*, B^*)$$

then, we obtain

$$|W_2(\tau)G_{y^*}(y_{0_2}^*(t, t; S_2^*(\tau), \tau) - W_1(\tau)G_{y^*}(y_{0_1}^*(t, t; S_1^*(\tau), \tau))| \leq P_3(M^*, B^*)\|W_1 - W_2\|_t + [P_1(M^*, B^*)t\|w_1 - w_2\|_t + P_2(M^*, B^*)\tau\|W_1 - W_2\|_t]R$$

and (62) hold.

Following [26] we obtain (63) and (65).

To prove (66) we use estimates obtained in [27].

To finish the thesis, we shall prove (67):

$$\int_0^t |g^*(\tau)| |N(S_2^*(t, t; y_{0_2}^*(\tau), \tau) - N(S_1^*(t, t; y_{0_1}^*(\tau), \tau))| d\tau \leq \|g^*\| \int_0^t \{ |N(S_2^*(t, t; y_{0_2}^*(\tau), \tau) - N(S_2^*(t, t; y_{0_1}^*(\tau), \tau))|$$

$$+ |N(S_2^*(t), t, y_{0_1}^*(\tau), \tau) - N(S_1^*(t), t, y_{0_1}^*(\tau), \tau)| \, d\tau \quad (74)$$

By the mean value theorem there exists  $c(\tau) \in (y_{0_1}^*(\tau), y_{0_2}^*(\tau))$  such that

$$\begin{aligned} & |N(S_2^*(t), t, y_{0_2}^*(\tau), \tau) - N(S_2^*(t), t, y_{0_1}^*(\tau), \tau)| \\ &= |N_\xi(S_2^*(t), t, c(\tau), \tau)| |y_{0_1}^*(\tau) - y_{0_2}^*(\tau)| \\ &\leq \frac{1}{4\sqrt{\pi}} \left\{ |S^*(t) - c(\tau)| \frac{\exp\left(\frac{-(S_2^*(t) - c(\tau))^2}{4(t-\tau)}\right)}{(t-\tau)^{3/2}} \right. \\ &\quad \left. + |S^*(t) + c(\tau)| \frac{\exp\left(\frac{-(S^*(t) + c(\tau))^2}{4(t-\tau)}\right)}{(t-\tau)^{3/2}} \right\} |y_{0_1}^*(\tau) - y_{0_2}^*(\tau)| \end{aligned}$$

then taking into account (52) and Lemma 2 we have

$$|N(S_2^*(t), t, y_{0_2}^*(\tau), \tau) - N(S_2^*(t), t, y_{0_1}^*(\tau), \tau)| \leq P_5(M^*, B^*) |y_{0_1}^*(\tau) - y_{0_2}^*(\tau)|$$

Furthermore, in a similar way we have that exists  $C(t) \in (S_1^*(t), S_2^*(t))$  such that

$$\begin{aligned} & |N(S_2^*(t), t, y_{0_1}^*(\tau), \tau) - N(S_1^*(t), t, y_{0_1}^*(\tau), \tau)| \\ &= |N_{y^*}(C(t), t, y_{0_1}^*(\tau), \tau)| |S_2^*(t) - S_1^*(t)| \leq P_6(M^*, B^*) |S_2^*(t) - S_1^*(t)| \end{aligned}$$

then (67) is obtained.  $\square$

**Theorem 6.** The map  $F: C_{R,\sigma} \rightarrow C_{R,\sigma}$  is well defined and it is a contraction map if  $\sigma$  satisfies the following inequalities:

$$\sigma \leq 1, \quad 2R\sigma \leq M^* < \frac{B^*}{3}, \quad R^2\sigma \leq 1 \quad (75)$$

$$H_1(\|\dot{g}^*\|_\sigma, a, b, B^*, \theta_f^*, \alpha^*, D, M^*, R, \sigma) \leq 1 \quad (76)$$

$$H_2(\|f^*\|, \|f^*\|, a, b, B^*, \theta_f^*, \alpha^*, D, M^*, R, \sigma) \leq 1 \quad (77)$$

where  $R$  is given by

$$R = 1 + \|f^*\| \left( a + bD + \frac{2}{|\alpha^* - \theta_f^*|} \right) \quad (78)$$

and

$$\begin{aligned} & H_1(\|\dot{g}^*\|_\sigma, a, b, B^*, \theta_f^*, \alpha^*, D, M^*, R, \sigma) \\ &= (a + bD) \left\{ (\theta_f^* + \alpha^*) C_1(B^*, R, M^*) \sigma + \frac{2}{\sqrt{\pi}} \|\dot{g}^*\|_\sigma \sqrt{\sigma} \right\} \\ &\quad + \left| \frac{1}{\alpha^* - \theta_f^*} \right| \left\{ (\theta_f^* + \alpha^*) \frac{R^2}{2\sqrt{\pi}} C_2(B^*, R) \sqrt{\sigma} + 4\sqrt{\frac{\sigma}{\pi}} \|\dot{g}^*\|_\sigma \right\} \quad (79) \end{aligned}$$

$$\begin{aligned} & H_2(\|\dot{g}^*\|_\sigma, a, b, B^*, \theta_f^*, \alpha^*, D, M^*, R, \sigma) \\ &= (a + bD) \left\{ \frac{2}{\sqrt{\pi}} \|f^*\| + (\theta_f^* + \alpha^*) \left( P_3(M^*, B^*) + RP_1(M^*, B^*) + \frac{RP_2(M^*, B^*)}{2} \right) \right. \\ &\quad \left. + \|\dot{g}^*\|_\sigma P_4(M^*, R) \right\} + \frac{2}{|\alpha^* - \theta_f^*|} \left\{ \frac{2\|f^*\|}{\sqrt{\pi}} + (\theta_f^* + \alpha^*) P_5(M^*, B^*) \right. \\ &\quad \left. + \|\dot{g}^*\|_\sigma P_6(M^*, B^*) \right\} \quad (80) \end{aligned}$$

Then there exists a unique solution on  $C_{R,\sigma}$  to the system of integral equations (38) and (39).

**Proof.** Firstly we demonstrate that  $F$  maps  $C_{R,\sigma}$  into itself, that is

$$\|F(\vec{w}^*)\|_\sigma = \max_{t \in [0, \sigma]} |F_1(w(t), W(t))| + \max_{t \in [0, \sigma]} |F_2(w(t), W(t))| \leq R$$

Taking into account Lemma 3 we have

$$|F_1(w(t), W(t))| \leq (a + bD) \left\{ \|f^*\| + (\theta_f^* + \alpha^*) C_1(B^*, R, M^*) t + 2\sqrt{\frac{t}{\pi}} \|\dot{g}^*\| \right\}$$

$$|F_2(w(t), W(t))| \leq \frac{2}{|\alpha^* - \theta_f^*|} \left\{ \|f^*\| + (\theta_f^* + \alpha^*) \frac{R}{4\sqrt{\pi}} C_2(B^*, R) \sqrt{t} + 2\sqrt{\frac{t}{\pi}} \|\dot{g}^*\| \right\}$$

and then

$$\|F(\vec{w}^*)\|_\sigma \leq \|f^*\| \left( a + bD + \frac{2}{|\alpha^* - \theta_f^*|} \right) + H_1(\|\dot{g}^*\|_\sigma, a, b, B^*, \theta_f^*, \alpha^*, D, M^*, R, \sigma)$$

where  $H_1$  is given by (79). Selecting  $R$  by (78) and  $\sigma$  such that (75) hold, we obtain  $\|F(\vec{w}^*)\|_\sigma \leq R$ . Now, we will prove that

$$\|F(\vec{w}_2^*) - F(\vec{w}_1^*)\|_\sigma \leq H_2(\|\dot{g}^*\|_\sigma, a, b, B^*, \theta_f^*, \alpha^*, D, M^*, R, \sigma) \|\vec{w}_2^* - \vec{w}_1^*\|_\sigma$$

where  $\vec{w}_1^* = (\vec{w}_1^1), \vec{w}_2^* = (\vec{w}_2^2) \in C_{R,\sigma}$ . Taking into account Lemma 5 we have

$$\begin{aligned} & \|F(\vec{w}_2^*) - F(\vec{w}_1^*)\|_\sigma = \max_{t \in [0, \sigma]} |F_1(w_2(t), W_2(t)) - F_1(w_1(t), W_1(t))| \\ &\quad + \max_{t \in [0, \sigma]} |F_2(w_2(t), W_2(t)) - F_2(w_1(t), W_1(t))| \\ &\leq \left\{ (a + bD) \left[ \frac{2}{\sqrt{\pi}} \|f^*\| + (\theta_f^* + \alpha^*) \left( P_3(M^*, B^*) + RP_1(M^*, B^*) + \frac{RP_2(M^*, B^*)}{2} \right) \right. \right. \\ &\quad \left. \left. + \|\dot{g}^*\|_\sigma P_4(M^*, R) \right] + \frac{2}{|\alpha^* - \theta_f^*|} \left[ \frac{2\|f^*\|}{\sqrt{\pi}} + (\theta_f^* + \alpha^*) P_5(M^*, B^*) \right. \right. \\ &\quad \left. \left. + \|\dot{g}^*\|_\sigma P_6(M^*, B^*) \right] \right\} \|\vec{w}_2^* - \vec{w}_1^*\|_\sigma \sqrt{\sigma} \\ &\leq H_2(\|\dot{g}^*\|_\sigma, a, b, B^*, \theta_f^*, \alpha^*, D, M^*, R, \sigma) \|\vec{w}_2^* - \vec{w}_1^*\|_\sigma \quad (81) \end{aligned}$$

By hypothesis (75)–(78) we have that  $F$  is a contraction.  $\square$

## 4. Conclusions

A one-dimensional free boundary problem is studied for a non-linear diffusion–convection equation whose diffusivity is heterogeneous in space as well as being non-linear. After using some non-linear transformations that simplify the transport equation to a form that is both linear and homogeneous in space, the transformed problem now has two free boundaries. It is shown that the problem is equivalent to a system of two integral equations. Then sufficient conditions for data was found applying the Banach contraction theorem, to prove existence and uniqueness, local in time of the solution.

## Acknowledgments

This paper has been partially sponsored by the Projects PIP No. 0460 of CONICET-UA and ANPCYT PICTO Austral 2008 No. 73, Rosario (Argentina). We appreciate the valuable suggestions by the anonymous referee which improve the paper.

## References

- [1] V. Alexiades, A.D. Solomon, Mathematical Modeling of Melting and Freezing Processes, Hemisphere, Taylor & Francis, Washington, 1983.
- [2] I. Athanassopoulos, G. Makrakakis, J.F. Rodrigues (Eds.), Free Boundary Problems: Theory and Applications, CRC Press, Boca Raton, 1999.
- [3] J.R. Cannon, The One-dimensional Heat Equation, Addison-Wesley, Menlo Park, 1984.

- [4] H.S. Carslaw, J.C. Jaeger, *Conduction of Heat in Solids*, Oxford University Press, London, 1959.
- [5] J.M. Chadam, H. Rasmussen (Eds.), *Free Boundary Problems Involving Solids*, Pitman Research Notes in Mathematics Series, vol. 281, Longman, Essex, 1993.
- [6] J. Crank, *Free and Moving Boundary Problems*, Clarendon Press, Oxford, 1984.
- [7] J.I. Diaz, M.A. Herrero, A. Liñan, J.L. Vazquez (Eds.), *Free Boundary Problems: Theory and Applications*, Pitman Research Notes in Mathematics Series, vol. 323, Longman, Essex, 1995.
- [8] A. Fasano, M. Primicerio (Eds.), *Nonlinear Diffusion Problems*, Lecture Notes in Mathematics, vol. 1224, Springer Verlag, Berlin, 1986.
- [9] N. Kenmochi (Ed.), *Free Boundary Problems: Theory and Applications*, I,II, Gakuto International Series: Mathematical Sciences and Applications, vols. 13 and 14, Gakkotosho, Tokyo, 2000.
- [10] V.J. Lunardini, *Heat Transfer with Freezing and Thawing*, Elsevier, Amsterdam, 1991.
- [11] A.C. Briozzo, M.F. Natale, D.A. Tarzia, The Stefan problem with the temperature-dependent thermal conductivity and a convective term with a convective condition at the fixed face, *Communications on Pure and Applied Analysis* 9 (5) (2010) 1209–1220.
- [12] M.F. Natale, D.A. Tarzia, Explicit solutions to the one-phase Stefan problem with temperature-dependent thermal conductivity and a convective term, *International Journal of Engineering Science* 41 (2003) 1685–1698.
- [13] C. Rogers, P. Broadbridge, On a nonlinear moving boundary problem with heterogeneity: application of reciprocal transformation, *Journal of Applied Mathematics and Physics (ZAMP)* 39 (1988) 122–129.
- [14] G. Bluman, S. Kumei, On the remarkable nonlinear diffusion equation, *Journal of Mathematical Physics* 21 (1980) 1019–1023.
- [15] A.C. Briozzo, M.F. Natale, D.A. Tarzia, Determination of unknown thermal coefficients for Storm's-type materials through a phase-change process, *International Journal of Non-Linear Mechanics* 34 (1999) 324–340.
- [16] P. Broadbridge, Non-integrability of non-linear diffusion–convection equations in two spatial dimensions, *Journal of Physics A—Mathematical and General* 19 (1986) 1245–1257.
- [17] P. Broadbridge, Integrable forms of the one-dimensional flow equation for unsaturated heterogeneous porous media, *Journal of Mathematical Physics* 29 (1988) 622–627.
- [18] J.H. Knight, J.R. Philip, Exact solutions in nonlinear diffusion, *Journal of Engineering Mathematics* 8 (1974) 219–227.
- [19] M.F. Natale, D.A. Tarzia, Explicit solutions for a one-phase Stefan problem with temperature-dependent thermal conductivity, *Bollettino dell'Unione Matematica Italiana* (8) 9-B (2006) 79–99.
- [20] R. Philip, General method of exact solution of the concentration-dependent diffusion equation, *Australian Journal of Physics* 13 (1960) 1–12.
- [21] M.F. Natale, D.A. Tarzia, The classical one-phase Stefan problem with temperature-dependent thermal conductivity and a convective term, *MAT-Serie* 15 (2008) 1–16.
- [22] G. Gripenberg, S.O. Londen, O. Staffans, *Volterra Integral and Functional Equations*, Cambridge University Press, Cambridge, 1990.
- [23] R.K. Miller, *Nonlinear Volterra Integral Equations*, W.A. Benjamin, Menlo Park, 1971.
- [24] A. Friedman, Free boundary problems for parabolic equations I. Melting of solids, *Journal of Mathematics and Mechanics* 8 (1959) 499–517.
- [25] L.I. Rubinstein, *The Stefan problem*, Translations of mathematical monographs, vol. 27, American Mathematical Society, Providence, 1971.
- [26] A.C. Briozzo, D.A. Tarzia, A one phase Stefan problem of a non-classical heat equation with a heat flux condition on the fixed face, *Applied Mathematics and Computation* 182 (2006) 809–819.
- [27] B. Sherman, A free boundary problem for the heat equation with prescribed flux at both fixed face and melting interface, *Quarterly of Applied Mathematics* 25 (1967) 53–63.