



Original article

## Native trees of the Northeast Argentine: Natural hosts of the *Cryptococcus neoformans*–*Cryptococcus gattii* species complex

Maria Emilia Cattana\*, María de los Ángeles Sosa, Mariana Fernández, Florencia Rojas, Magdalena Mangiaterra, Gustavo Giusiano

Departamento de Micología, Instituto de Medicina Regional, Universidad Nacional del Nordeste, Resistencia, Argentina

---

### ARTICLE INFO

**Article history:**

Received 11 September 2012

Accepted 4 June 2013

Available online 26 June 2013

**Keywords:**

*Cryptococcus gattii*

Ecological niche

Genotype

*Grevillea robusta*

*Peltophorum dubium*

*Tabebuia avellaneda*

---

### ABSTRACT

**Background:** In Argentina, information about epidemiology and environmental distribution of *Cryptococcus* is scarce. The city of Resistencia borders with Brazil and Paraguay where this fungus is endemic. All these supported the need to investigate the ecology of the genus and the epidemiology of cryptococcosis in this area.

**Aims:** The aim was to investigate the presence of species of *Cryptococcus neoformans*–*Cryptococcus gattii* complex and their genotypes in trees of the city of Resistencia.

**Methods:** One hundred and five trees were sampled by swabbing technique. The isolates were identified using conventional and commercial methods and genotyped by PCR-RFLP (Restriction Fragment Length Polymorphism).

**Results:** *Cryptococcus* was found in 7 out of the total trees. 6 out of 7 *Cryptococcus* isolates were identified as *C. neoformans* and one as *C. gattii*. *C. gattii* was isolated from *Grevillea robusta*. *C. neoformans* strains were isolated from *Tabebuia avellaneda* and *Peltophorum dubium*. Genotyping showed that all *C. neoformans* belonged to the VNI type and *C. gattii* belonged to the VGI type.

**Conclusions:** This represents the first study on the ecology of *Cryptococcus* spp. associated to trees from northeastern Argentina, and the first report describing *Grevillea robusta* as a host of members of this fungal genus. Another finding is the isolation of *C. neoformans* from *Tabebuia avellaneda* and *Peltophorum dubium*, both tree species native to northeastern Argentina.

© 2012 Revista Iberoamericana de Micología. Published by Elsevier España, S.L.U. All rights reserved.

## Árboles autóctonos del nordeste argentino. Huéspedes naturales de las especies del complejo *Cryptococcus neoformans*–*Cryptococcus gattii*

---

### RESUMEN

**Palabras clave:**

*Cryptococcus gattii*

Nicho ecológico

Genotipo

*Grevillea robusta*

*Peltophorum dubium*

*Tabebuia avellaneda*

**Antecedentes:** En Argentina la información sobre la epidemiología y la distribución ambiental de *Cryptococcus* es escasa.. Resistencia es una ciudad que limita con Brasil y Paraguay, donde este hongo es endémico. Esto apoya la necesidad de investigar la ecología de este género y la epidemiología de la criptococcosis en la región.

**Objetivos:** El objetivo del presente estudio fue investigar la presencia de especies del complejo *Cryptococcus neoformans* – *Cryptococcus gattii* y sus genotipos en árboles de la ciudad de Resistencia, situada en el nordeste argentino.

**Métodos:** Mediante la técnica del hisopo se tomaron muestras de 105 árboles. Los aislamientos se identificaron utilizando métodos convencionales y comerciales, y se genotipificaron mediante la prueba PCR-RFLP (Restriction Fragment Length Polimorphism).

**Resultados:** Se aisló *Cryptococcus* en 7 árboles. Se identificaron 6 aislamientos como *Cryptococcus neoformans* y uno como *Cryptococcus gattii*. Este último se aisló de *Grevillea robusta*. *Cryptococcus neoformans* se aisló de *Tabebuia avellaneda* y *Peltophorum dubium*. La genotipificación mostró que todos los aislamientos de *C. neoformans* pertenecían al tipo molecular VNI, y *C. gattii* al tipo molecular VGI.

---

\* Corresponding author.

E-mail addresses: [memiliacattana@yahoo.com.ar](mailto:memiliacattana@yahoo.com.ar), [emilycaty@hotmail.com](mailto:emilycaty@hotmail.com) (M.E. Cattana).

**Conclusiones:** El presente estudio es la primera investigación sobre la ecología del género *Cryptococcus* asociado a árboles del nordeste argentino, y la primera que describe *Grevillea robusta* como nicho ecológico de este género fúngico. Otro hallazgo es el aislamiento de *C. neoformans* de *Tabebuia avellaneda* y *Peltophorum dubium*, ambas especies de árboles originarias del nordeste argentino.

© 2012 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

The name *Cryptococcus* refers to a haploid and heterothallic group of encapsulated yeasts classified within the phylum Basidiomycota. Its teleomorph is *Filobasidiella*, a filamentous fungus belonging to the class Tremellomycetes.<sup>5</sup>

This genus is characterized by having a polysaccharide capsule that is its main factor of virulence. There are about 70 species of *Cryptococcus*, most of which live in the soil and are not harmful to humans.<sup>15</sup>

As currently accepted, the *Cryptococcus neoformans* complex contains two species that cause cryptococcosis in both humans and animals, a disease that causes significant morbidity and mortality in immunocompromised and immunocompetent patients.<sup>27,38</sup>

The differentiation into two species is due to genetic differences, the composition of the capsular polysaccharide structure, the teleomorph, the biochemical properties, the geographical distribution, the reservoir and the immunological state of the host to whom it affects.<sup>17,19,29,30,38</sup>

The species and varieties belonging to the *C. neoformans*–*C. gattii* complex are *C. gattii* (serotypes B and C) and *C. neoformans*, with its two varieties, *C. neoformans* var. *grubii* (serotype A) and *C. neoformans* var. *neoformans* (serotype D). There is also a hybrid of these two varieties, serotype AD. Molecular methods have allowed the discrimination of eight main genetic types: VNI and VNII (*C. neoformans* var. *grubii*, serotype A), VNIII (*C. neoformans* serotype AD), VNIV (*C. neoformans* var. *neoformans*, serotype D), VG1, VGII, VGIII, and VGIV (*C. gattii*, serotypes B and C).<sup>3,31</sup>

Since the infection is acquired by inhalation of infective propagules present in the environment, it is important to study the habitat of this fungus.<sup>38,40</sup>

Either one or both species of *Cryptococcus* have been found in different families and genera of trees from around the world.<sup>1,2,16,26,35,41</sup> *C. gattii* had been considered to be restricted to areas of tropical and subtropical climate, although there is evidence that it has expanded its distribution.<sup>6,11,16,18,19,24,42</sup>

In Argentina, although the incidence of cryptococcosis has increased 40- to 50-fold from the AIDS pandemic, the knowledge of the environmental distribution of *Cryptococcus* is scarce and epidemiological information is fragmented, scanty and based only on the communications from hospitals assisting HIV-positive patients.<sup>7,32,33</sup>

Currently *C. gattii* is endemic in Latin America. In Brazil, cryptococcosis by *C. gattii* is prevalent in immunocompetent patients, and is the 8th cause of meningitis, the majority of the cases related to children below 14 years of age.<sup>39,43</sup> Paraguay is among the Latin-American countries with high prevalence of this fungus, and outbreaks of infection with this agent have been reported in this country.<sup>8,24,25,43</sup> North-eastern Argentina borders with Brazil and Paraguay. Resistencia city is located in this region and the lack of knowledge about the presence of this fungus in the environment supports the need to investigate the ecology of the genus and the epidemiology of cryptococcosis in this zone.

The aim of this study was to investigate the presence and genotypes of members of the *Cryptococcus neoformans*–*Cryptococcus gattii* species complex in trees of the city of Resistencia.

## Materials and methods

### Study area

The study was performed in the city of Resistencia (27°27'05" S–58°59'12" W), located in northeastern Argentina. The city is located 51 m above sea level and is surrounded by rivers and lakes. The climate is subtropical with no dry season, with a mean annual rainfall of 1560 mm. The mean annual temperature is 21 °C, with extreme variations reaching 45 °C in summer and 0 °C in winter.<sup>37</sup>

Samples were taken from five parks, as well as from the Universidad Nacional del Nordeste campus and the Paseo de las Esculturas, all spaces that are centrally located in downtown. The 25 de Mayo Central Park has 4 ha, whereas the other four parks (Belgrano Park, España Park, 12 de Octubre Park, 9 de Julio Park) have only 1 ha. The latter are equidistant, about 600 m away from the central park. The campus is 1500 m to the south of the central park, whereas the Paseo de las Esculturas is 800 m to the northeast. The former has 5 ha whereas the latter has 0.50 ha (Fig. 1).

### Swabbing technique

Samples were collected by rubbing the inner of hollows or fissures of trees with a 5 mm of diameter sterile cotton-tipped swab moistened in sterile saline solution (0.85% NaCl) supplemented with chloramphenicol (10 mg/ml). Each swab was placed in a test tube with 3 ml of the solution.<sup>21</sup>

### Isolation


Without removing the swab, each tube was shaken manually for 5 min. The swab was then removed allowing the solution to settle for 10 min. The supernatant was diluted 1:10 in sterile saline solution. Then, 100 µl of the supernatant and 100 µl of the dilution were inoculated in separate plates of Pal medium<sup>34</sup> supplemented with biphenyl (0.1%). The inoculated plates were incubated at 32 °C.

### Biochemical and physiological identification of the isolates

After 48 h and up to 7 days, all brown colonies that grew on Pal medium were subcultured on Sabouraud medium. *Cryptococcus* isolates were identified according to the methodology of Kreger-van Rij<sup>20</sup> and with the commercial API ID32 method (bioMérieux, Marcy l'Etoile-France). *C. gattii* was identified by culturing the isolates on canavanine-glycine-bromothymol blue medium,<sup>23</sup> and by testing D-proline assimilation.<sup>10</sup>

### Molecular analysis

*Cryptococcus* isolates were genotyped by URA5 gene RFLP (Restriction Fragment Length Polymorphism) analysis. DNA extraction was performed according to the procedures described by Bosco Borgeat et al.<sup>4</sup> The PCR-RFLP reaction was performed as described by Meyer et al.<sup>31</sup> The PCR products were double digested with *Sau96I* (Fermentas) and *HhaI* (Fermentas) and separated in 3%



**Fig. 1.** Distribution of the seven parks sampled in Resistencia city.

agarose gel (Biodynamics, Argentina) electrophoresis at 100 V for 5 h. RFLP patterns were assigned visually by comparing them with the patterns obtained from the reference strains (*C. gattii*: CBS 10078 VG<sub>I</sub>; CBS 10082 VG<sub>II</sub>; CBS 10081 VG<sub>III</sub>; CBS 10101 VG<sub>IV</sub>. *C. neoformans* var. *grubii*: CBS 10085 VN<sub>I</sub>; CBS 10084 VN<sub>II</sub>. *C. neoformans* hybrid AD: CBS 10080 VN<sub>III</sub>. *C. neoformans* var. *neoformans*: CBS 10079 VN<sub>IV</sub>).

## Results

One hundred and five trees were sampled between March and December 2010. *Cryptococcus* was found in seven trees (6.7%).

Six out of the seven specimens were identified as *C. neoformans* (85.7%) and one as *C. gattii* (14.3%). All isolates were collected from different trees. Table 1 shows the tree species sampled and which of them showed the presence of *Cryptococcus*.

A second sampling was performed in positive trees obtaining the same RFLP genotypes.

Genotyping showed that all *C. neoformans* belonged to the variety *grubii* type VNI and that *C. gattii* belonged to the VGI type.

## Discussion

*Cryptococcus gattii* was first isolated by Ellis and Pfeiffer in 1990 from *Eucalyptus* trees in Australia and, for a long time, *Eucalyptus* spp. were pointed out as the ecological niche of *C. gattii*.<sup>12</sup> However, the isolation of *C. gattii* from *Eucalyptus* spp. outside Australia was considered as rare.<sup>41</sup> Our findings are consistent with the latter, since none of the *Eucalyptus* trees sampled showed the presence of *Cryptococcus* species. Affirming this, *C. gattii* has been found in 54 species of trees that are native to temperate, tropical and sub-tropical climates.<sup>41</sup> In the present study, *C. gattii* was isolated only from an old specimen of *Grevillea robusta*. This is the first report that refers to this species as a host for this fungus. There are many specimens of this species in the city of Resistencia and most of them are centenarians. The persistence of *C. gattii* in the same selected tree holes was confirmed by successive sampling. In agreement with other authors, this result shows that the ecological niche of *C. gattii* is not restricted to specific tree species.<sup>36,41</sup>

The natural habitat of *C. neoformans* has long been associated with nests of pigeons and other places with avian excreta.

**Table 1**  
Distribution of isolates and t

Tree species	n = 105	Specimens with positive culture	Species and genotypes of <i>Cryptococcus</i>
<i>Tipuana tipu</i>	49	1 <sup>a</sup>	<i>C. neoformans</i> var. <i>grubii</i> VNI
<i>Eucalyptus</i> sp.	10	–	–
<i>Tabebuia avellanedae</i>	8	1 <sup>b</sup>	<i>C. neoformans</i> var. <i>grubii</i> VNI
<i>Thuja</i> sp.	2	1 <sup>a</sup>	<i>C. neoformans</i> var. <i>grubii</i> VNI
<i>Grevillea robusta</i>	3	1 <sup>c</sup>	<i>C. gattii</i> VGI
<i>Ceiba insignis</i>	3	–	–
<i>Enterolobium contortisiliquum</i>	3	–	–
<i>Delonix regia</i>	1	–	–
<i>Bauhinia</i> sp.	2	–	–
<i>Ligustrum</i> sp.	2	–	–
<i>Peltophorum dubium</i>	8	3 <sup>c</sup>	<i>C. neoformans</i> var. <i>grubii</i> VNI
<i>Cinnamomum camphora</i>	1	–	–
<i>Taxodium distichum</i>	2	–	–
<i>Platanus</i> sp.	2	–	–
<i>Brachychiton</i> sp.	3	–	–
<i>Pinus</i> sp.	6	–	–

<sup>a</sup> España Park.

<sup>b</sup> Belgrano Park.

<sup>c</sup> 25 de Mayo Central Park.

accumulation.<sup>12,13,22,28</sup> However, the presence of this species has been confirmed in holes of tree trunks with obvious degradation of lignin.<sup>40</sup> In the present study, *C. neoformans* was found in *Tabebuia avellaneda* and *Peltophorum dubium*, both tree species native from Argentina. We found no references about these tree species as niche for any species of the genus *Cryptococcus*.

*Tipuana tipu*, a tree where *C. neoformans* was also found, is native from northwestern Argentina, in particular the Yungas and the subtropical forests of Bolivia. Because of its adaptability, *Tipuana tipu* has a wide distribution in Argentina, both by natural means and by the hand of man. In the city of Buenos Aires, it also has been reported harboring this yeast.<sup>36</sup>

The fact that all *C. neoformans* found belong to the VNI genotype is in agreement with reports from several countries indicating the VNI as the most frequent molecular type both as agent of cryptococcosis and in the environment.<sup>14,31,36</sup>

The *C. gattii* strain isolated from *Grevillea robusta* was type VGII. This finding agrees with that reported by Refojo et al. and Davel et al., who isolated *C. gattii*, though from other tree species in Buenos Aires city.<sup>9,36</sup> In other South American countries, such as Brazil and Colombia, the most common genotypes were VGII and VGIII.<sup>14,31,36,39,43</sup> This would demonstrate the geographic variation in the distribution of molecular types of *C. gattii*.

The study of the genetic relationship between environmental strains of *Cryptococcus* and those isolated from clinical specimens would be interesting to confirm that trees are a reservoir or source of infection.

This is the first study about ecological niches of *Cryptococcus* carried out in the northeast of Argentina. Further researches are necessary to find out which are the circulating genotypes in South America.

## Conflict of interest

The authors report no conflict of interest. The authors alone are responsible for the content and writing of the paper.

## Acknowledgement

Financial support for this work was provided by Fundación Alberto J Roemmers.

## References

- Bennett JE, Kwon-Chung KJ, Howard DH. Epidemiologic differences among serotypes of *Cryptococcus neoformans*. Am J Epidemiol. 1977;105:582–6.
- Boddy L, Frankland JC, West PV, editors. Ecology of saprotrophic basidiomycetes. British Mycological Society Symposium Series. 1st edn. London: Elsevier/Academic Press; 2008.
- Boekhout T, Theelen B, Diaz M, Fell JW, Hop WC, Abeln EC, et al. Hybrid genotypes in the pathogenic yeast *Cryptococcus neoformans*. Microbiology. 2001;147:891–907.
- Bosco Borgeat ME, Mazza M, Refojo N, Taverna M. Tipificación molecular de especies del género *Cryptococcus*. In: Merino LA, Giusiano G, editors. Manual de Métodos Moleculares para Estudios Microbiológicos. 1st edn. Buenos Aires: Asociación Argentina de Microbiología; 2011. p. 164–6.
- Bovers M, Hagen F, Boekhout T. Diversity of the *Cryptococcus neoformans*-*Cryptococcus gattii* species complex. Rev Iberoam Micol. 2008;25:S4–12.
- Bustamante Rufino B, Swine D. Aislamiento de *C. neoformans* var. *gattii* en dos pacientes peruanos. Rev Iberoam Micol. 1998;15:22–4.
- Canteros CE, Brudny M, Rodero L, Perrotta D, Davel G. Distribution of *Cryptococcus neoformans* serotypes associated with human infections in Argentina. Rev Argent Microbiol. 2002;34:213–8.
- Castañón Olivares LR, Arreguín Espinosa R, Ruiz Palacios, Santos G, López Martínez R. Frequency of *Cryptococcus* species and varieties in México and their comparison with some Latin American countries. Rev Latinoam Microbiol. 2000;42:35–40.
- Davel G, Abrantes R, Brudny M, Córdoba S, Rodero L, Canteros CE, et al. Primer aislamiento de *Cryptococcus neoformans* var. *gattii* en Argentina. Rev Argent Microbiol. 2003;35:110–2.
- Dufait R, Velho R, De Vroey C. Rapid identification of the two varieties of *Cryptococcus neoformans* by D-proline. Mykosen. 1987;30:483.
- Ellis DH. *Cryptococcus neoformans* var. *gattii* in Australia. J Clin Microbiol. 1987;25:430–1.
- Ellis DH, Pfeiffer TJ. Ecology, life cycle, and infectious propagule of *Cryptococcus neoformans*. Lancet. 1990;366:923–5.
- Ellis DH, Pfeiffer TJ. Natural habitat of *Cryptococcus neoformans* var. *gattii*. J Clin Microbiol. 1990;28:1642–4.
- Escandón P, Sánchez A, Martínez M, Meyer W, Castañeda E. Molecular epidemiology of clinical and environmental isolates of the *Cryptococcus neoformans* species complex reveals a high genetic diversity and the presence of the molecular type VGII mating type a in Colombia. FEMS Yeast Res. 2006;6:625–35.
- Fonseca A, Boekhout T, Fell JW. *Cryptococcus Vuillemin* (1901). In: Kurtzman CP, Fell JW, Boekhout T, editors. The yeasts, a taxonomic study. 5th edn. Elsevier B.V.; 2011. p. 1661–737.
- Gugnani HC, Mitchell TG, Litvintseva AP, Lengeler KB, Heitman J, Kumar A, et al. Isolation of *Cryptococcus gattii* and *Cryptococcus neoformans* var. *grubii* from the flowers and bark of eucaliptos trees in India. Med Mycol. 2005;43:565–9.
- Jarvis JN, Dromer F, Harrison T, Lortholary O. Managing cryptococcosis in the immunocompromised host. Curr Opin Infect Dis. 2008;21:596–603.
- Kidd SE, Hagen F, Tscharke RL, Huynh M, Bartlett KH, Fyfe M, et al. A rare genotype of *Cryptococcus gattii* caused the cryptococcosis outbreak on Vancouver Island (British Columbia, Canada). Proc Natl Acad Sci USA. 2004;101:17258–63.
- Kidd SE, Sorrell TC, Meyer W. Isolation of two molecular types of *Cryptococcus neoformans* var. *gattii* from insect frass. Med Mycol. 2003;41:171–6.
- Kreger-van Rij NJW, editor. The yeasts: a taxonomic study. Amsterdam: Elsevier Science Publishers; 1984.
- Krockenberger MB, Canfield PJ, Malik R. *Cryptococcus neoformans* in the koala (*Phascolarctos cinereus*): colonization by *C. neoformans* var. *gattii* and investigation of environmental sources. Med Mycol. 2002;40:263–72.
- Kwon-Chung KJ, Bennett J. Distribution of alpha and alpha mating types of *Cryptococcus neoformans* among natural and clinical isolates. Am J Epidemiol. 1978;4:337–9.
- Kwon-Chung KJ, Polacheck I, Bennett JE. Improved diagnostic medium for separation of *Cryptococcus neoformans* var. *neoformans* (serotypes A and D) and *Cryptococcus neoformans* var. *gattii* (serotypes B and C). J Clin Microbiol. 1982;15:535–7.
- Kwon-Chung KJ, Bennett JE. Epidemiologic differences between the two varieties of *Cryptococcus neoformans*. Am J Epidemiol. 1984;120:123–30.
- Kwon-Chung KJ. Cryptococcosis. In: Kwon-Chung KJ, Bennett JE, editors. Medical mycology. Philadelphia: Lea & Febiger; 1992. p. 397–446.
- Lazera MS, Cavalcanti MA, Trilles L, Nishikawa MM, Wanke B. *Cryptococcus neoformans* var. *gattii*—evidence for a natural habitat related to decaying wood in a pottery tree hollow. Med Mycol. 1998;36:119–22.
- Lin X, Heitman J. The biology of the *Cryptococcus neoformans* species complex. Annu Rev Microbiol. 2006;60:69–105.
- Littman ML, Borok R. Relation of the pigeon to cryptococcosis: natural carrier state, heat resistance and survival of *Cryptococcus neoformans*. Mycopathol Mycol Appl. 1968;35:922–33.
- Lizarazo-Niño J. Criptococosis en pacientes inmunocompetentes. Rev Neurol. 2005;40:5.
- Martín Mazuelos E, Aller García AI. Aspectos microbiológicos de la criptococosis en la era post-TARGA. Enferm Infect Microbiol Clin. 2010;28:40–5.
- Meyer W, Castañeda A, Jackson S, Huynh M, Castañeda E, IberoAmerican Cryptococcal Study Group. Molecular typing of IberoAmerican *Cryptococcus neoformans* isolates. Emerg Infect Dis. 2003;9:189–95.
- Mónaco LS, Tamayo Antabak N. Criptococosis en pacientes con SIDA: estudio de casos en el Hospital Paroissien en el período 1996–2007. Rev Argent Microbiol. 2008;40:218–21.
- Negrón R, Arechavala A, Robles AM, Bianchi M, Bava A, Helou S. Revisión clínica y evolución terapéutica de pacientes con criptococosis asociada al SIDA. Rev Iberoam Micol. 1995;12:12–5.
- Pal M, Mehrotra BS. Studies on the efficacy of sunflower seed agar for the isolation and identification of *Cryptococcus neoformans*. Arogya J Health Sci. 1982;8:74–9.
- Randhawa HS, Mussa AY, Khan ZU. Decaying wood in tree trunk hollows as a natural substrate for *Cryptococcus neoformans* and other yeast-like fungi of clinical interest. Mycopathologia. 2001;151:63–9.
- Refojo N, Perrotta D, Brudny M, Abrantes R, Hevia AI, Davel G. Isolation of *Cryptococcus neoformans* and *Cryptococcus gattii* from trunk hollows of living trees in Buenos Aires City, Argentina. Med Mycol. 2009;47:177–84.
- Rey W. Provincia del Chaco. In: Rey W, editor. Atlas total de la República Argentina. Buenos Aires: Centro editor de América Latina; 1992.
- Rippon JW. Criptococosis. In: Rippon JW, editor. Micología Médica. Hongos y Actinomicetos Patógenos. 3rd edn. México: Interamericana-McGraw-Hill; 1990. p. 629–59.
- Santos WR, Meyer W, Wanke B, Costa SP, Trilles L, Nascimento JL, et al. Primary endemic *Cryptococcosis gattii* by molecular type VGII in the state of Pará, Brazil. Mem Inst Oswaldo Cruz. 2008;103:813–8.

40. Sorrell TC, Ellis DH. Ecology of *Cryptococcus neoformans*. Rev Iberoam Micol. 1997;14:42–3.
41. Springer DJ, Chaturvedi V. Projecting global occurrence of *Cryptococcus gattii*. Emerg Infect Dis. 2010;16:14–20.
42. Torres-Rodríguez JM, Baró T, Morera Y, Alía C, López O, Hermoso de Mendoza M. Caracterización molecular de *Cryptococcus neoformans* var *gattii* causante de brotes epidémicos de criptococosis en cabras. Rev Iberoam Micol. 1999;16:164–5.
43. Trilles L, Lazéra Mdos S, Wanke B, Oliveira RV, Barbosa GC, Nishikawa MM, et al. Regional pattern of the molecular types of *Cryptococcus neoformans* and *Cryptococcus gattii* in Brazil. Mem Inst Oswaldo Cruz. 2008;103: 455–62.