

## Removal of cadmium by *Lactobacillus kefir* as a protective tool against toxicity

Esteban Gerbino<sup>1</sup>, Paula Carasi<sup>2</sup>, E. Elizabeth Tymczyszyn<sup>1</sup> and Andrea Gómez-Zavaglia<sup>1\*</sup>

<sup>1</sup> Center for Research and Development in Food Cryotechnology, CCT-CONICET La Plata, RA-1900, Argentina

<sup>2</sup> Laboratorio de Microbiología, Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Argentina

Received 21 October 2013; accepted for publication 3 March 2014; first published online 24 June 2014

The aim of this work was to evaluate the capacity of *Lactobacillus kefir* strains to remove cadmium cations and protect eukaryotic cells from cadmium toxicity. *Lb. kefir* CIDCA 8348 and JCM 5818 were grown in a 1/2 dilution of MRS broth supplemented with Cd(NO<sub>3</sub>)<sub>2</sub> ranging 0 to 1 mM. Growth kinetics were followed during 76 h at 30 °C by registering optical density at 600 nm every 4–10 h. The accumulated concentration of cadmium was determined on cultures in the stationary phase by atomic absorption. The viability of a human hepatoma cell line (HepG2) upon exposure to (a) free cadmium and (b) cadmium previously incubated with *Lb. kefir* strains was evaluated by determining the mitochondrial dehydrogenase activity. *Lb. kefir* strains were able to grow and tolerate concentrations of cadmium cations up to 1 mM. The addition of cadmium to the culture medium increased the lag time in all the concentrations used. However, a decrease of the total biomass (maximum Absorbance) was observed only at concentrations above 0.0012 and 0.0011 mM for strains CIDCA 8348 and JCM 5818, respectively. Shorter and rounder lactobacilli were observed in both strains upon microscopic observations. Moreover, dark precipitates compatible with intracellular precipitation of cadmium were observed in the cytoplasm of both strains. The ability of *Lb. kefir* to protect eukaryotic cells cultures from cadmium toxicity was analysed using HepG2 cells lines. Concentrations of cadmium greater than  $3 \times 10^{-3}$  mM strongly decreased the viability of HepG2 cells. However, when the eukaryotic cells were exposed to cadmium pre-incubated 1 h with *Lb. kefir* the toxicity of cadmium was considerably lower, *Lb. kefir* JCM 5818 being more efficient. The high tolerance and binding capacity of *Lb. kefir* strains to cadmium concentrations largely exceeding the tolerated weekly intake (TWI) of cadmium for food (2.5 µg per kg of body weight) and water (3 µg/l) addressed to human consumption, is an important added value when thinking in health-related applications.

**Keywords:** Cadmium, *Lb. kefir*, probiotics, bioaccumulation, HepG2 cells.

Cadmium is a toxic heavy metal with no reported biological functions. Its toxicity results from the coordination with biochemical ligands. As cadmium is considered a soft Lewis acid, it interacts easily with oxidisable soft ligands (i.e.: thiol groups) (Crichton, 2007). Complexation of cadmium with vital enzymes replaces physiological ions like zinc, calcium or nickel and inhibits the activity of enzymes that require active-site thiol groups (Carballo et al. 2012).

The main long-term effects of low-level cadmium exposure in humans are generally chronic obstructive pulmonary disease, emphysema and chronic renal tubular disease. Skeletal effects (i.e.: softening of bones resulting

from loss of minerals) and anaemia (resulting from cadmium interference with iron absorption) are also possible. Cadmium is transported by erythrocytes and albumin, and mainly stored in liver and kidneys, bound to metallothionein (50–75% of the body burden) (ATSDR, 2008).

Cadmium can occur as residue in food because of its environmental presence, as a result of human activities, including inadequate disposal electronic waste, mining or farming, and industrial production. Foodstuffs are the main source of cadmium exposure for the non-smoking general population (Satarug et al. 2010). According to the European Food Safety Association (EFSA), the tolerated weekly intake (TWI) for cadmium is 2.5 µg per kilogram of body weight. The TWI was defined as the estimated amount of cadmium that can be ingested weekly over a lifetime without

\*For correspondence; e-mail: angoza@qui.uc.pt

appreciable health risk (WHO 1989). The World Health Organization (WHO) established a guideline value for cadmium of 3 µg/l based on an allocation of 10% of the TWI to drinking-water (WHO 2006). The current average dietary exposure to cadmium for adults in Europe is close to the TWI, but in highly contaminated areas, it may be higher (EFSA 2012).

For this reason, all processes addressed to remove it are helpful. The capacity of different genera of lactic acid bacteria to adsorb heavy metals on their cell wall surface (biosorption) or accumulate them inside the cells (bioaccumulation) has gained interest in the last years and supports their use for several biotechnology processes (Halttunen et al. 2007; Mrvcic et al. 2012). In this regard, the GRAS (Generally Recognised as Safe) status of lactic acid bacteria and probiotics strengthens their relevance in cadmium removal, as they could be included in food matrices. This supports the application of probiotics in novel detoxification therapies (Urban & Kuthan, 2004; Zhai et al. 2013). In this context, Battikh et al. (2011) showed that a complex composed of *Lactobacillus acidophilus* and *Lb. bulgaricus* decreased the negative effects of cadmium on iron distribution in liver, kidney, leg and breast meat of chicken. It has also been published that the ability of some strains of *Streptococcus thermophilus* and *Lb. bulgaricus* to bind lead and cadmium has a beneficial effect against toxic heavy metals in rats (Salim et al. 2011). Furthermore, Bhakta et al. (2012) reported that the cadmium sorption by probiotic *Lb. reuteri*, may decrease the negative effects related to cadmium bioaccumulation in fish. Moreover, *Carnobacterium piscicola* and *Leuconostoc mesenteroides* subsp. Lysis are other species of lactic acid bacteria that can tolerate high concentrations of cadmium (Rial et al. 2011). Monachese (2012) reported a detailed analysis on the sequestration capacity of lead, cadmium and arsenic by different species of lactobacilli present in the gut microbiota (i.e.: *Lb. johnsonii*, *Lb. casei*, *Lb. plantarum* and *Lb. rhamnosus*).

*Lb. kefir* has well demonstrated probiotic properties (Golowczyc et al. 2007; Zheng et al. 2013) and is able to bind different heavy metals, including cadmium (biosorption) (Gerbino et al. 2011, 2012). In this work, we aim to get a deeper insight into the cadmium sequestering capacity of *Lb. kefir*. Strains of this species were incubated in presence of different concentrations of cadmium and the effect of this metal cation on growth (*lag* time and EC50) and morphology has been investigated. Finally, the capacity of *Lb. kefir* to protect HepG2 cells from cadmium toxicity was evaluated to support novel dietary strategies in detoxification of this harmful metal.

## Materials and methods

### Bacterial strains and growth conditions

Two strains of *Lb. kefir*, CIDCA 8348 isolated from kefir grains (Garrote et al. 2001), and JCM 5818, obtained from

the Japanese Collection of Microorganisms (Reiken, Japan), stored at – 80 °C, were used in this work. Frozen strains were activated under aerobic conditions in a 1/2 dilution of MRS broth (de Man et al. 1960) (Difco, Detroit, MI, USA) at 30 °C.

### Cadmium tolerance growth kinetics

A 2% w/v inoculum of *Lb. kefir* CIDCA 8348 or JCM 5818 ( $5 \times 10^8$  CFU/ml) was added to a 1/2 dilution of MRS broth placed in 96-well plates (Greiner Bio-one, Germany) and supplemented with concentrations of Cd(NO<sub>3</sub>)<sub>2</sub> ranging from 0 to 1 mM (Baker, Phillipsburg, New Jersey, USA). Strains were incubated 76 h at 30 °C. The optical density (OD) at 600 nm was determined every 4–10 h on a Synergy HT microplate reader (Bio-Tek Instruments, Winooski, Vermont, USA).

The increase of the *lag* time was determined as the difference between the *lag* time of *Lb. kefir* strains grown in presence of cadmium, and the *lag* time of strains grown in absence of cadmium (controls).

The total biomass obtained in each condition was referred to that of the controls in the stationary phase. The following equation was used:

$$\%OD = \frac{OD_{Cd^{2+}}}{OD_{ctrl}} \times 100 \quad (1)$$

where OD<sub>Cd<sup>2+</sup></sub> is the OD of microorganisms grown 55 h in presence of cadmium and OD<sub>ctrl</sub> is the OD of the controls in the stationary phase.

The effective concentration 50 (EC50), defined as the cadmium concentration that reduces 50% the biomass of a *Lb. kefir* culture in the stationary phase (maximum biomass obtained), was determined using GraphPad Prism 5 program (GraphPad Software Inc., San Diego, CA, 2007) as follows:

$$EC50 = \frac{\%OD + (\%OD_{max} - \%OD)}{1 + 10^{[(\log EC50 - X)S]}} \quad (2)$$

where EC50 is the effective concentration 50, %OD is the value obtained from Eq. (1), %OD<sub>max</sub> is the maximum value of %OD at 600 nm, X is the cadmium concentration and S is the slope of the plot.

### Bioaccumulation of cadmium

A 2% w/v inoculum of *Lb. kefir* strains was inoculated in 10 ml of 1/2 diluted MRS broth in presence of the EC50 for cadmium and incubated 48 h at 30 °C. Then, cells were washed twice in Milli Q water (Milli-Q plus; Millipore Cop., USA), harvested by centrifugation and pellets were resuspended in 1 ml cadmium-free ultrapure water. After centrifugation, cell pellets were collected and digested in concentrated HNO<sub>3</sub>-H<sub>2</sub>O<sub>2</sub> (García et al. 2006). The cadmium concentrations were determined in an atomic absorption spectrophotometer (SHIMADZU 6800). The values obtained were referred to the cadmium concentration added to the culture medium (100%). Results were expressed as percentage of bioaccumulation.

### Microscopy

For **optical microscopic** observations, microorganisms grown in presence of cadmium at the EC50 were fixed onto a microscope slide at 37 °C for 20 min and stained with crystal violet dye to enhance contrast in the microscopic image.

For **transmission electron microscopy** (TEM) observations, microorganisms grown in presence of cadmium at the EC50 were prepared according to Gerbino et al. (2011). Briefly, microorganisms were fixed in 20 g glutaraldehyde/l for 15 h at room temperature. Glutaraldehyde solution was prepared in phosphate-buffered saline (PBS) whose composition was: 0.144 g  $\text{KH}_2\text{PO}_4$ /l, 9 g  $\text{NaCl}$ /l, 0.795 g  $\text{Na}_2\text{HPO}_4$ /l, pH 7.2). Fixed cells were collected by centrifugation and washed three times with PBS. All samples were post-fixed with 10 g osmium tetroxide/l prepared in PBS, and dehydrated for 2 h. Samples were included in Epon 812 and sliced with a Sorvall MT 2B ultramicrotome. A JEM 1200 EX II transmission electron microscope (JEOL Ltd., Tokyo, Japan) was operated at 85 kV and photographed with a camera Erlangshen ES1000W, Model 785 (Gatan Inc., Pleasanton, California, USA) from Central Service Electronic Microscopy (Faculty of Veterinary Sciences, National University of La Plata). Microorganisms grown in 1/2 MRS broth without the addition of cadmium were used as controls.

### Eukaryotic cell cultures

HepG2 cells (human hepatoma cell line) were grown in Dulbecco's Modified Eagle's Medium (DMEM; Gibco BRL Life Technologies, Rockville, MD, USA) supplemented with 10% v/v inactivated (30 min/60 °C) foetal bovine serum (FBS, Gibco BRL Life Technologies, Rockville, MD, USA), 2 g  $\text{NaHCO}_3$ /l, 10 mg streptomycin/l and 10 IU penicillin G/ml and incubated at 37 °C in a 5% v/v  $\text{CO}_2$  atmosphere. The culture medium was changed every 2 d to attain a concentration of  $2 \times 10^5$  cells/ml. Cells were allowed to attach to the wells of polystyrene plates for 24 h before treatment with  $\text{Cd}(\text{NO}_3)_2$ , according to Fotakis & Timbrell (2006).

One millilitre of 48 h cultures of *Lb. kefir* CIDCA 8348 or JCM 5818 grown in 1/2 MRS (containing  $5 \times 10^8$  CFU/ml) was harvested and washed twice with ultra pure water (Milli-Q plus; Millipore Cop., USA). Bacterial pellets were resuspended in DMEM (pH=6) containing  $\text{Cd}(\text{NO}_3)_2$  at different concentrations of cadmium (0–0.03 mM) and incubated 1 h at 30 °C. Five hundred microlitres of the suspensions were added to HepG2 cell monolayers previously washed with PBS. Polystyrene plates were incubated 20 h at 37 °C in a 5% v/v  $\text{CO}_2$  atmosphere. Solutions of  $\text{Cd}(\text{NO}_3)_2$  ranging 0 to 0.03 mM were used as controls. A solution of 0.06 mM  $\text{KNO}_3$  was included to test the toxic effect of nitrates, and bacteria resuspended in DMEM that were not treated with  $\text{Cd}(\text{NO}_3)_2$  were used as negative controls.

### Mitochondrial dehydrogenase activity

The mitochondrial dehydrogenase activity of viable cells can be evaluated from the conversion of 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) to an insoluble purple formazan. The cell membranes are impermeable to the formazan product and therefore it accumulates in healthy cells. The MTT assay was performed as described by Mosmann (1983). Briefly, cells were incubated in serum free DMEM containing 0.8 mg MTT/ml for 4 h and washed with PBS. Dimethyl sulphoxide was added to remove formazan from cells and absorbance was recorded at 560 nm in a Synergy HT microplate reader (Bio-Tek Instruments, Winooski, Vermont, USA). Results were expressed as percentage of the viability of non-cadmium treated cells (100% viability).

### Statistical analysis


All the assays were conducted in triplicate, and average values were used for data analysis. The statistical data analysis was carried out using the statistical program GraphPad Prism 5 (GraphPad Software Inc., San Diego, CA, 2007).

## Results


The growth kinetics of *Lb. kefir* CIDCA 8348 and JCM 5818 in presence of different concentrations of  $\text{Cd}(\text{NO}_3)_2$  are shown in Fig. 1. Both *Lb. kefir* CIDCA 8348 (Fig. 1a) and JCM 5818 (Fig. 1b) were able to grow in presence of cadmium concentrations ranging 0 to 1 mM although with different yields in biomass and/or rates of growth. An increase of the lag time and a decrease of the maximum biomass and rate of growth (slope of the logarithmic phase) were observed as soon as the concentration of cadmium in the culture medium increased.

When plotting the increase of the lag time against the cadmium concentration in the culture medium, an abrupt increase of the lag time was observed for both strains at 0.0078 mM (Fig. 2). Below this value, the increase of the lag times for *Lb. kefir* JCM 5818 is lower than that of *Lb. kefir* CIDCA 8348, between 0.0078 and 0.25 mM, the behaviour is the opposite, and above 0.25 mM, strain *Lb. kefir* CIDCA 8348 again had a lower increase of the lag time.

The toxicity of cadmium cations was analysed in terms of the effective concentration 50 (EC50). In the context of this work, this parameter indicates the concentration of cadmium that inhibits 50% of the bacterial growth. A dose-response graph was constructed by plotting the %OD obtained after 55 h incubation (stationary phase, taken from Fig. 1a, b) v. the cadmium concentration in the culture media (Fig. 3). Using Eq. (2), the calculated EC50 was 0.0012 and 0.0011 mM for *Lb. kefir* CIDCA 8348 and JCM 5818, respectively (Table 1). These concentrations were then used for bioaccumulation assays and for microscopic




**Fig. 1.** Growth kinetics of *Lb. kefir*: CIDCA 8348 (a) and JCM 5818 (b) grown in presence of cadmium concentrations ranging 0–1 mM. ■ (0 mM); □ (0.0020 mM); ▽ (0.0078 mM); △ (0.0156 mM); ◆ (0.250 mM); ▲ (1 mM).



**Fig. 2.** Semilog plot showing the increase of the lag time vs the concentration of cadmium in the growth medium. The increase of the lag time was determined as the difference between the lag times of *Lb. kefir* incubated with a given concentration of cadmium and the lag time of strains incubated without cadmium (control). *Lb. kefir* CIDCA 8348 (◇); *Lb. kefir* JCM 5818 (●). Error bars indicate SD.

preparations. Table 1 also depicts the percentage of cadmium internalised by both strains, *Lb. kefir* JCM 5818 being the strain that incorporated it more efficiently.

Figure 4 shows the morphology of *Lb. kefir* CIDCA 8348 and JCM 5818 grown in presence or absence of cadmium



**Fig. 3.** Dose-response plot showing the percentage of total biomass (%OD) obtained for each condition of growth after 55 h incubation vs concentration of cadmium in the growth medium. %OD is referred to OD of strains grown in the absence of cadmium ( $OD_{max}$ ). *Lb. kefir* CIDCA 8348 (◇) and JCM 5818 (●).

**Table 1.** EC<sub>50</sub> and percentage of cadmium bioaccumulated by *Lb. kefir* CIDCA 8348 and JCM 5818

	<i>Lb. kefir</i>	
	CIDCA 8348	JCM 5818
EC <sub>50</sub> (mM)	0.012 ( $R^2$ 0.98)	0.011 ( $R^2$ 0.99)
Bioaccumulation (%)	72.5 ± 0.5	100 ± 0.5


cations. *Lb. kefir* CIDCA 8348 is an auto-aggregative strain (Fig. 4a) and although the cadmium treatment did not affect aggregation, rods became shorter and thicker (Fig. 4b). The non-aggregative strain *Lb. kefir* JCM 5818 (Fig. 4c) occurred as shorter rods after growing in presence of cadmium (Fig. 4d).

Electron microscopic pictures indicate that morphological features like shape and size were affected by the cadmium treatment (Fig. 5). Indeed, a slight shortening and rounding of bacterial cells was observed in both strains (Fig. 5b, d). Moreover, dark precipitates spread in the cytoplasm of both strains were noticed (Fig. 5b, d).

The ability of *Lb. kefir* to protect eukaryotic cells cultures from cadmium toxicity was analysed using HepG2 cells lines. Concentrations of cadmium higher than  $3 \times 10^{-3}$  mM strongly decreased the viability of HepG2 cells (Fig. 6). When the eukaryotic cells were exposed to cadmium pre-incubated with *Lb. kefir* strains, the toxicity of cadmium was considerably lower. *Lb. kefir* JCM 5818 was more efficient at protecting HepG2 cells as cadmium concentrations lower than  $3 \times 10^{-3}$  mM did not affect the viability of eukaryotic cells.

## Discussion

The resistance of microorganisms to heavy metal cations is very important in different scientific domains, namely


*Lb. kefir* CIDCA 8348*Lb. kefir* JCM 5818

**Fig. 4.** Microscopic observation (1000 $\times$ ) of *Lb. kefir* CIDCA 8348 and JCM 5818 grown in absence (a and c) and presence (b and d) of the EC50. Arrows indicate the regions where thickening of lactobacilli is more evident.


environmental and food sciences. Different mechanisms of detoxification have been described to explain the capacity of certain microorganisms to adsorb, tolerate and/or bioaccumulate toxic metals and prevent cellular damage (Cervantes et al. 2006; Halttunen et al. 2007; Mrvic et al. 2012). Detoxification mechanisms are particularly relevant in lactobacilli, as they may act as potential adjuncts for reducing the toxicity of cadmium in humans and animals.

The results obtained in this work showed for the first time that *Lb. kefir* strains can grow and tolerate cadmium concentrations up to 1 mM. Even when the addition of cadmium to the culture medium increased the lag time, only at concentrations above EC50, a decrease of the total biomass was observed (Figs. 1–3). The greater lag time indicates an increase in the time required by microorganisms for the adaptation to a harsh environment containing cadmium cations. Deleterious effects of cadmium include damage of DNA, inhibition of protein synthesis and competition for enzymatic reactive sites (Hynninen, 2010).

The microscopic observations reported in this work are compatible with intracellular precipitation of cadmium (Fig. 5). Different mechanisms have been proposed to explain the microbial tolerance to heavy metals. Most of them lead to reduced metal mobility, for example production of metal-binding peptides and proteins, organic or inorganic precipitation, intracellular metal sequestration, low bacterial transformation, and metal sorption by cell wall and other superficial structures (Adarsh et al. 2007; Panwichian et al. 2011). Formation of intracellular particles of cadmium was also reported for different lactobacilli and is a widespread phenomenon that could play a role in metal resistance because it decreases the harmful effect of cadmium. The formation of cadmium complexes with sulphur rich proteins or segregation into vesicles have been proposed as mechanisms (Prasad & Jha, 2010; Monachese, 2012). The morphological changes (Fig. 4) and cytoplasmic dark precipitates (Fig. 5) observed in both strains of *Lb. kefir* can be explained on this basis. Taking into account the probiotic properties of *Lb. kefir*, the high bioaccumulation capacity of both strains (Table 1) encourages their food use,



**Fig. 5.** Transmission electron microscopy of *Lb. kefir* CIDCA 8348 and JCM 5818 grown in absence (a and c) and presence (b and d) of the EC50. Arrows indicate precipitation of cadmium in the cytoplasm.



**Fig. 6.** Cytotoxicity of cadmium for HepG2 cells as measured by MTT assay. Percentage of viable cells vs cadmium concentration. HepG2 cells treated with suspensions of *Lb. kefir* CIDCA 8348 ( $\diamond$ ) or JCM 5818 ( $\bullet$ ) previously incubated with different concentrations of cadmium. HepG2 cells directly treated with different concentrations of cadmium ( $\blacksquare$ ). Error bars indicate sd.

thereby decreasing the concentration of free cadmium cations and preventing their toxicity in humans and animals.

HepG2 cells are generally used to investigate cadmium toxicity because, due to the high hepatotoxicity of cadmium, they provide a good approximation for subsequent studies in vivo (Fotakis & Timbrell, 2006). In our previous work, we reported that both *Lb. kefir* strains are able to detoxify water of cadmium (Gerbino et al. 2012). In this work, we showed that a prior interaction of cadmium with *Lb. kefir* decreases the toxicity of cadmium in HepG2 cells (Fig. 6). The higher efficiency of strain JCM 5818 is consistent with its major capacity to bioaccumulate cadmium cations (being a non-aggregative strain, a larger surface to interact with cadmium is available) (Table 1) and its higher binding capacity measured in mg cadmium per g dry biomass ( $q_{\max} = 61.6$  mg/g) (Gerbino et al. 2012). This way, a direct interaction of cadmium with the host microbiota containing different species of lactobacilli may result in a decrease of cadmium availability by immobilisation and/or uptake

of the metal. Zhai et al. (2013) reported that a strain of *Lb. plantarum* decreases the intestinal absorption of cadmium in mice and consequently, increases its faecal excretion in the first 24 h after acute oral exposure. The removal of cadmium by adsorbing it at the bacterial surface is one of the mechanisms proposed to explain this effect. In fact, the binding capacity is related to a protective effect against acute cadmium toxicity (Zhai et al. 2013). Both *Lb. kefir* strains used in this work are able to bind cadmium fast (after 1 h incubation) and efficiently (Gerbino et al. 2012). This, together with the capacity of lactobacilli to stimulate intestinal motility (Del Piano et al. 2010) supports the use of these strains to increase the faecal excretion of cadmium after acute exposure. The viability of eukaryotic cells exposed to *Lb. kefir* strains incubated with cadmium was not affected, indicating that adsorbed cadmium was not released in these experimental conditions.

Finally, it is noteworthy that the EC50 determined in this study (~3.55 mg/l) (Table 1) far exceeded the tolerated weekly intake of cadmium for food (EFSA, 2012) and water addressed for human consumption (3 µg/l) (WHO, 2006). The high tolerance and binding capacity of *Lb. kefir* strains to cadmium (even in concentrations that are incompatible with human life) is an important added value when thinking about health-related applications.

As consumption of harmful metals is not likely to decrease in the near future, the regular administration of formulations containing these strains appears to be an adequate and easy to implement strategy to prevent the harmful effects of cadmium in populations exposed to this toxic metal.

This work was supported by the Argentinean Agency for the Promotion of Science and Technology (Projects PICT/2008/145 and PICT/2011/0226), the Argentinean National Research Council (CONICET) (Project PIP 114-201101-00024). AGZ and EET are members of the research career CONICET (Argentina). EG and PC are postdoctoral and doctoral fellows from CONICET, respectively.

## References

- Adarsh VK, Mishra M, Chowdhury S, Sudarshan M, Thakur AR & Chaudhuri SR 2007 Studies on metal microbe interaction of three bacterial isolates from east Calcutta wetland. *Online Journal Biological Sciences* **7** 80–88
- Agency for Toxic Substances and Disease Registry (ATSDR) 2008 *Toxicological profile for Cadmium (Draft for Public Comment)*. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service. <http://www.atsdr.cdc.gov>. Agency for Toxic Substances and Disease Registry. Toxicological Profile for Cadmium
- Bhakta JN, Ohnishi K, Muneke Y, Iwasaki K & Wei MQ 2012 Characterization of lactic acid bacteria-based probiotics as potential heavy metal sorbents. *Journal of Applied Microbiology* **112** 1193–1206
- Battikh E, Safa A, Niccola MK & Aagha SI 2011 Effect of cadmium and *Lactobacillus* levels on iron concentration in different organs and meat of broiler chicken. *Journal of the University of Chemical Technology and Metallurgy* **46** 381–388
- Carballo R, Castiñeiras A, Domínguez-Martín A, García-Santos I & Niclós-Gutiérrez J 2012 Solid state structures of cadmium complexes with relevance for biological systems. In *Cadmium: From Toxicity to Essentiality. Series: Metal ions in Life Sciences*, Chapter 7, pp. 145–189 (Eds Sigel A, Sigel H & Sigel RKO). Springer ebooks.
- Cervantes C, Espino-Saldaña AE, Acevedo-Aguilar F, León-Rodríguez IL, Rivera-Cano ME, Avila-Rodríguez M, Wróbel-Kaczmarczyk K, Wróbel-Zasada K, Gutiérrez-Corona JF, Rodríguez-Zavala JS & Moreno-Sánchez R 2006 Interacciones microbianas con metales pesados. *Revista Latinoamericana Microbiología* **48** 203–210
- Crichton RR 2007 Metals in medicine and the environment. Cadmium. In *Biological Inorganic Chemistry. An Introduction*, Chapter 20, pp. 346–350. Oxford: Elsevier
- Del Piano M, Carmagnola S, Anderloni A, Andorno S, Ballarè M, Balzarini M, Montino F, Orsello M, Pagliarulo M, Sartori M, Tari R, Sforza F & Capurso L 2010 The use of probiotics in healthy volunteers with evacuation disorders and hard stools: a double-blind, randomized, placebo-controlled study. *Journal of Clinical Gastroenterology* **44** S30–34
- De Man JO, Rogosa M & Sharpe ME 1960 A medium for the cultivation of lactobacilli. *Journal of Applied Bacteriology* **23** 130–135
- European Food Safety Authority (EFSA) 2012 Cadmium dietary exposure in the European population. *EFSA Journal* **10** 2551. [37 pages; doi:10.2903/j.efsa.2012.2551. Available: [www.efsa.europa.eu/efsajournal](http://www.efsa.europa.eu/efsajournal)] [Accessed 5th October 2013]
- Fotakis G & Timbrell JA 2006 In vitro cytotoxicity assays: comparison of LDH, neutral red, MTT and protein assay in hepatoma cell lines following exposure to cadmium chloride. *Toxicology Letters* **160** 171–177
- García H, Senior A, El Zauahre M, Morán H, Acosta Y & Fernández N 2006 Análisis comparativo de dos técnicas de digestión para la determinación de metales pesados en lodos residuales. *Multiciencias* **6** 234–243
- Garrote GL, Abraham AG & De Antoni GL 2001 Chemical and microbiological characterisation of kefir grains. *Journal of Dairy Research* **68** 639–652
- Gerbino E, Mobili P, Tymczyszyn EE, Fausto R & Gómez-Zavaglia A 2011 FTIR spectroscopy structural analysis of the interaction between *Lactobacillus kefir* S-layers and metal ions. *Journal of Molecular Structure* **987** 186–192
- Gerbino E, Mobili P, Tymczyszyn EE, Frausto Reyes C, Araujo-Andrade C & Gómez-Zavaglia A 2012 Use of Raman spectroscopy and chemometrics for the quantification of metal ions attached to *Lactobacillus kefir*. *Journal Applied Microbiology* **112** 363–371
- Golowczyk MA, Mobili P, Garrote GL, Abraham AG & De Antoni GL 2007 Protective action of *Lactobacillus kefir* carrying S-layer protein against *Salmonella enterica* serovar Enteritidis. *International Journal of Food Microbiology* **118** 264–273
- Halttunen T, Salminen S & Tahvonen R 2007 Rapid removal of lead and cadmium from water by specific lactic acid bacteria. *International Journal Food Microbiology* **114** 30–35
- Hynninen A 2010 Zinc, cadmium and lead resistance mechanisms in bacteria and their contribution to biosensing. PhD Thesis. Finland: Department of Food and Environmental Sciences, Faculty of Agriculture and Forestry, University of Helsinki
- Monachese M 2012 Sequestration of lead, cadmium and arsenic by *Lactobacillus* species and detoxification potential. Master of Science Thesis. London, Ontario, Canada: University of Western Ontario
- Mosmann T 1983 Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* **65** 55–63
- Mrvic J, Stanzer D, Solic E & Stehlik-Tomas V 2012 Interaction of lactic acid bacteria with metal ions: opportunities for improving food safety and quality. *World Journal of Microbiology and Biotechnology* **28** 2771–2782
- Panwichian S, Kantachote D, Wittayaveerasak B & Mallavarapu M 2011 Removal of heavy metals by exopolymeric substances produced by resistant purple nonsulfur bacteria isolated from contaminated shrimp ponds. *Electronic Journal of Biotechnology* **14** 4

- Prasad K & Jha AK 2010 Biosynthesis of CdS nanoparticles: an improved green and rapid procedure. *Journal of Colloid Interface Science* **342** 68–72
- Rial D, Vázquez JA & Murado MA 2011 Effects of three heavy metals on the bacteria growth kinetics: a bivariate model for toxicological assessment. *Applied Microbiology and Biotechnology* **90** 1095–1109
- Salim AB, Badawy IH & Kassem SS 2011 Effect of lactic acid bacteria against heavy metals toxicity in rats. *Journal of American Science* **7** 264–274
- Satarug S, Garrett SH, Sens MA & Sens DA 2010 Cadmium, environmental exposure, and health outcomes. *Environmental and Health Perspectives* **118** 182–190
- Urban PL & Kuthan RT 2004 Application of probiotics in the xenobiotic detoxification therapy. *Nukleonika* **49** 43–45
- World Health Organization (WHO) 1989 Evaluation of Certain Food Additives and Contaminants. Thirty-third Report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series 776. Available: <http://www.who.int/ipcs/publications/jecfa/reports/en/index.html> [accessed 2nd October 2013]
- World Health Organization (WHO) 2006 *Guidelines for Drinking-water Quality*. Vol. 1, Recommendations–3rd ed. Available in: [http://www.who.int/water\\_sanitation\\_health/dwq/gdwq3rev/en/index.htm](http://www.who.int/water_sanitation_health/dwq/gdwq3rev/en/index.htm)
- Zhai Q, Wang G, Zhao J, Liu X, Tian F, Zhang H & Chen W 2013 Protective effects of *Lactobacillus plantarum* CCFM8610 against acute cadmium toxicity in mice. *Applied and Environmental Microbiology* **79** 1508–1515
- Zheng Y, Lu Y, Wang J, Yang L, Pan C & Huang Y 2013 Probiotic properties of *Lactobacillus* strains isolated from Tibetan kefir grains. *PLoS ONE* **8** e69868

*Journal of Dairy Research* (2014) **81** 287. © Proprietors of *Journal of Dairy Research* 2014  
doi:10.1017/S0022029914000351


## ERRATUM

### Removal of cadmium by *Lactobacillus kefir* as a protective tool against toxicity – ERRATUM

Esteban Gerbino, Paula Carasi, E. Elizabeth Tymczynsyn and Andrea Gómez-Zavaglia

doi:10.1017/S0022029914000314 Published Cambridge University Press, 24 June 2014

The y-axis of Fig. 6 should read % Viable cells (not Visible cells as printed).



#### Reference

Gerbino et al. 2014 Removal of cadmium by *Lactobacillus kefir* as a protective tool against toxicity. *Journal of Dairy Research* **81** 280–287