


Accepted Manuscript

Title: Oxfendazole flukicidal activity in pigs

Author: Pedro Ortiz Susana Terrones María Cabrera Cristian
Hoban Laura Ceballos Laura Moreno Candela Canton
Merixell Donadeu Carlos Lanusse Luis Alvarez



PII: S0001-706X(14)00107-7
DOI: <http://dx.doi.org/doi:10.1016/j.actatropica.2014.03.024>
Reference: ACTROP 3331

To appear in: *Acta Tropica*

Received date: 27-12-2013
Revised date: 17-2-2014
Accepted date: 28-3-2014

Please cite this article as: Ortiz, P., Terrones, S., Cabrera, M., Hoban, C., Ceballos, L., Moreno, L., Canton, C., Donadeu, M., Lanusse, C., Alvarez, L., Oxfendazole flukicidal activity in pigs, *Acta Tropica* (2014), <http://dx.doi.org/10.1016/j.actatropica.2014.03.024>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Oxfendazole flukicidal activity in pigs

¹Pedro Ortiz, ¹Susana Terrones, ¹María Cabrera, ¹Cristian Hoban, ²Laura Ceballos, ²Laura Moreno,
²Candela Canton, ³Meritxell Donadeu, ²Carlos Lanusse, ^{2*}Luis Alvarez

¹ Laboratorio de Inmunología, Facultad de Ciencias Veterinarias, Universidad Nacional de Cajamarca, Cajamarca, Perú.

²Laboratorio de Farmacología, Centro de Investigación Veterinaria de Tandil (CIVETAN), CONICET, Facultad de Ciencias Veterinarias, UNCPBA. Campus Universitario, 7000 Tandil, Argentina.

³GALVmed, Global Alliance for Livestock Veterinary Medicines, Edinburgh, Scotland, UK.

Running title: *Oxfendazole flukicidal activity in pigs*

*Corresponding Author: Laboratorio de Farmacología, Centro de Investigación Veterinaria de Tandil (CIVETAN), CONICET, Facultad de Ciencias Veterinarias, UNCPBA. Campus Universitario, 7000 Tandil, Argentina. TE: 54 249-4439850. e-mail address: lalvarez@vet.unicen.edu.ar

1 **Abstract**

2 Although oxfendazole (OFZ) is a well know broad-spectrum benzimidazole anthelmintic, the
3 assessment of its potential trematodicidal activity remains unexplored. OFZ administration at single
4 high doses has been recommended to control *Taenia solium* cysticercus in pigs. The current study
5 investigated the flukicidal activity obtained after a single high (30 mg/kg) oral dose of OFZ in pigs
6 harbouring a natural *Fasciola hepatica* infection. Sixteen (16) local ecotype pigs were randomly
7 allocated into two (2) experimental groups of 8 animals each named as follow: Untreated control and
8 OFZ treated, in which animals received OFZ (Synanthic[®], Merial Ltd., 9.06% suspension) orally at 30
9 mg/kg. At seven (7) days post-treatment, all the animals were sacrificed and direct adult liver fluke
10 counts were performed following the WAAVP guidelines. None of the animals involved in this
11 experiment showed any adverse event during the study. OFZ treatment as a single 30 mg/kg oral dose
12 showed a 100% efficacy against *F. hepatica*. In conclusion, the trial described here demonstrated an
13 excellent OFZ activity against *F. hepatica* in naturally infected pigs, after its administration at a single
14 oral dose of 30 mg/kg.

15

16 Key words: Oxfendazole, *Fasciola hepatica*, Efficacy, Pigs.

17

1 1. Introduction

2 Oxfendazole (OFZ) is a methyl-carbamate benzimidazole compound, indicated for the control of
3 gastrointestinal and lung nematodes and cestodes. OFZ was first marketed to be used in cattle, sheep
4 and horses. Additionally, OFZ has demonstrated activity against *Taenia solium* cysticercus in pigs after
5 its single oral administration at 30 mg/kg (Gonzales et al., 1996). OFZ treatment of *T. solium*-infected
6 pigs has been proposed as a tool to interrupt the transmission cycle of this parasite, protecting people
7 from neurocysticercosis, the most common parasitic infection of the human nervous system and the
8 most frequent preventable cause of epilepsy in endemic areas of Latin America, Africa and Asia
9 (Gonzales et al., 1996). The WHO estimated that globally, the parasite causes 50 million human cases
10 of taeniasis (infection with adult tapeworms) and cysticercosis, and 50,000 human deaths per year in
11 Africa, Asia and Latin America (WHO, 2012). In a recent study, OFZ orally administered to naturally
12 parasitized piglets at a single dose of 30 mg/kg was safe and highly efficacious (100%) against adult
13 stages of *Ascaris suum*, *Oesophagostomum* spp., *Trichuris suis* and *Metastrongylus* spp. (Alvarez et al.,
14 2013).

15
16 Fascioliasis, caused by the trematode liver fluke *Fasciola hepatica*, is the cause of considerable loss in
17 sheep and cattle production systems all over the world (Roberson and Courtney, 1995). Human
18 fascioliasis occurs as an accidental zoonotic disease in Africa, Western Europe and Latin America
19 (Mas-Coma et al., 2005). Considered a secondary zoonotic disease until the mid-1990s, human
20 fascioliasis is at present emerging or re-emerging in many countries, including increases of prevalence
21 and intensity and geographical expansion (Mas-Coma, 2005). Human fascioliasis is considered by the
22 WHO as an important human parasitic diseases, with estimates of 2.4 million (Rim et al., 1994) up to
23 17 million infected people (Hopkins, 1992). These figures could be even larger depending upon the
24 unknown situations in many countries, mainly in Asia and Africa (Mas-Coma, 2005). Sheep and cattle

1 are considered the main reservoir host species of *F. hepatica* (Hillyer et al., 1996; Mas-Coma et al.,
2 1997; 1999) but other species may provide a reservoir of infection. Pigs may play an important role in
3 fascioliasis transmission. For instance, fascioliasis has been reported in pigs from different
4 geographical areas, including Africa (El-Rafaie et al., 1984), Asia (Boes et al., 2000) and South
5 America (Mas-Coma et al., 1997). Valero and Mas-Coma (2000) found that there were no differences
6 in the infectivity of the metacercariae between sheep, cattle and pig isolates from the Bolivian
7 Altiplano, and in this geographical area *F. hepatica* adult development in the pig is similar to that
8 observed in sheep and cattle. The available epidemiological data suggest that pigs may play an
9 important role in fascioliasis transmission in hyperendemic areas and must, consequently, be taken into
10 account when applying control measures (Valero et al., 2001).

11

12 The main strategy for the effective control of fascioliasis is based on chemotherapy. However, there are
13 not anthelmintic compounds approved for the control of *F. hepatica* in pigs. The halogenated
14 benzimidazole triclabendazole has demonstrated to be active against liver flukes in pigs artificially
15 infected with *F. hepatica* metacercariae (Olaechea et al., 1989). However, at present triclabendazole is
16 not indicated to be used in pigs. In spite of its broad-spectrum activity against gastrointestinal
17 nematodes, OFZ is not indicated against *F. hepatica* (McKellar and Scott, 1990). However, some *in*
18 *vitro* ovicidal activity against *F. hepatica* eggs was previously reported (Alvarez et al., 2009), and it is
19 likely that a high dose of this benzimidazole compound may have flukicidal activity in domestic
20 animals, including pigs. In fact, in a recent study (Gomez-Puerta et al., 2012) performed in sheep,
21 flukicidal activity measured by the faecal egg count reduction test (FECRT), was reported after a single
22 OFZ dose. The potential of OFZ use in pigs at 30 mg/kg as a single oral dose against *F. hepatica*
23 should be investigated to search for a broader therapeutic indication for this anthelmintic compound,
24 when used as a single oral dose for the treatment of porcine cysticercosis. The goal of the current work

1 was to assess the flukicidal activity of a 9.06% OFZ suspension administered as a single oral dose (30
2 mg/kg) in pigs naturally infected with *F. hepatica*.

3

4

5

6 **2. Material and methods**

7 **2.1. Animals**

8 Local ecotype commercial pigs naturally parasitized with *F. hepatica* were involved in the current trial.
9 Pigs were fed *ad libitum* with a commercial balanced food and had free access to water. A 10 days
10 acclimatization period was allowed for the experimental animals. Animals were housed in two different
11 pens with concrete floors, protected from rain and prevailing winds, but without temperature control.
12 The animal phase of the current experiment was performed in the Faculty of Veterinary Science,
13 Universidad Nacional de Cajamarca, Cajamarca, Perú.

14

15 **2.2. Experimental design**

16 Sixteen pigs (28.0 ± 4.9 kg, 6-7 months old), naturally parasitized with *F. hepatica* were randomly
17 distributed into two groups (n= 8 each): Untreated control and OFZ treated group. Parasite infection
18 was confirmed by faecal egg counts (FEC) performed according Mooney et al., (2009). Briefly, 3 g of
19 faeces was added with water, and the contents was mixed thoroughly and poured through a tea strainer
20 to remove large debris. The filtrate was then allowed sediment for 10 min after which the supernatant
21 was siphoned off taking care not to disturb the sediment. The sediment was stained with two drops of
22 methylene blue and the entire sediment were assessed for liver fluke eggs using an optical microscope
23 (x40 magnification). The number of *F. hepatica* eggs observed was counted and the eggs per gram
24 (epg) of faecal material was calculated. In the OFZ treated group, treatment was performed by oral

1 administration of OFZ (Synanthic[®], OFZ 9.06%, Merial, France) at the dose of 30 mg/kg. Seven days
2 after treatment, animals were sacrificed and direct trematode counts of animals from the untreated
3 control and OFZ treated groups were performed following the WAAVP guidelines (Hennessy et al.,
4 2006). The total number of *F. hepatica* was recovered and counted according to Hennessy et al.,
5 (2006). Briefly, at necropsy, the gall bladder and liver of pigs was examined for living and dead *F.*
6 *hepatica*. After incising the gall bladder and bile ducts, the liver was cut along the large and small bile
7 ducts and hepatic veins and searched for flukes. Then the liver was cut into thin slices (0.5–1.0 cm in
8 width), soaked in warm (37 °C) saline in appropriate trays. The recovered liver flukes were transferred
9 to Petri dishes, filled with saline, and examined for vitality and appearance. The evaluation of adult
10 liver fluke burdens was done “blindly”. The OFZ efficacy was determined by the comparison of worm
11 burdens in treated versus untreated control animals. The following equation expresses the percentage of
12 efficacy (%E) against *F. hepatica* for the OFZ treated group (T) when compared with the untreated
13 control (C): %E= [(mean of S in C – mean of S in T) / mean of S in C] x 100. The geometric mean was
14 used as it most accurately represents the distribution of parasite populations within each group
15 (Hennessy et al., 2006). The criterion for efficacy was statistically significant differences in fluke
16 burdens between treated and untreated control groups and efficacy $\geq 90\%$. Nematode counts were
17 compared by non parametric unpaired test (Mann-Whitney) of log-transformed data. A value of $P < 0.05$
18 was considered statistically significant. The statistical analysis was performed using the Instat 3.0
19 Software (Graph Pad Software, CA, USA). Animal procedures and management protocols were carried
20 out in accordance with the Animal Welfare Policy (Act 087/02) of the Faculty of Veterinary Science,
21 Universidad Nacional de Cajamarca, Cajamarca, Peru.

22

1 3. Results

2 None of the animals involved in the current trial showed any adverse event during the study. This was
3 in agreement with a previously reported trial where the 9.06% OFZ formulation was orally
4 administered to pigs at 30, 90 and 150 mg/kg daily for three consecutive days, with a partial reduction
5 on feed intake during two to three days post-treatment in the groups treated with the highest OFZ doses
6 as the main clinical change on the health status of the treated pigs (Alvarez et al., 2012a). OFZ has a
7 large therapeutic index in the mammalian host, which explains the absence of appreciable toxic effect
8 in pigs treated with a single dose of 30 mg/kg. The individual FEC and adult liver fluke counts
9 obtained for the untreated control and OFZ treated group, and the efficacy observed after the OFZ
10 treatment are shown in Table 1.

11
12 A large variation in worm burdens was observed among experimental animals. In the control group, 4
13 to 140 adult flukes were recovered from the biliary duct. Furthermore, the FEC showed a large
14 variability ranging from 1 to 820 eggs/3 g of faeces. However, no statistical difference ($P > 0.05$) was
15 observed in FEC between the treated and untreated animals. Animal #4 in the untreated control group
16 had an unusually high egg count in 3 g of faecal material (820). The variation in liver fluke burdens
17 observed among experimental animals could be explained by the fact that they were obtained from
18 different farms, likely with different environmental contamination levels. Consequently, experimental
19 pigs could have been exposed to different metacercariae infection pressure. Even though the large
20 parasite infection variability observed in the current trial, a high efficacy (100%) against *F. hepatica* in
21 pigs was observed after the OFZ treatment at 30 mg/kg.

22

23

24

1 4. Discussion

2 Anthelmintic drugs are the main available tool to control parasitic infections, including liver flukes.
3 There are many compounds derived from different chemical families approved as flukicidal drugs for
4 use in sheep and cattle. However, available information on successful anthelmintic treatments against
5 liver flukes in pigs is scarce. Earlier work by Olaechea et al., (1989) demonstrated the activity of
6 triclabendazole at doses of 10 to 40 mg/kg, given to pigs at day 21 after the experimental infection with
7 100 *F. hepatica* metacercariae. However, no effective anthelmintic treatment is currently available for
8 removing liver flukes from pigs. The use of benzimidazole compounds as broad-spectrum
9 anthelmintics in all age groups of pigs is a common practice in different regions of the world
10 (Theodoropoulos et al., 2001; Beloeil et al., 2003). The oral OFZ single-dose therapy is a practical and
11 easy technique for pig deworming in extensive production systems, such as those observed in some
12 areas in developing countries. This OFZ treatment is useful for cysticercosis (Gonzales et al., 1997) and
13 adult nematode control (Alvarez et al., 2013). Additionally, the single OFZ oral administration at 30
14 mg/kg dose reached an outstanding flukicidal activity in the experimental work described here.

15
16 The benzimidazole anthelmintics active against *F. hepatica* include the methylcarbamate derivative
17 albendazole, and the halogenated thiol benzimidazole derivative, triclabendazole. The pro-
18 benzimidazole netobimin is also active against *F. hepatica*, but since it is metabolically converted to
19 albendazole in the rumen of treated sheep/cattle (Lanusse and Prichard, 1993), its activity must be
20 linked to albendazole. While albendazole activity is restricted to flukes older than 12 weeks (McKellar
21 and Scott, 1990), triclabendazole has been shown to have an excellent efficacy against both the mature
22 and immature adult stages of *F. hepatica* (Boray et al. 1983), which explain why this compound has
23 been the drug of choice for treating liver fluke infections in livestock for over 20 years. In previous
24 studies in sheep, OFZ at the oral dose of 5 and 15 mg/kg reached flukicidal efficacies of 14% and 20%,

1 respectively (Furmaga et al., 1982). This low efficacy could be related to a “pharmacodynamic
2 limitation” due to a lack drug-receptor affinity. However, a recent report demonstrated that none of the
3 sheep treated with OFZ (single oral dose, 30 mg/kg) showed *F. hepatica* eggs in faeces after 10 days of
4 treatment (Gomez-Puerta et al., 2012). The same oral dose was highly effective against adult flukes in
5 pigs, as show in the current reported trial. Thus, the limited efficacy observed in earlier efficacy trials
6 appears to be due to a pharmacokinetic-based limited parasite exposure and/or a lower OFZ receptor
7 affinity. As it has been described in nematodes (Alvarez et al., 2012b), the higher the OFZ
8 accumulation within the liver fluke, the greater the resultant clinical efficacy.

9
10 The increase on the albendazole dose was associated with enhancement in the plasma exposure of its
11 metabolites in sheep (Moreno et al., 2004; Alvarez et al., 2012b). In the same way, a 235% increment
12 in OFZ plasma concentration in pigs was observed at 5 days post-treatment after 90 mg/kg (5.7 ± 2.6
13 $\mu\text{g/mL}$) compared with 30 mg/kg dose ($1.7 \pm 1.1 \mu\text{g/mL}$; Alvarez et al, 2012a). It is clear that at least
14 under a certain dose range, the higher the OFZ dose given to pigs the greater the amount of drug
15 absorbed at the GI level. After OFZ treatment in pigs (30 mg/kg), an OFZ peak plasma concentration
16 of $5.40 \pm 0.65 \mu\text{g/mL}$ and an AUC of $209.9 \pm 33.9 \mu\text{g.h/mL}$ were reported (Moreno et al., 2012). This
17 OFZ plasma exposure is much larger than that observed in pigs, after the administration of the parent
18 thioether fenbendazole, which is rapidly and extensively converted *in vivo* into its active sulphoxide
19 metabolite OFZ (Lanusse and Prichard, 1993). After a fenbendazole oral dose (5 mg/kg) in pigs the
20 observed OFZ peak plasma concentration (C_{max}) and area under the plasma concentration vs time
21 curve (AUC) values were $0.66 \pm 0.22 \mu\text{g/mL}$ and $15.6 \pm 5.24 \mu\text{g.h/mL}$, respectively (Petersen and
22 Friis, 2000). The high OFZ plasma exposure observed after a single OFZ dose of 30 mg/kg in pigs
23 assures the trematode parasite being exposed to toxic drug concentrations for extended periods of time,

1 which may help to explain the high efficacy against *F. hepatica* in naturally infected pigs, described in
2 the current work.

3

4 **5. Conclusion**

5 OFZ administered as a single oral dose of 30 mg/kg to naturally parasitized pigs, was safe and highly
6 efficacious (100%) against adult stages of *F. hepatica*. The findings reported here may have a great
7 impact for liver fluke control in swine, particularly in those geographical areas where pigs play an
8 important role as reservoirs for human fascioliasis.

9

10 **6. Acknowledgments**

11 The authors would like to acknowledge Merial, France, who kindly provided the OFZ formulation
12 (Synanthic[®], OFZ 9.06%) used in the current work. The technical advice of Dr. Fermín Olaechea is
13 gratefully appreciated. This research was supported by the Agencia Nacional de Promoción Científica y
14 Tecnológica, Argentina and Consejo Nacional de Ciencia y Tecnología (CONCYTEC), Peru.

15

1 7. References

- 2 Alvarez, L., Moreno, G., Moreno, L., Ceballos, L., Shaw, L., Fairweather,
3 I., Lanusse, C., 2009. Comparative assessment of albendazole and triclabendazole ovicidal activity on
4 *Fasciola hepatica* eggs. *Vet. Parasitol.* 164, 211-216.
5
- 6 Alvarez, L., Domingue, G., Moreno, L., Ceballos, L., Bistoletti, M., Donadeu, M., Lanusse, C., 2012a.
7 Testing high oxfendazole doses to treat cysticercosis in pigs: a safety assessment. 12th International
8 Congress of the European Association for Veterinary Pharmacology and Toxicology, Noordwijk,
9 Netherlands.
10
- 11 Alvarez, L., Suarez, G., Ceballos, L., Moreno, L., Lanusse, C., 2012b. Dose-dependent systemic
12 exposure of albendazole metabolites in lambs. *J. Vet. Pharmacol. Therap.* 35, 365-372.
13
- 14 Alvarez, L., Saumell, C., Fusé, L., Moreno, L., Ceballos, L., Domingue, G., Donadeu, M., Dungu, B.,
15 Lanusse, C., 2013. Efficacy of a single high oxfendazole dose against gastrointestinal nematodes in
16 naturally infected pigs. *Vet. Parasitol.* 194, 70-74.
17
- 18 Beloeil, P.A., Chauvin, C., Fablet, C., Jolly, J.P., Eveno, E., Madec, F., Reperant, J.M., 2003.
19 Helminth control practices and infections in growing pigs in France. *Livest. Prod. Sci.* 81, 99-104.
20
- 21 Boes, J., Willingham, A., Fuhui, S., Xuguang, H., Eriksen, L., Nansen, P., Stewart, T., 2000.
22 Prevalence and distribution of pig helminths in the Dongting Lake Region (Hunan Province) of the
23 People's Republic of China. *J. Helminthol.* 74, 45-52.
24
- 25 Boray, J.C., Crowfoot, P.D., Strong, M.B., Allison, J.R., Schellenbaum, M., Von Orelli, M., Sarasin, G.,
26 1983. Treatment of immature and mature *Fasciola hepatica* infections in sheep with triclabendazole.
27 *Vet. Rec.* 113, 315-7.
28
- 29 El-Rafaie, S.A., Bassiouny, G.A., Marie, N.A.M. and Moris, E., 1984. Concomitant hepatic *Fasciola*
30 and hydatid infections in animals. *J. Egypt. Soc. Parasitol.* 14, 421-427.
31

- 1 Furmaga S, Gundalach JL, Sadzikowski A, Paciejewski S, 1982. (Systamex (oxfendazole) in the
2 treatment of parasitoses of sheep). Med. Weter. 38, 269-271.
3
- 4 Gonzalez, A.E., García, H.H., Gilman, R.H., Gavidia, C., Tsang, V.C., Bernal, T., Falcon, N., Romero,
5 M., Lopez-Urbina, M., 1996. Effective, single-dose treatment of porcine cysticercosis with
6 oxfendazole. Am. J. Trop. Med. Hyg. 54, 391-394.
7
- 8 Gonzalez, A.E., Falcon, N., Gavidia, C., Garcia, H. H., Tsang, V. C., Bernal, T., Romero, M., Gilman,
9 R.H., 1997. Treatment of porcine cysticercosis with oxfendazole: a dose-response trial. Vet. Rec. 141,
10 420-422.
11
- 12 Gomez-Puerta, L., Gavidia, L., Lopez-Urbina, M., Garcia, H.H., Gonzalez, A.E., 2012. Short Report:
13 Efficacy of a single oral dose of oxfendazole against *Fasciola hepatica* in naturally infected sheep. Am.
14 J. Trop. Med. Hyg., 86, 486-488.
15
- 16 Hennessy, D., Bauer, C., Boray, J., Conder, G., Dauschies, A., Johansen, M., Maddox-Hyttel, C.,
17 Roepstorff, A., 2006. World Association for the Advancement of Veterinary Parasitology
18 (W.A.A.V.P.): Second edition of guidelines for evaluating the efficacy of anthelmintics in swine. Vet.
19 Parasitol. 141,138-149.
20
- 21 Hillyer, G., Soler de Galanes, M., Buchón, P., Bjorland, J., 1996. Herd evaluation by enzyme-linked
22 immunosorbent assay for the determination of *Fasciola hepatica* infection in sheep and cattle from the
23 Altiplano of Bolivia. Vet. Parasitol. 61, 211-20.
24
- 25 Hopkins, D., 1992. Homing in on helminths. Am. J. Trop. Med. Hyg. 46, 626-634.
26
- 27 Lanusse, C., Prichard, R., 1993. Clinical pharmacokinetics and metabolism of benzimidazole
28 anthelmintics in ruminants. Drug Metab. Rev. 25, 235-279.
29

- 1 Mas-Coma, S., Rodriguez, M., Bargues, M., Valero, J., Coell, J., Gles, R., 1997. Secondary reservoir
2 role of domestic animal other than sheep and cattle in fascioliasis transmission in the Northern Bolivian
3 Altiplano. *Res. Rev. Parasitol.* 57, 39-46.
- 4
- 5 Mas-Coma, S., Anglés, R., Esteban, J.G., Bargues, M.D., Buchon, P., Franken, M., Strauss, W., 1999.
6 The northern Bolivian altiplano: a region highly endemic for human fascioliasis. *Trop. Med. Int. Health*
7 4, 454-467.
- 8
- 9 Mas-Coma, S., 2005. Epidemiology of fascioliasis in human endemic areas. *J. Helminthol.* 79, 207-
10 216.
- 11
- 12 Mas-Coma, S., Bargues, M., Valero, J., 2005. Fascioliasis and other plant-borne trematode zoonoses.
13 *Int. J. Parasitol.* 35, 1255-1278.
- 14
- 15 McKellar, Q., Scott, E., 1990. The benzimidazole anthelmintic agents-a review. *J. Vet. Pharmacol.*
16 *Therap.* 13, 223-247.
- 17
- 18 Mooney, L., Good, B., Hanrahan, J.P., Mulcahy, G., de Waal, T., 2009. The comparative efficacy of
19 four anthelmintics against a natural acquired *Fasciola hepatica* infection in hill sheep flock in the west
20 of Ireland. *Vet. Parasitol.* 164, 201-205.
- 21
- 22 Moreno, L., Echevarría, F., Muñoz, F., Alvarez, L., Sánchez, S., Lanusse, C., 2004. Dose-dependent
23 activity of albendazole against benzimidazole-resistant nematodes in sheep: relationship between
24 pharmacokinetics and efficacy. *Exp. Parasitol.* 106, 150-157.
- 25
- 26 Moreno, L., Lopez-Urbina, M.T., Farias, C., Domingue, G., Donadeu, M., Dungu, B., García, H.H.,
27 Gomez-Puerta, L.A., Lanusse, C., González, A.E., 2012. A high oxfendazole dose to control porcine
28 cysticercosis: Pharmacokinetics and tissue residue profiles. *Food Chem. Toxicol.* 50, 3819-3825.
- 29
- 30 Olaechea, F., Nansen, P., Christensen, N., 1989. Anthelmintic activity of triclabendazole against
31 *Fasciola hepatica* and *Echinostoma caproni* in mice and against *F.hepatica* in pigs. 14th Symposium,
32 Scandinavian Society of Parasitology, Helsingor, Denmark.

- 1 Petersen, M., Friis, C. 2000. Pharmacokinetic of fenbendazole following intravenous and oral
2 administration to pigs. *Am. J. Vet. Res.* 61, 573-576.
- 3
- 4 Rim, H., Farag, H., Sor Mai, S., Cross, H., 1994. Food-borne trematodes: ignored or emerging?
5 *Parasitol. Today.* 10, 207-209.
- 6
- 7 Roberson, E., Courtney, C., 1995. Anticestodal and antitrepatodal drugs. In: Adams, R., (Ed.)
8 *Veterinary Pharmacology and Therapeutics*, Iowa State University Press, Iowa, pp. 950-951.
- 9
- 10 Theodoropoulos, G., Theodoropoulou, E., Melissaropoulou, G., 2001. Worm control practices of pig
11 farmers in Greece. *Vet. Parasitol.* 97, 285-293.
- 12
- 13 Valero, M., Mas-Coma, S., 2000. Comparative infectivity of *Fasciola hepatica* metacercariae from
14 isolates of the main and secondary reservoir animal host species in the Bolivian Altiplano high human
15 endemic region. *Folia Parasitol. (Praha).* 47, 17-22.
- 16
- 17 Valero, M.A., Panova, M., Mas-Coma, S., 2001. Developmental differences in the uterus of *Fasciola*
18 *hepatica* between livestock liver fluke populations from Bolivian highland and European lowlands.
19 *Parasitol. Res.* 87, 337-342.
- 20
- 21 WHO, 2012. http://www.who.int/zoonoses/neglected_zoonotic_diseases/en/ (accessed 29.10.12).
- 22
- 23

1 **Table 1:** Individual pre-treatment (trial day -1) faecal egg counts (FEC), post-treatment fluke counts (trial day 7) and
 2 efficacy against *Fasciola hepatica*, obtained in naturally infected pigs from the untreated control group and the orally
 3 treated (30 mg/kg) with oxfendazole (OFZ).
 4
 5
 6

Animal #	FEC		Adult <i>F. hepatica</i> counts		FEC is expressed as egg in 3 g of faeces, estimated at trial day -1 (pre-treatment). No statistical difference (P>0.05, Mann-Whitney test) was observed
	Control	OFZ-treated	Control	OFZ-treated	
1	32	7	4	0	
2	2	1	4	0	
3	4	7	23	0	
4	820	13	140	0	
5	34	6	9	0	
6	14	48	5	0	
7	27	33	8	0	
8	1	85	8	0	
Geometric mean	15.1	12.3	8.3	0*	
Efficacy (%)	-	-	-	100	

21 between the Control and the OFZ treated group. *Statistically different (P< 0.05, Mann-Whitney test) adult fluke counts
 22 between the Control and the OFZ treated group.
 23
 24
 25
 26
 27

1 HIGHLIGHTS

- 2
- 3 Control measures in *F. hepatica*-endemic areas should include treatments in swine.
- 4 There are not available anthelmintic treatments to control *F. hepatica* in pigs.
- 5 The flukicidal activity of oxfendazole in pigs was assessed.
- 6 Oxfendazole (oral, 30 mg/kg) was 100% efficacious against *F. hepatica* in pigs.
- 7

Accepted Manuscript

Oxfendazole flukicidal activity in pigs

¹Pedro Ortiz, ¹Susana Terrones, ¹María Cabrera, ¹Cristian Hoban, ²Laura Ceballos, ²Laura Moreno, ²Candela Canton, ³Meritxell Donadeu, ²Carlos Lanusse, ²Luis Alvarez

¹ Laboratorio de Inmunología, Facultad de Ciencias Veterinarias, Universidad Nacional de Cajamarca, Cajamarca, Perú.

² Laboratorio de Farmacología, Centro de Investigación Veterinaria de Tandil (CIVETAN), CONICET, Facultad de Ciencias Veterinarias, UNCPBA. Campus Universitario, 7000 Tandil, Argentina.

³ GALVmed, Global Alliance for Livestock Veterinary Medicines, Edinburgh, Scotland, UK.

The current study investigated the flukicidal activity obtained after a single (30 mg/kg) oral dose of oxfendazole in pigs harbouring a natural *Fasciola hepatica* infection.

Animal Selection

(*Fasciola hepatica*-naturally infected pigs)




Trial design



Treatment



**Drug efficacy
assessment**



OXFENDAZOLE



Typical image of animal handling by the poorest people in Peru