

Solar UV-B radiation modifies the proportion of volatile organic compounds in flowers of field-grown grapevine (*Vitis vinifera* L.) cv. Malbec

Mariana Gil · Rubén Bottini · Mariela Pontin ·
Federico J. Berli · María Victoria Salomon ·
Patricia Piccoli

Received: 19 July 2013 / Accepted: 4 March 2014 / Published online: 14 March 2014
© Springer Science+Business Media Dordrecht 2014

Abstract Ultraviolet-B solar radiation (UV-B) is an environmental signal with biological effects in different plant tissues. Recent investigations reported dramatic changes of terpenes with a protective role in plant tissues submitted to biotic and abiotic stresses. This study examined the volatile organic compounds (VOCs) profile in flowers of *Vitis vinifera* L. cv. Malbec under filtered UV-B (or not). Gas chromatography–electron impact mass spectrometry analysis of flowers resulted in the identification of 12 VOCs, including eight sesquiterpenes, two aldehydes, and two ketones, being the oxygenated sesquiterpene farnesol the most abundant. The total amount of VOCs in flowers did not change irrespective UV-B had been filtered or not, suggesting those compounds have a protective role that is constitutive of the reproductive tissues. However UV-B increases the proportion of valencene, β -farnesene, α -panasinsene and hepatriacontanedione which would require further investigation.

Keywords Abiotic stress · Grape · Ultraviolet-B · Terpenes · VOCs

Introduction

Though ultraviolet-B radiation (UV-B; wavelength 280–315 nm) represents only a small portion of the total solar spectrum reaching the Earth's surface, it has large biological effects since it can activate the plant defense system leading to accumulation of secondary metabolites in different plant tissues (Teramura 2006), including grapevine leaves (Berli et al. 2010; Gil et al. 2012) and berries (Berli et al. 2011; Gil et al. 2013). Environmental UV-B levels are mainly regulated by season, latitude, altitude, time of day and cloudiness (McKenzie et al. 2003).

Plants synthesize and emit a large variety of volatile organic compounds (VOCs) with terpenoids and fatty-acid derivatives as the dominant classes (Pichersky and Gershenzon 2002). In grapevine, the evolution of these aromatic metabolites at the level of reproductive tissues has been explored mainly in grape berries of “floral” varieties like Muscat (Park et al. 1991; Coelho et al. 2006; Palomo et al. 2007). This is a biologically important aspect since VOCs can attract pollinators and may also protect reproductive tissues against pathogens (Pichersky and Gershenzon 2002; Lückner et al. 2004; Kegger and Pierik 2009). Different VOCs (especially monoterpenes and sesquiterpenes) with antimicrobial activity have been characterized in in vitro cultured grapevine (Escoriza et al. 2013). VOCs production in flowers of Arabidopsis may contribute to control bacterial infection and/or fungal infestations (Tholl et al. 2005). VOCs may also have a plant protective function towards abiotic stresses due to

Electronic supplementary material The online version of this article (doi:10.1007/s10725-014-9911-2) contains supplementary material, which is available to authorized users.

M. Gil · R. Bottini · M. Pontin · F. J. Berli ·
M. V. Salomon · P. Piccoli (✉)

Laboratorio de Bioquímica Vegetal, Instituto de Biología
Agrícola de Mendoza, Consejo Nacional de Investigaciones
Científicas y Tecnológicas-Universidad Nacional de Cuyo,
Almirante Brown 500, M5528AHB Chacras de Coria, Argentina
e-mail: ppiccoli@fca.uncu.edu.ar

M. Pontin
Estación Experimental Agropecuaria La Consulta, Instituto
Nacional de Tecnología Agropecuaria, CC8, 5567 La Consulta,
Argentina

their antioxidant activity that sequester free radicals so reducing oxidative damage (Aharoni et al. 2003; Wei and Shibamoto 2007) and inducing integrity and stability of membranes (Beckett et al. 2012). Therefore, one suspects that VOCs may reduce oxidative damage caused by UV-B in high altitude vineyard. If that occurs, VOCs will show considerably superior concentration in reproductive organs of grapes submitted to high UV-B.

In grapevine the pollen grains are site of terpene synthesis, which has been related with an increase in the activity of valencene synthase, enzyme responsible to produce sesquiterpenes in flowers of the cv. Cabernet Sauvignon (Martin et al. 2009). In the present work it was hypothesized that grapevine flowers increase the synthesis and/or accumulation of VOCs in response to UV-B in high altitude vineyards. The profile of VOCs in flowers of *Vitis vinifera* L. cv. Malbec field-grown plants from a high altitude vineyard exposed or not to relatively high ambient UV-B levels were monitored by gas chromatography–electron impact mass spectrometry (GC–EIMS).

Materials and methods

Plants of *V. vinifera* L. cv. Malbec from a high altitude vineyard at Gualtallary, Mendoza, Argentina (1450 m a.s.l., 69°15'W and 33°23'S) were exposed to two UV-B radiation regimens by covering the grapevines with specific plastic sheeting, from 15 days before flowering until sample collection. The experimental design and treatments were as previously reported in Berli et al. (2011). Solar UV-B radiation was cut-out to produce the minus UV-B treatment (–UV-B) using a clear polyester (100 µm), which absorbs more than 95 % of UV-B, and transmitted most of the solar radiation. Low density polyethylene (40 µm) transmitting most of the radiation from sunlight was used for the full UV-B treatment (+UV-B) and to minimize environmental differences between –UV-B and +UV-B treatments (no difference were found in air temperature). The plants in the +UV-B treatment were exposed to UV-B irradiances that reached up to 0.28 W m^{–2} during flowering (November). UV-B filtered and non-filtered plastics were positioned 2.5 m above ground level, covering the entire grapevine canopy. One inflorescence per experimental unit was sampled at flowering, stage 23 (Coombe 1995), collected at midday, immediately frozen with liquid nitrogen and kept at –80 °C until further analysis (n = 5).

Volatile organic compounds were determined according to Martin et al. (2009) with modifications. Ten flowers were randomly separated from inflorescences, weighed and immediately immersed in 0.5 mL of pentane in 1.5 mL glass vials with Teflon-coated screw caps, vortex-mixed vigorously and kept for 20 h at 4 °C in darkness. Then, aliquots of 1 µL of

the pentane fractions were injected in split–splitless mode into a Perkin-Elmer Elite-5MS, cross-linked methyl silicone capillary column (30 m length, 0.25 mm inner diameter, and 0.25 µm film thickness) fitted in a GC–EIMS (Clarus 500, PerkinElmer, Shelton, CT, USA). The GC program was: initial temperature at 45 °C for 1 min, followed by an increase of 2 °C min^{–1}–130 °C, then from 130–250 °C at a rate of 20 °C min^{–1} and held for 10 min at 250 °C. The ionization potential was 70 eV and a range of 40–500 atomic mass units was scanned. The identities of compounds were confirmed by comparison of their retention times and full scan mass spectra with those of authentic standards, and with mass spectra of the National Institute of Standards and Technology (NIST) library. Peak areas were referred to *n*-hexadecane standard for quantification of compounds.


The statistical evaluation was performed using the software Statgraphics Centurion XVI version 15.0.10 (Statpoint Technologies Inc., Warrenton, VA, USA). One-way ANOVA was used to test the variation in concentration of VOCs from flowers exposed to two UV-B radiation regimens. Significant differences were considered at probability of $P \leq 0.1$. Results are reported as a mean of five independent replicated assays with their standard error (SE).

Results and discussion

The chemical structures of the volatile compounds characterized by GC–EIMS in flowers are presented in Fig. 1 (for full mass spectra see Figures 1S–3S of Supplementary Material). Flowers emitted a mixture of compounds belonging to four chemical classes: sesquiterpenes, oxygenated sesquiterpenes, aliphatic aldehydes and ketones. The GC–EIMS analysis resulted in the identification of 12 compounds: the hydrocarbons sesquiterpenes (C₁₅H₂₄) α - β -farnesene, α - β -caryophyllene, valencene, α -panasinsene, and the oxygenated sesquiterpenes (C₁₅H₂₆O) nerolidol and farnesol; the aldehydes pentadecanal and 7, 10, 13-hexadecatrienal; and the ketones pentadecanone and 6, 28-heptatriacontadien-2-one (Table 1).

Irrespective of the UV-B treatments, the sesquiterpenes were the most abundant with an average of 81 % of the total while ketones represented 14 % and aldehydes 5 %. The total amounts of VOCs were not noticeably affected by UV-B. Among, the sesquiterpene, the perception of UV-B reduced the proportion of the oxygenated farnesol, the most abundant sesquiterpene, wherein –UV-B was 94.5 % and +UV-B 72.6 %, respectively. UV-B increased specifically, β -farnesene (representing 17.3 % of the total sesquiterpene in +UV-B and 2.36 % in –UV-B), valencene (4.4 vs. 0.8 %) and α -panasinsene (3.9 vs. 0.9 %); the proportion of the other compounds identified (except for 6, 28-heptatriacontadien-2-one) did not change significantly in this treatment. That is,

Fig. 1 Chemical structures of sesquiterpenes, oxygenated sesquiterpenes, aldehydes and ketones identified by GC–EIMS in grapevine flowers



+UV-B (as compared to –UV-B) increased β -farnesene (7.7-fold, $P_{(UV-B)} = 0.0557$), valencene (5.8-fold, $P_{(UV-B)} = 0.1236$), α -panasinsene (4.4-fold, $P_{(UV-B)} = 0.0872$), and 6, 28-heptatriacontadien-2-one (2.7-fold, $P_{(UV-B)} = 0.0356$).

Twelve VOCs were identified and quantified in grapevine flowers cv. Malbec in which sesquiterpenes predominated. These terpenoids have been previously reported in flowers and berries of *V. vinifera* L. (Lücker et al. 2004; Martin et al. 2009); they are synthesized by plants mainly as a defense mechanism against fungi (Moreira et al. 2003; Escoriza et al. 2013) and as insect pollinator attractants (Stranden et al. 2003). Previously, Gil et al. (2013) analyzed VOCs at different developmental stages in grape berries (at veraison, pre-harvest and harvest) and compounds differ from those identified in grape flowers. Four monoterpenes (limonene, pinene,

geraniol and eucalyptol), four aldehydes (hexanal, 2-heptenal, 2-pentenal and octanal), two alcohols (3-hexen-1-ol and 3-methyl pentanol) and one ketone (5-hepten-2-one) were detected. In the present study, and contrary to the hypothesis, UV-B did not stimulate the overall contents of VOCs in grapevine flowers. However, a differential synthesis of VOCs was detected; where the proportion of farnesol decreases while β -farnesene, valencene α -panasinsene, and 6, 28-heptatriacontadien-2-one increased. Anthers have been previously identified as the floral organs with the highest level of sesquiterpene volatiles in flowers of the cv. Cabernet Sauvignon (Martin et al. 2009). After flower opening, pollen is still well protected in the anther sacs and their walls attenuate UV-B radiation by at least 98 % (Flint and Caldwell 1983), and even the pollen grain wall is effective in attenuating

Table 1 VOCs assessed by GC–EIMS (ng mg⁻¹ flower FW) in grapevine flowers in field experiments (+UV-B and –UV-B). P_(UV-B): UV-B effect

	–UV-B	+UV-B	P _(UV-B)
Sesquiterpenes			
α-Farnesene	1.82 ± 2.06	3.30 ± 3.40	0.4283
β-Caryophyllene	1.19 ± 0.56	0.92 ± 0.35	0.3920
Valencene	2.03 ± 1.53	11.85 ± 8.6	0.1236
β-Farnesene	6.07 ± 6.88	46.94 ± 25.57	0.0557
α-Caryophyllene	0.31 ± 0.18	0.30 ± 0.22	0.8910
α-Panasinsene	2.41 ± 1.52	10.5 ± 7.66	0.0872
Nerolidol	0.37 ± 0.12	0.49 ± 0.61	0.7067
Farnesol	243.37 ± 55.03	197.28 ± 71.01	0.2844
Aldehydes			
Pentadecanal	0.61 ± 0.20	0.54 ± 0.06	0.4797
Hexadecatrienal	17.25 ± 9.10	17.85 ± 3.61	0.8951
Ketones			
Pentadecanone	46.63 ± 15.86	43.71 ± 16.08	0.7803
Heptatriacontadienone	0.275 ± 0.09	0.755 ± 0.34	0.0356

Values are mean ± SE

UV-B radiation (Uber 1939). Also, it has been reported a low responsiveness of grapevine flowers to UV-C, the most energetic type of UV radiation (Petit et al. 2009). Contrary to these reports, and in agreement with the study of Martin et al. (2009) in which sesquiterpene emission is light-dependent, we found differential responses by UV-B in some specific VOCs in flowers of grapevine cv. Malbec. Probably, the signal of UV-B is perceived directly by the flower and/or transmitted by other tissues. Gil et al. (2013) also found that in grapevine berries, monoterpenes associated with defense increased in response to +UV-B. The increase of β-farnesene, α-panasinsene and valencene may reduce oxidative damage in high altitude vineyards by UV-B and so requires further investigations. Especially because the augment of terpene synthesis in response to stressful environmental conditions seems to be crucial for the plant's tissues resistance to adverse environmental conditions (Berli et al. 2010; Beckett et al. 2012), defense towards pests (Moreira et al. 2003; Escoriza et al. 2013), and/or as pollinator attractants (Pichersky and Gershenzon 2002).

Acknowledgments This work was funded by FONCYT (PICT 2008 1666 and PID 0149-PAE 36789) to R.B., and (PICT 2007 02190) to P.P., CONICET (PIP 2008) to P.P., and Sepcyt-UNCuyo to R.B. and P.P. M.G. and M.V.S. are recipient of CONICET scholarships; R.B., F.B. and P.P. are members of CONICET. M.P. is member of INTA. The authors thank the helpful technical assistance of L. Bolcato in GC–EIMS.

References

- Aharoni A, Giri AP, Deuerlein S, Griepink F, De Kogel WJ, Verstappen FWA, Verhoeven HA, Jongmsa MA, Schwab W, Bouwmeester HJ (2003) Terpenoid metabolism in wild-type and transgenic plants. *Plant Cell* 15:2866–2884
- Beckett M, Loreto F, Velikova V, Brunetti C, Di Ferdinando M, Tattini M, Calfapietra C, Farrant J (2012) Photosynthetic limitations and volatile and non-volatile isoprenoids in the poikilochlorophyllous resurrection plant *Xerophyta humilis* during dehydration and rehydration. *Plant Cell Environ* 35: 2061–2074
- Berli FJ, Moreno D, Piccoli P, Hespagnol-Viana L, Silva MF, Bressan-Smith R, Cavagnaro JB, Bottini R (2010) Abscisic acid is involved in the response of grape (*Vitis vinifera* L.) cv. Malbec leaf tissues to ultraviolet-B radiation by enhancing ultraviolet-absorbing compounds, antioxidant enzymes and membrane sterols. *Plant, Cell Environ* 33:1–10
- Berli FJ, Fanzone M, Piccoli P, Bottini R (2011) Solar UV-B and ABA are involved in phenol metabolism of *Vitis vinifera* L. increasing biosynthesis of berry skin polyphenols. *J Agric Food Chem* 59:4874–4883
- Coelho E, Rocha SM, Delgadillo I, Coimbra MA (2006) Headspace-SPME applied to varietal volatile components evolution during *Vitis vinifera* L. cv. 'Baga' ripening. *Anal Chim Acta* 563: 204–214
- Coombe BJ (1995) Growth stages of the grapevine: adoption of a system for identifying grapevine growth stages. *Aust J Grape Wine Res* 1:104–110
- Escoriza G, Sansberro P, García Lampasona S, Gatica M, Bottini R, Piccoli P (2013) *In vitro* cultures of *Vitis vinifera* L. cv. Chardonnay synthesize the phytoalexin nerolidol upon infection by *Phaeoacremonium parasiticum*. *Phytopathol Mediterr* 52: 289–297
- Flint SD, Caldwell MM (1983) Influence of floral optical properties in the ultraviolet radiation environment of pollen. *Am J Bot* 70: 1416–1419
- Gil M, Pontin M, Berli FJ, Bottini R, Piccoli P (2012) Metabolism of terpenes in the response of grape (*Vitis vinifera* L.) leaf tissues to UV-B radiation. *Phytochemistry* 77:89–98
- Gil M, Bottini R, Berli FJ, Pontin M, Silva MF, Piccoli P (2013) Volatile organic compounds characterized from grapevine (*Vitis vinifera* L. cv. Malbec) berries increase at pre-harvest and in response to UV-B radiation. *Phytochemistry* 96:148–157
- Kegger W, Pierik R (2009) Biogenic volatile organic compounds and plant competition. *Trends Plant Sci* 15:126–131
- Lücker J, Bowen P, Bohlmann J (2004) *Vitis vinifera* terpenoid cyclases: functional identification of two sesquiterpene synthase cDNAs encoding (+)-valencene synthase and (–)-germacrene D synthase and expression of mono- and sesquiterpene synthases in grapevine flowers and berries. *Phytochem* 65:2649–2659
- Martin DM, Toub O, Chiang A, Lo BC, Ohse S, Lund ST, Bohlmann J (2009) The bouquet of grapevine (*Vitis vinifera* L. cv. Cabernet Sauvignon) flowers arises from the biosynthesis of sesquiterpene volatiles in pollen grains. *PNAS USA* 106:6881–6882
- McKenzie RL, Bjorn LO, Bais A, Ilyas M (2003) Changes in biologically active UV radiation reaching the earth's surface. *Photochem Photobiol Sci* 2:5–15
- Moreira IC, Lago JHG, Young MCM, Roque NF (2003) Antifungal aromadendrane sesquiterpenoids from the leaves of *Xylopia brasiliensis*. *J Brazil Chem Soc* 14:828–831
- Palomo ES, Diaz-Maroto MC, Vinas MAG, Soriano-Pérez A, Pérez-Coello MS (2007) Aroma profile of wines from Albillo and Muscat grape varieties at different stages of ripening. *Food Control* 18:398–403

- Park SK, Morrison JC, Adams DO, Noble AC (1991) Distribution of free and glycosidically bound monoterpenes in the skin and mesocarp of Muscat of Alexandria grapes during development. *J Agric Food Chem* 39:514–518
- Petit A, Baillieul F, Vaillant-Gaveau N, Jacquens L, Conreux A, Jeandet P, Clément C, Fontaine F (2009) Low responsiveness of grapevine flowers and berries at fruit set to UV-C irradiation. *J Exp Bot* 60:1155–1162
- Pichersky E, Gershenzon J (2002) The formation and function of plant volatiles: perfumes for pollinator attraction and defense. *Curr Opin Plant Biol* 5:237–243
- Stranden M, Liblikas I, König WA, Almaas TJ, Borg-Karlson AK, Mustaparta H (2003) (-)-Germacrene D receptor neurones in three species of heliothine moths: structure–activity relationships. *J Comp Physiol A* 189:563–577
- Teramura AH (2006) Effects of ultraviolet-B radiation on the growth and yield of crop plants. *Physiol Plant* 58:415–427
- Tholl D, Chen F, Petri J, Gershenzon J, Pichersky E (2005) Two sesquiterpene synthases are responsible for the complex mixture of sesquiterpenes emitted from *Arabidopsis* flowers. *Plant J* 42:757–771
- Über F (1939) Ultra-violet spectrophotometry of *Zea mays* pollen with the quartz microscope. *Am J Bot* 26:799–807
- Wei A, Shibamoto T (2007) Antioxidant activities and volatile constituents of various essential oils. *J Agric Food Chem* 55:1737–1742