

Duality and bosonization in Schwinger–Keldysh formulation

This content has been downloaded from IOPscience. Please scroll down to see the full text.

J. Stat. Mech. (2014) P09034

(<http://iopscience.iop.org/1742-5468/2014/9/P09034>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 163.10.1.238

This content was downloaded on 02/10/2014 at 18:49

Please note that [terms and conditions apply](#).

Duality and bosonization in Schwinger–Keldysh formulation

R E Gamboa Saraví, C M Naón and F A Schaposnik

Departamento de Física, Facultad de Ciencias Exactas, Universidad Nacional de La Plata and IFLP-CONICET, CC 67, 1900 La Plata, Argentina
E-mail: fidel@fisica.unlp.edu.ar

Received 31 July 2014

Accepted for publication 3 August 2014

Published 30 September 2014

Online at stacks.iop.org/JSTAT/2014/P09034

[doi:10.1088/1742-5468/2014/09/P09034](https://doi.org/10.1088/1742-5468/2014/09/P09034)

Abstract. We present a path-integral bosonization approach for systems out of equilibrium based on a duality transformation of the original Dirac fermion theory combined with the Schwinger–Keldysh time closed contour technique, to handle the non-equilibrium situation. The duality approach to bosonization that we present is valid for $D \geq 2$ space–time dimensions leading for $D = 2$ to exact results. In this last case we present the bosonization rules for fermion currents, calculate current–current correlation functions and establish the connection between the fermionic and bosonic distribution functions in a generic, non-equilibrium situation.

Keywords: bosonisation

Contents

1. Introduction	2
2. Generating functional for free Dirac fermions in the Schwinger–Keldysh time closed path	3
2.1. The model	3
2.2. Free fermions in the time contour	5
2.3. Incorporating sources	7
3. Duality and bosonization	7
4. Current–current correlations and the connection between fermionic and bosonic out of equilibrium distribution functions	12
5. Summary and discussion	14
Acknowledgments	15
Appendix A. Computation of the fermionic determinant: the Fujikawa Jacobian in Schwinger–Keldysh framework	15
References	20

1. Introduction

Bosonization is a powerful technique, widely used to analyze quantum systems in one spatial dimension [1]. In the context of relativistic quantum field theories (QFT), following the seminal papers by Coleman, Mandelstam and Witten [2–4], the bosonization procedure became a key tool in the solution of simplified models of strong interactions, such as QCD in $(1+1)$ dimensions. These developments also gave strength to the idea of duality, which is currently exploited with big success in the framework of string theories [5]. On the other hand, bosonization in condensed matter physics [6, 7] is an extremely important technique in the study of non-relativistic, one-dimensional systems (see [8] and references therein). Indeed, the Luttinger liquid state, which manifests experimentally in carbon nanotubes, polymer nanowires and quantum Hall edges, can be naturally understood in terms of bosonic collective modes, obtained analytically through bosonization. It is important to emphasize that the bosonization method was restricted to equilibrium physical situations until recently, when Gutman, Gefen and Mirlin in a significant work [9] built a bosonic theory for 1D fermions under non-equilibrium conditions. Starting from a model of free fermions of a definite chirality, they obtained an equivalent bosonic effective action containing terms of all orders in the fields. It should be stressed that this is an important achievement in view of the current, growing interest in non-equilibrium phenomena in nanoscopic systems [10].

In this work we present an out of equilibrium path-integral bosonization approach based in duality transformations proposed in [11, 12]. Our proposal is inspired by the ‘target space duality’ of string theory [13], and can be applied in any number of space dimensions, leading to exact results for Abelian and non-Abelian bosonization of massive and massless Dirac fermions in $1 + 1$ space–time dimension and perturbative answers in $d > 1$ space dimensions. We find that this path-integral bosonization framework is particularly appropriate when considering out of equilibrium systems using the Schwinger–Keldysh time closed contour technique [14–18] leading to very natural extensions of the Coleman–Mandelstam bosonization recipe [2].

The paper is organized as follows. In section 2 we introduce the path-integral generating functional for non-interacting Dirac fermions with dynamics governed by an action defined on the Schwinger–Keldysh time contour. We discuss boundary conditions in different branches of the contour and introduce the retarded, advanced and Keldysh fermionic Green functions. Then, in section 3, we present the path-integral bosonization approach based in duality transformations. After gauging the global symmetry of the original fermionic theory and introducing a Lagrange multiplier that constrains the auxiliary gauge field to be a pure gauge, we arrive at a bosonic dual and establish the correspondence between fermionic and bosonic currents. In section 4 we present the calculation of current–current correlation functions both in the bosonic and fermionic sectors and establish the connection between the fermionic and bosonic distribution functions for a generic, non-equilibrium situation. We summarize and discuss our results in section 5, where we also comment on possible extensions of our approach to $D > 2$ space–time dimensions.

2. Generating functional for free Dirac fermions in the Schwinger–Keldysh time closed path

2.1. The model

We start by considering a system of fermion fields ψ and ψ^\dagger in $D = d + 1$ space–time dimensions (later on we will specialize our results to the case $d = 1$). The Lagrangian density is given by

$$\mathcal{L}_F = \bar{\psi} i \not{\partial} \psi \quad (1)$$

Here $\not{\partial} \equiv \gamma^\mu \partial_\mu$, with γ^μ the Dirac matrices and $(x^\mu) \equiv (x^0 = t, \mathbf{x})$,

As is well-known, the Schwinger–Keldysh formulation [14, 15] enables us to extend the validity of a quantum field theory to the realm of non-equilibrium physics by defining the action $S[\psi, \bar{\psi}, s_\mu]$ along a closed time contour $\mathcal{C} = C_+ \cup C_- \cup C_\tau$ [16] represented in figure 1. Within this formulation, we start by considering the partition function

$$Z_F = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi e^{iS_F[\psi, \bar{\psi}]}, \quad (2)$$

with

$$S_F[\psi, \bar{\psi}] = \int_{\mathcal{C}} dt d^d x \bar{\psi} i \not{\partial} \psi, \quad (3)$$




Figure 1. Sketch of the closed time path contour. The upper branch of the path, C_+ , represents the usual time path, from $t = -\infty$ to $t = +\infty$. The lower branch C_- goes backward in time. Both branches are connected, in the remote past, by the curve C_τ .

where subscript \mathcal{C} indicates that the action is defined along a closed time path [16] so that in the path-integral defining Z_F the fermion fields should be integrated imposing the appropriate boundary conditions on each branch of the contour.

As indicated in figure 1, the contour is split into two branches, one from $t = -\infty$ to $t = +\infty$ (C_+), the other one completing the contour (C_-). In the remote past ($t = -\infty$) the forward and backward paths are connected by C_τ where $\tau \in (0, \beta)$ is imaginary time when evolving along C_τ and $\beta = 1/T$ is the inverse temperature. The precise form of this connection depends on the initial distribution, which can be assumed to be an equilibrium distribution. Fields in each one of the two contour sections C_+ and C_- will be denoted as ψ_+ , ψ_+^\dagger , and ψ_- , ψ_-^\dagger , respectively. Concerning fields in C_τ , they will be called ψ_τ , ψ_τ^\dagger . with this, action S_F will be written in the form

$$S_F[\psi, \bar{\psi}] = \int d^d x \int_{-\infty}^{\infty} dt (\mathcal{L}[\psi_+, \bar{\psi}_+] - \mathcal{L}[\psi_-, \bar{\psi}_-]) + \int_{C_\tau} dt d^d x \mathcal{L}_\tau \quad (4)$$

where

$$\mathcal{L}[\psi_\pm, \bar{\psi}_\pm] = \bar{\psi}_\pm i \not{\partial} \psi_\pm. \quad (5)$$

$$\mathcal{L}_\tau[\psi_\tau, \bar{\psi}_\tau] = \bar{\psi}_\tau i \not{\partial} \psi_\tau \quad (6)$$

The appropriate boundary conditions for the fermion fields are [16]

$$\begin{aligned} \psi_+(\mathbf{x}, t = +\infty) &= \psi_-(\mathbf{x}, t = +\infty) \\ \psi_+^\dagger(\mathbf{x}, t = +\infty) &= \psi_-^\dagger(\mathbf{x}, t = +\infty) \\ \psi_-(\mathbf{x}, t = -\infty) &= \psi_\tau(\mathbf{x}, t = 0) \\ \psi_-^\dagger(\mathbf{x}, t = -\infty) &= \psi_\tau^\dagger(\mathbf{x}, t = 0) \\ \psi_+(\mathbf{x}, t = -\infty) &= -\psi_\tau(\mathbf{x}, t = \beta) \\ \psi_+^\dagger(\mathbf{x}, t = -\infty) &= -\psi_\tau^\dagger(\mathbf{x}, t = \beta), \end{aligned} \quad (7)$$

where the minus sign in the two last equations is related to the Grassmann nature of fermionic variables. It is through these boundary conditions that the fermion fields ψ_\pm , ψ_τ are coupled. The role of the field ψ_τ is precisely to provide matching conditions for ψ_\pm and they will play no role in final expressions [17]. That is the reason why when we define the generating functional, in order to compute vacuum expectation values, we do not couple sources to ψ_τ fields and write

$$Z_F[s_\mu] = \int \mathcal{D}\bar{\psi}_+ \mathcal{D}\bar{\psi}_- \mathcal{D}\psi_+ \mathcal{D}\psi_- \exp(iS_F[\bar{\psi}, \psi, s]) \quad (8)$$

where

$$S_F[\bar{\psi}, \psi, s] = \int d^d x \int_{-\infty}^{\infty} dt (\mathcal{L}[\psi_+, \bar{\psi}_+, s_\mu^+] - \mathcal{L}[\psi_-, \bar{\psi}_-, s_\mu^-]) \quad (9)$$

Written in this form, one could think that the partition function (8) consists of two independent contributions which can be treated separately so that the partition function splits into two independent factors. However, as indicated above, the path-integration requires boundary conditions which in the present case are precisely those in equation (8) which do couple the \pm -fields.

2.2. Free fermions in the time contour

Let us now show how to take into account, within the path-integral formulation, the time boundary conditions. Let us denote ψ_a and $\bar{\psi}_a$ the spinors introduced in (4), with $a = +, -$. Thus, the free fermion action can be written in terms of the time contour matrix structure as

$$S_F[\psi, \bar{\psi}, 0] = \int d^d x dt \bar{\psi}_a D_{ab} \psi_b, \quad (10)$$

where

$$D = \begin{pmatrix} i\cancel{\partial} & 0 \\ 0 & -i\cancel{\partial} \end{pmatrix}.$$

As discussed above, the required boundary conditions (8) imply a correlation between $+$ and $-$ fields at the boundaries so that although D^0 is diagonal, its inverse operator (the Green function) could have non-vanishing off-diagonal components. Indeed, the time boundary terms lead to a Green function that is a non-diagonal matrix

$$G = \begin{pmatrix} G_{++} & G_{+-} \\ G_{-+} & G_{--} \end{pmatrix},$$

satisfying

$$D_x G_{xy} = \delta(t_x - t_y) \delta^d(\mathbf{x} - [y]) I_{4 \times 4}.$$

or, more explicitly, for the case of one spatial dimension

$$i \begin{pmatrix} 0 & \partial_0 + \partial_1 & 0 & 0 \\ \partial_0 - \partial_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -(\partial_0 + \partial_1) \\ 0 & 0 & -(\partial_0 - \partial_1) & 0 \end{pmatrix} \begin{pmatrix} 0 & G_{++}^R & 0 & G_{+-}^R \\ G_{++}^L & 0 & G_{+-}^L & 0 \\ 0 & G_{-+}^R & 0 & G_{--}^R \\ G_{-+}^L & 0 & G_{--}^L & 0 \end{pmatrix} = \delta(t_x - t_y) \delta(x - y) I_{4 \times 4}, \quad (11)$$

where the superscripts R and L indicate the chiral components of the fermionic Green functions. Note that when considering the usual equilibrium situation, the introduction of the contour is not mandatory and only the time forward branch is usually taken into account. Consequently, the Green function is just G_{++} , which is nothing but the time-ordered propagator. Let us mention, however, that in some applications the use of the closed time contour can be convenient even in equilibrium situations. In the present formulation G_{--} represents an anti-time-ordered propagator. The off-diagonal components G_{+-} and G_{-+} satisfy homogeneous differential equations and depend on the

distribution function. Moreover, it can be shown that only three of the four elements of G are independent. Indeed, one can verify that $G_{++} + G_{--} = G_{+-} + G_{-+}$. This suggests that the fermionic Green function can be written as a triangular matrix. In fact, following Larkin and Ovchinnikov [19], we perform the following rotation of fermion fields:

$$\Psi_1 = \frac{\psi_+ + \psi_-}{\sqrt{2}} \quad \Psi_2 = \frac{\psi_+ - \psi_-}{\sqrt{2}} \quad (12)$$

and

$$\bar{\Psi}_1 = \frac{\bar{\psi}_+ - \bar{\psi}_-}{\sqrt{2}} \quad \bar{\Psi}_2 = \frac{\bar{\psi}_+ + \bar{\psi}_-}{\sqrt{2}}. \quad (13)$$

These changes can be implemented introducing two matrices M and N , such that

$$\psi = M \Psi \quad \bar{\psi} = \bar{\Psi} N, \quad (14)$$

where Ψ and $\bar{\Psi}$ are now spinors constructed from Ψ_1, Ψ_2 , and

$$M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad N = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}.$$

In terms of these new fields the free action becomes

$$S_F[\Psi, \bar{\Psi}, 0] = \int dx dt \bar{\Psi} \mathcal{D} \Psi, \quad (15)$$

where now the free Dirac operator reads

$$\mathcal{D}^0 = \begin{pmatrix} i\partial\!\!\!/ & 0 \\ 0 & i\partial\!\!\!/ \end{pmatrix}. \quad (16)$$

The boundary conditions for spinors can be easily determined from those imposed on ψ_{\pm} and ψ_{\pm}^{\dagger} , equation (8).

Once again, as we did in (11), we can be more explicit by displaying the full matrix structure in the equation obeyed by the Green function in the $d = 1$ case,

$$\mathcal{D}_x^0 \mathcal{G}_{xy} = \delta(t_x - t_y) \delta(x - y) I_{4 \times 4},$$

where

$$\mathcal{D}^0 = i \begin{pmatrix} 0 & \partial_0 + \partial_1 & 0 & 0 \\ \partial_0 - \partial_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \partial_0 + \partial_1 \\ 0 & 0 & \partial_0 - \partial_1 & 0 \end{pmatrix} \quad (17)$$

and

$$\mathcal{G} = i \begin{pmatrix} 0 & \mathcal{G}_{\text{ret}}^R & 0 & \mathcal{G}_K^R \\ \mathcal{G}_{\text{ret}}^L & 0 & \mathcal{G}_K^L & 0 \\ 0 & 0 & 0 & \mathcal{G}_{\text{adv}}^R \\ 0 & 0 & \mathcal{G}_{\text{adv}}^L & 0 \end{pmatrix} \quad (18)$$

Here we have defined retarded, advanced and Keldysh components of the Green function as

$$\begin{aligned} \mathcal{G}_{\text{ret}}^i &= G_{++}^i - G_{+-}^i = G_{-+}^i - G_{--}^i, \\ \mathcal{G}_{\text{adv}}^i &= G_{++}^i - G_{-+}^i = G_{+-}^i - G_{--}^i, \quad i = R, L \\ \mathcal{G}_K^i &= G_{++}^i + G_{--}^i = G_{+-}^i + G_{-+}^i. \end{aligned} \quad (19)$$

2.3. Incorporating sources

Let us now include the source term appearing in (3), which, after the doubling of degrees of freedom, reads

$$\bar{\psi}_a s_{ab} \psi_b = \bar{\psi}_+ \not{s}_+ \psi_+ - \bar{\psi}_- \not{s}_- \psi_-, \quad (20)$$

where the vector source will be written as

$$s = \begin{pmatrix} \not{s}_+ & 0 \\ 0 & -\not{s}_- \end{pmatrix}.$$

Performing the rotation in fermion fields (14), we obtain

$$\bar{\psi}_a s_{ab} \psi_b = \bar{\Psi} \not{S} \Psi, \quad (21)$$

where the matrix \mathcal{S} takes the form

$$\not{S} = \begin{pmatrix} \frac{\not{s}_+ + \not{s}_-}{2} & \frac{\not{s}_+ - \not{s}_-}{2} \\ \frac{\not{s}_+ - \not{s}_-}{2} & \frac{\not{s}_+ + \not{s}_-}{2} \end{pmatrix}.$$

Note that, in contrast to Dirac operator \mathcal{D}^0 , \mathcal{S} is a non-diagonal matrix. At this point, it is convenient to define

$$\not{s}_c = \frac{\not{s}_+ + \not{s}_-}{2} \quad \not{s}_q = \frac{\not{s}_+ - \not{s}_-}{2}, \quad (22)$$

where the subscripts c and q stand for ‘classical’ and ‘quantum’ components of the vector fields (this terminology is habitual in the context of quantum non-equilibrium field theory, it refers to the role played by the fields when studying the classical equations of motion [18]). Now, the matrix \mathcal{S} can be written in the form

$$\not{S} = \not{s}_c \lambda^c + \not{s}_q \lambda^q \equiv \not{s}_a \lambda^a, \quad a = c, q \quad (23)$$

with

$$\lambda^c = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \lambda^q = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

At this point we have all the necessary ingredients to resume our main task.

3. Duality and bosonization

Taking into account that the Jacobian of the change of variables proposed in the previous sections is trivial (i.e. field independent), the fermionic path-integral in (2) can be written in terms of the fermion fields Ψ and $\bar{\Psi}$ introduced in equation (14). Once the source term is introduced, the same can be done for the generating functional which in terms of the new variables takes the form

$$Z_F[s_\mu] = \int \mathcal{D}\bar{\Psi} \mathcal{D}\Psi e^{iS_F[\Psi, \bar{\Psi}, S(s)]} \quad (24)$$

where

$$S_F = \int dt d^d x \bar{\Psi} (\mathcal{D}^0 + \not{S}(s)) \Psi. \quad (25)$$

Let us now introduce the functional bosonization procedure [12]. Although this formulation can be in principle used for arbitrary space–time dimensions, in order to make contact with the results of [9], from now on we will restrict our study to 1 + 1 dimensions ($d = 1$) which is besides the only case in which bosonization procedure is exact. Let us perform a local change of path-integral variables

$$\Psi \rightarrow g(x) \bar{\Psi}, \quad (26)$$

where $g(x)$ is composed of two successive $U(1)$ transformations

$$g(x) = g_c(x)g_q(x) \quad (27)$$

where

$$g_c(x) = \exp(i\alpha_c(x)\lambda^c) \quad (28)$$

$$g_q(x) = \exp(i\alpha_q(x)\lambda^q) \quad (29)$$

The boundary conditions on $\alpha = (\alpha_c, \alpha_q)$ should be chosen so that those on fermion fields, equation (8), remain valid. To obtain the appropriate boundary conditions for α we shall consider in what follows a general transformation depending on functions that will become eventually bosonic fields that we shall denote generically as ϕ .

Let us introduce a notation describing boundary conditions for the original fields ψ when written in terms of ψ_{\pm} and ψ_{τ}

$$\psi_{b_r}(x, t_i) = \epsilon_{b_s} \psi_{b_s}(x, t_i) \quad (30)$$

where index i denotes the different boundary points in the time contour ($t_i = -\infty, \infty, 0, \beta$) and ψ_{b_r}, ψ_{b_s} indicate the fields in different regions of the contour ($b_r, b_s s = +, -, 0\tau$). Concerning ϵ_{b_s} , it takes the value $\epsilon_s = +1$ for $s = \pm, 0$ and $\epsilon_s = -1$ for $s = \tau$ (see equation (8)).

Under a transformation $U_{b_r}[\phi_{r_i}(x, t)] = U_{b_r}(x, t)$ the fermion fields at the boundary transform according to

$$\psi_{b_r}(x, t_i) \rightarrow \psi_{b_r}(x, t_i)' = U_{b_r}(x, t_i) \psi_{b_r}(x, t_i) \quad (31)$$

Now, since $\psi_{b_r}(x, t_i)'$ should obey the same boundary conditions as $\psi_{b_r}(x, t_i)$, namely (30), one has that

$$U_{b_r}(x, t_i) = U_{b_s}(x, t_i) \quad (32)$$

where there is no ϵ_s factor as in the fermion case so as to preserve the signs implied by equation (30). These boundary conditions are then those satisfied by g and also by all bosonic fields to be introduced below.

One can always define a path-integral measure such that the Jacobians associated to transformations (28) and (29) are trivial [21–25] so that after the change of variables, the generating functional (24) becomes

$$Z_F[s_{\mu}] = \int \mathcal{D}\bar{\Psi} \mathcal{D}\Psi \exp \left(i \int d^2x \bar{\Psi} (\mathcal{D}^0 + \mathcal{G}(s) + ig^{-1}\not{\partial}g) \Psi \right). \quad (33)$$

We shall now introduce an auxiliary field \mathcal{B}_{μ}

$$\mathcal{B}_{\mu} = \mathcal{B}_{\mu}^a \lambda^a, \quad a = c, q \quad (34)$$

together with a delta function condition so that it represents a pure gauge,

$$\not{\mathcal{B}} + ig^{-1}\not{\partial}g = 0 \quad (35)$$

Now, instead of (33) we write the fermionic generating functional in the form

$$Z_F[s_\mu] = \int \mathcal{D}\mathcal{B}_\mu \mathcal{D}\bar{\Psi} \mathcal{D}\Psi \exp \left(i \int d^2x \bar{\Psi} (\mathcal{D}^0 + \mathcal{G}(s) + \mathcal{B}) \Psi \right) \delta(\mathcal{F}(\mathcal{B})), \quad (36)$$

where

$$\mathcal{F}(\mathcal{B}) = \epsilon_{\mu\nu} \partial^\mu \mathcal{B}^\nu. \quad (37)$$

imposes the pure gauge condition on \mathcal{B}_μ^a . As explained before, the bosonic field \mathcal{B}_μ^a obeys boundary conditions (32).

Taking into account that the fermionic path-integral is just the determinant of the Dirac operator, and the shift

$$\mathcal{B} + \mathcal{G}(s) \rightarrow \mathcal{B}, \quad (38)$$

one ends up with

$$Z_F[s_\mu] = \int \mathcal{D}\mathcal{B}_\mu \det(\mathcal{D}^0 + \mathcal{B}) \delta(\mathcal{F}(\mathcal{B}) - \mathcal{F}(\mathcal{S})). \quad (39)$$

We now introduce in the path-integral defining Z a scalar field $\Phi = \Phi_a \lambda^a i$ to represent the delta function constraint in the form

$$\delta(\mathcal{H}) = \int \mathcal{D}\Phi \exp \left(iC \operatorname{tr}_K \int d^2x \mathcal{H} \Phi \right) \quad (40)$$

with C a constant to be appropriately fixed below tr_K indicating a trace over matrices λ^a . Being Φ the bosonic field boundary conditions correspond to those given by equation (32).

Using the delta function representation (40) the generating functional $Z_F[s]$ takes the form

$$Z_F[s] = \int \mathcal{D}\Phi e^{iS_B[\Phi]} \exp \left(-iC \operatorname{tr}_K \int d^2x \mathcal{F}(\mathcal{S}[s]) \Phi \right), \quad (41)$$

with $S_B[\Phi]$ defined as

$$e^{iS_B[\Phi]} = \int \mathcal{D}\mathcal{B}_\mu \det(\mathcal{D}^0 + \mathcal{B}) \exp \left(iC \operatorname{tr}_K \int d^2x \mathcal{F}(\mathcal{B}) \Phi \right). \quad (42)$$

Given Z_F written as in (41) one can interpret $S_B[\Phi]$ as a bosonic action for the scalar field Φ and then establish an identity between the original fermionic generating functional (24) and a bosonic generating functional

$$Z_B[s] \equiv \int \mathcal{D}\Phi e^{iS_B[\Phi]} \exp \left(-iC \operatorname{tr}_K \int d^2x \mathcal{F}(\mathcal{S}[s]) \Phi \right), \quad (43)$$

that is

$$Z_F[s] = Z_B[s] \quad (44)$$

In order to find an explicit form for the bosonic action $S_B[\Phi]$ it is necessary to compute the fermionic determinant and then proceed to integrate over b . Once this is done one should obtain an identity between the original fermionic generating functional and what one can interpret as a bosonic generating functional $Z_B[\phi]$,

$$Z_B[s_\mu] = \int \mathcal{D}\Phi e^{iS_B[\Phi]} \exp \left(-iC \operatorname{tr}_K \int d^2x \mathcal{S}_\mu(s) \epsilon_{\mu\nu} \partial_\nu \Phi \right), \quad (45)$$

Thus, bosonization of the original fermionic theory has been achieved through the identity

$$Z_{\text{F}}[s_{\mu}] = Z_{\text{B}}[s_{\mu}], \quad (46)$$

This formula allows us to obtain, through functional derivation, the bosonic representation for fermionic currents.

In appendix A we present a careful evaluation of the fermionic determinant which takes the form

$$\det(\mathcal{D}^0 + \mathcal{B}) = \det \mathcal{D}^0 J_{\text{F}}^{\text{dec}}[\varphi] J_{\text{F}}^{\text{dec}}[\eta] \quad (47)$$

where the field \mathcal{B} has been written in terms of the general decomposition

$$\mathcal{B} = \not{\phi} \gamma_5 \varphi + \not{\phi} \eta \quad (48)$$

and $J_{\text{F}}^{\text{dec}}[\chi]$ is defined as (see appendix A)

$$J_{\text{F}}^{\text{dec}}[\chi] = \exp \frac{i}{2\pi} \int d^2x ((\partial_0 + \partial_1)\chi_q(\partial_0 - \partial_1)\chi_c + (\partial_0 + \partial_1)\chi_c(\partial_0 - \partial_1)\chi_q) \quad (49)$$

Now, inserting this result in (42) one has

$$e^{iS_{\text{B}}[\Phi]} = \det(\mathcal{D}^0) \int \mathcal{D}\eta J_{\text{F}}^{\text{dec}}[\eta] \mathcal{D}\varphi J_{\text{F}}^{\text{dec}}[\varphi] \exp \left(-i \text{C}tr_{\text{K}} \int d^2x \Phi \partial_{\mu} \partial^{\mu} \varphi \right) \quad (50)$$

We see that the η field is completely decoupled from Φ so that the integral over η just gives a trivial factor. Concerning the remaining φ integral, apart from the quadratic term in φ there is the linear term that couples φ with Φ .

$$e^{iS_{\text{B}}[\Phi]} = \mathcal{N} \int \mathcal{D}\varphi \exp \left(\frac{i}{2\pi} \int d^2x \left((\partial_0 + \partial_1)\varphi_q(\partial_0 - \partial_1)\varphi_c + (\partial_0 + \partial_1)\varphi_c(\partial_0 - \partial_1)\varphi_q - 2iC \Phi_q \partial_{\mu} \partial^{\mu} \varphi_q + \Phi_c \partial_{\mu} \partial^{\mu} \varphi_c \right) \right)$$

Then, integrating over φ one gets for $S_{\text{B}}[\Phi]$

$$e^{iS_{\text{B}}[\Phi]} = \exp \left(i\pi C^2 \text{tr}_{\text{K}} \int d^2x \lambda^q \partial_{\mu} \Phi \partial^{\mu} \Phi \right) \quad (51)$$

Inserting the result into the rhs of (41) we finally obtain

$$Z_{\text{B}}[s_{\mu}] = \mathcal{N} \int \mathcal{D}\Phi \exp \left(i\pi C^2 \text{tr}_{\text{K}} \int d^2x \lambda^q \partial_{\mu} \Phi \partial^{\mu} \Phi \right) \exp \left(-i \text{C}tr_{\text{K}} \int d^2x \mathcal{S}_{\mu} \epsilon^{\mu\nu} \partial_{\nu} \Phi \right), \quad (52)$$

where the factor \mathcal{N} , being independent of the fields and the sources, is a normalization factor that can be disregarded when computing correlation functions. At this point we also note that the constant C does not play any relevant role, as expected. Indeed, it becomes apparent that it can be eliminated from the action by rescaling the field. In order to facilitate comparison with previous results we redefine $2\sqrt{\pi}C\phi \rightarrow \phi$ obtaining

$$Z_{\text{B}}[s_{\mu}] = \mathcal{N} \int \mathcal{D}\phi \exp \left(\frac{i}{4} \text{tr}_{\text{K}} \int d^2x \lambda^q \partial_{\mu} \phi \partial^{\mu} \phi \right) \exp \left(-\frac{i}{2\sqrt{\pi}} \text{tr}_{\text{K}} \int d^2x \mathcal{S}_{\mu}[s] \epsilon^{\mu\nu} \partial_{\nu} \phi \right), \quad (53)$$

with \mathcal{N} a normalization constant.

From the generating functional (53) we see that the bosonic dual Lagrangian density takes the form

$$\mathcal{L}_B = \frac{1}{4} \text{tr}_K \lambda^q \partial_\mu \Phi \partial^\mu \Phi. \quad (54)$$

This bosonized Lagrangian can be written in a more illuminating way introducing the boson fields ϕ_+ and ϕ_- such that

$$\begin{aligned} \phi_+ &= \frac{\phi^c + \phi^q}{\sqrt{2}} \\ \phi_- &= \frac{\phi^c - \phi^q}{\sqrt{2}} \end{aligned} \quad (55)$$

In terms of the ϕ_\pm fields \mathcal{L}_B takes the form

$$\mathcal{L}_B = \frac{1}{2} (\partial_\mu \phi_+ \partial^\mu \phi_+ - \partial_\mu \phi_- \partial^\mu \phi_-). \quad (56)$$

which corresponds to a free boson model.

As was the case for fermions ψ_\pm , although the bosonic fields ϕ_\pm in the bosonic Lagrangian (56) ensemble decoupled, they are in fact correlated through the bosonic time boundary conditions

$$\begin{aligned} \phi_+(x, t = +\infty) &= \phi_-(x, t = +\infty) \\ \phi_-(x, t = -\infty) &= \phi_+(x, t = 0) \\ \phi_+(x, t = -\infty) &= \phi_+(x, t = \beta) \end{aligned} \quad (57)$$

The action associated to Lagrangian (56)

$$S_B[\phi_+, \phi_-] = \frac{1}{2} \int dx \int_{-\infty}^{\infty} dt (\partial_\mu \phi_+ \partial^\mu \phi_+ - \partial_\mu \phi_- \partial^\mu \phi_-), \quad (58)$$

can be also written in the form

$$S_B[\phi] = \frac{1}{2} \int_{\mathcal{C}} dt dx \partial_\mu \phi \partial^\mu \phi \quad (59)$$

where \mathcal{C} is the closed time contour introduced in the original fermionic action (3).

One can compute vacuum expectation values of fermion current correlations in terms of bosonic currents using the identity between fermionic and bosonic generating functionals established in (44). As an example, one has

$$\begin{aligned} \langle \bar{\psi}_\pm(x_1) \gamma_{\mu_1} \psi_\pm(x_n) \dots \bar{\psi}_\pm(x_n) \gamma_{\mu_n} \psi_\pm(x_n) \rangle_F &= \frac{1}{Z_F[s]} \frac{\delta^n Z_F[s]}{\delta s_\pm^{\mu_1}(x_1) \dots \delta s_\pm^{\mu_n}(x_n)} \Big|_{s=0} \\ &= \frac{1}{Z_B[s]} \frac{\delta^n Z_B[s]}{\delta s_\pm^{\mu_1}(x_1) \dots \delta s_\pm^{\mu_n}(x_n)} \Big|_{s=0} \\ &= (\mp(2\pi)^{-1/2})^n \langle \epsilon_{\mu_1 \nu_1} \dots \epsilon_{\mu_n \nu_n} \partial_{\nu_1} \phi_\pm(x_1) \dots \partial_{\nu_n} \phi_\pm(x_n) \rangle_B \end{aligned} \quad (60)$$

We can summarize our bosonization results in terms of the following fermion–boson correspondence

$$\mathcal{L}_F = \bar{\psi}_+ i \not{\partial} \psi_+ - \bar{\psi}_- i \not{\partial} \psi_- \rightarrow \mathcal{L}_B = \frac{1}{2} (\partial_\mu \phi_+ \partial^\mu \phi_+ - \partial_\mu \phi_- \partial^\mu \phi_-), \quad (61)$$

$$\bar{\psi}_+ \gamma^\mu \psi_+ \rightarrow -\frac{1}{\sqrt{2\pi}} \epsilon^{\mu\nu} \partial_\nu \phi_+, \quad (62)$$

$$\bar{\psi}_- \gamma^\mu \psi_- \rightarrow \frac{1}{\sqrt{2\pi}} \epsilon^{\mu\nu} \partial_\nu \phi_- . \quad (63)$$

In view of the connection (61), one can write a bosonization recipe for the action written in terms of the Keldysh contour \mathcal{C} in the form

$$S_F[\psi, \bar{\psi}] = \int_{\mathcal{C}} dt dx \bar{\psi} i \not{\partial} \psi \rightarrow S_B[\phi] = \frac{1}{2} \int_{\mathcal{C}} dt dx \partial_\mu \phi \partial^\mu \phi \quad (64)$$

To write this formula we have used the fact that fields on the C_τ contour are decoupled from sources thus playing no role in final expressions.

Equation (64) is formally identical to the Coleman–Luther–Mandelstam–Mattis bosonization recipe in the equilibrium case [1]. Concerning the bosonization recipe for currents equations (62)–(63), note that they both lead to the same conserved charge, $dQ/dt = 0$ since the relative sign of the currents is compensated by the opposite sign of the time-integral limit.

$$Q = \int dx^1 \bar{\psi} \gamma^0 \psi \rightarrow \int dx^1 \phi \partial_1 \phi \quad (65)$$

Let us conclude by stressing that we were able to arrive at the simple bosonization rules (61)–(63) implying a quadratic bosonic Lagrangian because one can compute the fermion determinant $\det(\mathcal{D}^0 + \mathcal{B})$ in a closed form, this being a key tool in our approach. This was the reason why we have considered a Lagrangian for both positive and negative chirality fermions (see with [9]) since, it is well-known [26] if just one chirality is considered, the Dirac operator does not lead to a well-defined eigenvalue problem and hence the definition of an associated Dirac determinant is not a well-defined problem.

4. Current–current correlations and the connection between fermionic and bosonic out of equilibrium distribution functions

Here, we shall evaluate vacuum expectation values of current correlations by using the bosonization recipe (61)–(63). More specifically, we shall consider the following correlation function

$$\langle (\bar{\psi}_+ \gamma^\mu \psi_+(x)) (\bar{\psi}_- \gamma^\nu \psi_-(y)) \rangle_F = -\frac{1}{2\pi} \langle \epsilon^{\mu\rho} \partial_\rho \phi_+(x) \epsilon^{\nu\sigma} \partial_\sigma \phi_-(y) \rangle_B, \quad (66)$$

where the lhs is to be computed for the original free fermionic model, and the rhs with the bosonic action for Lagrangian (56). The Fourier transform of the out of equilibrium right (R) and left (L) fermionic Green functions (20) are given by

$$\begin{aligned} \mathcal{G}_{\text{ret}}^{\text{R}}(k) &= \frac{1}{k^0 - k^1 + i\delta}, & \mathcal{G}_{\text{ret}}^{\text{L}}(k) &= \frac{1}{k^0 + k^1 + i\delta} \\ \mathcal{G}_{\text{adv}}^{\text{R}}(k) &= \frac{1}{k^0 - k^1 - i\delta}, & \mathcal{G}_{\text{adv}}^{\text{L}}(k) &= \frac{1}{k^0 + k^1 - i\delta} \\ \mathcal{G}_{\text{K}}^{\text{R}}(k) &= F(k^0)(\mathcal{G}_{\text{ret}}^{\text{R}}(k) - \mathcal{G}_{\text{adv}}^{\text{R}}(k)), & \mathcal{G}_{\text{K}}^{\text{L}}(k) &= F(k^0)(\mathcal{G}_{\text{ret}}^{\text{L}}(k) - \mathcal{G}_{\text{adv}}^{\text{L}}(k)), \end{aligned}$$

where $F(k^0)$ is the fermionic Wigner function, connected to the generic (not necessarily the one corresponding to equilibrium) fermionic distribution function $n_{\text{F}}(k^0)$

$$F(k^0) = 1 - 2n_{\text{F}}(k^0) \quad (67)$$

At thermodynamical equilibrium $n_F(k^0)$ coincides with the Fermi–Dirac distribution and then one has

$$F^{\text{eq}}(k^0) = \tanh\left(\frac{\beta k^0}{2}\right)$$

where $\beta = 1/kT$.

The free bosonic Green functions arising from the dual bosonic Lagrangian (56), with out of equilibrium bosonic distribution $n_B(k^0)$ can be expressed in terms of the inverse of the d'Alembertian operator Δ as

$$\begin{aligned}\Delta_{\text{ret}}(k) &= \frac{1}{k_0^2 + k^1 k_1 + i\delta} \\ \Delta_{\text{adv}}(k) &= \frac{1}{k_0^2 + k^1 k_1 - i\delta} \\ \Delta_K(k) &= B(k^0)(\Delta_{\text{ret}}(k) - \Delta_{\text{adv}}(k)),\end{aligned}\tag{68}$$

where

$$B(k^0) = 1 + 2n_B(k^0)\tag{69}$$

is the bosonic Wigner function. At equilibrium, $n_B(k^0)$ is the Bose–Einstein distribution and one has

$$B^{\text{eq}}(k^0) = \coth\left(\frac{\beta k^0}{2}\right)$$

We shall consider for definiteness the vacuum expectation values (66) for $\mu = \nu = 1$. Using

$$\langle \phi_+(x) \phi_-(y) \rangle_B = 2i\Delta_{+-}(x - y)\tag{70}$$

in the bosonic side we get

$$G_{+-}^{\text{R}}(z)G_{-+}^{\text{R}}(-z) + G_{+-}^{\text{L}}(z)G_{-+}^{\text{L}}(-z) = \frac{i}{\pi} \partial_0^x \partial_0^y \Delta_{+-}(z)\tag{71}$$

with $z = x - y$.

In terms of the physical Green functions (68) Δ_{+-} takes the form

$$\Delta_{+-} = \frac{1}{2}(\Delta_K + \Delta_{\text{adv}} - \Delta_{\text{ret}})\tag{72}$$

Then working in momentum space and using the explicit form of Green functions we get

$$\begin{aligned}\int_{-\infty}^{+\infty} dk_0 \int_{-\infty}^{+\infty} dk'_0 n_F(k^0)(1 - n_F(k'^0))(e^{-i(k_0 - k'_0)u} + e^{-i(k_0 - k'_0)v}) \\ = \frac{1}{2} \int_{-\infty}^{+\infty} dk_0 |k_0| (B(k^0) - 1)(e^{-ik_0 u} + e^{-ik_0 v}),\end{aligned}\tag{73}$$

where $u = z^0 + z^1$ and $v = z^0 - z^1$. Multiplying both sides by $e^{i\omega u}(e^{i\omega v})$ and integrating in u (v), we find

$$n_B(\omega) = \frac{1}{\omega} \int_{-\infty}^{+\infty} dk_0 n_F(k^0) [1 - n_F(k^0 - \omega)].\tag{74}$$

This is a remarkable equation that establishes the connection between the fermionic and bosonic distributions for a generic, non-equilibrium situation.

As a check, if one considers the case of equilibrium in which

$$n_F^{\text{eq}}(k_0) = 1/(e^{\beta k_0} + 1)$$

one can easily see that formula (74) leads to the correct equilibrium Bose–Einstein distribution function

$$n_B^{\text{eq}}(k_0) = 1/(e^{\beta k_0} - 1).$$

5. Summary and discussion

The path-integral approach originally developed for bosonization in $D = 2$ space–time dimensions as a particular case of a duality transformation [11, 12] has been very successful particularly because it allows us to make simple extensions to $D > 2$ dimensions. The duality process amounts to gauging the global symmetry of the original (fermionic) theory, and constraining the corresponding field strength to vanish by introducing a Lagrange multiplier that is finally identified as the dual bosonic field. In this way one can find, for example, that in the large mass limit of a $D = 3$ fermionic theory bosonization leads to a Chern–Simons model, this leading to the existence of operators of the Fermi theory which ought to exhibit fractional statistics [12]. Also in the case of fermions coupled to gravity, bosonization can be achieved and in this case the parity breaking that takes place in $D = 3$ Dirac fermion models can be exploited to simulate the effects of crystal defects on the electronic degrees of freedom of topological insulators in condensed matter physics [28–30].

We have presented in this work an extension of the above referred duality approach to bosonization to the case of out of equilibrium fermion models based in the Schwinger–Keldysh formalism. In this way starting from the generating functional $Z_F[s]$ for $D = 2$ Dirac fermions coupled to an external source s_μ written in terms of a path-integral in which the action S_F introduced in (3) is defined in a closed time contour we were able to show the identity of $Z_F[s]$ with $Z_B[s]$, bosonic generating functional in which the action (59) is also defined in the closed time contour. As in the fermionic case, the apparently decoupled boson field defined in each branch of the time contour are in fact connected through the boundary conditions (57).

Concerning the bosonization rules for currents, one finds in each branch of the closed time contour the usual result corresponding to a topologically conserved current $j_\pm^\mu \propto \epsilon^{\mu\nu} \partial\phi_\pm$ whose spatial integral coincides with the Pontryagin class, this showing the connection of bosonization with the existence of quantum anomalies, in the present case with the $D = 2$ chiral anomaly which arises through the non-triviality of the Fujikawa Jacobian [20].

By studying current–current correlations within our bosonization approach we have been able to find in a very simple way the relation between the fermionic n_F and bosonic n_B distribution functions for a generic non-equilibrium situation, equation (74). This relation agrees with the results of reference [9]. The connection leads to the correct Bose–Einstein distribution function in equilibrium if n_F is taken as the Fermi–Dirac distribution function.

Although we have considered the free Dirac fermion Lagrangian coupled to an external source, as in the equilibrium case the bosonization rules (61)–(64) are universal in the sense

that they hold for interacting fermion models [1]. Indeed, as first observed in the case of the massless and massive self-interacting Luttinger/Thirring model where bosonization was originally derived within the operator approach [2, 3, 6, 7], the bosonization rules for fermion currents are independent of the coupling constants of the theory thus coinciding with those for free fermions. Hence, a current–current self-interaction as in the Luttinger/Thirring model can be automatically bosonized using the free fermion recipe. This can be easily understood in the path-integral framework where the current–current interaction can be traded to a linear one using a Hubbard–Stratonovich transformation [23, 25]. The same holds in the case of fermions coupled to Abelian and non-Abelian gauge theories [21, 22–24]. We plan for future work the analysis of these models in the out of equilibrium case.

As stated in the introduction, out of equilibrium bosonization was first developed in reference [9] in a functional integral approach that differs from ours in the way the source terms are handled. The main difference lies in the fact that using the duality technique that we employed here, the external source can be factored out from the path-integral that allows us to calculate the bosonic action in a closed form and only remains linearly coupled to the dual boson field leading to the correct bosonized recipe. In spite of this difference, both approaches lead to the same relation between distribution functions.

We already mentioned that the duality approach to bosonization of fermions in equilibrium can be easily extended to $D > 2$ space–time dimensions so that the analysis presented here for the out of equilibrium case could be applied with no apparent problems to higher dimensions. Of particular relevance for condensed matter problems is the $D = 3$ case in connection with fractional statistics and when coupled to a gravitational background, in applications to study the problem of topological insulators.

The bosonization approach that we presented is based on duality transformations related to the global $U(1)$ symmetry of the original fermion theory. Since duality transformations can be easily extended to the non-Abelian case, one could develop without much effort an out of equilibrium non-Abelian bosonization along the lines of reference [12].

We expect to come back to all these issues in a future work.

Acknowledgments

This work was supported by CONICET, ANPCYT, CIC and UNLP, Argentina

Appendix A. Computation of the fermionic determinant: the Fujikawa Jacobian in Schwinger–Keldysh framework

In order to compute the fermionic determinant

$$\det(\mathcal{D}^0 + \mathcal{B}) = \int \mathcal{D}\bar{\Psi} \int \mathcal{D}\Psi \exp\left(i \int d^2x \bar{\Psi} (\mathcal{D}^0 + \mathcal{B}) \Psi\right) \quad (\text{A.1})$$

we follow [21] and perform a change in the path-integral fermionic variables

$$\bar{\Psi} = U \bar{\Psi}' \quad \bar{\Psi} = \bar{\Psi}' \bar{U}, \quad (\text{A.2})$$

where U and \bar{U} are given by

$$U = \exp i\gamma_5 \varphi \quad \bar{U} = \exp i\gamma_5 \varphi. \quad (\text{A.3})$$

In these expressions φ is a scalar field that belongs to the algebra generated by the λ^a matrices ($\varphi = \varphi_a \lambda^a$) and $\gamma_5 = \gamma_0 \gamma_1$, with the two-dimensional Dirac matrices

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

As explained in section 3, scalar fields like φ defined in the Keldysh closed time contour obey the following and should obey the boundary conditions (see equations (57))

$$\begin{aligned} \varphi_+(x, t = +\infty) &= \varphi_-(x, t = +\infty) \\ \varphi_-(x, t = -\infty) &= \varphi_\tau(x, t = 0) \\ \varphi_+(x, t = -\infty) &= \varphi_\tau(x, t = \beta) \end{aligned} \quad (\text{A.4})$$

Now one can always choose a gauge (the ‘decoupling gauge’ introduced in reference [22]) in which \mathcal{B} can be written in terms of a scalar field φ in the form

$$\mathcal{B}^{\text{dec}} = -i \bar{U} \not{\partial} U = \not{\partial} \gamma_5 \varphi, \quad (\text{A.5})$$

Boundary conditions for \mathcal{B}^{dec} coincide with those for φ given in equation (A.5)

After the change of variables (A.2)–(A.3) one then ends with

$$\det(\mathcal{D}^0 + \mathcal{B}^{\text{dec}}) = J_{\text{F}}^{\text{dec}}[\varphi] \det \mathcal{D}^0 \quad (\text{A.6})$$

where $J_{\text{F}}^{\text{dec}}[\varphi]$ is the Fujikawa Jacobian [20] associated to the change of variables

$$\mathcal{D} \bar{\Psi} \mathcal{D} \Psi = J_{\text{F}}^{\text{dec}} \mathcal{D} \bar{\Psi}' \mathcal{D} \Psi'. \quad (\text{A.7})$$

Note that the \mathcal{D}^0 determinant resulting from the $\bar{\Psi}', \Psi$ path-integration is a constant independent of \mathcal{B} . It should be stressed that Fujikawa’s method implies a gauge invariant regularization of the determinant so that the answer obtained in the decoupling gauge is independent of the gauge choice.

In the case of thermodynamical equilibrium, without the introduction of the closed time contour, J_{F} has been carefully evaluated [22–24]. Interestingly, we shall see that the main computational scheme remains valid when considering the closed time path.

The presence of Jacobian J_{F} is due to the non-invariance of the path-integral measure under the chiral factor $\exp(i\gamma_5 \varphi)$ in the change of variables (A.2). Even in the context of equilibrium QFTs, where, usually, only the forward time path is considered, the fermionic Jacobian J_{F} is directly connected with the chiral anomaly, as Fujikawa originally showed [20]. Its non-triviality is due to the necessity, in its computation, of a proper regularization that depends on the symmetries to be preserved on physical grounds [27]. This Jacobian is at the heart of the path-integral approach to fermion bosonization in two dimensional space–time fermion theories [1].

Let us consider an extended transformation U_τ , depending on a parameter τ ($\tau \in [0, 1]$):

$$U_\tau = \exp i\tau \gamma_5 \varphi. \quad (\text{A.8})$$

and define a τ -dependent operator \mathcal{D}^τ as

$$\mathcal{D}^\tau = \mathcal{D}^0 + (1 - \tau)\not{\partial}\gamma_5\varphi, \quad (\text{A.9})$$

such that for $\tau = 0$ it coincides with the original operator $\mathcal{D}^0 + \mathcal{B}$ and for $\tau = 1$ becomes the free operator. Next, we introduce the quantity

$$W(\tau) = \log \det \mathcal{D}^\tau, \quad (\text{A.10})$$

such that

$$- \int_0^1 \frac{dW(\tau)}{d\tau} d\tau = \log \mathcal{D}^\tau - \log \mathcal{D}^0. \quad (\text{A.11})$$

Then, in view of equations (A.6)–(A.11) we have

$$J_{\text{F}}^{\text{dec}} = \exp \left(- \int_0^1 \frac{dW(\tau)}{d\tau} d\tau \right). \quad (\text{A.12})$$

Once this Jacobian is calculated, equation (A.6) leads to an explicit form for the fermion determinant .

Now from equation (A.10), the integrand in the exponential argument can be written as

$$\frac{dW(\tau)}{d\tau} = \frac{d \log \det \mathcal{D}^\tau}{d\tau} = \text{Tr}(\mathcal{D}^\tau)^{-1} \frac{d\mathcal{D}^\tau}{d\tau}, \quad (\text{A.13})$$

where Tr denotes a trace over Lorentz, coordinate space and also over matrices λ^a . The calculation of $dW(\tau)/d\tau$ involves a divergence coming from the evaluation of the fermionic Green function $(\mathcal{D}^\tau)^{-1}(x, y)$ at the same space–time point. To see this we write the rhs of (A.13) in a more explicit way:

$$\frac{dW(\tau)}{d\tau} = \lim_{y \rightarrow x} \text{tr}_{\text{L}} \otimes \text{tr}_{\text{K}} \int d^2y B(x, y, \tau), \quad (\text{A.14})$$

where $B(x, y, \tau) = [\mathcal{D}^\tau]^{-1}(x, y)\not{\partial}_y\gamma_5\varphi(y)$. It is convenient to specify the equation obeyed by the Green function $[\mathcal{D}^\tau]^{-1}(x, y)$. In Lorentz space, both \mathcal{D}^τ and $[\mathcal{D}^\tau]^{-1}$ have an anti-diagonal matrix structure:

$$\mathcal{D}^\tau [\mathcal{D}^\tau]^{-1} = \begin{pmatrix} 0 & \mathcal{D}_{L,x}^\tau \\ \mathcal{D}_{R,x}^\tau & 0 \end{pmatrix} \begin{pmatrix} 0 & \mathcal{G}_R(x, y, \tau) \\ \mathcal{G}_L(x, y, \tau) & 0 \end{pmatrix} = \delta^2(x - y) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

where the subscripts L and R indicate the chiral (left and right) components of each fermion field. Of course, as explained before, \mathcal{G}_R and \mathcal{G}_L have, in turn, a triangular matrix structure in Keldysh space (see section 2). On the other hand, $[\mathcal{D}^\tau]^{-1}(x, y)$ is related to the free Green function as

$$[\mathcal{D}^\tau]^{-1}(x, y) \equiv \mathcal{G}(x, y) = \exp i(1 - \tau)\gamma_5\varphi(x) \mathcal{G}^0(x, y) \exp i(1 - \tau)\gamma_5\varphi(y),$$

with

$$\mathcal{G}^0(x, y) = \begin{pmatrix} 0 & \mathcal{G}_R^0(x, y) \\ \mathcal{G}_L^0(x, y) & 0 \end{pmatrix}.$$

Performing the corresponding matrix products (in Lorentz space), one can see that the integrand in (A.14) is a diagonal matrix:

$$B(x, y, \tau) = \begin{pmatrix} B_R(x, y, \tau) & 0 \\ 0 & B_L(x, y, \tau) \end{pmatrix},$$

where

$$B_{R,L}(x, y, \tau) = \pm e^{\pm i(1-\tau)\varphi(x)} \mathcal{G}_{R,L}^0(x, y) e^{\mp i(1-\tau)\varphi(y)} (\partial_0 \mp \partial_1)\varphi(y).$$

where the upper (lower) sign corresponds to B_R (B_L).

After performing the Lorentz trace (A.14) then reads

$$\frac{dW(\tau)}{d\tau} = \lim_{y \rightarrow x} \text{tr}_K \int d^2y (B_R(x, y, \tau) + B_L(x, y, \tau)). \quad (\text{A.15})$$

Here we note that, as anticipated, $\lim_{y \rightarrow x} B_{R,L}(x, y, \tau)$ is divergent. Indeed, in this limit the fermion functions $\mathcal{G}_{R,L}(x, x)$ represent products of fermion fields at the same point. We shall adopt a point-splitting regularization which amounts to introducing a two-vector ϵ and define the regularized Green functions through a *symmetric* $\epsilon \rightarrow 0$ limit:

$$\lim_{y \rightarrow x} \mathcal{G}_{R,L}(x, y) \equiv \lim_{\epsilon \rightarrow 0} \frac{\mathcal{G}_{R,L}(x, x + \epsilon) + \mathcal{G}_{R,L}(x, x - \epsilon)}{2}, \quad (\text{A.16})$$

with

$$\lim_{\epsilon \rightarrow 0} \frac{\epsilon^\mu \epsilon^\nu}{\epsilon^\rho \epsilon_\rho} = \frac{g^{\mu\nu}}{2}, \quad (\text{A.17})$$

and

$$\lim_{\epsilon \rightarrow 0} \frac{\epsilon^\mu}{\epsilon^\rho \epsilon_\rho} = 0, \quad (\text{A.18})$$

where $g^{\mu\nu}$ is the metric tensor. This leads to the following expressions:

$$\begin{aligned} \lim_{y \rightarrow x} B_{R,L}(x, y, \tau) &= \pm \lim_{\epsilon \rightarrow 0} (e^{\pm i(1-\tau)\varphi(x)} \frac{\mathcal{G}_{R,L}^0(x, x + \epsilon) + \mathcal{G}_{R,L}^0(x, x - \epsilon)}{2} e^{\mp i(1-\tau)\varphi(x)} \\ &\mp e^{\pm i(1-\tau)\varphi(x)} \frac{\mathcal{G}_{R,L}^0(x, x + \epsilon) - \mathcal{G}_{R,L}^0(x, x - \epsilon)}{2} \\ &\times e^{\mp i(1-\tau)\varphi(x)} \epsilon^\mu \partial_\mu \varphi(x)) (\partial_0 \mp \partial_1)\varphi(y). \end{aligned} \quad (\text{A.19})$$

At this point, in order to go further, we must recall that we have to perform the product of the λ^a in $\varphi = \varphi_a \lambda^a$ and in the triangular Green function matrices

$$\mathcal{G}_{R,L}^0 = \begin{pmatrix} \mathcal{G}_{R,L,\text{ret}}^0 & \mathcal{G}_{R,L,\text{K}}^0 \\ 0 & \mathcal{G}_{R,L,\text{adv}}^0 \end{pmatrix}. \quad (\text{A.20})$$

Now we have to specify the precise form of the Green functions. For simplicity we will use an equilibrium distribution at zero temperature. Interestingly, this choice does not imply any loss of generality, since we have confirmed that the regularization performed with Green functions corresponding to a generic out of equilibrium distribution yields the same result for J_F . Then we employ the functions

$$\mathcal{G}_{R,L,\text{ret}}^0(z, 0) = \frac{\theta(z_0)}{2\pi} \left(\frac{1}{z^0 \mp z^1 + i\delta} - \frac{1}{z^0 \mp z^1 - i\delta} \right), \quad (\text{A.21})$$

$$\mathcal{G}_{R,L,\text{adv}}^0(z, 0) = \frac{-\theta(-z_0)}{2\pi} \left(\frac{1}{z^0 \mp z^1 + i\delta} - \frac{1}{z^0 \mp z^1 - i\delta} \right), \quad (\text{A.22})$$

$$\mathcal{G}_{R,L,\text{K}}^0(z, 0) = \frac{-1}{2\pi} \frac{1}{z^0 \mp z^1 + i\delta}, \quad (\text{A.23})$$

with $\delta > 0$.

We find that the symmetric $\epsilon \rightarrow 0$ limit of $\frac{\mathcal{G}_{R,L}^0(x, x+\epsilon) + \mathcal{G}_{R,L}^0(x, x-\epsilon)}{2} \epsilon^\mu$ vanishes. Concerning $\frac{\mathcal{G}_{R,L}^0(x, x+\epsilon) - \mathcal{G}_{R,L}^0(x, x-\epsilon)}{2}$, only its Keldysh component will give a non-vanishing contribution, when contracted with the ϵ^μ factor, in the second term of (A.19):

$$\lim_{\epsilon \rightarrow 0} \frac{\mathcal{G}_{R,L}^0(x, x+\epsilon) - \mathcal{G}_{R,L}^0(x, x-\epsilon)}{2} \epsilon^\mu = \frac{g^{0\mu} \mp g^{1\mu}}{4\pi} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad (\text{A.24})$$

where here $g^{\mu\nu}$ is the metric tensor.

This result is valid for an arbitrary non-equilibrium $n_F(k_0)$. To see this we concentrate on the only component of the Green function that survives the symmetric limit, the Keldysh one. For simplicity we consider the R component which in the general situation takes the form (compare with (A.23))

$$\mathcal{G}_{R,K}^0(v, 0) = \frac{-1}{2\pi} \lim_{a \rightarrow 0} \int_0^\infty dk^0 e^{-ak^0} (1 - 2n_F(k_0)) \sin k^0 v, \quad (\text{A.25})$$

Here $v = z^0 - z^1$ is a light-cone coordinate and a an ultraviolet regulator. It is straightforward to get

$$\mathcal{G}_{R,K}^0(v, 0) = \frac{-1}{2\pi v} + O(v), \quad (\text{A.26})$$

which, when used in the lhs of (A.24) yields the same symmetric limit as in the equilibrium case.

Coming back to $dW(\tau)/d\tau$, we can now compute the regularized matrix products in (A.19) and take the Keldysh trace in (A.15), obtaining

$$\frac{dW(\tau)}{d\tau} = \frac{i(\tau - 1)}{\pi} \int d^2x ((\partial_0 + \partial_1)\varphi_q(\partial_0 - \partial_1)\varphi_c + (\partial_0 + \partial_1)\varphi_c(\partial_0 - \partial_1)\varphi_q). \quad (\text{A.27})$$

Replacing in (A.12) and integrating over τ we get

$$J_F^{\text{dec}}[\varphi] = \exp \frac{i}{2\pi} \int d^2x ((\partial_0 + \partial_1)\varphi_q(\partial_0 - \partial_1)\varphi_c + (\partial_0 + \partial_1)\varphi_c(\partial_0 - \partial_1)\varphi_q) \quad (\text{A.28})$$

or, writing the Jacobian in terms of $\mathcal{B}^d[\varphi]$

$$J_F^{\text{dec}} = \exp \left(-i \frac{1}{4\pi} \text{tr}_K \int d^2x \mathcal{B}_\mu^{\text{dec}} \lambda^q \mathcal{B}^{\text{dec}\mu} \right). \quad (\text{A.29})$$

This result has been obtained in the decoupling gauge in which B_μ was written in the form (A.5). Going from this gauge to an arbitrary one amounts to writing

$$\mathcal{B}^{\text{dec}} \rightarrow \mathcal{B} = -i \bar{U} \not{\partial} U + \not{\partial} \eta = \not{\partial} \gamma_5 \varphi + \not{\partial} \eta \quad (\text{A.30})$$

where the scalar field η satisfies the same boundary conditions as φ and so does \mathcal{B} . Then instead of (A.28) one has, in an arbitrary general case

$$J_F[\varphi, \eta] = J_F^{\text{dec}}[\varphi] \times J_F^{\text{dec}}[\eta] = \exp \left(-i \frac{1}{4\pi} \text{tr}_K \int d^2x \mathcal{B}_\mu \lambda^q \mathcal{B}^\mu \right). \quad (\text{A.31})$$

References

- [1] Stone M D 1993 *Bosonization* (Singapore: World Scientific)
- [2] Coleman S 1975 *Phys. Rev. D* **11** 2088
- [3] Mandelstam S 1975 *Phys. Rev. D* **11** 3026
- [4] Witten E 1984 *Commun. Math. Phys.* **92** 455
- [5] Núñez C, Olsen K and Schiappa R 2000 *J. High Energy Phys.* **JHEP07(2000)030**
- [6] Mattis D C 1974 *Phys. Rev. Lett.* **32** 714
- [7] Luther A and Peschel I 1974 *Phys. Rev. Lett.* **32** 992
- [8] Fradkin E 2013 *Field Theories in Condensed Matter Systems* 2nd edn (Cambridge: Cambridge University Press)
- Giamarchi T 2003 *Quantum Physics in One Dimension* (Oxford: Oxford Science Publications)
- Gogolin A O, Nersisyan A A and Tsvelik A M 2004 *Bosonization and Strongly Correlated Systems* (Cambridge: Cambridge University Press)
- [9] Gutman D B, Yuval Gefen and Mirlin A D 2010 *Phys. Rev. B* **81** 085436
- [10] Chen Y-F, Dirks T, Al-Zoubi G, Birge N O and Mason N 2009 *Phys. Rev. Lett.* **102** 036804
- Dirks T, Chen Y-F, Birge N-O and Mason N 2009 *Appl. Phys. Lett.* **95** 192103
- Altimiras C, le Sueur H, Gennser U, Cavanna A, Mailly D and Pierre F 2010 *Nat. Phys.* **6** 34
- Altimiras C, le Sueur H, Gennser U, Cavanna A, Mailly D and Pierre F 2010 *Phys. Rev. Lett.* **105** 226804
- Altimiras C, le Sueur H, Gennser U, Anthore A, Cavanna A, Mailly D and Pierre F 2012 *Phys. Rev. Lett.* **109** 026803
- [11] Burgess C P, Lutken C A and Quevedo F 1994 *Phys. Lett. B* **336** 18
- [12] Fradkin E and Schaposnik F A 1994 *Phys. Lett. B* **338** 253
- Le Guillou J C, Moreno E, Núñez C and Schaposnik F 1997 *Nucl. Phys. B* **484** 682
- [13] Giveon A, Porrati M and Rabinovici E 1994 *Phys. Rep.* **244** 77
- [14] Schwinger J 1961 *J. Math. Phys.* **2** 407
- [15] Keldysh L V 1964 *Zh. Eksp. Teor. Phys.* **47** 1515
- Keldysh L V 1965 *Sov. Phys.-JETP* **20** 1018
- [16] Babichenko V S and Kozlov A N 1986 *Solid State. Commun.* **59** 39
- [17] Kiselev M N and Oppermann R 2000 *Phys. Rev. Lett.* **85** 5631
- [18] Das A 1997 *Finite Temperature Field Theory* (Singapore: World Scientific)
- Kamenev A and Levchenko A 2009 *Adv. Phys.* **58** 197
- Kamenev A 2011 *Field Theories of Non-Equilibrium systems* (Cambridge: Cambridge University Press)
- Altland A and Simons B D 2010 *Condensed Matter Field Theory* (Cambridge: Cambridge University Press)
- [19] Larkin A I and Ovchinnikov Y N 1975 *Zh. Eksp. Teor. Fiz.* **68** 1915
- Larkin A I and Ovchinnikov Y N 1976 *Sov. Phys.—JETP* **41** 960
- [20] Fujikawa K 1979 *Phys. Rev. Lett.* **42** 1195
- Fujikawa K 1980 *Phys. Rev. D* **21** 2848
- Fujikawa K and Suzuki H 2004 *Path Integrals and Quantum Anomalies* (Oxford: Oxford Science Publications)
- [21] Roskies R and Schaposnik F A 1981 *Phys. Rev. D* **23** 558
- [22] Gamboa Saraví R E, Schaposnik F A and Solomin J E 1981 *Nucl. Phys. B* **185** 239
- [23] Furuya K, Gamboa Saraví R E and Schaposnik F A 1982 *Nucl. Phys. B* **208** 159
- [24] Gamboa Saraví R E, Schaposnik F A and Solomin J E 1984 *Phys. Rev. D* **30** 1353
- [25] Naón C M 1985 *Phys. Rev. D* **31** 2035
- [26] Alvarez-Gaume L and Ginsparg P H 1984 *Nucl. Phys. B* **243** 449
- [27] Cabra D and Schaposnik F 1989 *J. Math. Phys.* **30** 816
- [28] Hughes T L, Leigh R G and Fradkin E 2011 *Phys. Rev. Lett.* **107** 075502
- [29] Hughes T L, Leigh R G and Parrikar O 2013 *Phys. Rev. D* **88** 025040
- [30] Fradkin E, Moreno E F and Schaposnik F A 2014 *Phys. Lett. B* **730** 284
- Fujikawa K 1980 *Phys. Rev. D* **21** 2848
- Fujikawa K and Suzuki H 2004 *Path Integrals and Quantum Anomalies* (Oxford: Oxford Science Publications)