



Original Article

Alterations in biochemical parameters of the freshwater fish, *Alburnus mossulensis*, exposed to sub-lethal concentrations of Fenprothrin

Mahdi Banaee^{*1}, Antoni Sureda², Fazel Zohiery¹, Behzad Nematdoust Hagi¹, Daniela S. Garanzini³

¹Aquaculture Department, Natural Resource Faculty, Behbahan Khatam Alanbia University of Technology, Iran.

²Laboratori de Nutrició Comunitària i Estrès Oxidatiu; Department of Biology Science, Balearic Islands University, IllesBalears, Spain.

³Laboratorio de Ecotoxicología, Instituto de Investigaciones Marinas y Costeras (CONICET/ Universidad Nacional de Mar del Plata), Funes 3350 (B7602AYJ) - Mar del Plata – Argentina.

Abstract: Fenprothrin is a new pyrethroid insecticide used to control crop pests. The aim of this study was to evidence fenprothrin-induced oxidative stress and alterations in biochemical parameters in the freshwater fish, *Alburnus mossulensis*. Total antioxidant capacity, malondialdehyde (MDA), catalase activity (CAT), aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), creatine phosphokinase (CK), acetylcholinesterase (AChE) in the whole body extract were measured in *A. mossulensis* after exposure to sub-lethal concentrations of fenprothrin (approximately equal to 1, 2, 3, 5 and 10% of 96 h LC₅₀) for 15 days. The 24, 48, 72 and 96 h LC₅₀ of the fenprothrin for *A. mossulensis* was 562.28±45.19, 218.18±18.75, 136.18±11.90 and 121.38±11.84 µg/L at 24±2 °C. Exposure to 2.75, 5.50 and 12.6 µg/L fenprothrin significantly increased AST activity in fish. A significant increase in the ALP and LDH activities was observed in fish after a 15 day exposure to 1.25, 5.50 and 12.60 µg/L fenprothrin. Fenprothrin significantly induced lipid peroxidation and increased MDA levels in fish. Compared with the control group, total protein levels in fish decreased after exposure to 2.75, 5.50 and 12.60 µg/L fenprothrin on day 15. Total antioxidant capacity, AChE and CPK activities in fish exposed to fenprothrin were significantly lower than control group. There was a significant increase in the CAT and ALT activities in fish after exposure to 5.50 and 12.60 µg/L fenprothrin. In conclusion, fenprothrin has the potential to disrupt biochemical parameters in *A. mossulensis* and to induce oxidative stress.

Article history:

Received 11 December 2013

Accepted 12 March 2014

Available online 25 April 2014

Keywords:

Fenprothrin
Oxidative stress
Lipid peroxidation,
Biochemical parameters
Alburnus mossulensis

Introduction

The use of pyrethroid insecticides has recently increased compared to organochlorine pesticides due to their high selectivity and lower persistence in the environment (Goulding et al., 2013). Fenprothrin [2, 2, 3- tetramethylcyclopropanecarboxylic acid cyano (3-phenoxyphenyl) methyl ester] is a new pyrethroid insecticide used to control crop pests in tomatoes, tobacco, cotton farms and fruit orchards. Pyrethroid insecticides enter surface water via air drift, leaching from agricultural land and surface runoff during or after the application of pesticides

(Gan et al., 2005; Woudneh and Oros, 2006; Weston et al., 2009; Goulding et al., 2013). Although there is no report indicating the presence of fenprothrin in surface water and groundwater, the presence of other pyrethroid pesticides in sediment, surface water (Amweg et al., 2006; Holmes et al., 2008; Feo et al., 2010), deltaic regions of big rivers (Feo et al., 2010; Weston and Lydy, 2010) and estuaries (Woudneh and Oros, 2006) and agriculture wastewater (You et al., 2004) have been reported.

Pyrethroid pesticides are lipophilic compounds, which may be absorbed through gills, skin or

* Corresponding author: Mahdi Banaee
E-mail address: banaee@bkatu.ac.ir

alimentary ducts. Food poisoning with pyrethroids is another influential pathway of these pesticides on fish (Macneale et al., 2010). After entering the body, pyrethroid pesticides can be accumulated in fat tissues (Banaee, 2013). Pyrethroid pesticides in animal's body are rapidly hydrolyzed in the body of the exposed animals and their metabolites excreted. Carboxy esterases play an important role in the detoxification of these pesticides (Sogorb and Vilanova, 2002). Reactive oxygen species (ROS) produced during the detoxification process of pyrethroid pesticides are known as important agents that cause oxidative stress (Ansari et al., 2011; El-Demerdash, 2011). In addition, alterations in biochemical parameters (Das and Mukherjee, 2003; Parvez and Raisuddin, 2005), and hematologic factors (Saxena and Seth, 2002; Muranli and Güner, 2011), reproductive disorders (Moore and Waring, 2001), neurological disorders (Soderlund et al., 2002), histopathological changes (Velmurugan et al., 2007; Korkmaz et al., 2009; Kan et al., 2012), and mutations (Ansari et al., 2011; Beggel et al., 2011), and reduced survival rate of larvae and embryos (Köprücü and Aydın, 2004; Ural and Sağlam, 2005), are reported in several fish species exposed to various pyrethroid pesticides.

Werner and Moran (2008) suggested that acute toxicity of pyrethroids to fish is usually higher than 200 ng/L. However, synthetic pyrethroid insecticides are highly toxic to fish and aquatic invertebrates (Polat et al., 2002; Datta, A. Kaviraj, 2003; Ural and Sağlam, 2005; Feo et al., 2010). There is little information on the toxicity of fenprothrin to aquatic organisms.

Alburnus species belong to the Cyprinidae family that is found in most rivers in Iran. The wide distribution of the freshwater fish *Alburnus* sp. makes this species a potential and useful biomarker for monitoring aquatic ecosystems. Therefore, the aim of the present study was to evaluate the changes in some biochemical parameters of the freshwater fish *Alburnus mossulensis* after exposure to sub-lethal concentrations of fenprothrin.

Materials and Methods

Fish: Freshwater fish, *Alburnus mossulensis*, 7.36±1.60 g were netted from the Maroon River, Khuzestan Province, Iran. Fishes were kept in glass aquaria with dechlorinated water to acclimatize them to laboratory conditions (24±2°C, pH: 7.4±0.2, hardness 355±25 mg/L of CaCO₃ and 16 L: 8 D photoperiods) for two weeks prior to use. Specimens were fed with live food and commercial diet for ornamental fish.

Acute toxicity test: The acute toxicity test was performed according to semi-static methods described in the OECD procedure. Fishes were not fed 24 h before the experiments and during the acute toxicity test. The experiments consisted of a control group and five experimental groups. Acute test was performed to determine the appropriate toxicity range for the sub-lethal assay. 10 fish per group were exposed to different fenprothrin concentrations (0.0, 10, 50, 100, 250, 500 and 1000 µg/L fenprothrin, purity 30%) in 85 L aquarium. During the 96 h acute toxicity experiment, water in each aquarium was aerated and had the same conditions as the acclimation period. Test solutions were renewed every 24 h to maintain the chemical and the water quality. Every 24 h the dead fish were removed and the number of survivals was recorded. The experiment was repeated in triplicate. LC₅₀ values were calculated by the Probit Analysis test (Banaee et al., 2011).

Sub-lethal toxicity test: For sub-lethal toxicity tests, the concentrations of fenprothrin in water were maintained modestly below the 96 h LC₅₀ value. Based on this value, five sub-lethal concentrations 1, 2, 3, 5 and 10% of 96 h LC₅₀ were chosen for the freshwater fish (*Alburnus mossulensis*). The fish were divided into five treatments and a control group by triplicate (10 specimens per each aquarium). Test solutions in each aquarium were renewed every 24 h. On the other hand, the twenty percent of water was changed daily to reduce the build-up of metabolic wastes and to keep concentrations of fenprothrin near the nominal level. During the experiment, fishes

Table 1. Numerical value of lethal concentrations at different times

LC	Numerical value of lethal concentrations at different times			
	24 h	48 h	72 h	96 h
LC ₁₀	149.28±45.74	58.045±20.15	56.04±11.31	49.34±10.15
LC ₂₀	291.05±39.32	113.01±17.25	83.55±10.19	74.07±9.12
LC ₃₀	393.28±38.67	152.65±16.66	103.39±10.26	91.90±9.49
LC ₄₀	480.63±40.99	186.52±17.30	120.34±10.90	107.14±10.48
LC ₅₀	562.28±45.19	218.18±18.75	136.18±11.90	121.38±11.84
LC ₆₀	643.92±50.88	249.84±20.86	152.02±13.21	135.62±13.48
LC ₇₀	731.27±58.11	283.71 ±23.65	168.97±14.86	150.85±15.45
LC ₈₀	833.50±67.56	323.35±27.38	188.80±17.02	168.69±17.94
LC ₉₀	975.27±81.79	378.32±33.09	216.31±20.26	193.42±21.59
LC ₉₉	1311.97±118.12	508.87 ±47.86	281.64 ±28.61	252.15±30.75

were fed twice a day with commercial food, and the fish mortality was recorded.

At the end of the experimental period, on day 15, all fish per treatment were captured and anaesthetized with a clove powder extract. Then, they were sacrificed and washed with buffered normal saline. Fish were homogenized during two minutes in ice cold phosphate buffer (pH 7.4; 1:10, w/v) using a glass homogenizer and then centrifuged for 15 min at 15000 g at 4°C with a refrigerated centrifuge. Supernatants were immediately used to measure biochemical parameters by using spectrophotometric assays.

Biochemical Analysis: Lactate dehydrogenase (LDH) activity determination is based on measuring the conversion of pyruvate to L-lactate by monitoring the NADH oxidation. Aspartate aminotransferase (AST) was assayed in a coupled reaction with malate dehydrogenase in the presence of NADH. In the alanine aminotransferase (ALT) assay, the enzyme reacts with alanine and α -ketoglutarate to form glutamate and pyruvate. LDH converts pyruvate to lactate and NAD^+ . In the determination of creatine phosphokinase (CK) activity, the enzyme reacts with creatine phosphate and ADP to form ATP, which is coupled to the hexokinase/GDP reaction generating NADPH. All these activities were monitored by measuring the change in absorbance at 340 nm. Alkaline

phosphatase (ALP) assay is based on the enzyme-mediated conversion of *p*-nitrophenol phosphate to nitrophenol in an alkaline buffer at 405 nm (Moss and Henderson, 1999). Acetylcholinesterase (AChE) activity was determined by adding an adequate volume of the sample into a cuvette containing 0.1 M phosphate pH 8.0, and acetylcholine iodide (0.015 M) and dithiobisnitrobenzoic acid (0.01 M) as substrates. AChE activity was recorded during 180 s at 405 nm (Thomas, 1998). Levels of protein in liver tissue were determined by standard procedures used in clinical biochemistry laboratories based on manual biochemical kits (ParsAzemon Co, Iran) (Johnson et al., 1999).

Total antioxidant capacity was assayed according to the ferric reducing ability of plasma (FRAP) method. Briefly, the FRAP reagent contained 5 mL of a (10 mmol/L) TPTZ (2,4,6- tripyridyl- s- triazine) solution in 40 mmol/L HCL plus 5 mL of FeCl_3 (20 mmol/L) and 50 mL of acetate buffer, (0.3 mol/L, pH=3.6) and was prepared freshly. Aliquots of 100 μL supernatant were mixed with 3 mL FRAP reagent. The conversion rate of ferric tripyridyl-s-triazine (Fe^{3+} -TPTZ) complex to ferrous tripyridyl-s-triazine (Fe^{2+} -TPTZ) at pH 3.6 and temperature 25°C is directly proportional to the concentration of total antioxidant in the sample. Fe^{2+} -TPTZ has an intensive blue color that can be monitored up to 5 min at 593 nm by a UV/VIS spectrophotometer.

Table 2. Changes in the level of enzyme activities in the whole body of fish exposed to different concentrations of fenpropathrin

Concentration of Fenpropathrin ($\mu\text{g/L}$)	Enzyme activity levels (Unit per g protein tissue)					
	AST (U/g)	ALT (U/g)	ALP (U/g)	LDH (U/g)	CPK (U/g)	AChE (U/g)
0.00 $\mu\text{g/L}$	2.28 \pm 1.04 ^a	0.67 \pm 0.13 ^a	2.28 \pm 0.36 ^a	2.54 \pm 1.10 ^a	13.47 \pm 5.70 ^b	7.99 \pm 2.17 ^b
1.25 $\mu\text{g/L}$	2.33 \pm 0.97 ^a	0.51 \pm 0.14 ^a	4.21 \pm 2.03 ^{ab}	4.49 \pm 2.28 ^{ab}	5.45 \pm 0.80 ^a	5.66 \pm 2.14 ^a
2.75 $\mu\text{g/L}$	4.11 \pm 1.25 ^b	0.36 \pm 0.05 ^a	2.52 \pm 0.54 ^{ab}	4.64 \pm 2.26 ^{ab}	7.81 \pm 2.75 ^a	4.35 \pm 1.11 ^a
3.15 $\mu\text{g/L}$	3.55 \pm 1.60 ^{ab}	0.45 \pm 0.15 ^a	3.72 \pm 0.69 ^{ab}	4.05 \pm 1.34 ^{ab}	5.30 \pm 1.18 ^a	4.59 \pm 1.40 ^a
5.50 $\mu\text{g/L}$	6.00 \pm 1.11 ^c	2.40 \pm 0.50 ^c	4.88 \pm 1.29 ^b	5.94 \pm 3.07 ^b	4.65 \pm 0.93 ^a	4.29 \pm 0.48 ^a
12.60 $\mu\text{g/L}$	4.06 \pm 1.38 ^b	1.55 \pm 0.41 ^b	4.71 \pm 2.14 ^b	5.74 \pm 3.16 ^b	6.94 \pm 2.33 ^a	4.62 \pm 1.02 ^a

Calculations were performed using a calibration curve of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (100 to 1000 $\mu\text{M/L}$) (Iris et al., 1996).

Malondialdehyde (MDA) content was assessed by a modification of a thiobarbituric acid assay and was expressed as $\mu\text{mol/g}$ tissue (Placer et al., 1966). 500 μl of the supernatant was transferred to a Pyrex tube and mixed with 2500 μl trichloroacetic acid (20%) and 1000 μmL trichlorothiobarbituric acid (67%). The tubes were then placed in boiling water (100°C) for 15 min. After cooling, the mixtures were extracted with a solution containing 1000 μL of distilled water and 5000 μL *n*-butanol: pyridine (15: 1). The mixture was then centrifuged at 2000 g for 15 min at 4°C. The pink color produced by these reactions was measured spectrophotometrically at 532 nm to measure MDA levels. MDA concentration was calculated with an external standard of MDA. Tetraethoxypropane and absolute ethanol were used to prepare the MDA standards. Concentrations of MDA in whole body samples are expressed in μM per g protein.

Catalase activity was measured by an assay with hydrogen peroxide based on formation of its stable complex with ammonia molybdate. 0.2 ml of supernatant was incubated in 1 ml reaction mixture containing 65 mM hydrogen peroxide in 60 mM sodium-potassium phosphate buffer, pH 7.4 at 25°C for 4 min. The enzymatic reaction was stopped with 1 ml of 32.4 mM ammonium molybdate and concentration of the yellow complex of molybdate and hydrogen peroxide was measured at 405 nm. All chemical materials were obtained from Merck Co, Germany. All biochemical parameters were

measured in duplicate by UV/Vis Unico spectrophotometer (Model 2100).

Statistical analysis: Statistical analyses were performed using SPSS (Release 19 IBM) software. Data are presented as mean \pm SD. All the data were tested for normality (Kolmogorov-Smirnov test). Data were analyzed by one-way of variance analysis (ANOVA). The significant means were compared by Duncan test and a $P < 0.05$ was considered statistically significant.

Results

Numerical value of LC_{50} of fenpropathrin at 24, 48, 72 and 96 hours are presented in Table 1. Fish mortality was progressively elevated with increasing the concentration of fenpropathrin. Sub-lethal concentrations of fenpropathrin were determined according to 96h LC_{50} .

No mortality was observed in fish exposed to sub-lethal concentrations of fenpropathrin and control group during experimental periods. Loss of appetite, increased mucus secretion, increase in abnormal behavior, swimming in the surface of water and swimming vertically were the major changes observed in fish after exposure to the higher concentrations (5.50 and 12.60 $\mu\text{g/L}$) of fenpropathrin.

Changes in the enzymatic activities determined in the whole body of fish exposed to different concentrations of fenpropathrin are presented in Table 2.

AST activity in the whole body of fish exposed to 2.75, 5.50 and 12.60 $\mu\text{g/L}$ fenpropathrin was significantly higher than in the control group. A




Figure 1. Changes in the MDA levels in the whole body of fish exposed to different concentrations of fenpropathrin. Significant differences between values when compared with control groups were characterized by alphabet symbol ($P < 0.05$). Values represent mean \pm S.D.




Figure 2. Changes in the total antioxidant levels in the whole body of fish exposed to different concentrations of fenpropathrin. Significant differences between values when compared with control groups were characterized by alphabet symbol ($P < 0.05$). Values represent mean \pm S.D.

significant increase in ALT activity was observed in the whole body of fish exposed to 5.50 and 12.60 µg/L fenpropathrin when compared with control group. AChE and CPK activities significantly decreased in the whole body of fish after exposure to all concentrations of fenpropathrin. There was a significant increase in the ALP and LDH activities in the whole body of fish exposed to 1.25, 5.50 and 12.60 µg/L fenpropathrin when compared with control group.

Alterations in the malondialdehyde, total protein, total antioxidant levels and catalase activity in the whole body of fish exposed to different concentrations of fenpropathrin are presented in Figures 1 to 4.




Figure 3. Changes in the CAT activity in the whole body of fish exposed to different concentrations of fenpropathrin. Significant differences between values when compared with control groups were characterized by alphabet symbol ($P < 0.05$). Values represent mean \pm S.D.




Figure 4. Changes in the level of total protein in the whole body of fish exposed to different concentrations of fenpropathrin. Significant differences between values when compared with control groups were characterized by alphabet symbol ($P < 0.05$). Values represent mean \pm S.D.

A significant increase in MDA levels in the whole body of fish exposed to fenproparthin was observed in all treated groups when compared with control group (Fig. 1).

Total antioxidant capacity in the whole body of fish exposed to fenproparthin progressively decreased with increased concentrations of the pesticide (Fig. 2).

There was a significant increase in the CAT activity in the whole body of fish after exposure to 5.50 and 12.60 µg/L Fenproparthin respect to the control group (Fig. 3).

The results of this study showed that total protein levels in the whole body of fish exposed to 2.75, 5.50

and 12.60 µg/L fenpropathrin significantly decreased when compared with control group (Fig. 4).

Discussion

The present results show that fenpropathrin is highly toxic to freshwater fish, *A. mossulensis*. The toxicity of fenpropathrin on *A. mossulensis* increased with increasing the concentration and exposure time. When fishes were exposed to 10 µg/L fenpropathrin, only 3.34% died after 96 h, whereas all the fishes (100%) died after 48 h when were exposed to a concentration of 1000 µg/L fenpropathrin. In addition, the 24, 48, 72 and 96 h LC₅₀ values of fenpropathrin of *A. mossulensis* were calculated as 562.3±45.2, 218.2±18.8, and 136.2±11.9 and 121.4±11.8 µg/L, respectively.

Imbalance, vertical swimming and swimming in the water, loss of appetite, bleeding at the base of the fins and eye balls were important clinical symptoms observed in fish exposed to high concentrations of fenpropathrin. Behavior disorders in fish exposed to fenpropathrin maybe associated to the neurotoxicity potential of pyrethroid pesticides. Neurological disorders in animals are often attributed to a dysfunction in ion channels of neurons, particularly sodium ions after exposure to pyrethroid pesticides (Sogorb and Vilanova, 2002; Banaee, 2013). However, a decrease in AChE activity may also play a role on the behavioral changes in fish after exposure to fenpropathrin. Reduced AChE activity in the whole body of *A. mossulensis* exposed to fenpropathrin may be due to changes in this enzyme function (Tayebati et al., 2009; El-Demerdash, 2011). Decreases in the AChE activity levels in different tissues of fish were observed after exposure to chlorpyrifos (Halappa and David, 2009; Sharbidre et al., 2011), diazinon (Banaee et al., 2011), methyl parathion (Sharbidre et al., 2011), monocrotophos (Rao, 2006), and atrazine (Santos and Martinez, 2012), permethrin and deltamethrin (Goulding et al., 2013). Although, there is no information on the detoxification mechanism of fenpropathrin in fish, free radicals that are produced during the biotransformation of this pesticide in liver tissue may

be the most important cause of oxidative stress in *A. mossulensis*. Increases in MDA levels in the whole body extract of fish may be an important bio-indicator of lipid peroxidation in different tissues of freshwater fish, after exposure to fenpropathrin. An increase in MDA levels were reported in different tissues of fish exposed to diazinon (Oruç and Usta, 2007; Isik and Celik, 2008), deltamethrin (Yonar and Sakin, 2011), methyl parathion (Monteiro et al., 2006; Isik and Celik, 2008; Sharbidre et al., 2011), chlorpyrifos (Sharbidre et al., 2011), carbamazepine (Li et al., 2010), and trazine (Paulino et al., 2012).

A decrease in total antioxidant capacity was an important detrimental response of the fish antioxidant defense system to increased free radicals after exposure to fenpropathrin. Similar results were observed in rainbow trout and carp after exposure to diazinon and cyfluthrin (Sepici-Dinçel et al., 2009; Banaee et al., 2013). The overproduction of free radicals during pesticide detoxification may be associated with a decrease in the hepatic total antioxidant capacity (Monteiro et al., 2006; Banaee et al., 2013). Impairment in the synthesis of enzymatic and non-enzymatic antioxidants may be the most important factor in reducing the levels of cellular total antioxidant. Therefore, the decline in the cellular antioxidant capacity makes the cells more vulnerable to oxidative stress damage. A decrease in non-enzymatic antioxidant levels were observed in different tissue of fish exposed to chlorpyrifos (Sharbidre et al., 2011) and andatrazine (Santos and Martinez, 2012).

The elevated MDA levels and total antioxidant capacity are indicative of increased oxidative stress in fish. The results in the present study indicated that the imbalance between oxidants and antioxidants in cells was the most important factor in causing oxidative stress in fish after exposure to fenpropathrin. Similar results were observed in the sheepshead minnow (*Cyprinodon variegatus*) (Harper et al., 2008), *Channa punctata* (Sayeed et al., 2003; Atif et al., 2005), medaka (*Oryzias latipes*) (Ingeborg et al., 2002) exposed to bifenthrin esfenvalerate, respectively.

Catalase plays an important role in the elimination of hydrogen peroxide in cells (Banaee et al., 2013). Hydrogen peroxide may play a role in increasing MDA levels in the fish exposed to fenprothrin. Moreover, an increase in catalase activity in the whole body of fish after exposure to high concentrations of fenprothrin may be effective in the removal of hydrogen peroxide produced in cells during detoxification process. An increase in CAT activity in different tissues of fish exposed to atrazine (Paulino et al., 2012) was reported.

AST and ALT enzyme activities are important in cellular nitrogen metabolism, oxidation of amino acids, and liver gluconeogenesis (Banaee, 2013). In stress situations, increased activity of liver enzymes such as AST and ALT has stimulatory effects on gluconeogenic mechanism (Banaee, 2012; 2013). Thus, increased levels of these aminotransferase activities may have played an important role in energy supply for fish exposed to fenprothrin. Similar changes were observed in Labeorohita, after exposure to fenvalerate (Prusty et al., 2011). Increase in ALT and AST activities were observed in plasma, liver and kidney of *Oreochromis mossambicus*, *Cyprinus carpio* and Korean rockfish (*Sebastes schlegeli*) after exposure to monocrotophos (Rao, 2006), bifenthrin (Velisek et al., 2009) and cypermethrin (Jee et al., 2005).

The results showed that the protein of the whole body was reduced, which may be related with the increased transaminase activities. The increased activity of these enzymes may lead to protein breakdown to provide energy and regeneration of tissue damages. Fish muscle breakdown under stress situation may be one of the main reasons for decreasing tissue protein. Decreased total protein levels were reported in *Oreochromis mossambicus*, *O. niloticus*, *Cyprinus carpio*, and *Oncorhynchus mykiss* exposed to endosulfan (Kumar et al., 2011), malathion (Patil and David, 2008), diazinon (Banaee et al., 2008; Banaee et al., 2011; Banaee et al., 2013), lindane (Saravanan et al., 2011), and cypermethrin (Korkmaz et al., 2009), bifenthrin (Velisek et al., 2009) and cypermethrin (Jee et al., 2005).

Reduction of muscle mass of fish may decrease CPK activity in fish after exposure to fenprothrin. Rosalki (1998) believed that damage to connective tissues and a reduction of muscle mass are the main reasons for the decrease in the activity of the CPK enzyme.

After fish exposure to pesticides, LDH may increase to supply energy to fish. LDH is an enzyme that participates in the anaerobic pathway of carbohydrate metabolism. The increase of LDH activity is a diagnostic index widely used to recognize increases of anaerobic metabolism resulting from depletion of energy under anaerobic and environmental stress conditions (Banaee, 2012; 2013). The increase of LDH activity in the whole body extract of fish was a physiological mechanism to provide more energy to deal with the effects fenprothrin on the freshwater fish, *A. mossulensis*. Similar changes in LDH activity were observed in crayfish exposed to endosulfan (Banaee and Ahmadi, 2011). Increased LDH activity in the gill and brain of tilapia, *Oreochromis mossambicus* after exposure to monocrotophos was reported by Rao (2006). In contrast, Tripathi and Shasmal (2011) showed an inhibitory effect of chlorpyrifos on LDH activities in different tissues of the fish.

ALP plays a significant role in phosphate hydrolysis and in membrane transport and it also acts as a good bio-indicator of stress in biological systems. Increased ALP activity in the whole body extract of fish may be due to the effects of fenprothrin on transphosphorylation activity. Increases in ALP activity were reported in the whole body extract of crayfish after exposure to endosulfan (Banaee and Ahmadi, 2011). Rao (2006) reported that the ALP activities were increased in plasma, gills and kidneys of tilapia exposed to monocrotophos.

It can be concluded that fenprothrin was highly toxic to *A. mossulensis*. Exposure to sub-lethal concentrations of fenprothrin resulted in significant biochemical alterations and behavioral changes which may be potentially disruptive for the survivability of *A. mossulensis*. This fact should be taken into consideration when this pesticide is used

for pest control in agriculture fields surrounding freshwater ecosystems. In conclusion, measuring oxidative stress biomarker and other biochemical parameters in the present study was useful for monitoring the sub-lethal effects of fenprothrin on freshwater fish.

Acknowledgment

The authors are grateful to the laboratory technician, Mr. Ranjbar, for his cooperation and assistance throughout the research.

References

Amweg E.L., Weston D.P., You J., Lydy M.J. (2006). Pyrethroid insecticides and sediment toxicity in urban creeks from California and Tennessee. *Environmental Science and Technology*, 40: 1700-1706.

Ansari R.A., Rahman S., Kaur M., Anjum S., Raisuddin S. (2011). In vivo cytogenetic and oxidative stress-inducing effects of cypermethrin in freshwater fish, *Channa punctata* Bloch. *Ecotoxicology and Environmental Safety*, 74: 150-156.

Atif F., Parvez S., Pandey S., Ali M., Kaur M., Rehman H., Khan H.A., Raisuddin S. (2005). Modulatory effect of cadmium exposure on deltamethrin-induced oxidative stress in *Channa punctata* (Bloch). *Archive Environment Contamination Toxicology*, 49: 371-377.

Banaee M. (2012). Adverse effect of insecticides on various aspects of fish's biology and physiology: Insecticides – Basic and Other Applications Book, Edited by Sonia Soloneski and Marcelo Larramendy, Published by InTech, Chapter, 6: 101-126.

Banaee M. (2013). Physiological dysfunction in fish after insecticides exposure: Insecticides often Undesired but still so Important, Edited by Stanislav Trdan, Published by InTech, Chapter 4: 103-142.

Banaee M., Ahmadi, K. (2011). Sub-lethal Toxicity Impacts of Endosulfan on Some biochemical Parameters of the Freshwater Crayfish (*Astacus leptodactylus*). *Research Journal of Environmental Sciences*, 5(11): 827-835.

Banaee M., Mirvagefi A.R., Rafei G.R., Majazi Amiri, B. (2008). Effect of sub-lethal Diazinon Concentrations on Blood Plasma Biochemistry. *International Journal of Environmental Research*, 2(2): 189-198.

Banaee M., Sureda A., Mirvaghefi A.R., Ahmadi K. (2011). Effects of Diazinon on Biochemical Parameters of Blood in Rainbow Trout (*Oncorhynchus mykiss*). *Pesticide Biochemistry and Physiology*, 99: 1-6.

Banaee M., Sureda A., Mirvaghefi A.R., Ahmadi K. (2013). Biochemical and histological changes in the liver tissue of Rainbow trout (*Oncorhynchus mykiss*) exposed to sub-lethal concentrations of diazinon. *Fish Physiology and Biochemistry*, 39: 489-501.

Beggel S., Connon R., Werner I., Geist J. (2011). Changes in gene transcription and whole organism responses in larval fathead minnow (*Pimephales promelas*) following short-term exposure to the synthetic pyrethroid bifenthrin. *Aquatic Toxicology*, 105: 180-188.

Das B.K., Mukherjee S.C. (2003). Toxicity of cypermethrin in *Labeo rohita* fingerlings: biochemical, enzymatic and haematological consequences. *Comparative Biochemistry and Physiology, Part C*, 134: 109-121.

Datta M., Kaviraj A. (2003). Acute toxicity of the synthetic pyrethroid deltamethrin to freshwater cafish *Clarias gariepinus*. *Bulletin of Environmental Contamination and Toxicology*, 70: 296-299.

El-Demerdash F.M. (2011). Lipid peroxidation, oxidative stress and acetylcholinesterase in rat brain exposed to organophosphate and pyrethroid insecticides. *Food and Chemical Toxicology*, 49: 1346-1352.

Feo M.L., Ginebreda A., Eljarrat E., Barceló D. (2010). Presence of pyrethroid pesticides in water and sediments of Ebro River Delta. *Journal of Hydrology*, 393: 156-162.

Gan J., Lee S.J., Liu W.P., Haver D.L., Kabashima J.N. (2005). Distribution and persistence of pyrethroids in runoff sediments. *Journal of Environmental Quality*, 34: 836-841.

- Goulding A.T., Shelley L.K., Ross P.S., Kennedy C.J. (2013). Reduction in swimming performance in juvenile rainbow trout (*Oncorhynchus mykiss*) following sublethal exposure to pyrethroid insecticides. *Comparative Biochemistry and Physiology, Part C*, 157: 280–286.
- Halappa R., David M. (2009). Behavioural responses of the fresh water fish, *Cyprinus carpio* (Linnaeus) following sublethal exposure to chlorpyrifos. *Turkish Journal of Fisheries and Aquatic Sciences*, 9: 233-238.
- Harper H.E., Penington P.L., Hoguet J., Fulton M.H. (2008). Lethal and sublethal effects of the pyrethroid, bifenthrin, on grass shrimp (*Palaemonetes pugio*) and sheepshead minnow (*Cyprinodon variegatus*). *Journal of Environmental Health, Part B*. 43: 476-483.
- Holmes R.W., Anderson B.S., Phillips B.M., Hunt J.W., Crane D.B., Mekebri A., Connor V. (2008). Statewide investigation of the role of pyrethroid pesticides in sediment toxicity in California's urban waterways. *Environmental Science and Technology*, 42: 7003-7009.
- Ingeborg W., Juergen G., Mark O., Philipp R., David H. (2002). Effects of dietary exposure to the pyrethroid pesticide esfenvalerate on medaka (*Oryzias latipes*). *Marine Environmental Research*, 54: 609-614.
- Iris F., Benzie F., Strain J.J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical Biochemistry*, 239: 70–76
- Isik I., Celik I. (2008). Acute effects of methyl parathion and diazinon as inducers for oxidative stress on certain biomarkers in various tissues of rainbowtrout (*Oncorhynchus mykiss*). *Pesticide Biochemistry and Physiology*, 92: 38–42.
- Johnson A.M., Rohlf E.M., Silverman L.M. (1999). Proteins. In: Burtis CA, Ashwood ER, editors. *Tietz Textbook of Clinical Chemistry*. 3rd ed. Philadelphia: W.B. Saunders Company; p. 477-540.
- Kan Y., Cengiz E.I., Ugurlu P., Yanar M. (2012). The protective role of vitamin E on gill and liver tissue histopathology and micronucleus frequencies in peripheral erythrocytes of *Oreochromis niloticus* exposed to deltamethrin. *Environmental Toxicology and Pharmacology*, 34: 170–179.
- Köprücü K., Aydın R. (2004). The toxic effects of pyrethroid deltamethrin on the common carp (*Cyprinus carpio* L.) embryos and larvae. *Pesticide Biochemistry and Physiology*, 80: 47–53.
- Korkmaz N., Cengiz E.I., Unlu E., Uysal E., Yanar M. (2009). Cypermethrin-induced histopathological and biochemical changes in Nile tilapia (*Oreochromis niloticus*), and the protective and recuperative effect of ascorbic acid. *Environmental Toxicology and Pharmacology*, 28: 198–205.
- Kumar N., Prabhu P.A.J., Pal A.K., Remya S., Aklakur M., Rana R.S., Gupta S., Raman R.P., Jadhao S.B. (2011). Anti-oxidative and immunohematological status of Tilapia (*Oreochromis mossambicus*) during acute toxicity test of endosulfan. *Pesticide Biochemistry and Physiology*, 99: 45–52.
- Li Z H., Velisek J., Zlabek V., Grabic R., Machova, J., Kolarova J., Randak T. (2010). Hepatic antioxidant status and hematological parameters in rainbow trout, *Oncorhynchus mykiss*, after chronic exposure to carbamazepine. *Chemico-Biological Interactions*, 183: 98–104.
- Macneale K.H., Kiffney P.M., Scholz N.L. (2010). Pesticides, aquatic food webs, and the conservation of Pacific salmonids. *Frontiers in Ecology and the Environment*, 8: 475-482.
- Monteiro D.A., de Almeida J.A., Rantin F.T., Kalinin A.L. (2006). Oxidative stress biomarkers in the freshwater characid fish, *Brycon cephalus*, exposed to organophosphorus insecticide Folisuper 600 (methyl parathion). *Comparative Biochemistry and Physiology, Part C*, 143: 141–149.
- Moore A., Waring C.P. (2001). The effects of a synthetic pyrethroid pesticide on some aspects of reproduction in Atlantic salmon (*Salmo salar* L.). *Aquatic Toxicology*, 52: 1-12.
- Moss D.V., Henderson A.R. (1999). Clinical enzymology In: Burtis CA, Ashwood ER, editors. *Tietz Textbook of Clinical Chemistry*. 3rd ed. Philadelphia: W.B. Saunders Company; p. 617-721.

- Muranli F.D.G., Güner U. (2011). Induction of micronuclei and nuclear abnormalities in erythrocytes of mosquito fish (*Gambusia affinis*) following exposure to the pyrethroid insecticide lambda-cyhalothrin. *Mutation Research*, 726: 104–108.
- Oruç E.Ö., Usta D. (2007). Evaluation of oxidative stress responses and neurotoxicity potential of diazinon in different tissues of *Cyprinus carpio*. *Environmental Toxicology and Pharmacology*, 23: 48–55.
- Parvez S., Raisuddin S. (2005). Protein carbonyls: novel biomarkers of exposure to oxidative stress-inducing pesticides in freshwater fish *Channa punctata* (Bloch). *Environmental Toxicology and Pharmacology*, 20: 112–117.
- Patil V.K., David M. (2008). Behaviour and respiratory dysfunction as an index of malathion toxicity in the freshwater fish, *Labeo rohita* (Hamilton). *Turkish Journal of Fisheries and Aquatic Sciences*, 8: 233-237.
- Paulino M.G., Sakuragui M.M., Fernandes M.N. (2012). Effects of atrazine on the gill cells and ionic balance in a neotropical fish, *Prochilodus lineatus*. *Chemosphere*, 86: 1–7.
- Placer Z.A., Cushman L., Johnson B.C. (1966). Estimation of products of lipid peroxidation (malonyl dialdehyde) in biological fluids. *Analytical Biochemistry*, 16: 359-364.
- Polat, H., Erkoc, F.U., Viran, R., and Kocak, O. (2002). Investigation of acute toxicity of beta-cypermethrin on guppies, *Poecilia reticulata*, *Chemosphere*, 49: 39–44.
- Prusty A.K., Kohli M.P.S., Sahu N.P., Pal A.K., Saharan N., Mohapatra S., Gupta SK. (2011). Effect of short term exposure of fenvalerate on biochemical and haematological responses in *Labeo rohita* (Hamilton) fingerlings. *Pesticide Biochemistry and Physiology*, 100: 124–129.
- Rao J.V. (2006). Biochemical alterations in euryhaline fish, *Oreochromis mossambicus* exposed to sub-lethal concentrations of an organophosphorus insecticide, monocrotophos. *Chemosphere*, 65: 1814–1820.
- Rosalki S.B. (1998). Low serum creatine kinase activity. *Clinical Chemistry*, 44(5): 905.
- Santos T.G., Martinez C.B.R. (2012). Atrazine promotes biochemical changes and DNA damage in a Neotropical fish species. *Chemosphere*, 89: 1118–1125.
- Saravanan M., Prabhu Kumar K., Ramesh M. (2011). Haematological and biochemical responses of freshwater teleost fish *Cyprinus carpio* (Actinopterygii: Cypriniformes) during acute and chronic sublethal exposure to lindane. *Pesticide Biochemistry and Physiology*, 100: 206-211.
- Saxena K.K., Seth N. (2002). Toxic effects of cypermethrin on certain hematological aspects of fresh water fish, *Channa punctatus*. *Bulletin of Environmental Contamination and Toxicology*, 69: 364–369.
- Sayeed I., Parvez S., Pandey S., Bin-Hafeez B., Haque R., Raisuddin S. (2003). Oxidative stress biomarkers of exposure to deltamethrin in freshwater fish, *Channa punctatus* (Bloch). *Ecotoxicology and environmental safety*, 56: 295-301.
- Sepici-Dinçel A., Çağlan Karasu Benli A., Selvi M., Sarıkaya R., Şahin D., Özkul I.A., Erkoç F. (2009). Sublethal cyfluthrin toxicity to carp (*Cyprinus carpio* L.) fingerlings: biochemical, hematological, histopathological alterations. *Ecotoxicology and Environmental Safety*, 72: 1433–1439.
- Sharbidre A.A., Metkari V., Patode P. (2011). Effect of methyl parathion and chlorpyrifos on certain biomarkers in various tissues of guppy fish, *Poecilia reticulata*. *Pesticide Biochemistry and Physiology*, 101: 132-141.
- Soderlund D.M., Clark J.M., Sheets L.P., Mullin L.S., Piccirillo V.J., Sargent D., Stevens J.T., Weiner M.L. (2002). Mechanisms of pyrethroid neurotoxicity: implications for cumulative risk assessment. *Toxicology*, 171: 3–59.
- Sogorb M.A., Vilanova E. (2002). Enzymes involved in the detoxification of organophosphorus, carbamate and pyrethroid insecticides through hydrolysis. *Toxicology Letters*, 128: 215–228.
- Tayebati S.K., Di Tullio M.A., Ricci A., Amenta F. (2009). Influence of dermal exposure to the

pyrethroid insecticide deltamethrin on rat brain microanatomy and cholinergic/dopaminergic neurochemistry. *Brain Research*, 1301: 180-188.

Thomas L. (1998). *Clinical laboratory diagnostics*. 1st ed Frankfurt: TH-Books Verlagsgesellschaft; p. 65-71.

Tripathi, G., Shasmal, J., (2010). Reparation of chlorpyrifos-induced impairment by thyroxine and vitamin C in fish. *Ecotoxicology and Environmental Safety*, 73: 1397-1401.

Ural M.Ş., Sağlam N. (2005). A study on the acute toxicity of pyrethroid deltamethrin on the fry rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792). *Pesticide Biochemistry and Physiology*, 83: 124-131.

Velisek J., Svobodova Z., Machova J. (2009). Effects of bifenthrin on some haematological, biochemical and histopathological parameters of common carp (*Cyprinus carpio*). *Fish Physiology and Biochemistry*, 35(4): 583-590.

Velmurugan B., Selvanayagam M., Cengiz E.I., Unlu E. (2007). Histopathology of lambda-cyhalothrin on tissues (gill, kidney, liver and intestine) of *Cirrhinus mrigala*. *Environmental Toxicology and Pharmacology*, 24: 286-291.

Werner I., Moran K. (2008). Effects of pyrethroid insecticides on aquatic organisms. In: Gan, J., Spurlock, F., Hendley, P., Weston, D. (Eds.), *Synthetic Pyrethroids: Occurrence and Behavior in Aquatic Environments*. ACS Symposium Series, vol. 991. American Chemical Society, Washington, D.C., pp. 310-334.

Weston D.P., Lydy M.J. (2010). Urban and agricultural sources of pyrethroid insecticides to the Sacramento-San Joaquin Delta of California. *Environmental Science and Technology*, 44: 1833-1840.

Weston D.P., Holmes R.W., Lydy M.J. (2009). Residential runoff as a source of pyrethroid pesticides to urban creeks. *Environmental Pollution*, 157: 287-294.

Woudneh M.B., Oros D.R. (2006). Pyrethroids, pyrethrins, and piperonyl butoxide in sediments by high-resolution gas chromatography/high-resolution

mass spectrometry. *Journal of Chromatography A*, 1135: 71-77.

Yonar M.E., Sakin F. (2011). Ameliorative effect of lycopene on antioxidant status in *Cyprinus carpio* during pyrethroid deltamethrin exposure. *Pesticide Biochemistry and Physiology*, 99: 226-231.

You J., Weston D.P., Lydy M.J. (2004). A Sonication Extraction Method for the analysis of pyrethroid, organochlorine pesticides from sediment by gas chromatography with electron-capture detection. *Archives of Environmental Contamination and Toxicology*, 47: 141-147.