

Demersal and pelagic species of fish and squid from the Patagonian shelf

Elena Beatríz Eder¹, María Rosa Marin¹, Mirtha Noemí Lewis¹

¹ *Centro para el Estudio de Sistemas Marinos CESIMAR-CENPAT-CONICET, Boulevard Almirante Brown 2915, U9120ACD, Puerto Madryn, Chubut, Argentina*

Corresponding author: *Elena B. Eder* (eder@cenpat-conicet.gob.ar)

Academic editor: *D. Bloom* | Received 16 January 2017 | Accepted 28 March 2017 | Published 13 April 2017

<http://zoobank.org/AE401D3C-E5D9-4055-A487-DB09C19B1905>

Citation: Eder EB, Marin MR, Lewis MN (2017) Demersal and pelagic species of fish and squid from the Patagonian shelf. *ZooKeys* 668: 139–145. <https://doi.org/10.3897/zookeys.668.11826>

Resource citation: Eder EB, Marin MR, Lewis MN (2016) Demersal and pelagic species of fish and squid from the Patagonian shelf. v1.7. ArOBIS Centro Nacional Patagónico. Dataset/Samplingevent. <http://arobis.cenpat-conicet.gob.ar:8081/resource?r=argentina-fishes&v=1.7>

Abstract

The dataset contains 2007 records of occurrence of 39 species of fish and 2 species of squid distributed on the Patagonian continental shelf and slope. This dataset describes a new and revised version of the original data published through OBIS with individual morphometrics. Specimens are representative of pelagic, demersal, demersal-pelagic, demersal-benthic and benthic habits and they were collected by commercial fishing vessels in autumn (May–June, 2001, 51 catches), winter (July–August, 2001, 38 catches) and summer (January–February, 2002, 112 catches). The sampling was carried out with bottom trawls at a depth range of 73–370 m. The survey was located between 39°–52°S and 55°–65°W.

Keywords

Occurrence, Patagonia, teleosts, elasmobranchs, cephalopods, Southwestern Atlantic Ocean, demersal habitat, pelagic habitat

Data published through

GBIF: <http://arobis.cenpat-conicet.gob.ar:8081/resource?r=argentina-fishes>

Introduction

The Patagonian continental shelf, in the Atlantic margin of South America, is characterized by a remarkable productivity accounted by its oceanographic and bathymetric features, promoting ‘hotspots’ of biodiversity and attracting marine top predators, such as sea birds and marine mammals. The objective of this dataset is to provide basic information (location and depth capture, and individual morphometrics) of potential prey species of marine top predators from the Patagonian shelf, representative of pelagic, demersal, demersal-pelagic, demersal-benthic and benthic habits as useful information for ecological and biogeographical studies.

Taxonomic coverage

General taxonomic coverage description: All specimens were identified to species level. The dataset included 39 marine species of fish, representative of half the best-known fish species from the Argentine continental shelf (55.7 %, Cosseau and Perrota 2013), and 2 species of squid, that were captured during three seasons, autumn (May–June, 2001), winter (July–August, 2001) and summer (January–February, 2002). Autumn and summer records presented a high diversity of species (80.5 and 78 %, respectively), while only 46% of the species were present in winter (Figure 1).

Taxonomic ranks

Kingdom: Animalia

Phylum: Chordata

Class: Chondrichthyes

Order: Chimaeriformes, Carcharhiniformes, Squaliformes, Torpediniformes, Rajiformes.

Family: Callorhynchidae, Scyliorhinidae, Squalidae, Torpedinidae, Rajidae

Genera: Callorhynchus, Schroederichthys, Squalus, Discopyge, Bathyraja, Dipturus, Psammobatis.

Species: *Callorhynchus callorhynchus*, *Schroederichthys bivius*, *Squalus acanthias*, *Discopyge tshudii*, *Bathyraja brachyurops*, *Bathyraja albomaculata*, *Bathyraja scaphiops*, *Bathyraja macloviana*, *Dipturus chilensis*, *Psammobatis scobina*, *Psammobatis normani*.

Class: Actinopterygii




Figure 1. Distribution of occurrences of species during three seasons. Occurrence was calculated in each season as: (number of localizations where the species was present/ total number of localizations)*100.

Order: Anguilliformes, Gadiformes, Ophidiiformes, Pleuronectiformes, Perciformes, Scorpaeniformes.

Family: Congridae, Macrouridae, Moridae, Merlucciidae, Gadidae, Ophidiidae, Paralychnthyidae, Achiropsettidae, Scombridae, Centrolophidae, Stromateidae, Pinguipedidae, Carangidae, Serranidae, Bramidae, Cheilodactylidae, Bovichtidae, Nototheniidae, Scorpaenidae, Congiopodidae, Triglidae, Zoarcidae, Psychrolutidae.

Genera: Bassanago, Coelorrhynchus, Salilota, Austrophyctic, Macruronus, Merlucius, Micromesistius, Genypterus, Paralichthys, Mancopsetta, Scomber, Seriolella, Stromateus, Pseudopercis, Parona, Acanthistius, Brama, Nemadactylus, Cottoperca, Dissostichus, Patagonotothen, Sebastes, Congiopodus, Prionotus, Iluocoetes, Cottunculus.

Species: *Bassanago albescens*, *Coelorrhynchus fasciatus*, *Salilota australis*, *Austrophyctic marginata*, *Macruronus magellanicus*, *Merluccius hubbsi*, *Merluccius australis*, *Micromesistius australis*, *Genypterus blacodes*, *Paralichthys patagonicus*, *Mancopsetta maculata*, *Scomber japonicus*, *Seriolella punctata*, *Stromateus brasiliensis*, *Pseudopercis semifasciata*, *Parona signata*, *Acanthistius patachonicus*, *Brama brama*, *Nemadactylus bergi*, *Cottoperca gobio*, *Dissostichus eleginoides*, *Patagonotothen ramsayi*, *Pseudopercis*, *Sebastes oculata*, *Congiopus peruvianus*, *Prionotus nudigula*, *Iluocoetes fimbriatus*, *Cottunculus granulatus*.

Phylum: Mollusca

Class: Cephalopoda

Order: Oegopsida

Family: Ommastrephidae, Onychoteuthidae.

Genera: *Illex*, *Moroteuthis*.

Species: *Illex argentinus*, *Moroteuthis ingens*.

Spatial coverage

General spatial coverage: The Argentine continental shelf, in the Atlantic margin of South America (Figure 2), is bounded by the line of the coast and the continental slope and is the most extensive submerged plain in the world, with an area of 930.000 km²

(Cousseau and Perrotta 2013). It is characterized by gentle slopes and low-relief. The shelf waters are of sub-Antarctic origin diluted by continental flows and modified by exchanges of mass and heat with the atmosphere. The main distribution of the species of this dataset is:

- The internal and external sectors of the shelf off the Province of Buenos Aires (35° S) to Patagonia (48° S).
- The southern part of the Patagonian-Fuegian shelf and Malvinas/Falkland Islands, along the outer shelf to the North, up to approximately 42° S.
- Deeper waters near the continental slope.

Coordinates: 52°S and 39°S Latitude; 65°W and 55°W Longitude.

Temporal coverage: May 11, 2001–June 26, 2001, July 21, 2001–August 31, 2001, January 11, 2002–February 27, 2002.

Methods

Study extent description

The sampling area was located on the Patagonian continental shelf (Figure 2), within 52 to 39°S and 65 to 55°W. This is the largest submerged plain in the Southern Hemisphere (930.000 km²), characterized by relatively shallow depth (mostly 150 m deep). Productive zones are associated with major water mass transitions, currents nutrient flow connected to upwelling and bathymetric features (Cosseau and Perrotta 2013). The different water masses promote the recycling of nutrients accounting for its high productivity (Podesta et al. 1991, Carreto et al. 1995). The productivity of this area generates 'hotspots' of biodiversity and foraging of marine top predators is regularly observed (Croxall and Wood 2002, Campagna and Croxall 2003, Bastida and Rodríguez 2005). The specimens of this work were obtained through autumn (May-June, 2001), winter (July-August, 2001) and summer (January-February).

Sampling description

Specimens of fish and squid (commercial target and by catch) were taken daily by commercial fishing vessels operating with bottom trawls during autumn (May-June, 2001, 51 catches), winter (July-August, 2001, 38 catches) and summer (January-February, 2002, 112 catches). The fishing company provided the associated data of each fishing haul: date, hour, location (decimal Latitude and Longitude of the position while pulling the net) and depth of the catch (the maximum depth reached by the net). Specimens were frozen on board, and identified at species level at the Ichthyology laboratory of Centro Nacional Patagónico, Puerto Madryn, Argentina.




Figure 2. Spatial coverage (sampling locations).

The taxonomical identification of species was made by the specialists Dr A. Gosztonyi and Dr M. Re and the scientific names and their current accurate spelling were also reviewed using suitable literature (Brunetti et al. 1998; Cousseau and Perrotta 2013) and the WoRMS web site (<http://www.marinespecies.org/>).

Sex (when possible) and morphometric measures were taken for each specimen: wet mass (g) and wet mass of viscera (g, empty stomach); total and standard length (cm) for fish; mantle, head and fin length and width (cm) for squid; left and right fin length (cm) and maximum fin width (cm) for skates.

Method step description

Step1: Sampling locality and depth were recorded in each season.

Step2: Specimens were sent to the lab for species and sex identification and morphometric measurements.

Datasets

The data is published on a Creative Commons Attribution Non Commercial (CC-BY-NC) 4.0 license.

Dataset description

The star schema used to arrange the data has a DwC Event Core (seasons and catches) and two extensions: Occurrence (species) and ExtendedMeasurementOrFacts (individual morphometrics) (<http://bdj.pensoft.net/articles.php?id=10989>). The Darwin Core terms included in each file are:

Event core: type, eventID, parentEventID, samplingProtocol, eventDate, locationID, waterBody, locality, minimumDepthInMeters, decimalLatitude, decimalLongitude.

Occurrence (extension): modified, institutionCode, collectionCode, basisOfRecord, occurrenceID, catalogNumber, sex, occurrenceStatus, eventID, identifiedBy, scientificNameID, scientificName, kingdom, phylum, class, order, family, genus, specificEpithet, scientificNameAuthorship.

ExtendedMeasurementOrFacts (extension): occurrenceID, measurementType, measurementTypeID, measurementValue, measurementUnit, measurementUnitID, measurementDeterminedDate, measurementDeterminedBy.

Object name: Darwin Core Archive Demersal and pelagic species of fish and squid from the Patagonian shelf

Character encoding: UTF-8

Format name: Darwin Core Archive format

Format version: 1.0

Distribution: <http://arobis.cenpat-conicet.gob.ar:8081/archive.do?r=argentina-fishes>

Publication date of data: 2016-11-25

Language: English

Licenses of use: This work is licensed under a Creative Commons Attribution Non Commercial (CC-BY-NC) 4.0 License (<http://creativecommons.org/licenses/by-nc/4.0/legalcode>).

Metadata language: English

Date of metadata creation: 2016-09-07

Hierarchy level: Dataset

Acknowledgements

We are indebted to Harengus S.A. fishing company for the samples provided, and to Centro Nacional Patagónico (CENPAT-CONICET) for institutional support. We thank Dr. A. Gosztonyi and Dr. M. Re for their help with species identification, R. Mazzanti for technical assistance, Dr. M. Degradi and Dr. G. Garaffo for the language assistance and Dr G. Lovrich, Dr. D. Brosens and an anonymous reviewer for their valuable comments.

References

- Bastida R, Rodríguez D (2005) Marine Mammals of Patagonia and Antarctica. Vazquez Mazzini, Buenos Aires, 207 pp.
- Brunetti NE, Ivanovic ML, Beatriz E (1998) Calamares Ommastrephidae (Cephalopoda, Ommastrephidae). In: Boschi EE (Ed.) El Mar Argentino y sus recursos pesqueros, Tomo 2. Los moluscos de interés pesquero: cultivos y estrategias reproductivas de bivalvos y equinoideos. Contribución Instituto Nacional de Investigación y Desarrollo Pesquero No. 1044, Mar del Plata, Argentina, 37–68.
- Campagna C, Croxall J (2003) Application of the landscape species approach to the Patagonian Large Marine Ecosystem (The Sea and Sky Project). UICN-WPC, Durban.
- Carreto JI, Lutz VA, Carignan MO, Cucchi Colleoni AD, De Marco SG (1995) Hydrography and chlorophyll-a in the transect from the coast to the shelf break in the Argentinean Sea. *Continental Shelf Research* 15: 315–336. [https://doi.org/10.1016/0278-4343\(94\)E0001-3](https://doi.org/10.1016/0278-4343(94)E0001-3)
- Cosseau MB, Perrota G (2013) Peces marinos de Argentina. Biología, distribución, pesca. Instituto Nacional de Investigación y Desarrollo Pesquero, Mar del Plata, Argentina, 193 pp.
- Croxall JP, Wood AG (2002) The importance of the Patagonian Shelf to top predator species breeding at South Georgia. *Aquatic Conservation* 12: 101–118. <https://doi.org/10.1002/aqc.480>
- Podesta GP, Brown OB, Evans RH (1991) The annual cycle of satellite-derived sea surface temperature in the Southwestern Atlantic Ocean. *American Meteorological Society* 4: 457–467. [https://doi.org/10.1175/1520-0442\(1991\)004<0457:tacosd>2.0.co;2](https://doi.org/10.1175/1520-0442(1991)004<0457:tacosd>2.0.co;2)