

# Apoptosis induced by a snake venom metalloproteinase from *Bothrops alternatus* venom in C2C12 muscle cells

Soledad Bustillo<sup>1</sup> · Andrea C. Van de Velde<sup>1</sup> · Verónica Matzner Perfumo<sup>1</sup> · Claudia C. Gay<sup>1</sup> · Laura C. Leiva<sup>1</sup>

© Springer Science+Business Media New York 2017

**Abstract** In this study, the apoptosis inducing effects of baltergin as well as its influence on cell adhesion and migration on muscles cells in vitro were studied. Morphological analysis made by scanning electron and phase contrast microscopy demonstrated typical futures of programmed cell death, apoptosis. This mechanism was confirmed by fluorescence staining, molecular analysis of endonuclease activity and increased mRNA expression level of two representative genes (p53 and bax). On the other hand, baltergin exert an inhibition effect on myoblast cell adhesion and migration in vitro probably through a mechanism that involves the interaction of this enzyme with cell integrins. In conclusion, our results suggest that the absence of appropriate extracellular matrix contacts triggers anoikis. Therefore, this is the first report that demonstrated the mechanism of programmed cell death triggered by baltergin, a PIII metalloprotease isolated from *Bothrops alternatus* venom, in a myoblast cell line.

**Keywords** Baltergin · Metalloproteinase · *Bothrops alternatus* · Anoikis · Apoptosis

## Introduction

Snake venom metalloproteinases (SVMPs) play a relevant role in the pathogenesis of *Bothrops* intoxication. The functional diversity of SVMPs is in part due to the structural organization of different combinations of catalytic,

disintegrin, disintegrin-like and cysteine-rich domains, which categorizes SVMPs in 3 classes of precursor molecules (PI, PII and PIII) further divided in 11 subclasses. This heterogeneity is currently correlated to genetic accelerated evolution and post-translational modifications [1].

According to proteomic studies, SVMPs consists about 43% of *Bothrops alternatus* venom [2]. Baltergin is a member of P-III class of SVMPs isolated from this venom which is responsible to induce relevant local effects such as hemorrhage, edema and myotoxicity as well as systemic bleeding in mice [3]. Previous studies demonstrated cytotoxic and apoptotic effects induced by *B. alternatus* whole venom on muscle C2C12 cell line [4]. In particular, this metalloproteinase together with a non-toxic acidic PLA<sub>2</sub> isolated from the same venom, evidenced a synergistic effect on cell detachment of the mentioned muscle cell line and an endothelial cell line (tEnd) [5, 6]. This loss of adhesion from culture substrate, occurred without involving a cytolytic effect, as indicated by the lack of release of cytosolic LDH to supernatants in cells exposed to the toxins.

The use of skeletal muscle myoblasts/myotubes from C2C12 cell line as targets for venom toxins has been proposed as a useful in vitro model to study their myotoxic mechanism(s), as it correlates with the muscle-damaging activity observed in vivo [7]. Therefore, this cell culture is an appropriate cell model system to evaluate the myotoxicity caused by snake venoms or isolated toxins as proven by several studies [4, 8–13].

In anchorage-dependent cells, cell-to-cell and cell-to-matrix contacts are necessary for the maintenance of cell survival [14]. Linked with cell adherence status, anoikis has been described as a specific type of apoptosis caused by inappropriate cell-ECM interactions and all of the features that characterize apoptosis, including nuclear fragmentation, membrane blebbing and externalization of

✉ Soledad Bustillo  
solebustillo@yahoo.es

<sup>1</sup> Universidad Nacional del Nordeste (UNNE), Corrientes, Argentina

phosphatidyl serine (PS), are observed during this process [15].

It has been reported earlier that different snake venom metalloproteases (SVMPs) induce apoptosis. In particular, class PIII of these enzymes has demonstrated to induce this type of cell death principally on endothelial cells as a consequence of loss of adhesion [16–20]. However, the mechanism of cell death triggered by baltergin was not yet studied.

Thus, in order to achieve a better understanding on the role of metalloproteinase activity in envenomation, we analyzed the events involved in cell death induced by baltergin on muscle cells and its pharmacological potential as a structural model for drug design for cancer therapy.

## Materials and methods

### Venom and toxin

*B. alternatus* crude venom was supplied by the Serpentarium of Corrientes, Argentina and kept at  $-20^{\circ}\text{C}$  after vacuum dried. The metalloproteinase, baltergin was isolated from *B. alternatus* venom as described by Gay et al. [3]. In order to assay the isolated enzyme for purity, electrophoresis was performed on 12% polyacrylamide slab gels following the method of Laemmli [21].

### Cell culture

C2C12 (CRL-1772, ATCC) is a subclone murine skeletal muscle cell line derived from mouse myoblast cells obtained from normal adult C3H leg muscle. This cell line rapidly differentiates and produces extensive contracting myotubes that express characteristic muscle proteins [22]. Cytolysis was assessed in undifferentiated myoblasts. Cells were grown in  $25\text{ cm}^2$  flasks using Dulbecco's minimum essential medium (DMEM—Gibco Argentina), supplemented with 5% fetal bovine serum (FBS—Natocor Argentina), Penicillin–Streptomycin (Invitrogen) in a humidified atmosphere with 5%  $\text{CO}_2$  at  $37^{\circ}\text{C}$ .

### Phase contrast microscopy (PCM)

C2C12 cells were seeded in 6-well plates,  $1 \times 10^6$  cells per well in the same growth medium (DMEM-FBS 5%). When monolayers reached 80–90% confluence, culture medium was washed out and  $200\text{ }\mu\text{g/mL}$  of the purified enzyme was added (3 mL/well). After 4 h incubation, at  $37^{\circ}\text{C}$  and 5%  $\text{CO}_2$  in humid atmosphere, the cell morphological changes were investigated qualitatively using a phase contrast microscope (Axiovert 40<sup>®</sup>, Carl Zeiss Argentina). The photos were taken with a digital camera (Canon CCD

2272 $\times$ 1704, Argentina). In addition, the effect of baltergin inhibited catalytically with Na<sub>2</sub>EDTA was assayed. Briefly,  $100\text{ }\mu\text{L}$  of enzyme (4 mg/mL) was preincubated with the same volume of Na<sub>2</sub>EDTA (100 mM) for 1 h at  $37^{\circ}\text{C}$ . Excess reagent was eliminated by gel filtration on Sephadex G-25. The inhibited fraction, baltergin-Na<sub>2</sub>EDTA, was collected and protein concentration was determined by Bradford and adjust to  $200\text{ }\mu\text{g/mL}$  [23]. Moreover, the effectiveness of inhibition was checked before the exposure to cells by the determination of the residual proteolytic activity [24].

### Scanning electron microscopy (SEM)

Cells were grown on coverslips until 70–80% confluence and then incubated with PBS (control) or baltergin ( $200\text{ }\mu\text{g/mL}$ ) during 4 h at  $37^{\circ}\text{C}$ . Cells were then fixed at room temperature for 60 min in 2.5% glutaraldehyde (pH 7.4). Coverslips were air dried and dehydrated with increasing concentrations of acetone, dried by the  $\text{CO}_2$  critical-point method, sputtercoated with gold/palladium and examined with a JEOL 5800 LV Scanning Electron Microscope (SEM).

### Fluorescence staining

Myoblast cells were grown on cover slips and treated with baltergin metalloproteinase ( $200\text{ }\mu\text{g/mL}$ ) for 4 h at  $37^{\circ}\text{C}$ . PBS was used instead of baltergin for control assays. After incubation, cultured myoblasts were washed twice with PBS and gently mixed with a mixture of acridine orange (AO) ( $1\text{ }\mu\text{g/mL}$ ) and ethidium bromide (EB) ( $1\text{ }\mu\text{g/mL}$ ) dye solution for one minute as described by Spector et al. [25]. Coverslips were applied to the slides; afterwards, the sections were observed and photographed under a fluorescence microscope (Axioskop 40<sup>®</sup>/Axioskop 40 FL<sup>®</sup>, Carl Zeiss, Argentina).

### Adhesion assay

For adhesion assay, C2C12 cells ( $2.5\text{--}3 \times 10^4$ /well) were preincubated for 30 min at room temperature with Baltergin ( $10\text{--}300\text{ }\mu\text{g/mL}$ ) or culture medium (control group) and then added to 96-well plate previously coated with FBS:PBS (1:4) for 1 h at  $37^{\circ}\text{C}$  followed blocked with 1% heat-inactivated BSA (Sigma- $1\text{ }\mu\text{g}/\mu\text{L}$ ) in PBS. After 1.5 h, non-adherent cells were removed by careful aspiration and washed with PBS. Adherent cells were fixed with methanol:glacial acetic acid (3:1 ratio), and stained with 0.5% crystal violet in 20% (v/v) methanol. After dye release, adherent cells were microscopically observed and recorded photographically. The dye was released from the cells by addition of ethanol:glacial acetic acid (3:1 ratio).

The optical density of the released dye solution was read at 620 nm and the percentage of cell adhesion was determined by comparing the resulting absorbances with the mean absorbance of control groups (considered as 100% of adhesion) [26]. For the assessment of cytolysis, the release of the cytosolic enzyme lactate dehydrogenase (LDH) was used as an indicator of membrane damage [5]. Aliquots (40  $\mu$ L) of the supernatant in culture wells were collected, and LDH activity was determined by using a UV kinetic assay kit (LDH BR CE, Spain). Cytolysis was expressed as percentage, using as 0% and 100% reference values the LDH activity of supernatants from cells exposed to medium alone, or to 0.1% (v/v) Triton X-100, respectively. All assays were carried out in triplicate wells.

### Migration assay

Cell migration was measured by Wound-healing assay. Briefly, cells were grown to 90% confluence in a 6-well plate at 37 °C, 5% CO<sub>2</sub> incubator. A wound was created by scratching cells with a sterile pipette tip, cells were washed twice with PBS to remove floating cells and then incubated in culture medium in the absence or presence of baltergin (25 and 50  $\mu$ g/mL) for 24 h. Also, the effect of the same concentrations of enzyme but catalytically inhibited with Na<sub>2</sub>EDTA were evaluated. The inhibition process was performed as it was previously described in item 3. Cell migration into the wound surface was determined under an inverted microscope. Wound widths were measured with Image-Pro software and percentage of cell migration was calculated by comparing with control groups (considered as 100% of migration).

### DNA fragmentation assay

For the detection of apoptotic DNA cleavage,  $1 \times 10^6$  cells were grown on 6-well plates at 37 °C, 5% CO<sub>2</sub>. Cells were treated with baltergin (200  $\mu$ g/mL) for 4 h and then trypsinized and lysed with lysis buffer (Tris-HCl 50 mM, pH 8.0, EDTA 10 mM, SDS 0.5%, proteinase K 0.5 mg/mL) for 20 min. DNA extraction was made with phenol/chloroform/isoamyl alcohol solution (25:24:1), precipitated with 0.1 volumes of sodium acetate (3 M) and 2.5 volumes of ethanol (96%) and resuspended in buffer Tris-HCl 10 mM pH 8.0, EDTA 1.0 mM pH 8.0 with RNase (20 mg/mL Invitrogen). DNA was visualized by electrophoresis on a 1.5% agarose gel (Biodynamics) containing ethidium bromide (Promega).

### RNA isolation and RT-PCR

To determine the expressions of p53 and bax mRNAs,  $1 \times 10^6$  cells were grown on 6-well plates at 37 °C, 5% CO<sub>2</sub>

and treated with baltergin (200  $\mu$ g/mL) for 4 h. Total RNA was isolated from the cells using Trizol reagent (Invitrogen) according to manufacturer's instructions. Two micrograms of RNA, 0.5  $\mu$ L of Oligo dt (Biodynamics) and 9.5  $\mu$ L of dimethyl pyrocarbonate-treated water (DEPC-H<sub>2</sub>O) were added together, and the mixture was heated at 70 °C for 5 min. One microliter of Moloney Murine Leukemia Virus Reverse Transcriptase (M-MLV RT) (Promega), 5  $\mu$ L of M-MLV RT buffer (Promega), 1.125  $\mu$ L of dNTPs (10 mM) (Fermentas), 0.625  $\mu$ L de RNasa Out (Invitrogen) were then added to the mixture, and the final volume was adjusted to 25  $\mu$ L with the DEPC-H<sub>2</sub>O. Subsequently, the reaction mixture was incubated at 42 °C for 1 h. PCR amplification was performed in a 25  $\mu$ L reaction volume containing 3  $\mu$ L of the appropriate cDNA, 2  $\mu$ L of dNTPs (Fermentas), 1  $\mu$ L of the 10  $\mu$ M primer set (Fermentas), 0.12  $\mu$ L of GoTaq<sup>®</sup> DNA polymerase (5 unites/ $\mu$ L) (Promega), 5  $\mu$ L de buffer of reaction buffer (Green GoTaq<sup>®</sup> buffer) (Promega), 1  $\mu$ L of Cl<sub>2</sub>Mg (Promega) and 12.88  $\mu$ L of H<sub>2</sub>O. The sequences of mouse p53 primers were 5'-GATGACTGCCATGGAGGAGT-3' (a 20-mer sense oligonucleotide) and 5'-CTCGGGTGGCTCATAAGGTA-3' (a 20-mer anti-sense oligonucleotide). The sequences of mouse bax primers were 5'-AGATGAACTGGATAGCAATATGGA-3' (a 24-mer sense oligonucleotide) and 5'-CCA CCCTGGTCTTGGATCCAGACA-3' (a 24-mer anti-sense oligonucleotide). The sequences of the internal control cyclophilin primers were 5'-ACCCACCGTGTCTTCCGAC-3' (a 20-mer sense oligonucleotide) and 5'-CATTTGCCATGGACAAGATG-3' (a 20-mer antisense oligonucleotide). The expected sizes of the PCR products were 664 bp for p53, 270 bp for bax, and 291 bp for cyclophilin. The PCR procedure was carried out using a thermocycler (Thermo) under the following conditions: initial denaturation at 94 °C for 5 min, followed by 25 amplification cycles, each consisting of denaturation at 94 °C for 30 s, annealing at 58 °C for 30 s, and extension at 72 °C for 30 s, followed by additional extension step at the end of the procedure at 72 °C for 5 min. Amplification products were separated on 1.5% agarose gel (Biodynamics) and visualized by ethidium bromide staining (Promega). Finally a photographic record of the amplified products was obtained and bands were analyzed by densitometry with computer software Uvi Pro gel documentation system (UV Tech) to calculate relative gene expression.

### Statistical analysis

Data represent the mean  $\pm$  standard deviation (SD) of at least four replications. Statistical significance was tested by one-way ANOVA and Tukey (HSD) and p-values inferior to 0.05 were considered significant.

## Results

### Phase contrast microscopy (PCM)

Cell morphology analysis by PCM showed that untreated C2C12 cells were homogeneously distributed on cultured field; they exhibited normal, elongated and tapered morphology (Fig. 1a). After 4 h incubation with 200  $\mu\text{g}/\text{mL}$  of baltergin cell rounding, chromatin condensation and some areas devoid of cells were observed (Fig. 1b). The inhibition of catalytical activity by  $\text{Na}_2\text{EDTA}$  decreased the number of cells with morphological features of apoptosis but not abolished completely the effect (Fig. 1c).

### Scanning electron microscopy (SEM)

To assess the baltergin-induced change in cell morphology, cells were examined by scanning electron microscopy. As shown in Fig. 2b, c, d, numerous membranes blebbings, decrease in cells density, shrinkage of cell volume, apoptotic bodies and dispersed aggregates of the supernatant were induced with 200  $\mu\text{g}/\text{mL}$  of this toxin.

### Fluorescence staining

In order to determine whether baltergin induced morphological alterations that could be attributed to an apoptotic mechanism, myoblast cells were incubated with 200  $\mu\text{g}/\text{mL}$  for 4 h and then stained with the nucleic acid-binding fluorochromes, acridine orange and ethidium bromide. Control untreated cells exhibited a green fluorescence, due to exclusion of ethidium bromide but not of acridine orange. Viable cells showed a light green nucleus with intact structure and presented punctuate orange red fluorescence in


the cytoplasm, representing lysosomes stained by acridine orange (Fig. 3a). After 4 h of incubation with Baltergin, typical features of apoptosis were observed. Apoptotic cells exhibited a bright green nucleus (showing condensation of chromatin), dense green areas and evident membrane blebbing (Fig. 3b).

### Adhesion assay


Baltergin (10–300  $\mu\text{g}/\text{mL}$ ) inhibited cellular adhesion in a concentration-dependent manner (Fig. 4), being significantly different from the control at all concentrations tested. The highest amount of enzyme (300  $\mu\text{g}/\text{mL}$ ) preincubated with myoblast cells, inhibited cell adhesion by 32% with respect to control assays. This effect was not related to cytolysis, since no significant LDH release to supernatants was detected even at the highest concentration tested. Thus, the enzyme induced endothelial cell detachment without cytolysis, i.e. without plasma membrane disruption.

### Migration assay

Wound healing assay was performed to evaluate the inhibitory effect of baltergin on the migratory features of C2C12 cells. The wound scratch in control cells was almost completely closed after 24 h of incubation (Fig. 5b). However, treatment with 25 and 50  $\mu\text{g}/\text{mL}$  baltergin resulted in the suppression of wound healing in a concentration-dependent manner, decreasing respect to controls by 15 and 48% respectively (Fig. 5c, e, g). This inhibitory effect on migration was found to be inferior with the catalytically inhibited enzyme, compared with the observed with baltergin alone, for both concentrations assayed (Fig. 5d, f, g).



**Fig. 1** Morphological changes in myoblast cell cultures under phase contrast microscopy **a** Control, **b** Baltergin, 200  $\mu\text{g}/\text{mL}$  for 4 h, **c** Baltergin- $\text{Na}_2\text{EDTA}$ , 200  $\mu\text{g}/\text{mL}$  for 4 h




**Fig. 2** C2C12 cells observed at SEM after baltergin treatment (4 h). **a** Control myoblasts **b** Baltergin 200 µg/mL ( $\times 270$ ) **c** Baltergin 200 µg/mL ( $\times 500$ ) **d** Baltergin 200 µg/mL ( $\times 4500$ ). Apoptotic bodies (*arrows*)

### DNA fragmentation assay


To ascertain that baltergin induces apoptosis, DNA

fragmentation that reflects the endonuclease activity was analyzed. As shown in Fig. 6, the treatment for 4 h with 100 µg/mL of enzyme showed fragmentation of DNA but



**Fig. 3** Acridine orange and ethidium bromide fluorescence staining. **a** Control **b** Representative photomicrograph showing cell shrinkage, chromatin condensation and the formation of blebs on the cell surface

associated with apoptosis. Myoblasts were grown on coverslides and treated for 4 h minutes with 200 µg/mL of baltergin



**Fig. 4** Effect of baltergin on C2C12 cells adhesion. Myoblasts were added to culture wells precoated with FBS:PBS (1:4) in the presence or absence of baltergin (12.5–300 µg/mL). After 1.5 h incubation, the non-adherent cells were washed away and the adherent cells were stained with Cristal Violet. Cell adhesion is expressed as % of the untreated control. Each bar shows the mean  $\pm$ SD of three independent experiments that were performed in triplicate. \* $p < 0.05$  compared with the control


incubation with 200 µg/mL resulted in the formation of fragments that could be seen as the characteristic ladder pattern by electrophoresis.

#### Effect of baltergin on the expression of p53 and bax mRNA

In the present study, we investigated the mRNA expression level of two representative genes (p53 and bax) after treatment with baltergin using the RT-PCR technique. The copy number of each gene was normalized to the reference gene (cyclophilin) to give relative expression levels, and the change in gene expression was compared with the untreated control samples (Fig. 7a). Our results demonstrated that the mRNA expression level of both genes, increased significantly following treatment with the enzyme as compared to control cells. In particular, levels of bax and p53 mRNA following the treatment with 200 µg/mL for 4 h were increased to 1.19 and 1.51 respectively (Fig. 7b, c).

#### Discussion

Snake venom metalloproteinases (SVMPs) degrade various components of the basement membrane and are also able to hydrolyze cell membrane proteins, such as integrins and cadherins, involved in cell–matrix and cell–cell adhesion [27]. Molecular approaches have been previously



**Fig. 5** Effect of baltergin on migration in C2C12 myoblast cells. **a–d** Photographs of wounds in monolayers of cells: control 0 h (**a**) and 24 h (**b**), baltergin 25  $\mu\text{g}/\text{mL}$  (**c**) and 50  $\mu\text{g}/\text{mL}$  (**e**), baltergin- $\text{Na}_2\text{EDTA}$  25  $\mu\text{g}/\text{mL}$  (**d**) and 50  $\mu\text{g}/\text{mL}$  (**f**). **g** Quantitative analysis


of cell migration. Each bar shows the mean  $\pm$ SD of three independent experiments that were performed in triplicate. \* $p < 0.05$  compared with control 0 h

performed with high molecular weight metalloproteinases from a number of Viperidae species in order to elucidate the complex integrin-disintegrin interactions. SVMPs containing disintegrin-like domains (PIII/PIIIb class) may play a role in targeting the protein to a particular site in cells such as platelets, and endothelial cells, as well as in integrins, extracellular matrix and other substrates.

Recently, a two-step model was explained by Gutierrez et al. [28], for the mechanism of action of hemorrhagic SVMPs. Briefly, in a first step SVMPs bind to and hydrolyze critical structural components of the basement membrane of capillary vessels that results in the mechanical weakening of this scaffold structure. As a consequence, in a second step, the biophysical hemodynamic forces normally operating in the microcirculation induce a distention of the


vessel wall until the capillary is eventually disrupted, with the consequent extravasation of blood [28].

Baltergin is a member of P-III class of SVMPs isolated from *B. alternatus* venom which is responsible to induce relevant local effects. Previous works from our group demonstrated in vivo myotoxic effects of this metalloproteinase [3]. After intramuscular injection in murine gastrocnemium muscle, the metalloproteinase induced a late increase in serum creatine kinase, characteristic of muscle damage induced by SVMPs [29]. Also, we evaluated the capacity of this protein to induce cellular death and interestingly, this metalloproteinase showed a rapid dose-dependent detachment of myoblast (C2C12) and endothelial (t-End) cells but without cytolysis [5, 6]. In the present study, C2C12 cell line was also used. These cells are an appropriate in vitro



**Fig. 6** Electrophoretic examination of genomic DNA of C2C12 cells. Genomic DNA was extracted and analyzed by electrophoresis on the 1.5% agarose gel containing ethidium bromide. Line 1: Control, Line 2: baltergin 100 µg/mL, Line 3: baltergin 200 µg/mL

model system and a suitable target for the cytotoxic action of crude venoms or isolated enzymes, as proven by previous studies [8–11, 13, 30]. Baltergin induces apoptotic morphological changes in these myoblast cells. Under the phase-contrast microscope, baltergin-treated cells showed cell shrinkage, cytoplasm condensation, and irregularity in shape. Besides, dual acridine orange/ethidium bromide (AO/EB) fluorescent staining was used to identify apoptosis-associated changes of cell membranes. Fluorescence staining evidenced that cell membrane remains intact after toxin treatment exhibiting green but not red/orange fluorescence, indicative of acridine orange retention and ethidium



**Fig. 7** Effect of baltergin (200 µg/mL) on mRNA expressions of p53 and bax in C2C12 cells. The levels of p53 and bax mRNA were detected using the reverse transcription-polymerase chain reaction (RT-PCR) analysis. Cyclophilin mRNA was used as the internal control. **a** Cyclophilin control, **b** Bax mRNA expression, **c** p53 mRNA expression

bromide exclusion. Morphological changes associated with apoptosis were observed, principally, membrane blebbing and apoptotic bodies. The scanning electron microscopy also confirms these changes with the detection of cell shrinkage and once more membrane blebbing and the presence of apoptotic bodies.

On the other hand, cell adhesion is a critical event in many biological phenomena such as development, differentiation, signals transduction, maintenance of tissue



structure, wound healing, and tumor metastasis. Moreover, a vital step in the invasive cell processes is adhesion, cell migration and angiogenesis [60]. Therefore, two other approaches to this study were the evaluation of the in vitro toxin effect on cell adhesion and migration processes. The inhibition of adhesion promoted by baltergin was similar to those reported in other studies where metalloproteinases containing disintegrin-like domains, or disintegrins, inhibited cellular adhesion [31–34]. Our results demonstrated that baltergin induced a disruption of focal adhesion signaling, followed by loss of cell anchorage without involving a cytolytic effect, as indicated by the lack of release of cytosolic LDH to supernatants in cells exposed to the enzyme.

Besides, this toxin was also able to inhibit cell migration in a dose dependent manner. In this assay, the inhibition of baltergin with Na<sub>2</sub>EDTA showed the partial role of the catalytic activity in this effect. Thus, probably, other non-catalytic mechanisms could be involved.

Integrins are cell surface receptors that play critical roles in both processes (adhesion and migration). For this reason, it is possible that these effects were due in part to the specific binding or interaction of this toxin to  $\alpha\beta$  integrins and their subtypes. Previously, it has been shown that classical RGD-disintegrins induce apoptosis by interfering with  $\alpha\beta$  attachment to ECM proteins [16, 35, 36]. The absence of appropriate ECM contacts triggers apoptosis, and this programmed mechanism of cell death in response to lack of adhesion or inappropriate adhesion has been termed *anoikis* [15].

It is known that this mechanism of programmed cell death involves the activation of endonucleases resulting in the cleavage of genomic DNA into well-defined fragments that appear as the characteristic ladder pattern upon agarose gel electrophoresis [37]. To provide evidence supporting the involvement of apoptosis in the baltergin-induced cytotoxicity, the DNA fragmentation assay was performed and the distinctive ladder pattern characteristic of apoptotic cell death was detected.

Consistent with the induction of apoptosis, this toxin increased the expression of proapoptotic proteins. Apoptosis can be triggered through either the intrinsic and/or the extrinsic pathways. The intrinsic pathway is mediated through a mitochondria-dependent mechanism that is regulated by members of the Bcl-2 family. The p53 protein is a fundamental mediator in cellular processes of growth arrest and apoptosis and directly activates the proapoptotic Bcl-2 proteins. Our results showed that baltergin up-regulated the expression of p53 and Bax, one of the key members of the Bcl-2 family which convert from harmless monomers into deadly oligomers that form pores in the mitochondrial outer membrane [38].

Snake venom components captivate medical interest as potential molecules for the treatment of tumors [1, 16,

39–42]. The design of anti-migratory compounds is a particularly promising approach because these compounds not only delay tissue invasion and the formation of metastases by cancer cells emerging from primary tumor sites, but also restore a certain level of sensitivity to apoptosis and/or autophagy in these slowly migrating cells [43]. In this context many toxins from snake venoms have been investigated as a possible treatment for cancer; among these compounds are PLA<sub>2</sub>s, SVMPs, disintegrins, C-type lectins, etc [30, 41, 44–49]. In particular, SVMPs are promising prototypes because of their capacity of degrading various proteins of the basement membrane, as well as cell components involved in cell-matrix and cell–cell adhesion [50], as it was demonstrated in this work for baltergin.

This is the first report that demonstrated the mechanism of programmed cell death triggered by a PIII metalloprotease isolated from *B. alternatus* venom. The present data show that baltergin causes apoptosis, specifically anoikis, of myoblast cells directly related to the detachment effect and probably due to the interaction of this toxin with integrins.

**Acknowledgements** Financial support to these studies was received from Secretaría General de Ciencia y Técnica, Universidad Nacional del Nordeste (UNNE) (CF01-2013 and CF02-2013). Verónica Matzner Perfumo and Andrea Van de Velde are recipients of fellowships from ‘Facultad de Ciencias Exactas y Naturales y Agrimensura. UNNE’ and ‘Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)’, respectively. The authors thank Lic. Laura Rey from ‘Centro de Producción de Sueros Antiofídicos’ (CEPSAN; Corrientes, Argentina) for providing *Bothrops alternatus* venoms.

#### Compliance with ethical standards

**Conflict of interest** The authors declare that there are no conflicts of interest regarding this manuscript.

#### References

1. Moura-da-Silva AM, Furlan MS, Caporino MC, Grego KF, Portes-Junior JA, Clissa PB, Valente RH, Magalhaes GS (2011) Diversity of metalloproteinases in *Bothrops neuwiedi* snake venom transcripts: evidences for recombination between different classes of SVMPs. *BMC Genet* 12:94
2. Ohler M, Georgieva D, Seifert J, von Bergen M, Arni RK, Genov N, Betzel C (2010) The venomomics of *Bothrops alternatus* is a pool of acidic proteins with predominant hemorrhagic and coagulopathic activities. *J Proteome Res* 9:2422–2437
3. Gay CC, Leiva LC, Marunak S, Teibler P, Acosta de Perez O (2005) Proteolytic, edematogenic and myotoxic activities of a hemorrhagic metalloproteinase isolated from *Bothrops alternatus* venom. *Toxicon* 46:546–554
4. Bustillo S, Lucero H, Leiva LC, Acosta O, Kier Joffé EB, Gorodner JO (2009) Cytotoxicity and morphological analysis of cell death induced by *Bothrops* venoms from the northeast of Argentina. *J Venom Anim Toxins Incl Trop Dis* 15:28–42
5. Bustillo S, Garcia-Denegri ME, Gay C, Van de Velde AC, Acosta O, Angulo Y, Lomonte B, Gutierrez JM, Leiva L (2015)

- Phospholipase a enhances the endothelial cell detachment effect of a snake venom metalloproteinase in the absence of catalysis. *Chem-Biol Interact* 240:30–36.
6. Bustillo S, Gay CC, García Denegri ME, Ponce-Soto LA, de Kier Joffe EB, Acosta O, Leiva LC (2012) Synergism between baltergin metalloproteinase and Ba SPII RP4 PLA<sub>2</sub> from *Bothrops alternatus* venom on skeletal muscle (C2C12) cells. *Toxicol* 59:338–343.
  7. Angulo Y, Lomonte B (2005) Differential susceptibility of C2C12 myoblasts and myotubes to group II phospholipase A2 myotoxins from crotalid snake venoms. *Cell Biochem Funct* 23:307–313
  8. Omran MAA, Fabb SA, Dickson G (2004) Biochemical and morphological analysis of cell death induced by Egyptian cobra (*Naja haje*) venom on cultured cells. *J Venom Anim Toxins incl Trop Dis* 10:219–241.
  9. Damico DC, Nascimento JM, Lomonte B, Ponce-Soto LA, Joazeiro PP, Novello JC, Marangoni S, Collares-Buzato CB (2007) Cytotoxicity of Lachesis muta muta snake (bushmaster) venom and its purified basic phospholipase A2 (LmTX-I) in cultured cells. *Toxicol* 49:678–692
  10. Lomonte B, Angulo Y, Rufini S, Cho W, Giglio JR, Ohno M, Daniele JJ, Geoghegan P, Gutierrez JM (1999) Comparative study of the cytolytic activity of myotoxic phospholipases A2 on mouse endothelial (tEnd) and skeletal muscle (C2C12) cells in vitro. *Toxicol* 37:145–158
  11. Bonfim VL, Ponce-Soto LA, Novello JC, Marangoni S (2006) Cytotoxic action in myoblasts and myotubes (C2C12) and enzymatic characterization of a new phospholipase A2 isoform (Bj-V) from *Bothrops jararacussu* venom. *Protein Pept Lett* 13:707–713
  12. Lomonte B, Angulo Y, Santamaria C (2003) Comparative study of synthetic peptides corresponding to region 115–129 in Lys49 myotoxic phospholipases A2 from snake venoms. *Toxicol* 42:307–312
  13. Villalobos JC, Mora R, Lomonte B, Gutierrez JM, Angulo Y (2007) Cytotoxicity induced in myotubes by a Lys49 phospholipase A2 homologue from the venom of the snake *Bothrops asper*: evidence of rapid plasma membrane damage and a dual role for extracellular calcium. *Toxicol In Vitro* 21:1382–1389
  14. Grossmann J (2002) Molecular mechanisms of “detachment-induced apoptosis–Anoikis”. *Apoptosis* 7:247–260
  15. Valentijn AJ, Zouq N, Gilmore AP (2004) Anoikis. *Biochem Soc Trans* 32:421–425
  16. Tanjoni I, Weinlich R, Della-Casa MS, Clissa PB, Saldanha-Gama RF, de Freitas MS, Barja-Fidalgo C, Amarante-Mendes GP, Moura-da-Silva AM (2005) Jararhagin, a snake venom metalloproteinase, induces a specialized form of apoptosis (anoikis) selective to endothelial cells. *Apoptosis* 10:851–861
  17. Masuda S, Hayashi H, Atoda H, Morita T, Araki S (2001) Purification, cDNA cloning and characterization of the vascular apoptosis-inducing protein, HV1, from *Trimeresurus flavoviridis*. *Eur J Biochem* 268:3339–3345
  18. Trummel K, Tonismagi K, Siigur E, Aaspollu A, Lopp A, Sillat T, Saat R, Kasak L, Tammiste I, Kogerman P, Kalkkinen N, Siigur J (2005) A novel metalloprotease from *Vipera lebetina* venom induces human endothelial cell apoptosis. *Toxicol* 46:46–61
  19. You WK, Seo HJ, Chung KH, Kim DS (2003) A novel metalloprotease from *Gloydius halys* venom induces endothelial cell apoptosis through its protease and disintegrin-like domains. *J Biochem* 134:739–749
  20. Baldo C, Tanjoni I, Leon IR, Batista IF, Della-Casa MS, Clissa PB, Weinlich R, Lopes-Ferreira M, Lebrun I, Amarante-Mendes GP, Rodrigues VM, Perales J, Valente RH, Moura-da-Silva AM (2008) BnP1, a novel P-I metalloproteinase from *Bothrops neuwiedi* venom: biological effects benchmarking relatively to jararhagin, a P-III SVMP. *Toxicol* 51:54–65
  21. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680–685
  22. Yaffe D, Saxel O (1977) Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. *Nature* 270:725–727
  23. Hammond JB, Kruger NJ (1988) The Bradford method for protein quantitation. *Methods Mol Biol* 3:25–32
  24. Wang WJ, Huang TF (2002) Purification and characterization of a novel metalloproteinase, acurhagin, from *Agkistrodon acutus* venom. *Thromb Haemost* 87:641–650
  25. Specter DL, Robert DG, Lesli AL (1998) *Cells, a laboratory manual, subcellular localization of genes and their product*. New York, USA
  26. Paludo KS, Gremski LH, Veiga SS, Chaim OM, Gremski W, de Freitas Buchi D, Nader HB, Dietrich CP, Franco CR (2006) The effect of brown spider venom on endothelial cell morphology and adhesive structures. *Toxicol* 47:844–853
  27. Gutiérrez JM, Rucavado A, Escalante T, Díaz C (2005) Hemorrhage induced by snake venom metalloproteinases: biochemical and biophysical mechanism involved in microvessel damage. *Toxicol* 45:997–1011
  28. Gutierrez JM, Escalante T, Rucavado A, Herrera C (2016) Hemorrhage caused by snake venom metalloproteinases: a journey of discovery and understanding. *Toxins* 8:93.
  29. Andriao-Escarso SH, Soares AM, Rodrigues VM, Angulo Y, Diaz C, Lomonte B, Gutierrez JM, Giglio JR (2000) Myotoxic phospholipases A(2) in bothrops snake venoms: effect of chemical modifications on the enzymatic and pharmacological properties of bothropstoxins from *Bothrops jararacussu*. *Biochimie* 82:755–763
  30. Araya C, Lomonte B (2007) Antitumor effects of cationic synthetic peptides derived from Lys49 phospholipase A2 homologues of snake venoms. *Cell Biol Int* 31:263–268
  31. Costa EP, Santos MF (2004) Jararhagin, a snake venom metalloproteinase-disintegrin, stimulates epithelial cell migration in an in vitro restitution model. *Toxicol* 44:861–870
  32. Cominetti MR, Ribeiro JU, Fox JW, Selistre-de-Araujo HS (2003) BaG, a new dimeric metalloproteinase/disintegrin from the *Bothrops alternatus* snake venom that interacts with  $\alpha 5b1$  integrin. *Arch Biochem Biophys* 416:171–179
  33. Maria DA, da Silva MG, Correia Junior MC, Ruiz IR (2014) Antiproliferative effect of the jararhagin toxin on B16F10 murine melanoma. *BMC Complement Altern Med* 14:446
  34. Gomes MS, Naves de Souza DL, Guimaraes DO, Lopes DS, Mamede CC, Gimenes SN, Ache DC, Rodrigues RS, Yoneyama KA, Borges MH, de Oliveira F, Rodrigues VM (2015) Biochemical and functional characterization of Bothropoidin: the first haemorrhagic metalloproteinase from *Bothrops pauloensis* snake venom. *J Biochem* 157:137–149
  35. Wu WB, Peng HC, Huang TF (2003) Disintegrin causes proteolysis of beta-catenin and apoptosis of endothelial cells. Involvement of cell–cell and cell–ECM interactions in regulating cell viability. *Exp Cell Res* 286:115–127
  36. Yeh CH, Peng HC, Huang TF (1998) Accutin, a new disintegrin, inhibits angiogenesis in vitro and in vivo by acting as integrin  $\alpha v\beta 3$  antagonist and inducing apoptosis. *Blood* 92:3268–3276
  37. Lee MH, Jang MH, Kim EK, Han SW, Cho SY, Kim CJ (2005) Nitric oxide induces apoptosis in mouse C2C12 myoblast cells. *J Pharmacol Sci* 97:369–376
  38. Westphal D, Kluck RM, Dewson G (2014) Building blocks of the apoptotic pore: how Bax and Bak are activated and oligomerize during apoptosis. *Cell Death Differ* 21:196–205

39. Kamiguti AS, Hay CR, Theakston RD, Zuzel M (1996) Insights into the mechanism of haemorrhage caused by snake venom metalloproteinases. *Toxicon* 34:627–642
40. Calderon LA, Sobrinho JC, Zaqueo KD, de Moura AA, Grabner AN, Mazzi MV, Marcussi S, Nomizo A, Fernandes CF, Zuliani JP, Carvalho BM, da Silva SL, Stabeli RG, Soares AM (2014) Antitumoral activity of snake venom proteins: new trends in cancer therapy. *BioMed Res Int* 2014:203639.
41. Yang RS, Tang CH, Chuang WJ, Huang TH, Peng HC, Huang TF, Fu WM (2005) Inhibition of tumor formation by snake venom disintegrin. *Toxicon* 45:661–669
42. Correa MC Jr, Maria DA, Moura-da-Silva AM, Pizzocaro KF, Ruiz IR (2002) Inhibition of melanoma cells tumorigenicity by the snake venom toxin jararhagin. *Toxicon* 40:739–748
43. Lefranc F, Brotchi J, Kiss R (2005) Possible future issues in the treatment of glioblastomas: special emphasis on cell migration and the resistance of migrating glioblastoma cells to apoptosis. *J Clin Oncol* 23:2411–2422
44. Bazaa A, Luis J, Srairi-Abid N, Kallech-Ziri O, Kessentini-Zouari R, Defilles C, Lissitzky JC, El Ayeb M, Marrakchi N (2009) MVL-PLA2, a phospholipase A2 from *Macrovipera lebetina transmediterranea* venom, inhibits tumor cells adhesion and migration. *Matrix Biol* 28:188–193
45. Costa TR, Menaldo DL, Oliveira CZ, Santos-Filho NA, Teixeira SS, Nomizo A, Fuly AL, Monteiro MC, de Souza BM, Palma MS, Stabeli RG, Sampaio SV, Soares AM (2008) Myotoxic phospholipases A(2) isolated from *Bothrops brazili* snake venom and synthetic peptides derived from their C-terminal region: cytotoxic effect on microorganism and tumor cells. *Peptides* 29:1645–1656
46. da Silva RJ, da Silva MG, Vilela LC, Fecchio D (2002) Antitumor effect of *Bothrops jararaca* venom. *Mediators Inflamm* 11:99–104
47. de Carvalho DD, Schmitmeier S, Novello JC, Markland FS (2001) Effect of BJcuL (a lectin from the venom of the snake *Bothrops jararacussu*) on adhesion and growth of tumor and endothelial cells. *Toxicon* 39:1471–1476
48. Gebrim LC, Marcussi S, Menaldo DL, de Menezes CS, Nomizo A, Hamaguchi A, Silveira-Lacerda EP, Homs-Brandeburgo MI, Sampaio SV, Soares AM, Rodrigues VM (2009) Antitumor effects of snake venom chemically modified Lys49 phospholipase A2-like BthTX-I and a synthetic peptide derived from its C-terminal region. *Biologicals* 37:222–229
49. Souza DH, Iemma MR, Ferreira LL, Faria JP, Oliva ML, Zingali RB, Niewiarowski S, Selistre-de-Araujo HS (2000) The disintegrin-like domain of the snake venom metalloprotease alternagin inhibits  $\alpha$ 2b1 integrin-mediated cell adhesion. *Arch Biochem Biophys* 384:341–350
50. Moura-da-Silva AM, Baldo C (2012) Jararhagin, a hemorrhagic snake venom metalloproteinase from *Bothrops jararaca*. *Toxicon* 60:280–289