

NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 9. Issue 2. Pages 145-292. 2014
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS**PROFESSOR ALEJANDRO F. BARRERO**

Department of Organic Chemistry,
University of Granada,
Campus de Fuente Nueva, s/n, 18071, Granada, Spain
afbarre@ugr.es

PROFESSOR ALESSANDRA BRACA

Dipartimento di Chimica Bioorganica e Biofarmacia,
Università di Pisa,
via Bonanno 33, 56126 Pisa, Italy
braca@farm.unipi.it

PROFESSOR DEAN GUO

State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry
University of Wollongong
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry
The University of Alabama in Huntsville
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Institute of Natural Medicine
Institute of Natural Medicine, University of Toyama,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@inm.u-toyama.ac.jp

PROFESSOR DAVID E. THURSTON

Department of Pharmaceutical and Biological Chemistry,
The School of Pharmacy,
University of London, 29-39 Brunswick Square,
London WC1N 1AX, UK
david.thurston@pharmacy.ac.uk

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Roberto G. S. Berlinck
São Carlos, Brazil

Prof. Anna R. Bilia
Florence, Italy

Prof. Maurizio Bruno
Palermo, Italy

Prof. César A. N. Catalán
Tucumán, Argentina

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Fatih Demirci
Eskişehir, Turkey

Prof. Dominique Guillaume
Reims, France

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Hisahiro Hagiwara
Niigata, Japan

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F., Mexico

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Vladimir I Kalinin
Vladivostok, Russia

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Göttingen, Germany

Prof. Marie Laccaille-Dubois
Dijon, France

Prof. Shoen-Sheng Lee
Taipei, Taiwan

Prof. Imre Mathe
Szeged, Hungary

Prof. Ermino Murano
Trieste, Italy

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Luca Rastrelli
Fisciano, Italy

Prof. Stefano Serra
Milano, Italy

Prof. Monique Simmonds
Richmond, UK

Dr. Bikram Singh
Palampur, India

Prof. John L. Sorensen
Manitoba, Canada

Prof. Johannes van Staden
Scottsville, South Africa

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2014 subscription price: US\$2,395 (Print, ISSN# 1934-578X); US\$2,395 (Web edition, ISSN# 1555-9475); US\$2,795 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Nutritional and Functional Properties of Aqueous and Hydroalcoholic Extracts from Argentinean Propolis

Fátima C. Danert^a, Catiana Zampini^{a,b,c,#}, Roxana Ordoñez^{a,b,c}, Luis Maldonado^d, Enrique Bedascarrasbure^d and María Inés Isla^{a,b,c,#}

^aINQUINOA (Instituto de Química del Noroeste Argentino). CONICET; ^bFacultad de Bioquímica, Química y Farmacia. Ayacucho 471; ^cFacultad de Ciencias Naturales e IML. Miguel Lillo 205. Universidad Nacional de Tucumán. 4000. San Miguel de Tucumán. Tucumán. Argentina; ^dEstación Experimental Agropecuaria Famaillá. Instituto Nacional de Tecnología Agropecuaria. Ruta provincial 301, km 32, Famaillá, Tucumán, Argentina.

#Both authors have the same participation

misla@tucbbs.com.ar

Received: October 26th, 2012; Accepted: December 16th, 2013

Bee propolis is a natural product extensively used as an ingredient in functional foods in amounts that may confer health benefits. The aim of this study was to investigate the nutritional and functional properties (antioxidant activity) of aqueous and ethanolic extracts of propolis samples from Santiago del Estero province, northern Argentina. All propolis extracts contained macronutrients (glucose, fructose, sucrose and proteins), micronutrients (Na, K, Ca, P and Mg) and trace elements (Fe). Spectrophotometric, TLC and HPLC-DAD analyses showed the presence of several phenolic, flavonoid and non-flavonoid compounds, but in all cases the flavonoids prevailed. The PCA of polyphenolic content provided a clear separation of propolis in Group 1 (SE1, 2, 3, 4 and 7) and Group 2 (SE5 and 6) associated with phenolic compound content and collection regions. Two compounds, pinocembrin and chrysin, which could be used as chemical markers of Santiago del Estero propolis, were identified in all samples. Propolis samples extracted with water presented better radical scavenging ability than ethanolic extracts, independent of the antioxidant method (scavenging activity of ABTS^{•+}, DPPH[•], HO[•] and O₂^{•-} and β-carotene bleaching test). Such results correlated closely with the levels of total phenols and flavonoids in samples. The results justify the use of Argentine propolis as a functional dietary supplement.

Keywords: Argentine propolis, Macronutrients, Micronutrients, Functional compounds.

Bee propolis is a natural product extensively used as a phytochemical ingredient in functional foods in amounts that may confer health benefits. Since the chemical composition of propolis depends on the phytogeographical characteristics of the collection site, and as Argentina has a wide biodiversity, propolis obtained in different regions (northwest, northeast, Cuyo, center and Patagonia) at different times of the year is expected to have different chemical composition and, consequently, different biological activities. Several flavonoids and chalcones were identified in Argentine propolis, principally in northwestern Argentine propolis [1a-g]. These phytochemicals have antioxidant, anti-inflammatory, antibacterial and antifungal properties [1a-g, 2a-f]. However, up to now, the nutritional properties of propolis have not been analyzed, despite the fact that it is included in the Argentine Food Code.

Propolis ethanolic extract (PEE) is well known and has attracted much interest. Ethanol is the most commonly used solvent because the lipophilic compounds of propolis are easy to extract with it. However, this method has some drawbacks such as a strong residual flavor, adverse reactions and the intolerance to alcohol of some people [3a]. Propolis aqueous extract (PAE) has been featured in few reports, although PAE and its main constituents (including caffeoylquinic acids) have greater antioxidative effects and greater inhibitory activity against some enzymes than PEE and its constituents [3b]. The aim of this study was to investigate the nutritional and functional properties (antioxidant activity) of aqueous and ethanolic propolis extracts from Santiago del Estero province, Northern Argentina.

According to Sforcin & Bankova [4a], a critical step in propolis study is the selection and extraction of the propolis specimens that

Table 1: Content of macronutrients of SE-propolis by HPLC and dry weight of soluble principles extracted with ethanol and water.

Samples	Ethanolic extracts				Protein	Dry weight
	Total Sugars*	Sucrose**	Glucose**	Fructose**		
SE 1	5.2 ± 0.3	1.0 ± 0.05	1.7 ± 0.08	2.4 ± 0.1	20.4 ± 1.1	55.0 ± 0.03
SE 2	6.4 ± 0.02	1.1 ± 0.06	2.1 ± 0.1	1.3 ± 0.07	22.3 ± 3.4	51.2 ± 0.02
SE 3	5.9 ± 0.4	1.0 ± 0.05	1.4 ± 0.07	1.0 ± 0.05	29.8 ± 2.7	51.2 ± 0.03
SE 4	6.5 ± 0.04	1.0 ± 0.05	1.3 ± 0.07	1.0 ± 0.05	15.9 ± 1.3	30.0 ± 0.02
SE 5	10.8 ± 0.1	1.0 ± 0.05	1.0 ± 0.05	0.9 ± 0.05	7.1 ± 1.7	46.2 ± 0.02
SE 6	7.7 ± 0.04	1.3 ± 0.07	2.2 ± 0.1	2.9 ± 0.1	13.5 ± 2.7	41.2 ± 0.02
SE 7	5.7 ± 0.6	1.0 ± 0.05	1.5 ± 0.08	1.0 ± 0.05	20.8 ± 0.6	45.0 ± 0.02
Aqueous extracts						
SE 1	1.7 ± 0.1	0.1 ± 0.01	1.0 ± 0.05	1.1 ± 0.06	0.6 ± 0.06	5.7 ± 0.003
SE 2	1.5 ± 0.04	1.0 ± 0.05	1.0 ± 0.05	0.9 ± 0.05	0.9 ± 0.01	4.8 ± 0.01
SE 3	2.0 ± 0.3	0.1 ± 0.01	0.1 ± 0.01	1.0 ± 0.05	1.1 ± 0.03	4.7 ± 0.01
SE 4	2.1 ± 0.01	0.1 ± 0.01	1.0 ± 0.05	1.0 ± 0.05	1.4 ± 0.2	9.6 ± 0.01
SE 5	1.7 ± 0.1	2.5 ± 0.12	1.0 ± 0.05	1.0 ± 0.05	0.8 ± 0.02	3.3 ± 0.01
SE 6	2.1 ± 0.1	0.1 ± 0.01	0.1 ± 0.01	0.1 ± 0.01	0.7 ± 0.1	4.8 ± 0.01
SE 7	1.9 ± 0.1	ND	0.9 ± 0.05	0.9 ± 0.05	1.0 ± 0.1	5.8 ± 0.01

* all data were expressed as g/100 g of propolis. ND: not detected.

will be used. Propolis samples from different regions of Santiago del Estero, Argentina, were selected for this study. Water and 80° ethanol were used as extraction solvents. Water dissolved a small part of the constituents of propolis, about 3-10% of its weight, whereas 80° ethanol dissolved 30-50%, depending on the sample (Table 1). To our knowledge, propolis sugar content (sucrose, fructose and glucose) has been investigated in propolis from Kenya, Tanzania (≤ 1%) [4b], Greece, Cyprus (0.7 to 49.0 %) [4c] and Malta (0.8 to 37.2%) [4d], but not in Argentine propolis. All propolis from the SE exhibited the presence of sugars, with values from 5.2 to 10.8% for PEE and 1.5 to 2.1% for PAE (Table 1). Glucose, fructose and sucrose are believed to originate from either nectar or honey, introduced occasionally by bees, or hydrolyzed flavonoid glycosides and non-flavonoid glycosides [4b]. Mucilages were listed among potential sources of sugars in propolis [4e]. Until

now, the hypothesis that some plant mucilages are additional sources of sugars seems better-founded because there is numerous proof that bees collect propolis from plant materials that contain flavonoid aglycones, but not glycosides [4b,4e].

Table 2: Phenolic compounds and flavonoid content in aqueous and ethanolic propolis extracts from different regions of Santiago del Estero.

PEE	Phenolic compounds g GAE/100 g prop	Flavonoids g QE/100 g prop	Phenolic non flavonoids g GAE/100 g prop
SE 1	34.8 ± 2.6	66.0 ± 1.5	17.5 ± 1.2
SE 2	37.9 ± 1.7	49.7 ± 1.6	15.3 ± 1.5
SE 3	31.8 ± 2.7	56.8 ± 2.5	23.0 ± 1.7
SE 4	40.9 ± 3.4	34.6 ± 0.9	13.1 ± 1.4
SE 5	27.3 ± 1.9	21.3 ± 0.1	5.7 ± 0.6
SE 6	27.8 ± 0.4	29.8 ± 0.6	9.0 ± 0.8
SE 7	42.4 ± 1.1	46.6 ± 0.8	14.2 ± 1.0

PAE	Phenolic compounds g GAE/100g prop	Flavonoids g QE/100g prop	Phenolic non flavonoids g GAE/100g prop
SE 1	0.9 ± 0.04	2.1 ± 0.04	0.8 ± 0.01
SE 2	1.5 ± 0.05	3.1 ± 0.4	0.8 ± 0.02
SE 3	1.3 ± 0.01	3.5 ± 0.2	1.1 ± 0.014
SE 4	2.2 ± 0.2	4.3 ± 0.3	0.9 ± 0.002
SE 5	0.9 ± 0.01	3.0 ± 0.3	0.6 ± 0.009
SE 6	1.0 ± 0.03	2.5 ± 0.1	0.6 ± 0.002
SE 7	1.4 ± 0.02	3.2 ± 0.4	1.0 ± 0.02

The protein content in PEE was between 7.1 and 29.1%, while in PAE the values were between 0.6 and 1.4% (Table 1). This is the first report about protein content in propolis samples. Micronutrients are involved in numerous biochemical processes and an adequate intake of some of them relates to the prevention of diseases [5a]. Mineral element characterization was carried out. Among the macroelements, K was the main mineral with 313.0-655.0 mg/100 g, followed by Mg (51.6-227.7 mg/100 g), P (29.7-77.4 mg/100 g) and Ca (12.0-60.4 mg/100 g). Na presented values between 24.9-136.6 mg/100 g. Regarding microelements, Fe presented high values (32.1-75.0 mg/100 g).

TLC analysis showed the presence of several phenolic, flavonoid (F) and non-flavonoid (NF) compounds, but in all cases the flavonoids were the major bands. Quantitative assays confirmed that flavonoids are the main phenolic compounds in all PEE (Table 2). PEE possessed significantly higher amounts of total phenols and flavonoids than aqueous extracts. The results indicated that the content of total phenols and flavonoids varied according to the collection zone. SE3 and SE1 propolis had the highest non flavonoid and flavonoid contents, respectively (Table 2). According to PCA, the PC1 divides SE propolis into group 1 (SE1, 2, 3, 4 and 7) and group 2 (SE5 and 6); both groups were significantly different mainly in their polyphenolic compound content (CCF). PC2 associates SE propolis 2, 4 and 7 with the phenolic compounds content, and SE1 and SE3 with F and NF. These two components explain 86.1% of the total variability of SE propolis (Figure 1).

The HPLC-DAD chromatographic pattern of all PEE was different in quantity and absorption intensity of peaks. SE3 extracts showed the major peak numbers with UV-spectra characteristic of flavonoids [5b], while the other extracts showed lower numbers of constituents. These results reveal a major structural diversity for SE3 propolis. The peaks at 19.6 and 21 min were identified as pinocembrin and chrysin, respectively, and were detected in all Santiago del Estero propolis extracts. To identify each peak, PEE was co-eluted with reference compounds. These same compounds were identified previously in ethanolic extract of Argentine propolis from other regions [1a,1d] and in propolis from other countries [4c].

A rapid TLC fingerprinting of Argentine propolis, together with reference substances, confirmed two main components (chrysin and pinocembrin) from their R_f values and UV fluorescence after spraying with NP. The UV profiles in HPLC analysis of aqueous extracts displayed similar results as those observed in quantitative




Figure 1: PCA of propolis from Santiago del Estero, Argentina.

Table 3: SC₅₀ values (µgGAE/mL) of aqueous extracts (PAE) and ethanolic extracts (PEE) of propolis from different regions of Santiago del Estero.

Sample	ABTS ^{•+} scavenging		DPPH [•] scavenging		β-carotene bleaching	
	PEE	PAE	PEE	PAE	PEE	PAE
SE1	6.0±0.4	1.1±0.06	36.0±2.7	10.0±1.1	10.8±0.8	2.2±0.2
SE2	5.7±1.0	0.8±0.06	20.0±1.5	6.8±0.5	4.5±0.3	1.8±0.1
SE3	2.5±0.1	0.8±0.1	9.0±0.7	9.0±0.8	2.8±0.2	1.7±0.1
SE4	9.5±0.4	0.8±0.08	20.0±1.5	8.4±0.5	7.6±0.6	2.4±0.2
SE5	8.3±0.6	0.8±0.08	29.0±2.2	12.5±0.6	7.2±0.5	1.8±0.1
SE6	7.1±0.4	0.8±0.03	34.0±2.5	12.8±2.1	3.6±0.3	2.2±0.2
SE7	5.5±0.3	0.6±0.2	17.0±1.3	6.8±0.6	6.5±0.5	2.1±0.2

analysis of total phenolic and non-flavonoid phenolics. PAE showed lower peak numbers than PEE and the highest intensities tended to be crowded in places with low retention times.

PAE presented the highest scavenging activity with the lowest SC₅₀ values (Table 3). All aqueous extracts were effective ABTS^{•+} scavengers in a concentration-dependent manner in SE samples from 0.5 to 2.5 µg GAE/mL (r²: 0.99) with similar SC₅₀ values and around 1 µg GAE/mL. On the other hand, SC₅₀ values for ethanolic extracts were between 2.5 and 10 µg GAE/mL. SE3 ethanolic extracts were the most active ABTS^{•+} scavenger. The results demonstrated that all extracts had marked electron donor properties for neutralizing free radicals by forming stable products.

SC₅₀ values for aqueous and ethanolic extracts, BHT, and Trolox on the DPPH radical were 4 to 15 µg GAE/mL (r²: 0.985), 9 to 36 µg GAE/mL (r²: 0.813), 11.0 µg/mL (r²: 0.994) and 10 mM (r²: 0.970), respectively. SE3 ethanolic extract was the most active free radical scavenger. All extracts were able to protect linoleic acid from oxidation with IC₅₀ values of 1.7-2.4 µg GAE/mL for PAE and 3.0-11.0 µg GAE/mL for PEE. The hydroxyl radical is the most reactive as it can attack and damage almost every molecule found in living cells. OH[•] reactions include the ability to interact with DNA purine and pyrimidine. It can also abstract hydrogen atoms from biological molecules, including thiols, leading to the formation of sulfur radicals able to combine with oxygen to generate oxysulfur radicals, a number of which damage biological molecules. Only aqueous extracts were assayed as the reactants used to determine the scavenging capacity of HO[•] and O₂^{•-} can react with most of the organic solvents and substances employed for buffer preparations. SC₅₀ values of HO[•] and O₂^{•-} for aqueous extracts were 10 to 100 µg GAE/mL and 53 to 275 µg GAE/mL, respectively (data not shown).

In conclusion, SE propolis from Argentina includes nutrients such as sugars, proteins, and minerals that are essential in ATP- energy producing metabolism. Non-nutrients such as phenolic compounds, principally flavonoids, were found to play important roles in oxidative stress as antioxidants that scavenge reactive species related to various lifestyle-related diseases, e.g., obesity, type 2 diabetes or cardiovascular diseases.

Experimental

Propolis samples: Several samples from different areas of Santiago del Estero, Northwestern Argentina, were analyzed: Departamento Capital, El Palomar (SE1), Departamento Figueroa (SE2 and SE7), Departamento Banda (SE3 and SE5), Departamento Copo (SE4), and Departamento Robles (SE6). The samples were stored at -20°C until use.

Sample processing: Two g of ground propolis was extracted with 25 mL distilled water for 5 min in an ultrasonic bath at room temperature. The suspension was boiled at 80°C for 2 h and centrifuged at 5000 g. The residue was re-extracted under the same conditions and the supernatants were combined to reach a final volume of 100 mL of propolis aqueous extract (PAE).

Two g of ground propolis was extracted with 15 mL 80° ethanol and heated at 70°C for 35 min. The preparation was centrifuged at 8000 g for 10 min. The insoluble material was re-extracted with ethanol and the supernatants combined to produce propolis ethanolic extract (PEE) [6a].

Thin-layer chromatography (TLC): PEE and PAE (3 µg and 5 µg, respectively) were spotted onto TLC plates (Kieselgel 60 F254 0.2 mm, Merck), which were developed using chloroform: ethyl acetate (80:20). The separated components were visualized under ultraviolet light (254 and 365 nm, UV Lamp Model UVGL-58 Mineralight Lamp, USA), followed by spraying with NP reagent [6b].

Chemical profile by HPLC analysis: Chromatography was performed using a HPLC system consisting of a Waters 1525 Binary HPLC Pumps system with a 1500 Series Column Heater, (Rheodyne Inc., Cotati, CA). A Waters 2998 photodiode array detector (PAD) was used to analyze the extracts. A XBridge™ C18 column (4.6 x 150 mm, 5 µm; Waters corporation, Milford, MA) at 40°C and a gradient solvent system consisting of solvent A (methanol) and solvent B (9% acetic acid) (conditions: 25–45% A from 0 to 10 min and kept at 45% A from 10 to 20 min; 45–70% A from 20 to 40 min; 70–75% A from 40 to 50 min; 75–100% A from 50 to 55 min and kept at 100% A from 55 to 60 min) were used for separation. The flow rate was set at 0.8 mL/min. The sample injection volume was 20 µL. PAD acquisitions were performed in the range 190–500 nm, and chromatograms were integrated at 280 nm.

Sugar determination by spectrophotometric method and HPLC: The phenol–sulfuric acid method [6c] was used for determination of total neutral sugars in PAE and PEE. Results were expressed as g of glucose/100 g dry weight (DW). The samples were also analyzed by HPLC (Waters) with an automatic injector {ALLIANCE Waters 2695, detector IR Waters 410 and column sugar pack (Waters)}. The elution solvents were water containing EDTA-Na. The flow rate was 0.5 mL/min. Temperature: 85°C. The retention times of identified compounds were checked by co-injection with commercial standards.

Protein determination: Soluble protein concentration was determined by the method of Bradford [6d]. Results were expressed as mg of bovine serum albumin (BSA) /100 g DW.

Minerals: Na, K, Ca, Fe and Mg were determined by atomic absorption spectroscopy using a Perkin Elmer 3110 spectrophotometer (acetylene air flame and hollow-cathode lamps).

P was determined using an atomic absorption spectrometer 600 (Perkin Elmer). Results are expressed as mg/100 g propolis.

Determination of total phenolic and non-flavonoid phenolic content: Total phenolic content of the samples was determined using the Folin–Ciocalteu method [7a]. Results are expressed in g gallic acid equivalents (GAE) per 100 g DW. Non-flavonoid phenols were measured by determination of total phenol content remaining after precipitation of the flavonoids with acidic formaldehyde [7b]. Results are expressed in g GAE/100 g DW.

Total flavonoids: Total flavonoid content was determined with aluminum chloride [7c] and expressed as g quercetin equivalents (QE) per 100 g DW.

ABTS free radical scavenging activity: The antioxidant capacity assay was carried out by an improved ABTS^{•+} method [8a]. ABTS^{•+} solution (1 mL) was added to PAE (0.625 to 5 µg GAE/mL) and PEE (2.5 to 10 µg GAE/mL). Absorbance was read at 734 nm at 1 min and 6 min. SC₅₀ was determined (concentration of total phenolic compound able to scavenge 50% of ABTS^{•+}). Results were expressed as µg GAE/mL. BHT and Trolox were used as reference antioxidants.

DPPH free radical scavenging activity: DPPH radical scavenging activity was measured [8b]. DPPH solution (1.5 mL of 300 µM in 96° ethanol) was added to 0.5 mL of PAE (containing 5 to 20 µg of GAE/mL) and PEE (containing 2.5 to 10 µg of GAE/mL) and shaken for 20 min at room temperature. Absorbance was measured at 515 nm. The percentage inhibition $[(A_0 - A_1/A_0) \times 100]$ was plotted against phenolic compounds content and SC₅₀ values were determined (concentration of total phenolic compounds able to scavenge 50% of DPPH free radical). BHT and Trolox were used as reference radical scavengers.

β-Carotene bleaching assay: Antioxidant activity was determined according to the β-carotene bleaching method [8b]. The initial absorbance at 470 nm was registered at zero time (t₀) and at 120 min. Antioxidant activity (AA%) was calculated as percent inhibition relative to control. IC₅₀ values denote the sample concentration required to inhibit 50% β-carotene bleaching.

Hydroxyl radical scavenging: The deoxyribose assay as described by Chobot [8c], with a slight modification, was applied to measure HO[•] scavenger capacity. The aqueous extract was dissolved in a KH₂PO₄/KOH buffer solution (50 mM, pH 7.4) to yield final concentrations from 10 to 200 µg GAE/mL; 50 µL of a 10.4 mM 2-deoxy-D-ribose solution, 100 µL of FeCl₃ (50 µM) and 100 µL of 52 µM EDTA were added. To start the Fenton reaction, 50 µL of 10 mM H₂O₂ and 50 µL of 1.0 mM ascorbic acid were added. The mixture was incubated at 37°C for 60 min. Then, 500 µL of 1% 2-thiobarbituric acid dissolved in 3% trichloroacetic acid (w/v) was added to each vial and maintained at 100°C for 20 min. To extract the reaction product of malondialdehyde (MDA) and thiobarbituric acid, 700 µL of *n*-butanol was added, and the mixture was vigorously vortexed. The *n*-butanol layers of the tubes, each 600 µL, were separated and the absorbance was determined at 532 nm. Assays were performed in triplicate. Reaction mixtures without the test compound served as positive control (100% MDA). The negative control contained the full reaction mixture except 2-deoxy-D-ribose. Controls without either EDTA or ascorbic acid were performed. Inhibition (I) of deoxyribose degradation in percent was calculated.

Superoxide anion radical ($O_2^{\cdot-}$) scavenging capacity: The $O_2^{\cdot-}$ were generated in 120 μ L of sodium phosphate buffer (19 mM, pH 7.4), which contained 40 μ L NADH (2 mM), 20 μ L of NBT (1 mM), 40 μ L of PMS (60 μ M), PEE and PAE at different concentrations or distilled water and ethanol for controls. The color reaction was detected at 550 nm using a Microplate reader [8d]. SC_{50} values denote the μ g GAE/mL required to scavenge 50% of superoxide free radicals.

Statistical analysis: Sampling and analyses were performed in triplicate, and the data are presented as mean \pm standard deviation.

References

- [1] (a) Agüero M, González M, Lima B, Svetaz L, Sanchez M, Sacchino S, Feresin G, Schmeda-Hirschmann G, Palermo J, Tapia A. (2010) Argentinean propolis from *Zuccagnia punctata* Cav. (Caesalpinieae) exudates: Phytochemical characterization and antifungal activity. *Journal of Agricultural and Food Chemistry*, **58**, 194–201; (b) Chaillou L, Nazareno M. (2009) Bioactivity of propolis from Santiago del Estero, Argentina, related to their chemical composition. *LWT - Food Science Technology*, **42**, 1422–1427; (c) Kumazawa S, Ahna M, Fujimoto T, Kato M. (2010) Radical-scavenging activity and phenolic constituents of propolis from different regions of Argentina. *Natural Product Research*, **24**, 804–812; (d) Isla MI, Paredes Guzman J, Nieva Moreno MI, Koo H, Park YK. (2005) Some chemical composition and biological activity of northern Argentine propolis. *Journal of Agricultural and Food Chemistry*, **53**, 1166–1172; (e) Isla MI, Carrasco Juárez B, Nieva Moreno MI, Zampini I, Ordoñez R, Sayago J, Cuello S, Alberto MR, Bedescarrabure E, Alvarez A, Coccini F, Maldonado LM. (2009) Effect of seasonal variations and collection form on antioxidant activity of propolis from San Juan, Argentina. *Journal of Medicinal Food*, **12**, 1334–1342; (f) Nieva Moreno MI, Zampini IC, Ordoñez RM, Vattuone M, Isla MI. (2005) Evaluation of the cytotoxicity, mutagenicity and antimutagenicity of propolis from Amaicha del Valle, Tucumán, Argentina. *Journal of Agricultural and Food Chemistry*, **53**, 8957–8962; (g) Vera N, Solórzano E, Ordoñez R, Maldonado L, Bedascarrabure E, Isla MI. (2011) Chemical composition of Argentinean propolis collected in extreme regions and its relation with antimicrobial and antioxidant activities. *Natural Product Communications*, **6**, 823–827.
- [2] (a) Isla MI, Nieva Moreno MI, Vattuone MA, Sampietro AR. (2001) Antioxidant activity of Argentine propolis extracts. *Journal of Ethnopharmacology*, **76**, 165–170; (b) Lima B, Tapia A, Luna L, Fabani MP, Schmeda-Hirschmann G, Podio NS, Wunderlin DA, Feresin G, Main GE. (2009) Flavonoids, DPPH activity, and metal content allow determination of the geographical origin of propolis from the Province of San Juan (Argentina). *Journal of Agricultural Food Chemistry*, **57**, 2691–2698; (c) Lozina L, Peichoto M, Acosta O, Granero G. (2010) Standardization and organoleptic and physicochemical characterization of 15 Argentinean propolis. *Latin American Journal of Pharmacology*, **29**, 102–110; (d) Nieva Moreno MI, Isla MI, Cudmani NG, Vattuone MA, Sampietro AR. (1999) Screening of antibacterial activity of Amaicha del Valle (Tucumán, Argentina) propolis. *Journal of Ethnopharmacology*, **68**, 97–102; (e) Nieva Moreno MI, Isla MI, Vattuone MA, Sampietro AR. (2000) Comparison of the free radical-scavenging activity of propolis from several regions. *Journal of Ethnopharmacology*, **71**, 109–114; (f) Ordoñez R, Zampini IC, Nieva Moreno MI, Isla MI. (2011) Potential application of Argentine propolis to control some phytopathogenic bacteria. *Microbiology Research*, **166**, 578–584.
- [3] (a) Konishi S, Sawaya A, Custódio AR, Cunha I, Shimizu M. (2004) Influence of solubilising agents on antimicrobial activity of propolis extracts and hydro alcoholic spray formula. *Mensagem Doce*, **75**, 22–25; (b) Matsui T, Ebuchi S, Fujise T, Abesundara KJ, Doi S, Yamada H, Matsumoto K. (2004) Strong antihyperglycemic effects of water-soluble fraction of Brazilian propolis and its bioactive constituent, 3, 4, 5-tri-*O*-caffeoylquinic acid. *Biological & Pharmaceutical Bulletin*, **27**, 1797–1803.
- [4] (a) Sforzin JM, Bankova V. (2011) Propolis: Is there a potential for the development of new drugs? (Review). *Journal of Ethnopharmacology*, **133**, 253–260; (b) Qian W L, Khana Z, Watson DG, Fearnley J. (2008) Analysis of sugars in bee pollen and propolis by ligand exchange chromatography in combination with pulsed amperometric detection and mass spectrometry. *Journal of Food Composition and Analysis*, **21**, 78–83; (c) Kalogeropoulos N, Konteles SJ, Troullidou EM, Karathanos V. (2009) Chemical composition, antioxidant activity and antimicrobial properties of propolis extracts from Greece and Cyprus. *Food Chemistry*, **116**, 452–461; (d) Popova M, Trusheva B, Antonova D, Cutajar S, Mifsud D, Farrugia C, Tsvetkova I, Najdenski H, Bankova V. (2011) The specific chemical profile of Mediterranean propolis from Malta. *Food Chemistry*, **126**, 1431–1435; (e) Bankova VS, Christov RS, Tejera AD. (1998) Lignans and other constituents of propolis from the Canary Islands. *Phytochemistry*, **49**, 1411–1415.
- [5] (a) Palmer J, Venkateswaran V, Fleshner NE, Klotz LH, Cox ME. (2008) The impact of diet and micronutrient supplements on the expression of neuroendocrine markers in murine Lady transgenic prostate. *Prostate*, **68**, 345–353; (b) Mabry TJ, Markham KR, Thomas MB. (1970) *The Systematic Identification of Flavonoids*. Springer-Verlag, Berlin.
- [6] (a) Norma IRAM-INTA 15935-2. (2008) Extractos de Propóleos. Instituto Argentino de Normalización -Buenos Aires, Argentina; (b) Wagner H, Blatt S, Zgainsky EM. (1984) *Plant drug analysis*. Springer, Berlin; (c) Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. (1956) Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, **28**, 350–356; (d) Bradford MM. (1976) Rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, **72**, 248–254.
- [7] (a) Singleton VL, Orthofer R, Lamuela-Raventos RM. (1999) Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, **299**, 152–178; (b) Isla MI, Salas A, Danert FC, Zampini IC, Ordoñez RM. (2013) Analytical methodology optimization to estimate the content of non-flavonoid phenolic compounds in Argentine propolis extracts. *Pharmaceutical biology*. in press; (c) Woisky R, Salatino A. (1998) Analysis of propolis: some parameters and procedures for chemical quality control. *Journal of Apiculture Research*, **37**, 99–105.
- [8] (a) Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology Medicinal*, **26**, 1231–1237; (b) Ordoñez AA, Gómez D, Vattuone MA, Isla MI. (2006) Antioxidant activity of *Sechium edule* (Jacq) Swartz. *Food Chemistry*, **97**, 452–458; (c) Chobot V. (2010) Simultaneous detection of pro- and antioxidative effects in the variants of the deoxyribose degradation assay. *Journal of Agricultural Food Chemistry*, **58**, 2088–2094; (d) Valentão P, Fernandes E, Carvalho F, Andrade PB, Seabra RM, Bastos ML. (2001) Antioxidant activity of *Centaurium erythraea* infusion evidenced by its superoxide radical scavenging and xanthine oxidase inhibitory activity. *Journal of Agricultural and Food Chemistry*, **49**, 3476–3479.

Cyclic Dipeptides Produced by Marine Sponge-Associated Bacteria as Quorum Sensing Signals Gennaro Roberto Abbamondi, Salvatore De Rosa, Carmine Iodice and Giuseppina Tommonaro	229
Isolation of the Tetrapeptide Apicidins G, H and I from the Fungus <i>Fusarium semitectum</i> Suciati and Mary J. Garson	233
The Co-identity of Lipiarmycin A3 and Tiacumicin B Angelo Bedeschi, Piera Fonte, Giovanni Fronza, Claudio Fuganti and Stefano Serra	237
PSY-I, a <i>Taxus chinensis</i> var. <i>mairei</i> Extract, Inhibits Cancer Cell Metastasis by Interfering with MMPs Zao-qian Zheng, Ying-Ying Fu, Bo-Heng Li, Mei-Ling Zhang, Xiu-Li Yang, Chuan-wei Xin, Jia-na Shi, Yin Ying and Ping Huang	241
Antimicrobial Activity of Endophytic Fungi Isolated from <i>Swietenia macrophylla</i> Leaves Darah Ibrahim, Chong Chai Lee and Lim Sheh-Hong	247
Aroma Profile of Star Anise and the Structure-odor Relationship of Anethole Toshio Hasegawa, Haruna Seimiya, Takashi Fujihara, Noriko Fujiwara and Hideo Yamada	251
The Essential Oil of <i>Populus balsamifera</i> Buds: its Chemical Composition and Cytotoxic Activity Marianne Piochon-Gauthier, Jean Legault, Muriel Sylvestre and André Pichette	257
Comparative Chemical Study and Cytotoxic Activity of <i>Uvariadendron angustifolium</i> Essential Oils from Benin Jean-Pierre Noudogbessi, Magali Gary-Bobo, Aristide Adomou, Elvis Adjalian, Guy Alain Alitonou, Félicien Avlessi, Marcel Garcia, Dominique C. K. Sohounhloue and Chantal Menut	261
Composition of Essential Oil from <i>Tagetes minuta</i> and its Cytotoxic, Antioxidant and Antimicrobial Activities Nasser A. Awadh Ali, Farukh S. Sharopov, Ali G. Al-kaf, Gabrielle M. Hill, Norbert Arnold, Saeed S. Al-Sokari, William N. Setzer and Ludger Wessjohann	265
Chemical Composition of the Essential Oil of <i>Croton bonplandianus</i> from India Rajesh K. Joshi	269
Chemical Composition of <i>Angelica panicii</i> Essential Oil Determined by Liquid and Headspace GC-MS Techniques Strahinja R. Simonović, Vesna P. Stankov-Jovanović, Violeta D. Mitić, Marija D. Ilić, Goran M. Petrović and Gordana S. Stojanović	271
Chemical Description and Essential Oil Yield Variability of Different Accessions of <i>Salvia lavandulifolia</i> Jaime Usano-Aleman, Jesús Palá-Paúl, Manuel Santa-Cruz Rodríguez and David Herraiz-Peñalver	273
Antimicrobial Constituents and Synergism Effect of the Essential Oils from <i>Cymbopogon citratus</i> and <i>Alpinia galanga</i> Sarin Tadtong, Rith Watthanachaiyingcharoen and Narisa Kamkaen	277
<i>In vitro</i> Antibacterial Activity of <i>Libanotis montana</i> Essential Oil in Combination with Conventional Antibiotics Dragoljub L. Miladinović, Budimir S. Ilić, Tatjana M. Mihajilov-Krstešev, Jovana L. Jović and Marija S. Marković	281
<u>Review/Account</u>	
Modernization of Ayurveda: A Brief Overview of Indian Initiatives Ambarish Mukherjee, Mousumi Banerjee, Vivekananda Mandal, Amrithesh C. Shukla and Subhash C. Mandal	287

Natural Product Communications

2014

Volume 9, Number 2

Contents

<u>Original Paper</u>	<u>Page</u>
Volatile Glycosides from the Leaves of <i>Morus alba</i> with a Potential Contribution to the Complex Anti-diabetic Activity Attila Hunyadi, Ibolya Herke, Katalin Veres, Anna Erdei, András Simon and Gábor Tóth	145
A New Eudesmane Sesquiterpene from <i>Dichrocephala integrifolia</i> Fang Qin, Yi-Bing Wu, Rui-xia Guo, Mei Dong, Françoise Sauriol, Qing-Wen Shi, Yu-Cheng Gu and Hiromasa Kiyota	149
Evaluation of the Anti-melanoma Activities of Sarcophine, (+)-7α,8β-Dihydroxydeepoxysarcophine and Sarcophytolide from the Red Sea Soft Coral <i>Sarcophyton glaucum</i> Pawel T. Szymanski, Safwat A. Ahmed, Mohamed M. Radwan, Sherief I. Khalifa and Hesham Fahmy	151
Steroidal Aglycones from Stems of <i>Marsdenia tenacissima</i> that Inhibited the Hedgehog Signaling Pathway Lin Zhang, Feng-yang Chen, Shi-fang Xu, Yi-ping Ye and Xiao-yu Li	155
Chemical Constituents, and their Cytotoxicity, of the Rare Wood Decaying Fungus <i>Xylaria humosa</i> Sirirath Sodngam, Sasiphimol Sawadsitang, Nuttika Suwannasai and Wiyada Mongkolthanaruk	157
Alkaloids from <i>Habranthus tubispathus</i> and <i>H. jamesonii</i>, two Amaryllidaceae with Acetyl- and Butyrylcholinesterase Inhibition Activity Valeria Cavallaro, Natalia P. Alza, María G. Murray and Ana P. Murray	159
Further Characterization of Foliar Flavonoids in <i>Crossostephium chinense</i> and their Geographic Variation Ayumi Uehara, Junichi Kitajima, Goro Kokubugata and Tsukasa Iwashina	163
Hawaiian Propolis: Comparative Analysis and Botanical Origin Saori Inui, Takahiro Hosoya and Shigenori Kumazawa	165
Nutritional and Functional Properties of Aqueous and Hydroalcoholic Extracts from Argentinean Propolis Fátima C. Danert, Catiana Zampini, Roxana Ordoñez, Luis Maldonado, Enrique Bedascarrasbure and María Inés Isla	167
Anti-trypanosomal Phenolic Derivatives from <i>Baccharis uncinella</i> Simone dos S. Grecco, Maria Júlia P. Félix, João Henrique G. Lago, Érika G. Pinto, André G. Tempone, Paulete Romoff, Marcelo José P. Ferreira and Patricia Sartorelli	171
Polyphenols in Representative <i>Teucrium</i> Species in the Flora of R. Macedonia: LC/DAD/ESI-MSⁿ Profile and Content Ilija Mitreski, Jasmina Petreska Stanoeva, Marina Stefova, Gjoshie Stefkov and Svetlana Kulevanova	175
In vitro Inhibitory Effects of <i>Limonium contortirameum</i> and <i>L. virgatum</i> Extracts from Sardinia on α-Amylase, α-Glucosidase and Pancreatic Lipase Marzia Foddai, Violet Kasabri, Giacomo L. Petretto, Emanuela Azara, Angela Sias, Fatma U. Afifi, Giovanna Delogo, Mario Chessa and Giorgio Pintore	181
Search for Skin-whitening Agent from <i>Prunus</i> Plants and the Molecular Targets in Melanogenesis Pathway of Active Compounds Kazuya Murata, Keisuke Takahashi, Haruka Nakamura, Kimihisa Itoh and Hideaki Matsuda	185
In Silico Prediction of Tyrosinase and Adenylyl Cyclase Inhibitors from Natural Compounds Pedro Fong, Henry H. Y. Tong and Chi M. Chao	189
Molecular Docking and Reaction Kinetic Studies of Chrysin Binding to Serum Albumin Bingli Jiang, Anran Zhao, Jianhua Miao, Pengfei Chang, Hailin Chen, Weigao Pan and Cuiwu Lin	195
Anthocyanins from the Flowers of Nagai Line of Japanese Garden Iris (<i>Iris ensata</i>) Kaori Kitahara, Yoshinori Murai, Sang Woo Bang, Junichi Kitajima, Tsukasa Iwashina and Yukio Kaneko	201
New Chromone and Triglyceride from <i>Cucumis melo</i> Seeds Sabrin R. M. Ibrahim	205
Five New Acylphloroglucinol Glycosides from the Leaves of <i>Eucalyptus robusta</i> Qian-Yi Guo, Xiao-Jun Huang, Bing-Xin Zhao, Yu-Qing Jian, Shi-Lin Luo, Ying Wang and Wen-Cai Ye	209
Utric acid and Triacylglycerides Production by the Cultured Lichen Mycobiont of <i>Ramalina celastri</i> Alejandra T. Fazio, Mónica T. Adler and Marta S. Maier	213
A New Lignan Glycoside from <i>Chamaecyparis obtusa</i> var. <i>breviramea</i> f. <i>crippsi</i> Jian Xu, Guang-Zhi Zeng, Ke-Li Chen, Yi-Mei Liu, Zhang-Hua Sun, Ning-Hua Tan and Yu-Mei Zhang	215
Synthesis of 2-Acetyl-1,4-Dimethoxynaphthalene, A Potential Intermediate for Disubstituted Naphtho[2,3,c]pyran-5,10-dione Kimberly Chinae, Willian Vera and Ajoy K. Banerjee	217
Anthraquinone and Naphthoquinone Derivatives from the Roots of <i>Coptosapelta flavescens</i> Wipapan Kongyen, Vatcharin Rukachaisirikul, Souwalak Phongpaichit, Nongyao Sawangjaroen, Phruksa Songsing and Hattaya Madardam	219
Methicillin-resistant <i>Staphylococcus aureus</i>, Vancomycin-resistant <i>Enterococcus faecalis</i> and <i>Enterococcus faecium</i> active Dimeric Isobutyrylphloroglucinol from <i>Ivesia gordonii</i>	221
Biological Studies of Turmeric Oil, Part 3: Anti-Inflammatory and Analgesic Properties of Turmeric Oil and Fish Oil in Comparison with Aspirin James N. Jacob and Dinesh K. Badyal	225

Continued inside backcover