

## REVIEW

## Neutrophil apoptosis in the context of tuberculosis infection



Mercedes Alemán\*

IMEX-CONTICET-ANM, Buenos Aires, Argentina

## ARTICLE INFO

## Article history:

Received 20 November 2014

Accepted 24 March 2015

## Keywords:

Apoptosis

Neutrophils

Tuberculosis

## SUMMARY

Polymorphonuclear neutrophils comprise two-thirds of peripheral blood leukocytes and are key components of innate immunity as a first line of defense against bacterial and fungal pathogens. Their microbicidal mechanisms are essential for bacterial killing, the enhancement of inflammatory reactions and also comprise signaling molecules which have been implicated in signal transduction cascades. In tuberculosis, the number of neutrophils increases in the affected lung. In addition, they become activated and apoptotic due to the bacterial burden. As apoptosis is promoted by reactive oxygen species (ROS) during phagocytosis, the advantages and benefits to the host as well as the strategies displayed by the pathogen to avoid or retard apoptosis are discussed in this review.

© 2015 Elsevier Ltd. All rights reserved.

**1. Neutrophils and apoptosis**

The influx of neutrophils to the lung is one of the first events in the pathogenesis of tuberculosis [1] playing a main role in the acute phase of the disease [2]. Neutrophils display microbicidal mechanisms that include the release of proteolytic enzymes, antimicrobial peptides, and the rapid production of ROS [3]. Since the excessive neutrophil activation might cause severe tissue damage and inflammatory diseases [4], certain mechanisms have been evolved in order to resolve inflammation. Programmed cell death in neutrophils, or apoptosis, constrains the release of inflammatory mediators and toxic granule constituents [5] through the recognition and phagocytosis of the bacilli by macrophages, a process that is essential to resolve inflammation [6]. In addition, it has recently been reported a new-ROS dependent cell death mechanism, which leads to neutrophil extracellular traps (NETs) and induces NET-mediated killing [7].

**2. Neutrophil apoptosis and tuberculosis**

As the disease evolves, the number of neutrophils increases in the bronchoalveolar lavage fluid of the affected lung [8]. Although

this mobilization is important for microbicidal function, the powerful mediators such as oxidants and elastase which help neutrophils to be effective killers, can also injure endothelial cells and produce structural damage [9]. In line with this, it has been demonstrated that neutrophils from patients with active tuberculosis display an increased spontaneous apoptosis in vitro [10,11] which is related to the activation state of these cells in bloodstream [12]. Pathogen-induced cell death often involves the modulation of apoptosis and, accordingly, the activation of neutrophils induced by *Mycobacterium tuberculosis* (*Mtb*) accelerates neutrophil apoptosis in vitro [12–14].

Cytokines and cytokine-producing cells are present in pleural effusions from people with malignant and infectious diseases [15]. Patients with tuberculous pleuritis have elevated amounts of tumor necrosis factor (TNF)- $\alpha$  [16,17], transforming growth factor- $\beta$  [17,18], interferon- $\gamma$  [11,16,19], interleukin (IL)-8 [20], IL-6 [21], IL-18 [19], and IL-10 [17], which may modulate the lifetime of neutrophils. In this context it has been described that, after extravasation at the infected lung, the expression of those receptors associated with activation/apoptosis (CD11b, CD64, TNF-R55, and FasL) is up-regulated in neutrophils [22]. Therefore, the cytokine milieu in the pleural space appears to influence the signaling pathways on activated neutrophils, leading to apoptosis and inhibiting proinflammatory capacity.

A large body of evidence has now indicated that neutrophils participate as a bridge between innate and adaptive immune

\* Pacheco de Melo 3081, Buenos Aires, 1425 Argentina. Tel.: +54 11 4 8055695; fax: +54 11 4 803 9475.

E-mail address: [melealeman@hotmail.com](mailto:melealeman@hotmail.com).

responses, for example by acquiring DC markers [22,23], or by leading to specific T lymphocyte responses by crosspresentation [24]. In this regard, phagocytosis of apoptotic bodies promotes a more efficient fusion of the phagosome with the lysosome which results in the digestion of the pathogen and promoting antigen presentation to T cells [25]. Therefore, we can consider the inflammatory milieu as a place subjected to a delicate balance between the infected and uninfected neutrophils, limiting inflammation and/or the generation of specific immune activity.

### 3. Neutrophils and Toll-like receptors

The ability to discriminate between self and non-self is a process in which pattern-recognition receptors such as the Toll like receptors (TLRs) respond to evolutionary conserved non-self proteins [26]. TLRs are members of the IL-1R superfamily whose signaling pathways involve mitogen-activated protein kinase (MAPK) cascades, including c-Jun N-terminal kinase, p38, and extracellular signal-regulated kinases (ERK) MAPK. NF-κB is a heterodimer composed of p50 and p65 subunits which, in resting cells, remains in the cytoplasm because of its interaction with inhibitory κB (IκB) proteins. However, the phosphorylation events which follow TLR stimulation, lead to polyubiquitination and degradation of IκB, allowing nuclear translocation of NF-κB subunits to bind DNA and to trigger transcription of pro-inflammatory cytokine genes [27]. Moreover, the response of neutrophils to cytokines [28] and their activation [29] also involves p38 and ERK MAPK pathway. Human neutrophils express an array of TLRs like TLRs 1, 2, 4, 5, 6, 7, 8, 9, and 10 [30] involved in interleukin-8 (IL-8) secretion, an essential event in neutrophil migration to the lung [31].

### 4. *Mtb* and TLR signaling in neutrophils

Recognition of mycobacterial components by TLRs is a key step in initiating innate immune responses upon mycobacterial infection. In particular, TLR2 appears to be critical for sensing mycobacteria inducing a crucial proinflammatory signal, TNF- $\alpha$  [32].

Signaling pathways triggered by TLRs have been previously implicated in the regulation of neutrophil lifespan [33,34] involving tyrosine phosphorylation events [35]. In fact, it has been shown that *Mtb* activates p38 MAPK in neutrophils in vitro via the TLRs pathway inducing its activation and apoptosis [14,36]. These findings are coherent with high expression of the activated form of this kinase in peripheral blood as well as in pleural effusion neutrophils from tuberculosis patients [22]. Pathogens unable to bind directly to TLRs likely activate signaling complexes via lipid raft reorganization involving numerous cell surface proteins [37]. In this context, a variety of pathogens interact with lipid rafts which are involved in the attachment, cell entry, and intracellular survival of several microorganisms such as *Brucella suis* [38], *Chlamydia trachomatis* [39], and *Mycobacterium bovis* [40]. In addition, membrane lipid rafts are signaling platforms enriched in cholesterol that are required for innate immune responses [41] mediating the efficiency of oxidase coupling to receptors in the neutrophil membrane. On the other hand, TLR2 cooperates with other receptors in lipid rafts on macrophages and DC membrane, for the production of inflammatory cytokines in response to fungal wall-derived β-glucans [42]. Furthermore, it has also been reported that TLR2/lectin-1 cooperation induces ROS production, together with the rapid and sustained phosphorylation of p38 and Syk in monocytes [43,44], and neutrophils [45,46] in response to mycobacteria.

### 5. Neutrophils and reactive oxygen species

Phagocytes produce ROS through activation of nicotinamide adenine dinucleotide phosphate reduced (NADPH) oxidase that is assembled at the plasma membrane. Although the ability of ROS to kill *Mtb* has been demonstrated only in mice [47], the increase in superoxide and hydrogen peroxide could affect important signaling enzymes necessary for physiologic cell signaling. For instance, ROS participate in TLR2 signaling pathways together with a sustained phosphorylation of p38 suggesting an essential role of ROS in TLR2 signaling pathways [48]. In addition, neutrophil apoptosis is dependent on the generation of ROS during the phagocytosis of *Mtb* [13,14] involving the activation of p38. Therefore, *Mtb*-induced respiratory burst in neutrophils can also regulate the inflammatory response by induction of apoptosis.

It has been demonstrated that ROS activate NF-κB by an indirect mechanism [49] through the oxidation of thioredoxin (Trx) that in turn release the transcription factor complex protein-1, which is an upstream kinase in the c-JunN-terminal and p38 kinases [50], permitting its activation and downstream signaling [51]. Human neutrophils express several caspases, cysteine-dependent enzymes and thus potentially redox sensitive, which are key enzymes in apoptosis [52]. In this context, it has been described that *Mtb*-induced apoptosis in neutrophils is dependent on caspase-3, which is dependent on ROS and it has also been shown to be rapidly activated by *Mtb* [14].

### 6. Neutrophils and *Mtb* immune evasion

The ability of *Mtb* to infect and cause disease depends on its capacity to evade killing by phagocytic cells and to achieve this, *Mtb* is equipped with numerous strategies of immune evasion. For instance, it has been reported that virulent mycobacteria are able to escape from fused phagosomes and multiply [53]. In addition, *Mtb*-infected macrophages appear to be diminished in their ability to present antigens to T cells, which leads to persistent infection [54] and even more, *Mtb* has developed mechanisms to prevent both DC migration and antigen presentation [55,56]. In this way, antigen presenting cells contribute to defective T cell proliferation and function by the production of cytokines including TGF-β, IL-10 [57] or IL-6 [58]. In addition, neutrophils impair *Mtb*-induced DC maturation, thereby limiting their capacity to stimulate T cells [24].

Several pathogens could avoid neutrophil killing by limiting phagocytosis [59,60] and/or decreasing ROS production [61–64]. In line with this, *Mtb* could employ CD11b for entry, which do not induce respiratory burst [65]. Additionally, inhibition of ROS production also delays apoptosis in neutrophils and therefore, anti-apoptotic properties of a pathogen could be considered as a mechanism of immune evasion in which neutrophil survival is controlled by the bacilli allowing its persistence inside the host cell. This is coherent with the fact that attenuated strains of *Mtb* have been found to be much more effective than virulent strains at inducing apoptosis [66]. In this manner, it has been recently described that some *Mtb* strains fail to induce apoptosis due to an inability to stimulate ROS production and the induction of anti-apoptotic pathways [13]. This strategy has been also described for other pathogens like *Anaplasma phagocytophilum* which fails to trigger ROS and delays neutrophil apoptosis [67].

Other pathogen abilities rely on their capacity to secrete superoxide dismutase and catalase which are ROS antagonists [68], as well as sulphatides, LAM and phenolic-glycolipid I (PGL-I) which are potent oxygen radical scavengers [69]. In terms of host genetics, it has been proposed that there is a direct association between gene polymorphisms in patients with tuberculosis (i.e. polymorphisms in TLR2), and their associated mycobacterial isolates [70].


## 7. Conclusions

Tuberculosis remains the single largest infectious disease with ten million new TB cases and 2 million deaths which are estimated to occur yearly, more than any time in history [71]. The intracellular replication of the pathogen and its spread from the lungs to other sites occur before the development of adaptive immune responses, thus, innate immune mechanisms play a key role during the early stages of *Mtb* infection. Therefore, modulation of ROS production

**Funding:** This work was supported by grants from the Agencia Nacional de Promoción Científica y Tecnológica, ANPCyT (PICT 1497/12), Consejo Nacional de Investigaciones Científicas y Técnicas, CONICET (PIP112-200801-01476) and Fundación Alberto J Roemmers.

**Competing interests:** The author declare to have no competing interests.

**Ethical approval:** Not required.



would simultaneously control bacterial and neutrophil survival that may be beneficial to *Mtb* for establishing a niche where bacteria evade host immunity [46,72,73]. In contrast to other pathogens, *Mtb* lacks more classical virulence factors such as toxins and therefore, epidemiologic fitness of a strain can be influenced by the genetic background of host and pathogen, host/pathogen interactions, and the environment [74,75].

Virulence can be defined as the ability to cause active disease in humans. So, differences between genotypes may be slight but they may lead to remarkable phenotypic variation responsible for a differential modulation of host immune response impacting in the onset and progression of the disease. The variable incidence of epidemiologic and clinical outcomes of the disease could explain why some *Mtb* strains are more virulent than others and vary in their ability to cause disease in humans [76]. Thus, an evolutionary compensation which restores bacterial fitness may play a role in the epidemiological persistence of certain strains allowing them to acquire drug resistance-conferring mutations.

## Acknowledgments

I specially thank Dr Juan Ignacio Basile for helpful comments and critical reading of the manuscript.

A schematic representation of the molecules implicated in mycobacterial recognition by neutrophils. TLR2/dectin-1 cooperates to induce ROS production together with the rapid and sustained phosphorylation of p38 and Syk, inducing activation and apoptosis in neutrophil. *Mtb* could employ CD11b for entry, which do not induce respiratory burst. The MAPK can directly phosphorylate various transcription factors to initiate the expression of proinflammatory cytokine genes.

## References

- [1] Riedel DD, Kaufmann SHE. Chemokine secretion by human polymorphonuclear granulocytes after stimulation with *Mycobacterium tuberculosis* and lipoarabinomannan. *Infect Immun* 1997;65:4620–3.
- [2] Condros R, Rom WN, Liu YM, Schluger NW. Local immune responses correlate with presentation and outcome in tuberculosis. *Am J Respir Crit Care Med* 1998;157:729–35.
- [3] May ME, Spagnuolo PJ. Evidence for activation of a respiratory burst in the interaction of human neutrophils with *Mycobacterium tuberculosis*. *Infect Immun* 1987;55:2304–7.
- [4] Cook DN, Pisetsky DS, Schwartz DA. Toll-like receptors in the pathogenesis of human disease. *Nat Immunol* 2004;5:975–9.
- [5] DeLeo FR. Modulation of phagocyte apoptosis by bacterial pathogens. *Apoptosis: Int J Program Cell Death* 2004;9(4):399–413.
- [6] Cox G, Crossley J, Xing Z. Macrophage engulfment of apoptotic neutrophils contributes to the resolution of acute pulmonary inflammation in vivo. *Am J Respir Cell Mol Biol* 1995;12:232–7.
- [7] Fuchs TA, Abed U, Goosmann C, Hurwitz R, Schulze I, Wahn V, Weinrauch Y, Brinkmann V, Zychlinsky A. Novel cell death program leads to neutrophil extracellular traps. *J Cell Biol* 2007;176:231–41.

- [8] Ozaky T, Nakahira S, Tani K, Ogushi F, Yasuoka S, Ogura T. Differential cell analysis in bronchoalveolar lavage fluid from pulmonary lesions of patients with tuberculosis. *Chest* 1992;2:54–9.
- [9] Brown AE, Holzer TJ, Andersen BR. Capacity of human neutrophils to kill *Mycobacterium tuberculosis*. *J Infect Dis* 1987;156:985–9.
- [10] Alemán M, Beigier-Bompadre M, Borghetti C, de la Barrera S, Abbate E, Istariz M, Sasiain MC. Activation of peripheral blood neutrophils from patients with active advanced tuberculosis. *Clin Immunol* 2001;100:87–95.
- [11] Hirsch CS, Toosi Z, Johnson JL, Luzze H, Ntambi L, Peters P, McHugh M, Okwera A, Joloba M, Mugenyi P, Mugerwa RD, Terebuh P, Eline JJ. Augmentation of apoptosis and interferon- $\gamma$  production at sites of *Mycobacterium tuberculosis* infection in human tuberculosis. *J Infect Dis* 2001;183:779–88.
- [12] Alemán M, García A, Saab M, de la Barrera S, Finiazzi M, Abbate E, Sasiain MC. *Mycobacterium tuberculosis*-induced activation accelerates apoptosis in peripheral blood neutrophils from patients with active tuberculosis. *Am J Respir Cell Mol Biol* 2002;27:583–92.
- [13] Romero MM, Balboa L, Basile JI, López B, Ritacco V, de la Barrera SS, Sasiain MC, Barrera L, Alemán M. Clinical isolates of *Mycobacterium tuberculosis* differ in their ability to induce respiratory burst and apoptosis in neutrophils as a possible mechanism of immune escape. *Clin Dev Immunol* 2012;2012. <http://dx.doi.org/10.1155/2012/152546>.
- [14] Perskvist N, Long M, Stendahl O, Zheng L. *Mycobacterium tuberculosis* promotes apoptosis in human neutrophils by activating caspase-3 and altering expression of Bax/BclxL via an oxygen-dependent pathway. *J Immunol* 2002;168:6358–65.
- [15] Hoheisel G, Izicki G, Roth M, Chan CH, Leung JC, Reichenberger F, Schauer J, Perruchoud AP. Compartmentalization of proinflammatory cytokines in tuberculous pleurisy. *Respir Med* 1998;92:14–7.
- [16] Barnes PF, Fong SJ, Brennan PJ, Twomey PE, Mazumder A, Modlin RL. Local production of tumor necrosis factor and IFN- $\gamma$  in tuberculous pleuritis. *J Immunol* 1990;145:149–54.
- [17] Olobo JO, Geletu M, Eguale T, Hiwot K, Aderaye G, Britton S. Circulating TNF-a, TGF- $\beta$ , and IL-10 in tuberculous patients and healthy contacts. *Scand J Immunol* 2001;53:85–91.
- [18] Maeda J, Ueki N, Ohkawa T, Iwahashi N, Nakano T, Hada T, Higashino K. Local production and localization of transforming growth factor- $\beta$  in tuberculous pleurisy. *Clin Exp Immunol* 1993;92:32–8.
- [19] Song CH, Lee JS, Nam, Kim J-M, Suhr J-W, Jung S-S, Na M-J, Paik T-H, Kim H-J, Park J-K, Jo E-K. IL-18 production in human pulmonary and pleural tuberculosis. *Scand J Immunol* 2002;56:611–8.
- [20] Pace E, Gjormakaj M, Melis M, Profits M, Spatafora M, Vignola AM, Bonsignore G, Mody C. Interleukin-8 induces lymphocyte chemotaxis into the pleural space: role of pleural macrophages. *Am J Respir Crit Care Med* 1999;159:1592–9.
- [21] Xirouchaki N, Tzanakis N, Bourous D, Kyriakou D, Karkavitsas N, Alexandrakis M, Siafakas NM. Diagnostic value of interleukin-1 $\alpha$ , interleukin-6, and tumor necrosis factor in pleural effusions. *Chest* 2002;121:815–20.
- [22] Alemán M, de la Barrera S, Schierloh P, Alves L, Yokobori N, Baldini M, Abbate E and Sasiain MC: In tuberculous pleural effusions activated neutrophils undergo apoptosis and acquire dendritic cell-like phenotype. *J Infect Dis* 2005;192:399–409.
- [23] Iking-Konert C, Csekö C, Wagner C, Stegmaier S, Andrassy K, Hänsch GM. Transdifferentiation of polymorphonuclear neutrophils: acquisition of CD83 and other functional characteristics of dendritic cells. *J Mol Med* 2001;79:464–74.
- [24] Alemán M, de la Barrera S, Schierloh P, Yokobori N, Baldini M, Musella R, Abbate E, Sasiain M. Spontaneous or *Mycobacterium tuberculosis*-induced apoptotic neutrophils exert opposite effects on the dendritic cell-mediated immune response. *Eur J Immunol* 2007;37:1524–37.
- [25] Schaible UE, Winau F, Sieling PA, Fischer K, Collins HL, Brinkmann V, Kaufmann SHE. Apoptosis facilitates antigen presentation to T lymphocytes through MHC-I and CD1 in tuberculosis. *Nat Med* 2003;9:1039–46.
- [26] Medzhitov R. Toll-like receptors and innate immunity. *Nat Rev Immunol* 2001;1:135–45.
- [27] Kawai T, Akira S. Signaling to NF-( $\kappa$ B) by Toll-like receptors. *Trends Mol Med* 2007;13(11):460–9.
- [28] Kobayashi SD, Voyich JM, Braughton KR, DeLeo FR. Down-regulation of proinflammatory capacity during apoptosis in human polymorphonuclear leukocytes. *J Immunol* 2003;170:3357–68.
- [29] Brumell JH, Burkhardt AL, Bolen JB, Grinstein S. Endogenous reactive oxygen intermediates activate tyrosine kinases in human neutrophils. *J Biol Chem* 1996;271:1455–61.
- [30] Sabroe I, Jones EC, Usher LR, Whyte MKB, Dower SK. Toll-like receptor (TLR)2 and TLR4 in human peripheral blood granulocytes: a critical role for monocytes in leukocyte lipopolysaccharide responses. *J Immunol* 2002;168:4701–10.
- [31] Hayashi F, Means TK, Luster AD. Toll-like receptors stimulate human neutrophil function. *Blood* 2003;102:2660–9.
- [32] Jo EK, Yang CS, Choi CH, Harding CV. Intracellular signalling cascades regulating innate immune responses to Mycobacteria: branching out from Toll like receptors. *Cell Microbiol* 2007;9:1087–98.
- [33] Ward C, Chilvers ER, Lawson MF, Pryde JG, Fujihara S, Farrow SN, Haslett C, Rossi AG. NF- $\kappa$ B activation is a critical regulator of human granulocyte apoptosis in vitro. *J Biol Chem* 1999;274:4309–18.
- [34] Alvarado-Kristensson M, Porn-Ares MI, Grethe S, Smith D, Zheng L, Andersson. p38 mitogen-activated protein kinase and phosphatidylinositol 3-kinase activities have opposite effects on human neutrophil apoptosis. *FASEB J* 2002;16:129–31.
- [35] Yousefi S, Green DR, Blaser K, Simon HU. Protein-tyrosine phosphorylation regulates apoptosis in human eosinophils and neutrophils. *Proc Natl Acad Sci U S A* 1994;91:10868–72.
- [36] Alemán M, Schierloh P, de la Barrera S, Musella RM, Saab MA, Baldini M, Abbate E, Sasiain MC. *Mycobacterium tuberculosis* triggers apoptosis in peripheral neutrophils involving TLR2 and p38-mitogen protein kinase in tuberculosis patients. *Infect Immun* 2004;72:5150–8.
- [37] Triantaifilou M, Triantaifilou K. Lipopolysaccharide recognition: CD14, TLRs and the LPS-activation cluster. *Trends Immunol* 2002;23:301–4.
- [38] Naroeni A, Porte F. Role of cholesterol and ganglioside GM1 in entry and sort-term survival of *Brucella suis* in murine macrophages. *Infect Immun* 2002;70:1640–4.
- [39] Jutras I, Abrami L, Dautry-Varsat A. Entry of the *Lymphogranuloma Venereum* strain of *Chlamydia trachomatis* into host cells involves cholesterol-rich membrane domains. *Infect Immun* 2003;71(1):260–6.
- [40] Gatfield J, Pieters J. Essential role for cholesterol in entry of mycobacteria into macrophages. *Science* 2000;288(5471):1647–51.
- [41] Manes S, del Real G, Martinez AC. Pathogens: raft hijackers. *Nat Rev Immunol* 2003;3:557–68.
- [42] Gantner BN, Simmons RM, Canavera SJ, Canavera SJ, Akira S, Underhill DM. Collaborative induction of inflammatory responses by dectin-1 and Toll-like receptor 2. *J Exp Med* 2003;197:1107–17.
- [43] Shin D-M, Yang C-S, Yu J-M, Lee J-Y, Kim KH, Shin SJ, Takahara K, Lee SJ, Jo E-K. *Mycobacterium abscessus* activates the macrophage innate immune response via a physical and functional interaction between TLR2 and dectin-1. *Cell Microbiol* 2008;10(8):1608–21.
- [44] Yadav M, Schorey JS. The beta-glucan receptor dectin-1 functions together with TLR2 to mediate macrophage activation by mycobacteria. *Blood* 2006;108:3168–75.
- [45] Zhang X, Majlessi L, Deriaud E, Leclerc C, Lo-Man R. Coactivation of syk kinase and MyD88 adaptor protein pathways by bacteria promotes regulatory properties of neutrophils. *Immunity* 2009;31:761–71.
- [46] Romero MM, Basile JI, López B, Ritacco V, Barrera L, Sasiain M, Alemán M: Outbreaks of *Mycobacterium tuberculosis* MDR strains differentially induce neutrophil respiratory burst involving lipid rafts, p38 MAPK and Syk. *BMC Infect Dis* doi: 10.1186/1471-2334-14-262
- [47] Flesch I, Kaufmann SH. Mycobacterial growth inhibition by interferon-gamma-activated bone marrow macrophages and differential susceptibility among strains of *Mycobacterium tuberculosis*. *J Immunol* 1987;138:4408–13.
- [48] Yang C-S, Shin D-M, Lee H-M, Son JW, Lee SJ, Akira S, Gougerot-Pocidalo M-A, El-Benna J, Ichijo H, Jo E-K. ASK1-p38 MAPK-p47phox activation is essential for inflammatory responses during tuberculosis via TLR2-ROS signalling. *Cell Microbiol* 2008;10(3):741–54.
- [49] Janssen-Heininger YM, Poynter ME, Baeuerle PA. Recent advances towards understanding redox mechanisms in the activation of nuclear factor kappaB. *Free Radic Biol Med* 2000;28:1317–27.
- [50] Shaulian E, Karin M. AP-1 as a regulator of cell life and death. *Nat Cell Biol* 2002;4:E131–6.
- [51] Saitoh M, Nishitoh H, Fujii M, Takeda K, Tobiume K, Sawada Y, Kawabata M, Miyazono K, Ichijo H. Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1. *EMBO J* 1998;17:2596–606.
- [52] Yamashita K, Takahashi A, Kobayashi S, Hirata H, Mesner Jr PW, Kaufmann SHE, Yonehara S, Yamamoto K, Uchiyama T, Sasada M. Caspases mediate tumor necrosis factor- $\alpha$ -induced neutrophil apoptosis and down-regulation of reactive oxygen production. *Blood* 1999;93(2):674–85.
- [53] Moreira AL, Wang J, Tsanova-Berkova L, Hellmann W, Freedman VH, Kaplan G. Sequestration of *Mycobacterium tuberculosis* in tight vacuoles *in vivo* in lung macrophages of mice infected by the respiratory route. *Infect Immun* 1997;65:305–8.
- [54] Hmama Z, Gabathuler R, Jefferies WA, deJong G, Reiner NE. Attenuation of HLA-DR expression by mononuclear phagocytes infected with *Mycobacterium tuberculosis* is related to intracellular sequestration of immature class II heterodimers. *J Immunol* 1998;161:4882–93.
- [55] Wolf AJ, Linas B, Trevejo-Nunez GJ, Kincaid E, Tamura T, Takatsu K, Ernst JD. *Mycobacterium tuberculosis* infects dendritic cells with high frequency and impairs their function *in vivo*. *J Immunol* 2007;179(4):2509–19.
- [56] Balboa L, Romero MM, Yokobori N, Schierloh P, Geffner L, Basile JI, Musella RM, Abbate E, de la Barrera S, Sasiain MC, Alemán M. *Mycobacterium tuberculosis* impairs dendritic cell response by altering CD1b, DC-SIGN and MR profile. *Immunol Cell Biol* 2010;88:716–26.
- [57] Rojas RE, Balaji KN, Subramanian A, Boom WH. Regulation of human CD4+ ab T cell receptor positive (TCR+) and gd (TCR +T-cell responses to *Mycobacterium tuberculosis* by interleukin-10 and transforming growth factor b. *Infect Immun* 1999;67:6461–72.
- [58] van Heyningen TK, Collins HL, Russell DG. IL-6 produced by macrophages infected with *Mycobacterium* species suppresses T cell responses. *J Immunol* 1997;158:303–7.
- [59] Cornelius GR. Yersinia type III secretion: send in the effectors. *J Cell Biol* 2002;158:401–8.
- [60] Cunningham MW. Pathogenesis of group A streptococcal infections. *Clin Microbiol Rev* 2000;13:470–511.

- [61] Lei B, DeLeo FR, Hoe NP, Graham MR, Macie SM, Cole RL, Liu M, Hill HR, Low DE, Federle MJ, Scott JR, Musser JM. Evasion of human innate and acquired immunity by a bacterial homologue of CD11b that inhibits opsonophagocytosis. *Nat Med* 2001;7:1298–304.
- [62] Mott J, Rikihisa Y. Human granulocytic ehrlichiosis agent inhibits superoxide anion generation by human neutrophils. *Infect Immun* 2000;68:6697–703.
- [63] Mott J, Rikihisa Y, Tsunawaki S. Effect of *Anaplasma phagocitophila* on NADPH oxidase components in human neutrophils and HL-60 cells. *Infect Immun* 2002;70:7009–18.
- [64] Siemsen D, Kirpotina LN, Jutila MA, Quinn MT. Inhibition of the human neutrophil NADPH oxidase by *Coxelia burnetii*. *Microbes Infect* 2009;11:671–9.
- [65] Schlesinger LS, Bellinger-Kawahara CG, Payne NR, Horwitz MA. Phagocytosis of *Mycobacterium tuberculosis* is mediated by human monocyte complement receptors and complement component C3. *J Immunol* 1990;144(7):2771–80.
- [66] Keane J, Remold HG, Kornfeld H. Virulent *Mycobacterium tuberculosis* strains evade apoptosis of infected alveolar macrophages. *J Immunol* 2000;164(4):2016–20.
- [67] Björjesson DL, Kobayashi SD, Whitney AR, Voyich JM, Argue CM, DeLeo FR. Insights into pathogen immune evasion mechanisms: *Anaplasma phagocytophylum* fails to induce an apoptosis differentiation program in human neutrophils. *J Immunol* 2005;174:6364–72.
- [68] Andersen P, Askgaard D, Ljungqvist L, Bennedsen J, Heron I. Proteins released from *Mycobacterium tuberculosis* during growth. *Infect Immun* 1991;59:1905–10.
- [69] Chan J, Fujiwara T, Brennan P, McNeil M, Turco SJ, Sibley JC, Snapper M, Aisen P, Bloom B. Microbial glycolipids: possible virulence factors that scavenge oxygen radicals. *Proc Natl Acad Sci U.S.A* 1989;86:2453–7.
- [70] Caws M, Thwaites G, Dunstan S, Hawn TR, Lan NTN, Thuong NTT, Stepniewska K, et al. The influence of host and bacterial genotype on the Development of Disseminated Disease with *Mycobacterium tuberculosis*. *PLoS Pathog* 2008;4(3). <http://dx.doi.org/10.1371/journal.ppat.1000034>. e1000034.
- [71] WHO. Global tuberculosis control-surveillance, planning, financing. Geneva, Switzerland: WHO; 2010 (Contract No: WHO/HTM/TB/2008.393).
- [72] Chackerian A, Alt JM, Perera TV, Dascher CC, Behar SM. Dissemination of *Mycobacterium tuberculosis* is influenced by host factors and precedes the initiation of T-cell immunity. *Infect Immun* 2002;70(8):4501–9.
- [73] Hingley-Wilson SM, Sambandamurthy VK, Jacobs WR. Survival perspectives from the world's most successful pathogen, *Mycobacterium tuberculosis*. *Nat Immunol* 2003;4(10):949–55.
- [74] Luciani F, Sisson SA, Jiang H, Francis AR, Tanaka MM. The epidemiological fitness cost of drug resistance in *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U.S.A* 2009;106:14711–5.
- [75] O'Sullivan DM, McHugh TD, Gillespie SH. Mapping the fitness of *Mycobacterium tuberculosis* strains: a complex picture. *J Med Microbiol* 2010;59:1533–5.
- [76] Malik NJ, Godfrey-Faussett P. Effects of genetic variability of *Mycobacterium tuberculosis* strains on the presentation of disease. *Lancet Infect Dis* 2005;5(3):174–83.