



## Tick-borne Rickettsiales: Molecular tools for the study of an emergent group of pathogens



Eliana C. Guillemi <sup>a,\*</sup>, Laura Tomassone <sup>b</sup>, Marisa D. Farber <sup>a</sup>

<sup>a</sup> Inst. de Biotecnología, INTA Castelar. Los Reseros y N. Repetto, 1686 Hurlingham, Buenos Aires, Argentina

<sup>b</sup> Dipartimento di Scienze Veterinarie, Università degli Studi di Torino, Largo Braccini 2, 10095 Grugliasco, Torino, Italy

### ARTICLE INFO

#### Article history:

Received 5 February 2015

Received in revised form 9 October 2015

Accepted 9 October 2015

Available online xxxx

#### Keywords:

*Rickettsia*

*Ehrlichia*

*Anaplasma*

Blood-meal analysis

### ABSTRACT

The use of molecular techniques in recent years has enhanced the sensitivity and specificity of the diagnosis of Rickettsiales, a bacterial order which includes significant emerging and re-emerging pathogens of humans and animals. Molecular detection enables the accurate identification at the species level, providing additional information on the epidemiology and course of the clinical cases. Moreover, PCR and enzyme restriction analysis of the vector blood meal can be employed to study the tick feeding source and possibly identify pathogen's reservoir. Here, we review the molecular tools available for the identification and characterization of tick-borne bacteria from the genera *Rickettsia*, *Ehrlichia* and *Anaplasma* and for the study of ticks feeding behavior. We summarize the significant criteria for taxonomic identification of Rickettsiales species and propose a procedure algorithm for the classification of bacterial isolates as members of this order.

© 2015 Elsevier B.V. All rights reserved.

### Contents

1. Introduction . . . . .	87
2. Tick-borne disease diagnosis based on sample type . . . . .	88
3. Anaplasmataceae family members: identification by means of available molecular tools . . . . .	88
3.1. End point PCR . . . . .	88
3.2. Real-time PCR . . . . .	90
3.3. Tools for studying intraespecific variation at the population level . . . . .	90
4. Rickettsiaceae family members: identification by means of available molecular tools . . . . .	90
4.1. End point PCR . . . . .	90
4.2. Real-time PCR . . . . .	91
5. Middle and high-throughput tools for the study of Rickettsiales . . . . .	91
6. Suggested algorithms for taxonomic characterization and identification of novel species . . . . .	93
7. Vertebrate reservoir host identification through blood meal analysis . . . . .	94
8. Conclusions . . . . .	94
Acknowledgments . . . . .	95
References . . . . .	95

### 1. Introduction

The order Rickettsiales is composed of obligate intracellular Gram-negative bacteria infecting eukaryotic cells (Kang et al., 2014). It consists

of two families, namely Rickettsiaceae and Anaplasmataceae. The Rickettsiaceae family is integrated by bacteria from the genera *Rickettsia* and *Orientia*, that occupy an intracytoplasmic compartment in the host cell, while the Anaplasmataceae family is composed of the genera *Ehrlichia*, *Anaplasma*, *Neorickettsia* and *Wolbachia* which occupy an intravacuolar compartment within infected host cells (Dumler et al., 2001). Importantly, *Ehrlichia* spp., *Anaplasma* spp. and several *Rickettsia* spp., are tick-borne pathogens.

\* Corresponding author.

E-mail address: [guillemi.eliana@inta.gob.ar](mailto:guillemi.eliana@inta.gob.ar) (E.C. Guillemi).

*Rickettsia* is the most significant genus within the family Rickettsiaceae. A modern classification based on whole-genome analysis data (Fuxelius et al., 2007; Gillespie et al., 2007, 2008), divides the genus in four groups: the ancestral group (AG), the typhus group (TG), the spotted fever group (SFG), and the transitional group (TRG). AG comprises *R. bellii* and *R. canadensis*, which have no records of pathogenicity. TG consists of *R. typhi* and *R. prowazekii*. The SFG is integrated by several species, including *R. rickettsii*, *R. parkeri*, *R. conorii*, *R. massilliae*. Finally, TRG comprises *R. akari*, *R. australis* and *R. felis*. A large number of these organisms are transmitted by hard ticks (*Ixodidae* family) and are associated with emerging life-threatening human zoonoses (Anderson et al., 1992; Buller et al., 1999). Rickettsiae from the TG are not tick-borne, but are transmitted by infected flea, lice and mites (Azad, 1988; Durden and Traub, 2002; Gross, 1996).

The genus *Ehrlichia* belongs to the Anaplasmataceae family and comprises six species (*E. canis*, *E. chaffeensis*, *E. muris*, *E. ruminantium*, *E. ewingii* and *E. ovis*) from which *E. chaffeensis* and *E. ewingii* are recognized as human zoonotic pathogens (Breitschwerdt et al., 1998; Goldman et al., 1998). Recent reports suggest that some other *Ehrlichia* species might be human pathogens, such as *E. canis*, the etiologic agent of canine monocytic ehrlichiosis (Perez et al., 2006), and organisms closely related to *E. muris*, a rodent and deer associated ehrlichia (Nefedova et al., 2008; Pritt et al., 2011). Finally, *E. ruminantium*-like organisms have been implicated in human infections in South Africa (Louw et al., 2005) and in the USA, where the Panola Mountain Ehrlichia (PME) is reported (Reeves et al., 2008).

The genus *Anaplasma* (Anaplasmataceae family) currently comprises six species: *A. phagocytophilum*, *A. platys*, *A. marginale*, *A. centrale*, *A. bovis* and *A. ovis* (Dumler et al., 2001). Among these species, *A. phagocytophilum* causes febrile disease in humans and animals and *A. platys* is considered as a low risk pathogen for humans (Otranto et al., 2009), while *A. marginale* is a highly prevalent pathogen of livestock worldwide, causing important losses to production.

Due to the public health implications of bacteria from the order Rickettsiales, a proper identification protocol is needed. In this sense, in recent years the number of reports of rickettsiosis and ehrlichiosis has increased (Dumler, 2010; Nicholson et al., 2010). Remarkably, several studies in South America revealed the occurrence in new areas (Abarca et al., 2012; André et al., 2010; Cicuttin and Nava, 2013; García-García et al., 2010; Martínez et al., 2008; Miranda et al., 2011, 2012; Romer et al., 2014; Tomassone et al., 2008).

Diagnosis using serology faced the difficulty that a detectable level of antibodies against *Rickettsia* sp. does not appear in blood until the day 7 or 10 after the onset of the disease (Leitner et al., 2002). In the case of *Ehrlichia* genus, attempts to early identify bacteria by serologic methods, even applying the most widely used techniques, tends to fail in 56–67% of the cases (Childs et al., 1999). For this reason, the diagnosis in the acute phase must rely on the detection of the bacteria in the clinical specimens. Furthermore, direct observation of the *Ehrlichia* morula in stained blood smears also tends to result false negative (22%) (Childs et al., 1999). For *Anaplasma* species, direct observation requires a trained operator and it is useful only for early stages of infection. Similarly, *Rickettsia* species circulate in blood in low numbers even during the acute phase of the disease (Breitschwerdt et al., 1990; Parola et al., 2005). In this sense, the molecular methods become important for diagnosis because, in addition to improving sensitivity, they may give additional information regarding the epidemiology and course of the clinical case.

Up to now, several articles have reviewed the diagnostic methods for the identification of bacteria from the order Rickettsiales (Allison and Little, 2013; Fenollar and Raoult, 2004; Sparagano et al., 1999). The nowadays popularization of PCR together with the affordable molecular biology consumables due to a wider offer of alternative kits, give rise to an overwhelming number of diagnostic test reports. In this regard, here we propose an integral revision of the available molecular tools for tick-borne Rickettsiales so as to introduce the need to define a core of

reliable techniques allowing results comparison among laboratories. Furthermore, we also include a section referring to identification of novel species and the challenge of building up a consensus for taxonomy validation. Finally, we introduce the approaches for the study of ticks' feeding behavior.

## 2. Tick-borne disease diagnosis based on sample type

When studying tick-borne diseases, we can analyze a diversity of samples. These can be ticks, sampled from the host (feeding ticks) or collected in the environment (questing ticks), or host tissues (blood or other tissues). Samples of various origins can be a source of different information, and contribute to the knowledge of epidemiological aspects regarding the pathogens and their vectors. In example, the detection of the *Rickettsia* spp. DNA in the host blood can generally be achieved only during rickettsemia and is thus an evidence of recent infection (Horta et al., 2007). The finding of a rickettsial agent in a tick can be linked to its reservoir/vector/amplifier host role, or simply to the blood meal that has been taken on an infected vertebrate host. Each result has then to be carefully evaluated and interpreted, according to the known epidemiology of the studied disease.

Different molecular analyses can be performed to detect multiple pathogens in the extracted nucleic acid samples. The detection of coinfections is of crucial medical concern because interactions among pathogens can alter host susceptibility, infection length and clinical symptoms, increasing the disease severity. From an epidemiological point of view, coinfections may modify the infection risk (Andersson et al., 2013; Lommano et al., 2012; Vaumourin et al., 2013). Finally, the extracted DNA can be submitted to a 'blood meal analysis', in order to study the tick feeding behavior and gather information on potential reservoirs of tick-borne agents.

## 3. Anaplasmataceae family members: identification by means of available molecular tools

### 3.1. End point PCR

Many PCR protocols targeting the 16S rRNA gene were developed for the identification of the genera *Ehrlichia*, *Anaplasma* and their species. In Table 1, 16S rRNA gene based protocols for the identification if these genera are summarized. Whatever the protocol used, once the genus is detected (Dawson et al., 1994; Eddlestone et al., 2007a; Munderloh et al., 1996; Parola et al., 2000; Wen et al., 2002), subsequent PCRs are required to identify the species of major medical importance. In most cases, the fragments obtained for genus identification can be employed as template for a second step reaction. This is the case of the primers designed for *E. chaffeensis* (Anderson et al., 1992), *E. canis* (Murphy et al., 1998; Wen et al., 1997) and *E. ewingii* (Murphy et al., 1998), that can be used in a second round of amplification employing as template the 479 bp product generated with ECC and ECB oligonucleotides (Dawson et al., 1994). Another nested protocol was proposed by Inokuma et al. (2001a), who designed *Ehrlichia* sp. specific primers and additional *E. muris* and *A. platys* oligonucleotides for a specific second step amplification. For *A. platys* detection, a single step PCR was also proposed by Inokuma et al. (2000).

Finally, Teshale et al. (2015) developed a semi-nested PCR protocol to simultaneously detect the genera *Ehrlichia* and *Anaplasma*, followed by fragment length polymorphism (RFLP) that enables speciation. However, RFLPs protocols are acknowledged as being time-consuming, and exact genotyping cannot always be achieved (Rasmussen, 2012).

In addition to 16S rRNA, other target genes were used for both *Ehrlichia* and *Anaplasma* identification. Dumler et al. (2001) reorganized the genera in Anaplasmataceae family based on nucleotide sequences of 16S rRNA and *groEl* genes. Shortly after, Inokuma et al. (2001b) reinforced this new organization employing a well supported *gltA*-based phylogenetic tree. This last gene has been previously validated for

**Table 1**

Primers for Anaplasmataceae family identification targeting 16S rRNA gene.

Organism	Primers	Sequence	Amplicon size	Reference
<i>Anaplasma</i> and <i>Ehrlichia</i>	GEPs	5'-CTGGCCGCAAGCYTAACACATGCAAGTCGAACCGA-3'	432 bp	Eddlestone et al. (2007a)
	GEpas	5'-CTTCTTCTRGGTACCGCTATTATCTTCCYATG-3'		
<i>Ehrlichia</i> sp.	EHR16SD	5'-GGTACCYACAGAAGAAGTCC-3'	345 bp	Parola et al. (2000)
	EHR16SR	5'-TAGCACTCATCGTTACAGC-3'		
<i>Ehrlichia</i> sp.	PER1	5'-TTTATCGCTTATTAGATGAGCCTATG-3'	451 bp	Munderloh et al. (1996)
	PER2	5'-CTCTACACTAGGAATTCCGCT-3'		
<i>Ehrlichia</i> sp.	Eh-out1	5'-TTGAGAGTTTGATCTGGCTCAGAACG-3'	650 bp	Wen et al. (2002)
	Eh-out2	5'-CACCTCTACACTAGGAATTCCGCTATC-3'		
	Eh-gs1	5'-GTAATAACTGTATAATCCCTG-3'	280 bp	
	Eh-gs2	5'-GTACCGTCAATTATCTTCCCTA-3'		
<i>Ehrlichia</i> sp.	ECC	5'-AGAACGAAACGCTGGCGCAAGC-3'	479 bp	Dawson et al. (1994)
	ECB	5'-CGTATTACCGCGGCTGCTGCCA-3		
<i>E. chaffeensis</i>	HE1	5'-CAATTGCTTATAACCTTTGGTTAAAAT-3'	389 bp	Anderson et al. (1992)
	HE3	5'-TATAGGTACCGTCATTATCTTCCCTAT-3'		
<i>E. ewingii</i>	EE52	5'-CGAACAACTCTAAATAGTCTCTGAC-3'	394 bp	Murphy et al. (1998)
	HE3	5'-TATAGGTACCGTCATTATCTTCCCTAT-3'		
<i>E. canis</i>	ECAN	5'-CAATTATTTAGCCTCTGGCTATAGGA-3'	389 bp	
	HE3	5'-TATAGGTACCGTCATTATCTTCCCTAT-3'		
<i>E. canis</i>	ECA	5'-AACACATCGAACGCAACCGA-3'	410 bp	Wen et al. (1997)
	HE3	5'-TATAGGTACCGTCATTATCTTCCCTAT-3'		
<i>Ehrlichia</i> sp.	fD1	5'-AGAGTTGATCTGGCTCAG-3'	760 bp	Inokuma et al. (2001a)
	EHR16SR	5'-TAGCACTCATCGTTACAGC-3'		
<i>E. muris</i>	MURIS	5'-GATAGCTACCCATAGCTTTTTAGC-3'	414 bp	
	GA1UR	5'-GAGTTGCGGGACTCTCT-3'		
<i>A. platys</i>	PLATYS-F	5'-AAGTCGAACGGATTGTC-3'	506 bp	
	PLATYS-R	5'-CTTAACTTACCGAAC-3'		
<i>A. platys</i>	PLATYS	5'-GATTTTGTCTAGCTTGTATG-3'	678 bp	Inokuma et al. (2000)
	EHR16SR	5'-TAGCACTCATCGTTACAGC-3'		

identification and phylogenetic analysis of *Rickettsia* (Roux et al., 1997) and *Bartonella* species (Birtles and Raoult, 1996; Joblet et al., 1995), since *gltA* allows a better discrimination among closely related species due to its higher polymorphisms in comparison to 16S rRNA genes. In Table 2, PCR protocols based on *gltA* and *groEl* genes are summarized.

When searching for a defined species in the Anaplasmataceae family, a broader range of PCR protocols are available, targeting many different gene fragments. For studying *E. canis*, Stich et al. (2002) developed a nested PCR targeting the p30 open reading frame (ORF) resulting in a specific reaction that amplifies an exclusive fragment from *E. canis*.

that is even different from its orthologous *p28* from *E. chaffeensis*. The sensitivity of both the first and the second step were higher than for the nested PCR based on the 16S rRNA gene (100 and 1000 times respectively). However, only by applying the two steps it was possible to identify *E. canis* carriers in a canine survey (Stich et al., 2002). For studying *E. chaffeensis*, Wagner et al. (2004) amplified a fragment from the *p28* ORF, which is the most studied transmembrane protein family in *E. chaffeensis* (Crocquet-Valdes et al., 2011; Ohashi et al., 2001). The one-step reaction developed on P28 resulted 1000 times more sensitive than the nested protocol amplifying a 389-pb fragment of the 16S rRNA gene.

**Table 2**Primers based on *gltA* and *groEl* genes for Anaplasmataceae family identification.

Target Gene	Organism	Primers	Sequence	Amplicon size	Reference
<i>gltA</i>	<i>Anaplasma</i> and <i>Ehrlichia</i>	F1b EHR-CS779R	5'-GATCATGARCARAATGCTTC-3' 5'-GCNCCMCCATGMGCTC-3'	119 bp	Inokuma et al. (2002)
	<i>E. chaffeensis</i> ,	F4b	5'-CCGGTTTTATGCTACTGC-3'	776 bp	Inokuma et al. (2001b)
	<i>E. muris</i> ,	R1b	5'-CGATGACCAAACCCAT-3'		
	<i>E. ruminantium</i> , <i>A. phagocytophilum</i> , <i>A. marginale</i>				
	<i>E. chaffeensis</i> ,	EHR-CS136F	5'-TYATGTCYACTGCTGCKTG-3	650 bp	Inokuma et al. (2001b)
	<i>E. canis</i> , <i>E. muris</i> ,	EHR-CS778R	5'-GCNCCMCCATGMGCTGG-3		
	<i>E. ruminantium</i> ,				
	<i>A. phagocytophilum</i> , <i>A. marginale</i> ,				
	<i>A. centrale</i>				
	<i>A. platys</i>	PLA-CSM136F PLA-CS1359R	5'-TTGAAAAAGTAAGCGGAGC-3' 5'-AACACAGGTTATGACAC-3'	1495 bp	Inokuma et al. (2002)
		groEL-643 s groEL-1236 as	5'-ACT GAT GGT ATG CAR TTT GAY CG-3' 5'-TCT TTR CGT TCY TTG ACY TCA ACTTC-3'	600 bp	Barber et al. (2010)
	<i>E. canis</i>	gro-E-canis163s gro-E-canis573as	5'-AAA TGT AGT TGT AAC GGG TGA ACA G-3' 5'-AGA TAA TAC CTC ACG CTT CAT AGA CA-3'	410 bp	Otranto et al. (2010)
	<i>A. platys</i>	PLA-HS475F PLAT-HS1198R	5'-AAGCGAAAGAACGAGCTTCA-3' 5'-CATAGCTGAAGTCAGGAC-3'	724 bp	Inokuma et al. (2002)
	<i>A. platys</i>	GroAplats-35 s GroAplats-550as	5'-AGCGTAGTCGATTCTCAGTTT-3' 5'-TCGCCGTTAGCAGAGATGGTAG-3'	516 bp	Beall et al. (2008)
	<i>A. phagocytophilum</i> ,	EphplgroEL(569)F	5'-ATGGTATGCACTTGTGATCCG-3'	624 bp	Alberti et al. (2005)
	<i>A. platys</i>	EphplgroEL(1193)R	5'-TCTACTCTGCTTGTGCGTC-3'		
	<i>A. phagocytophilum</i>	EphplgroEL(569)F	5'-ATGGTATGCACTTGTGATCCG-3'	574 bp	
		EphgroEL(1142)R	5'-TTGAGTACAGAACACCCGGAA-3'		
	<i>A. platys</i>	EphplgroEL(569)F EplgroEL(1084)R	5'-ATGGTATGCACTTGTGATCCG-3' 5'-CATAGCTGAAGTGGAGGAC-3'	516 bp	

For the molecular detection of *E. ruminantium*, diagnostic tests targeting *pCS20* sequences have extensively been used (Faburay et al., 2007; Peter et al., 1995). In the case of *A. marginale*, *msp5* (major surface protein 5 gene) is the target locus of choice for diagnosis purposes, given that detection of persistently infected cattle is important to control the movement of infected cattle into and from disease-free regions (Torioni de Echaide et al., 1998; OIE Terrestrial Manual, 2015). The *ankA* gene is a useful target for *A. phagocytophilum* detection (Beall et al., 2008). It is an exclusive gene for the *Anaplasma* genus and is also considered one of the major components involved in the pathogenesis of *A. phagocytophilum* (Rikihisa, 2010, 2011).

### 3.2. Real-time PCR

In recent years, a wide range of real-time protocols (RT-PCR) for the identification of Anaplasmataceae species have been developed. They have the advantage of not requiring amplicon sequencing to confirm the positive result and allow the analysis of large sample numbers. Furthermore, they enable quantification of DNA molecules and the reduction of false positive results, since the reaction tubes can remain closed for the interpretation of the results, thus decreasing the possibility of contamination risk.

Available RT-PCR protocols can employ TaqMan or SYBR Green technologies and are focused on diverse genes such as *sodB* (Fe superoxide dismutase), for detection of at least 5 *Ehrlichia* spp. copies per reaction (Quroollo et al., 2014); *dsb* (disulfide bond formation protein gene) for *E. chaffeensis*, *E. ewingii* and *E. canis* (Doyle et al., 2005); *homP* (hypothetical outer membrane protein gene) (Stoffel et al., 2014) and *msp2* (Courtney et al., 2004; Scorpio et al., 2004) for *E. chaffeensis*, *msp1b* for *A. marginale* (Carelli et al., 2007) and TRP120 for *E. chaffeensis*, which encodes an important immunodominant antigen of this pathogen (Yu et al., 1996; Paddock and Childs, 2003).

In particular, the reaction designed by Bell and Patel (2005) is able to discriminate *A. phagocytophilum*, *E. chaffeensis* and *E. ewingii*, based on nucleotide differences found in the amplicon through a melting curve. This assay shows a considerably higher sensitivity and specificity when compared to standard PCR protocols (Anderson et al., 1992; Childs et al., 1999; Chu, 1998), and allows the simultaneous identification of the three pathogens. The protocol proposed by Sirigireddy et al. in 2005, is a multiplex real-time PCR for the identification of up to five different zoonotic risk bacteria species from the Anaplasmataceae family (*E. chaffeensis*, *E. canis*, *E. ewingii*, *A. phagocytophilum* and *A. platys*). This molecular tool targets the 16S rRNA gene using oligonucleotides specific for the genus *Ehrlichia* and *Anaplasma*, and implements species-specific TaqMan probes. This assay has an analytical sensitivity of 100 copies of each pathogen and is able to identify coinfections, when the concentration difference between organisms is more than 100 times. In the year 2007, Carelli et al. developed a RT-PCR specific for *A. marginale* which was later improved by Decaro et al. (2008) for the simultaneous detection and quantification of *A. marginale* and *A. centrale* in a duplex reaction, for pathogenesis studies on bovine acute anaplasmosis. Also, the three main taxonomic loci, *groEL*, 16S rRNA and *gltA*, were employed for RT-PCR protocols in the Anaplasmataceae family. Their most important characteristics are summarized in Table 3.

### 3.3. Tools for studying intraespecific variation at the population level

In order to improve the discrimination power between *E. chaffeensis* populations, Sumner et al. (1999) developed the intraspecific molecular marker VLPT (Variable-length PCR target). This methodology consists of a nested PCR targeting an ORF encoding a 44-KD polypeptide with four tandem repeats of 90 bp each. In this region, three to six repetitive units can be detected, so that the *E. chaffeensis* populations can be characterized based on the variable length of the obtained amplicons.

Multilocus Sequence Typing (MLST) is a widely accepted method for molecular characterization of bacteria. It enables the genotypic characterization of isolates and the study of the global dispersion of some new variants of pathogens (Mayer et al., 2002). In addition, the data obtained by MLST strategy apply to evolutionary and population studies (Jolley et al., 2000). A MLST scheme has been developed for *E. ruminantium* (Adakal et al., 2009), *A. phagocytophilum* (Huhn et al., 2014) and *A. marginale* (Guillemi et al., 2015). This methodology consists of amplifying and sequencing a set of single copy genes and the further identification of single nucleotides polymorphisms (SNPs) between genotypes.

In Fig. 1 we summarize as an algorithm the available molecular protocols for the study of this group of pathogens, together with Rickettsiaceae family (see below).

## 4. Rickettsiaceae family members: identification by means of available molecular tools

### 4.1. End point PCR

The serological typing and identification remained the reference diagnostic method for Rickettsiaceae (Philip et al., 1978) until Roux et al. (1997) analyzed and compared citrate synthase (*gltA*) sequences from different *Rickettsia* species. They also compared *gltA* with 16S rDNA sequences and found that the average rate of sequence change in *gltA* was faster than in 16S rRNA, which indicates that *gltA* was less conserved in rickettsiae. This result would suggest that for discrimination within the *Rickettsia* genus, *gltA* sequences may be quite valuable. In the same publication, the authors also suggested two other molecular candidates: genes encoding rOmpA and rOmpB proteins. These molecules are specific for the SFG rickettsiae and the members of the *Rickettsia* genus respectively, and have been shown to exhibit high degrees of difference between species employing PCR-RFLP (Eremeeva et al., 1994; Regnery et al., 1991; Roux et al., 1996) or sequencing (Roux et al., 1996) than other molecules (*gltA*, 17-kDa protein, 16S rRNA). Accordingly, Fournier et al. (1998) amplified the whole *ompA* gene (excluding the central region containing the tandem repeat units) of 21 SFG rickettsiae isolates and confirmed that the intra-SFG variability in *ompA* was markedly higher than that of 16S rRNA or *gltA* genes. Later, by amplifying and sequencing the *ompB* gene of 24 representatives of the *Rickettsia* genus, Roux and Raoult (2000) observed that the study of this protein, which has common epitopes to both SFG and TG rickettsiae, required to be analyzed in combination with *gltA* and *ompA* to obtain a reliable phylogeny for all the bacteria in the *Rickettsia* genus.

Since these genes resulted to have a high discrimination power, many PCR protocols have been described using them as targets (Table 4). Even, Regnery et al. (1991) developed an identification and species discrimination system, based on the amplification of two fragments from the *gltA* gene, followed by an endonuclease restriction of the second amplicon. Through this PCR scheme followed by RFLP, *Rickettsia* species and genotypes can be differentiated. Kidd et al. (2008) developed a PCR with primers targeting a small region of the *ompA* gene for the identification of SFG rickettsiae. The sensitivity of this reaction was improved (up to 5 DNA copies) when the primers were adapted to a real-time PCR (see below). This PCR protocol appeared to be more sensitive than the assay amplifying the largest portion of the *ompA* gene described by Roux et al. (1996), and has enough variability in this smaller region to distinguish species based on sequencing. A protocol targeting the 17-kD antigen gene, that implies a nested PCR followed by RFLP analysis, was designed for *R. conorii*. Although this technique is not specific for this species (since the nested PCR also amplifies DNA from *R. australis*, *R. rickettsii*, *R. akari*, *R. typhi*, and *R. prowazekii*), it enables differentiation of SFG from TG rickettsiae. For this purpose, a RFLP test based on the digestion of the 214-bp PCR product with the restriction enzyme *Bf1* is necessary. The SFG rickettsiae amplicon contains one *Bf1* site, resulting in two fragments, while the TG rickettsiae

**Table 3**

RT-PCR protocols for Anaplasmataceae family identification and quantification.

Target gene	Organism	Primers/probes	Discriminatory system	Quantification	Reference
16S rRNA	<i>E. chaffeensis</i>	sense 5'-AACACATGCAAGTCGAACGG-3'antisense 5'-CCCCGCAGGGATTATAACA-3'	SYBR Green	Yes	Li et al. (2002)
	<i>E. chaffeensis</i>	ECH16S-17 5'-CGCGGAACGCTAACACATG-3'	TaqMan probes	Yes	Loftis et al. (2003)
		ECH16S-97 5'-CCCGTCTGCCACTAACAAATTATT-3'			
		ECH16S-38 5'-6-carboxyfluorescein-AGTCGAACGGACAATTGCTTAAACCTTTGGT-3'			
	<i>A. platys</i>	EP-963F: 5'-CGCAGTTCGCGTGGATCT-3'	TaqMan probes	Yes	Eddlestone et al. (2007b)
		EP-1029R: 5'-CCCAACATCTCACGACAGCA-3'			
		EP16Sprobe: 5'-FAM-TGACGACAGCCATGCGACGACCTG-TAMRA-3'			
	<i>A. phagocytophilum</i>	Ehrl/Anapl_F 5'-CTCAGAACGAAACGGCTTG-3'	TaqMan probes	No	Sirigireddy and Ganta (2005)
	<i>A. platys</i>	Ehrl_R 5'-CATTTCTAATGCCATTCC-3'			
	<i>E. chaffeensis</i>	Anapl_R 5'-CATTTCTAGTGGCTATCCC-3'			
	<i>E. ewingii</i>	Ehrl/Anapl_R 5'-GTATTACCGCGGCTGGCAC-3'			
	<i>E. canis</i>	E. ewi_long.F 5'CTCAGAACGAAACGCTGGCGCAAGCTAACACATGCGAACGA			
		ACAATTCTAAATAGTCTGACTATTAGATAGTTAGTGGCACAC-3'			
		A. pla_long.F 5'CTCAGAACGAAACGCTGGCGCAAGCTAACACATGCGAACGA			
		CGAACCGATTTTGTCTGAGCTTGCTATGATAAAAATTAGTGGCAGAC-3'			
		Ehrl/Anapl_capture 5'-Biotin-CCCCCCCCCCCCGACTCTTCTRGGTACCGTC-3'			
		E. chaf 5'-TET/CTTATAACCTTTGGTATAAATAATTGTTAG/TAMRA-3'			
		E. can 5'-FAM/TATAGCCTCGGCTATAGGAAATTGTTAG/TAMRA-3'			
		E. ewi 5'-ROX/CTAAATAGTCTGACTATTAGATAGTTAG/BQH2-3'			
		A. pla 5'-FAM/CGGATTITGTCTGAGCTTGCTATGAT/DABCYL-3'			
		A. pha 5'-TET/TTGCTATAAGAATAATTAGTGGCAGACG/DABCYL-3'			
		Ehrl/Anaplcommon 5'-ROX/TAACACATGCGAACGGA/BQH2-3'			
gltA	<i>A. phagocytophilum</i>	ApF: 5'-TTTGGCCGCTGAATACAGT-3'	TaqMan probes	No	Henningsson et al. (2015)
		ApR: 5'-TCTCGAGGGAATGATCTAACAGT-3'			
		ApM: 5'- FAM -TGCCTGAACAAGTTATG-3'			
	<i>A. ovis</i>	sense primer: 5'-AGGTACCGGGTATCGCTGTGA-3'	TaqMan probes	Yes	Chi et al. (2015)
		anti-sense primer: 5'-AGGTTGGATCTGCCCTGTGA-3'			
		probe: 5'-(FAM)ACATTACAGGCACACCTCTGGCATGC(BHQ1)-3'			
groEL	<i>A. centrale</i>	AC-For 5'-CTATACACGGCTTGATCTC-3'	TaqMan probes	Yes	Decaro et al. (2008)
		AC-Rev: 5'-CGCTTTATGATGTTGATGC-3'			
		AC-Pb 5'-TexasRed-ATCATCATCTTCCCTTACCTCGTBHQ2-3'			
	<i>A. phagocytophilum</i>	ESP-F 5'-TACTCAGAGTGTCTCAATGT-3'	Melting curve	No	Bell and Patel (2005)
	<i>E. chaffeensis</i>	ESP-R 5'-GCATACCATCAGTTTCAAC-3'			
	<i>E. ewingii</i>	Ec/e-FL 5'-ATTTCAGCTAATGGAGATAAGAATATAG- FITC-3'			
		Aph-FL 5'-GTCAGCAATGGAGACAAGACATAGGA- FITC-3'			
		ESP-RD 5'-Red640-GTAAGATTGCACAGTGTGTTCAAGAACGTA-Phosphate-3'			

amplicon contains two sites, which generate three fragments (Leitner et al., 2002).

Fournier and Raoult (2004) developed a suicide PCR for the detection of both TG and SFG rickettsiae from skin biopsy specimens. A suicide PCR consists of single-use primers; in this case oligonucleotides were selected from conserved regions flanking 300 to 600-bp variable fragments of genes present in both *R. conori* and *R. prowazekii* genomes (targeting, each year of the study, a new gene that had never been amplified before in the laboratory). The positive PCR products were sequenced in both directions. This technique, based on single-use primers targeting single-use DNA fragments, presents a specificity of 100% and a sensitivity of 68%, but is 1.5 times more sensitive than the conventional PCR by Regnery et al. (1991). The authors suggest that suicide PCR should be reserved for use on patients suspected of having a rickettsiosis but with negative regular PCR results.

For species-specific identification, few PCR protocols are available (Raoult et al., 2001; Tzianabos et al., 1989). In an attempt to increase the sensitivity and avoid false positive amplifications, Raoult et al. (2001) developed a suicide PCR. This scheme targets the *ompA* gene from *R. africae* and is based on three sets of two pairs of single-use primers that amplify non-overlapping fragments in a nested protocol. The main disadvantage of the suicide PCR is that once a pair of primers is used in a laboratory, it cannot be re-used.

#### 4.2. Real-time PCR

Because of the intracellular habitat of Rickettsiae and their slow generation time, the use of traditional techniques to quantify the number of viable Rickettsiae in a sample is laborious, inaccurate, and tedious (Eremeeva et al., 2003). On the other hand, RT-PCR has significant


advantages over end-point PCR methods. For instance, RT-PCR requires shorter assay times; it does not require gel electrophoresis and has the capacity of analyzing large numbers of samples with fully automated equipment. Moreover, RT-PCR can be used as a quantitative method and for the identification of *Rickettsia* species on samples contaminated with other microorganisms.

As it was mentioned above for Anaplasmataceae, RT-PCR protocols can employ TaqMan or SYBR Green chemistries and are focused on diverse genes. In the case of Rickettsiaceae family, the genes usually targeted are: *gltA*, *ompA* and *ompB* (Table 5).

#### 5. Middle and high-throughput tools for the study of Rickettsiales

In addition to the molecular protocols described for the identification of one or a group of pathogens, dedicated tools for the simultaneous identification of this group of pathogens have been developed. In this sense, the Reverse Line Blot Hybridization (RLBH), DNA microarray and next generation sequencing techniques can be mentioned.

The RLBH methodology enables the simultaneous detection of different bacterial species from diverse samples. It consists on a PCR assay (nested or unique) that is then coupled to a hybridization reaction. Each species can be identified by a species-specific oligonucleotide that acts as a probe and that it is covalently bound to a membrane where the hybridization reaction, with PCR product, will take place. The name of the method arises from the reversed arrangement of the components. A device called "line blotter" is required to carry out this methodology, and enables the evaluation of different samples applying different probes that are immobilized in a transverse sense to the line where the amplification product will be loaded. Furthermore, the oligonucleotides employed in the amplification step are biotinilated in the



**Fig. 1.** Suggested algorithms for Rickettsiales identification and characterization.

5' terminus giving rise to a labeled PCR product, which enables the hybridization signal recognition through a coupled detection reaction using streptavidin-peroxidase, followed by a chemiluminescence developed reaction, visualized by exposing the membrane to X-ray film. The

membrane with the specific oligonucleotides in different lines is placed in a transversal sense in relation to that in which the PCR products are loaded, so that the different species simultaneously amplified by PCR can be specifically detected in the intersection of the oligonucleotide

**Table 4**  
Primers for identifying SFG Rickettsiae.

Rickettsia group	Primers sequence	Target Gene	Amplicon size	Reference
<i>Rickettsia</i> spp.	120-M59: 5'-CCGCAGGGTTGGTAACGTG-3' 120-807: 5'-CCTTITAGATTCACGCCCTAA-3'	<i>ompB</i>	833 bp	Roux and Raoult (2000)
<i>Rickettsia</i> spp.	CS-78 5'-GCAAGTATCGGTGAGGATGTAAT-3' CS-323 5'-GCTTCTTAAATTCAATAATCAGGAT-3'	<i>gltA</i>	401 bp	Labruna et al. (2004)
<i>Rickettsia</i> spp.	RpCS.415 5'-GCTATTATGCTTGCCTGT-3' RpCS.1220 5'-TGCATTCTTCCATTGTGC-3'	<i>gltA</i>	806 bp	de Sousa et al. (2006)
<i>Rickettsia</i> spp.	RCK/23-5-F: 5'-GATAGGTCRGRGTGGAAGCAC-3' RCK/23-5-R: 5'-TCGGGAYGGGATCGTGTTC-3'	23S-5S	388 bp	Jado et al. (2006)
SFG <i>Rickettsia</i>	RpCS.877p 5'-GGGGCCTCTCACGGCG-3' RpCS.1258n 5'-ATTGCGAAAAGTACAGTGAAACA-3'	<i>gltA</i>	381 bp	Regnery et al. (1991)
	Rrl90.70p 5'-ATGGCGAATATTCTCCAAAA-3' Rrl90.602n 5'-AGTCAGCATCGCTCCCCCT-3'	<i>ompA</i>	532 bp	
SFG <i>Rickettsia</i>	Rr 190.70p 5'-ATGGCGAATATTCTCCAAAA-3' 190-701 5'-CTTCCCTTAATGCCAGCATCT-3'	<i>ompA</i>	631 bp	Roux et al. (1996)
SFG <i>Rickettsia</i>	SLO1F: 5'-CACCACTCAACCGCAG-3' SLO1R: 5'-GCCGGGGCTGCAGATTG-3'	<i>ompA</i>	489 bp	Raoult et al. (2002)
SFG <i>Rickettsia</i>	107F 5'-GCTTATTACCAACCTCAAC-3' 299R 5'-TRATCACCAACCGTAAGTAAAT-3'	<i>ompA</i>	209–212 bp	Kidd et al. (2008)
SFG <i>Rickettsia</i>	Rc.rompB.4.362p: 5'-GTCAGGGTTACTCTCTGATGC -3' Rc.rompB.4.836n: 5'-CGTAGCTCCATCTAGCATCAG-3'	<i>ompB</i>	475 bp	Choi et al. (2005)
	Rc.rompB.4.496p: 5'-CCAATGGCAGGACTTAGCTACT-3' Rc.rompB.4.762n: 5'-AGGCTGGCTGATACACGGAGTAA-3'		267 bp	

**Table 5**

RT-PCR protocols for Rickettsiaceae family identification.

Organism	Target gene	Primers/probes	Discriminatory system	Quantitative	Reference
<i>Rickettsia</i> spp.	<i>gltA</i>	CS-5 5'-GAGAGAAAAATTATCCAAATGTTGAT -3' CS-6 5'-AGGGTCTTCGTGCATITCTT -3' probe 5'-6-FAM dCATTGTGCCATCCAGCTACCGT BHQ1-3'	Fluorogenic probe	No	Labruna et al. (2004)
<i>Rickettsia</i> spp.	<i>gltA</i>	gltAF 5'-TCGCAAATGTTCACGGTACTT-3' gltAR 5'-TCGTGCATITCTTCATTGTCATTC-3'	Melting curve (SYBR Green)	No	Stenos et al. (2005)
	<i>ompB</i>	ompBF 5'-CGACGTTAACGGTCTCATTC-3' ompBR 5'-ACCGGTTCTTGTAGTTTCGTC-3'		No	Paris et al. (2008)
SFG	<i>ompA</i>	RR190.547F 5'-CCTGCCGATAATTACAGGTTA -3' RR190.701R 5'-GTTCCGTTAACGGCAGCAT -3'	SYBR Green	Yes	Eremeeva et al. (2003)
<i>R. felis</i> <i>R. conorii</i> <i>R. typhi</i>	<i>gltA</i>	CS877F 5'-GGGGCCCTGCTACGGCGG-3' CS1258R 5'-ATTGCAAAAGTACAGTGAACA-3'	Melting curve (SYBR Green)	Yes	Rolain et al. (2002)

and the PCR product lines. This membrane can be re-used up to 20 times and has the major advantage of enabling the simultaneous analysis of up to 43 samples in the same membrane (Kong and Gilbert, 2006). RLH membranes have been developed to identify pathogens from the Anaplasmataceae family (Bekker et al., 2002; Tomassone et al., 2008). The PCR step employs oligonucleotides which target a conserved region of the 16S rRNA of *Anaplasma* and *Ehrlichia*, containing a hyper-variable region with polymorphisms specific for each of the known species in the two genera (Bekker et al., 2002). After amplification, the product is loaded in the membrane including specific oligonucleotides for *A. phagocytophilum*, *A. marginale*, *A. centrale*, *A. ovis*, *E. chaffeensis*, *E. ruminantium*, *Ehrlichia* sp. *ommatjenne* and *E. canis*. Generic and species-specific probes were also designed for eleven *Rickettsia* species by Jado et al. (2006). The PCR required in the first step targets the 23S-5S internal transcribed spacer (ISR), and the resulting amplicons are then hybridized to a membrane containing the generic (*Rickettsia* catch-all, SFG and TG) and species-specific probes (*R. akari*, *R. bellii*, *R. slovaca*, *R. conorii*, *R. aeschlimannii*, *R. rickettsii*-*R. sibirica*, *R. helvetica*, *R. felis*, *R. australis*, *R. prowazekii*, *R. typhi*). For *R. rickettsii* and *R. sibirica*, a common probe was designed due to the high homology of their 23S-5S ISR sequences, and amplicon sequencing is needed for discriminating between the two species. RLH is a useful tool for diagnostic and epidemiologic surveys and enables the detection of new microbial species. Although being a simple methodology of easy interpretation, it is specific enough to detect single-base-pair changes and showed high sensitivity for Rickettsiales identification (Schouls et al., 1999). Jado et al. (2006) report a sensitivity of one plasmid copy per reaction, which was determined in the presence of foreign DNAs from different clinical and environmental samples. Moreover, RLH membranes can be designed combining probes for different pathogens. In this sense, specific genus and species probes for Rickettsiaceae and Anaplasmataceae can be placed in the same membrane so that simultaneous detection of bacteria from the order can be carried out in the same reaction.

The DNA microarray is another tool based on molecular hybridization. El-Ashker et al. (2015) have developed a DNA microarray chip for the simultaneous identification of cattle tick-borne pathogens including *Anaplasma* species. As it was described for the RLH, after a first step of amplification of a 16S rRNA biotinylated amplicon, the PCR fragments are then hybridized and incubated with streptavidin-peroxidase conjugate for further visualization through a Chip Scanner. Nevertheless, the validation of this microarray is still in progress and has not been widely adopted.

The molecular approaches described above provide a way to gauge diversity in a microbial community, but considering that it is likely that ticks are host to a larger diversity of, as yet non-described, microorganisms, those molecular tools are not an efficient way to determine the presence or absence of any microbes in a sample, nor they provide robust information on relative abundances of a given microbe. In this sense, the advent of next generation sequencing techniques (NGS) promises to surmount many of the limitations in current molecular approaches to microbial diversity studies (Clay and Fuqua, 2010).

## 6. Suggested algorithms for taxonomic characterization and identification of novel species

Classification of members of the genus *Rickettsia* was based on mouse serotyping (Philip et al., 1978), Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) or pulsed-field gel electrophoresis (PFGE) (Fournier et al., 2003) until Fournier et al. (2003) established objective guidelines for isolate classification based on the sequencing of specific genes. These guidelines enable the classification of an isolate as member or not of this genus at various taxonomic levels. Previously, DNA taxonomy has been demonstrated to be useful for bacterial classification purposes (Bouyer et al., 2001; Niebylski et al., 1997; Tautz et al., 2003). The complete published sequences of the 16S rRNA (*rrs*), *gltA*, *ompB* and *gene D* (encoding the surface-exposed high-molecular-weight protein PS120), and the 5'end of the *ompA* gene of all validated *Rickettsia* species (except for *R. peacockii*) were compared. Then, based on the percentage of homology of each gene in relation to the known *Rickettsiae*, the unknown bacteria can be identified at the different taxonomic levels (Fournier et al., 2003).

In the last few years there have been reports of isolates that were characterized as *Ehrlichia* spp. by molecular methods, but failed to assign a species due to scarce sequence similarities against validated *Ehrlichia* species. In this regard, a novel tick-transmitted *Ehrlichia* called Panola Mountain was reported based on phylogenetic reconstruction from fragments of the 16S rRNA, *gltA*, *map1*, *map2*, and *ribonuclease III* genes. This organism was first detected in a goat and *A. americanum* ticks from a park in the metropolitan area of Atlanta, GA, U.S.A (Loftis et al., 2006). On the other hand, *Ehrlichia* sp. AvBat was proposed as a novel species by Socolovschi et al. (2012) after analyzing phylogenetic trees based on the *gltA* and 16S rRNA genes. Later, using a different set of target regions, Cruz et al. (2012) and O'Nion et al. (2015) reported the isolation of potentially novel *Ehrlichial* species from the hemolymph of *Rhipicephalus microplus* engorged females that had been collected from naturally infested cattle in Brazil and from horses in Nicaragua, respectively. Even, *Candidatus* status genera were proposed after phylogenetic reconstruction of diverse gene sequences, like "*Candidatus Xenohaliotis*" (Friedman et al., 2000) and "*Candidatus Neoehrlichia*" (Schouls et al., 1999).

This findings remark the need of a clear and agreed algorithm for new *Ehrlichia* species identification, similar to the one available for *Rickettsia* genus (Fournier et al., 2003).

A useful tool for the identification of new species of both Anaplasmataceae and Rickettsiaceae is the RLH method. As a first step, a conserved region can be amplified by PCR, the targets are the 23S-5S internal transcribed spacer for *Rickettsia* genus, and the 16S rRNA gene for *Ehrlichia* and *Anaplasma* genera. Then by applying RLH, the hypervariable region can be evaluated, since the membrane contains specific probes with polymorphisms for each of the known species of the three genera. If analyzing a sample, the common fragment is detected but no species-specific probes are identified, evaluation of possible new species should be considered through different target

genes. For *Ehrlichia* and *Anaplasma* genera, three target genes are suggested (Dumler et al., 2001; Inokuma et al., 2001a, 2001b) while for *Rickettsia* spp. a scheme of five target genes is proposed (Fournier et al., 2003) (Fig. 2).

## 7. Vertebrate reservoir host identification through blood meal analysis

Understanding the dynamics of pathogen transmission from infected wild hosts is critical for the study of tick-borne-pathogen epidemiology. Ticks can acquire pathogens by two main transmission ways: vertical (transovarial), in which the pathogen is transmitted from the female to offspring during egg development; and horizontal, in which pathogens are acquired during the blood meal feeding from reservoir or infected host (Allan et al., 2010). Moreover, transmission can occur between infected and susceptible ticks feeding simultaneously in close proximity on the same host (cofeeding transmission); such mechanism may enable the transmission of a pathogen even between ticks feeding on a non-competent host (Randolph et al., 1996). Efforts to identify reservoirs have historically been laborious, requiring in many cases the capture of potential hosts, their experimental infection with the organism of interest and the subsequent evaluation of their efficiency in the transmission process under controlled conditions (xenodiagnosis; LoGiudice et al., 2003). Even with such complex procedures, it is not always possible to assess the true importance of each vertebrate in the pathogens maintenance (Brisson et al., 2008). Xenodiagnostic procedures are essential to confirm the capacity of an infected vertebrate to transmit the pathogen to the vector, and of being infected by a vector. However, the molecular identification of the host DNA in the arthropod blood meal, together with the pathogen detection in the tick, may be a valid tool for the identification of possible reservoir sources of blood pathogens. This is particularly valuable in the study of the epidemiology of pathogens whose transmission cycles are not completely known, as

in the case of several Rickettsiales. Unfortunately, diverse factors can interfere with the detection of vertebrates DNA in vector samples. Among such factors, we have the scarce host blood volume available in the tick (due to degradation or the incapacity of the tick to engorge until repletion on certain hosts). The choice of the genome region that will be the target for molecular detection is also critical (Kent, 2009), since degradation of host DNA occurs after a period of time and only short DNA fragments are found.


Molecular protocols for detection of host DNA are in constant development, with the use of new targets and primers (Wodecka et al., 2014). One available tool is DNA sequencing of amplicons obtained from vertebrate conserved genes like the mitochondrial cytochrome b (CytB) (Pierce et al., 2009; Che Lah et al., 2015) or vertebrate rRNA genes. This option is accurate and specific but expensive when a great amount of samples are analyzed (Kent, 2009); moreover, it is not always possible to identify multiple blood meals within a vector. An alternative is the use of PCR-RFLP (Kirstein and Gray, 1996; Wodecka et al., 2014), which employs restriction endonucleases enzymes for the fragmentation of amplicons obtained from vertebrate conserved genes. This digestion results in differentially sized segments which are resolved visually by gel electrophoresis. The disadvantage of this technique is that, for a correct identification, a library of restriction patterns from known vertebrates is required (Kent, 2009). Another available tool is the PCR amplification of vertebrate DNA with universal primers, followed by RLBH; 18S rRNA gene (Allan et al., 2010; Pichon et al., 2003, 2005) and 12S rDNA gene (Humair et al., 2007) are generally used as targets. Recently, a new sensitive method for tick blood meal analysis has been reported: a non-targeted, genome-free, proteomics-based, spectral-matching methodology (Önder et al., 2013). This tool employs direct spectral matching to determine the vertebrate species from which the tick acquired the blood. A great advantage is that heavily degraded DNA can be detected, even when several months from the last meal have passed. Accordingly, further advances in the field are expected with the development of next-generation sequencing (NGS). This technique may enable the simultaneous identification of the vertebrate blood and pathogen infection within the arthropod vectors; in addition, it may allow the identification of multiple blood meals and requires a small amount of vertebrate reservoir DNA. The standardization of the methodologies and a decrease in costs are needed to make NGS a commonly used tool for the analysis of blood meal.

## 8. Conclusions

Rickettsiosis, ehrlichiosis, and anaplasmosis are increasingly reported in humans and animals worldwide, and tools for their recognition and prevention are needed. Although several techniques have been developed in the recent years, there is a need for a reorganization of the diagnostic procedures to make them more uniform across laboratories and geographic areas.

In our experience, the approach of choice should start with the identification of pathogens from Rickettsiaceae and Anaplasmataceae families at the genus level in host and tick samples, employing the main target genes validated for each family. Then, for positive reactions, further analysis can be carried out. For this purpose, groups and species can be characterized by sequencing the amplified products or applying specie-specific real-time or end point PCR protocols. For *Ehrlichia* and *Anaplasma* species, isolate characterization through specific molecular markers is also feasible. These methodologies are useful for molecular epidemiology surveys as they enable genotypic characterization of isolates and the study of the origin of new variants of pathogens.

Although NGS is not widely used for diagnostic purpose so far (Lefterova et al., 2015), it is promising for microbial diversity studies, including the study of the tick microbiome. Moreover, the same tick sample examined for pathogen identification can provide data for the study of the tick feeding behavior, through the blood meal analysis. This approach is critical for understanding tick-borne-diseases as it brings



**Fig. 2.** Suggested algorithms for taxonomic characterization and identification of novel *Rickettsia*, *Anaplasma* and *Ehrlichia* species.

information related to vertebrates that may act as reservoir or amplifier hosts (Labruna, 2009), contributing to the dispersion of tick-borne pathogens. The identification of new reservoir/amplifier hosts is a challenge in a constantly changing environment, where the increasing contact rate between wild and domestic animals, and animals and human beings, leads to the emergence and re-emergence of infectious diseases.

## Acknowledgments

We thank, Dr. Julia Sabio y García for the English language editing and Luis Fernandez for providing bibliography.

## References

- Abarca, K., López, J., Acosta-Jamett, G., Lepe, P., Soares, J.F., Labruna, M.B., 2012. A third *Amblyomma* species and the first tick-borne rickettsia in Chile. *J. Med. Entomol.* 49 (1), 219–222.
- Adakal, H., Meyer, D.F., Carasco-Lacombe, C., Pinarello, V., Allègre, F., Huber, K., Stachurski, F., Morand, S., Martinez, D., Lefrançois, T., Vachiery, N., Frutos, R., 2009. MLST scheme of *Ehrlichia ruminantium*: genomic stasis and recombination in strains from Burkina-Faso. *Infect. Genet. Evol.* 9 (6), 1320–1328.
- Alberti, A., Zobba, R., Chessa, B., Addis, M.F., Sparagano, O., Pinna Parpaglia, M.L., Cubeddu, T., Pintori, G., Pittau, M., 2005. Equine and canine *Anaplasma phagocytophilum* strains isolated on the island of Sardinia (Italy) are phylogenetically related to pathogenic strains from the United States. *Appl. Environ. Microbiol.* 71 (10), 6418–6422.
- Allan, B.F., Goessling, L.S., Storch, G.A., Thach, R.E., 2010. Blood meal analysis to identify reservoir hosts for *Amblyomma americanum* ticks. *Emerg. Infect. Dis.* 16 (3), 433–440.
- Allison, R.W., Little, S.E., 2013. Diagnosis of rickettsial diseases in dogs and cats. *Vet. Clin. Pathol.* 42 (2), 127–144.
- Anderson, B.E., Sumner, J.W., Dawson, J.E., Tzianabos, T., Greene, C.R., Olson, J.G., Fishbein, D.B., Olsen-Rasmussen, M., Holloway, B.P., George, E.H., 1992. Detection of the etiologic agent of human ehrlichiosis by polymerase chain reaction. *J. Clin. Microbiol.* 30 (4), 775–780.
- Andersson, M., Bartkova, S., Lindestad, O., Råberg, L., 2013. Co-infection with "Candidatus neoehrlichia mikurensis" and *Borrelia afzelii* in *Ixodes ricinus* ticks in southern Sweden. *Vector Borne Zoototic Dis.* 13 (7), 438–442.
- André, M.R., Adania, C.H., Machado, R.Z., Allegretti, S.M., Felipe, P.A., Silva, K.F., Nakaghi, A.C., 2010. Molecular and serologic detection of *Ehrlichia* spp. in endangered Brazilian wild captive felids. *J. Wildl. Dis.* 46 (3), 1017–1023.
- Azad, A., 1988. Farhang Relationship of Vector Biology and Epidemiology of Louse and Flea-Borne Rickettsioses. Walker David Biology of Rickettsial Diseases. CRC Press, Inc., Boca Raton, Florida, USA, pp. 51–61.
- Barber, R.M., Li, Q., Diniz, P.P., Porter, B.F., Breitschwerdt, E.B., Claiborne, M.K., Birkenheuer, A.J., Levine, J.M., Levine, G.J., Chandler, V., Kenny, P., Nghiêm, P., Wei, S., Greene, C.E., Kent, M., Platt, S.R., Greer, K., Schatzberg, S.J., 2010. Evaluation of brain tissue or cerebrospinal fluid with broadly reactive polymerase chain reaction for *Ehrlichia*, *Anaplasma*, spotted fever group Rickettsia, *Bartonella*, and *Borrelia* species in canine neurological diseases (109 cases). *J. Vet. Intern. Med.* 24 (2), 372–378.
- Beall, M.J., Chandrashekhar, R., Eberts, M.D., Cyr, K.E., Diniz, P.P., Mainville, C., Hegarty, B.C., Crawford, J.M., Breitschwerdt, E.B., 2008. Serological and molecular prevalence of *Borrelia burgdorferi*, *Anaplasma phagocytophilum*, and *Ehrlichia* species in dogs from Minnesota. *Vector Borne Zoototic Dis.* 8 (4), 455–464.
- Bekker, C.P., de Vos, S., Taoufik, A., Sparagano, O.A., Jongejan, F., 2002. Simultaneous detection of *Anaplasma* and *Ehrlichia* species in ruminants and detection of *Ehrlichia ruminantium* in *Amblyomma variegatum* ticks by reverse line blot hybridization. *Vet. Microbiol.* 89 (2–3), 223–238.
- Bell, C.A., Patel, R., 2005. A real-time combined polymerase chain reaction assay for the rapid detection and differentiation of *Anaplasma phagocytophilum*, *Ehrlichia chaffeensis*, and *Ehrlichia ewingii*. *Diagn. Microbiol. Infect. Dis.* 53 (4), 301–306.
- Birtles, R.J., Raoult, D., 1996. Comparison of partial citrate synthase gene (glta) sequences for phylogenetic analysis of *Bartonella* species. *Int. J. Syst. Bacteriol.* 46, 891–897.
- Bouyer, D.H., Stenos, J., Croquet-Valdes, P., Moron, C., Vsevolod, P., Zavala-Velasquez, J.E., Foil, L., Stothard, D., Azad, A., Walker, D., 2001. *Rickettsia felis*: molecular characterization of a new member of the spotted fever group. *Int. J. Syst. Evol. Microbiol.* 51, 339–347.
- Breitschwerdt, E.B., Levy, M.C., Davidson, M.G., Walker, D.H., Burgdorfer, W., Curtis, B.C., Babineau, C.A., 1990. Kinetics of IgM and IgG responses to experimental and naturally acquired *Rickettsia rickettsii* infection in dogs. *Am. J. Vet. Res.* 51 (8), 1312–1316.
- Breitschwerdt, E.B., Hegarty, B.C., Hancock, S.I., 1998. Sequential evaluation of dogs naturally infected with *Ehrlichia canis*, *Ehrlichia chaffeensis*, *Ehrlichia equi*, *Ehrlichia ewingii*, or *Bartonella vinsonii*. *J. Clin. Microbiol.* 36 (9), 2645–2651.
- Brisson, D., Dykhuizen, D.E., Ostfeld, R.S., 2008. Conspicuous impacts of inconspicuous hosts on the Lyme disease epidemic. *Proc. Biol. Sci.* 275 (1631), 227–235.
- Buller, R.S., Arens, M., Hmiel, S.P., Paddock, C.D., Sumner, J.W., Rikhisa, Y., Unver, A., Gaudreault-Keener, M., Manian, F.A., Liddell, A.M., Schmulewitz, N., Storch, G.A., 1999. *Ehrlichia ewingii*, a newly recognized agent of human ehrlichiosis. *N. Engl. J. Med.* 341 (3), 148–155.
- Carelli, G., Decaro, N., Lorusso, A., Elia, G., Lorusso, E., Mari, V., Ceci, L., Buonavoglia, C., 2007. Detection and quantification of *Anaplasma marginale* DNA in blood samples of cattle by real-time PCR. *Vet. Microbiol.* 124 (1–2), 107–114.
- Che Lah, E.F., Yaakop, S., Ahamed, M., Md Nor, S., 2015. Molecular identification of blood meal sources of ticks (Acarina, Ixodidae) using cytochrome b gene as a genetic marker. *Zookeys* (478), 27–43.
- Chi, Q., Liu, Z., Li, Y., Yang, J., Chen, Z., Yue, C., Luo, J., Yin, H., 2015. Development of a real-time PCR assay for detection and quantification of *Anaplasma ovis* infection. *Transbound. Emerg. Dis.* 60 (Suppl 2), 119–124.
- Childs, J.E., Sumner, J.W., Nicholson, W.L., Massung, R.F., Standaert, S.M., Paddock, C.D., 1999. Outcome of diagnostic tests using samples from patients with culture-proven human monocytic ehrlichiosis: implications for surveillance. *J. Clin. Microbiol.* 37 (9), 2997–3000.
- Choi, Y.J., Lee, S.H., Park, K.H., Koh, Y.S., Lee, K.H., Baik, H.S., Choi, M.S., Kim, I.S., Jang, W.J., 2005. Evaluation of PCR-based assay for diagnosis of spotted fever group rickettsiosis in human serum samples. *Clin. Diagn. Lab. Immunol.* 12 (6), 759–763.
- Chu, F.K., 1998. Rapid and sensitive PCR-based detection and differentiation of aetiologic agents of human granulocytotropic and monocyte-tropic ehrlichiosis. *Mol. Cell. Probes* 12, 93–99.
- Cicuttin, G., Nava, S., 2013. Molecular identification of *Rickettsia parkeri* infecting *Amblyomma triste* ticks in an area of Argentina where cases of rickettsiosis were diagnosed. *Mem. Inst. Oswaldo Cruz* 108 (1), 123–125.
- Clay, K., Fuqua, C., 2010. The tick microbiome: diversity, distribution and influence of the internal microbial community for a blood-feeding disease vector. The State of the Science workshop: "Critical Needs and Gaps in Understanding Prevention, Amelioration, and Resolution of Lyme and Other Tick-Borne Diseases: The Short-Term and Long-Term Outcomes" Washington, D.C.
- Courtney, J.W., Kostelnik, L.M., Zeidner, N.S., Massung, R.F., 2004. Multiplex real-time PCR for detection of *Anaplasma phagocytophilum* and *Borrelia burgdorferi*. *J. Clin. Microbiol.* 42 (7), 3164–3168.
- Crocquet-Valdes, P.A., Thirumalapura, N.R., Ismail, N., Yu, X., Saito, T.B., Stevenson, H.L., Pietsch, C.A., Thomas, S., Walker, D.H., 2011. Immunization with *Ehrlichia* P28 outer membrane proteins confers protection in a mouse model of ehrlichiosis. *Clin. Vaccine Immunol.* 18 (12), 2018–2025.
- Cruz, A.C., Zweigarth, E., Ribeiro, M.F., da Silveira, J.A., de la Fuente, J., Grubhoffer, L., Valdés, J.J., Passos, L.M., 2012. New species of *Ehrlichia* isolated from Rhipicephalus (Boophilus) microplus shows an ortholog of the *E. canis* major immunogenic glycoprotein gp36 with a new sequence of tandem repeats. *Parasit. Vectors* 5, 291.
- Dawson, J.E., Stallknecht, D.E., Howerth, E.W., Warner, C., Biggie, K., Davidson, W.R., Lockhart, J.M., Nettles, V.F., Olson, J.G., Childs, J.E., 1994. Susceptibility of white-tailed deer (*Odocoileus virginianus*) to infection with *Ehrlichia chaffeensis*, the etiologic agent of human ehrlichiosis. *J. Clin. Microbiol.* 32 (11), 2725–2728.
- de Sousa, R., Barata, C., Vitorino, L., Santos-Silva, M., Carrapato, C., Torgal, J., Walker, D., Bacellar, F., 2006. *Rickettsia sibirica* isolation from a patient and detection in ticks, Portugal. *Emerg. Infect. Dis.* 12 (7), 1103–1108.
- Decaro, N., Carelli, G., Lorusso, E., Lucente, M.S., Greco, G., Lorusso, A., Radogna, A., Ceci, L., Buonavoglia, C., 2008. Duplex real-time polymerase chain reaction for simultaneous detection and quantification of *Anaplasma marginale* and *Anaplasma centrale*. *J. Vet. Diagn. Investig.* 20 (5), 606–611.
- Doyle, C.K., Labruna, M.B., Breitschwerdt, E.B., Tang, Y.W., Corstvet, R.E., Hegarty, B.C., Bloch, K.C., Li, P., Walker, D.H., McBride, J.W., 2005. Detection of medically important *Ehrlichia* by quantitative multicolor TaqMan real-time polymerase chain reaction of the dsb gene. *J. Mol. Diagn.* 7 (4), 504–510.
- Dumler, J.S., 2010. Fitness and freezing: vector biology and human health. *J. Clin. Invest.* 120 (9), 3087–3090.
- Dumler, J.S., Barbet, A.F., Bekker, C.P., Dasch, G.A., Palmer, G.H., Ray, S.C., Rikihisa, Y., Rurangirwa, F.R., 2001. Reorganization of genera in the families Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, descriptions of six new species combinations and designation of *Ehrlichia equi* and 'HGE agent' as subjective synonyms of *Ehrlichia phagocytophila*. *Int. J. Syst. Evol. Microbiol.* 51 (Pt 6), 2145–2165.
- Durden, L.A., Traub, R., 2002. Fleas (Siphonaptera). In: Mullen, G.R., Durden, L.A. (Eds.), Medical and Veterinary Entomology. Academic Press, New York. NY, pp. 103–125.
- Eddlestone, S.M., Diniz, P.P., Neer, T.M., Gaunt, S.D., Corstvet, R., Cho, D., Hosgood, G., Hegarty, B., Breitschwerdt, E.B., 2007a. Doxycycline clearance of experimentally induced chronic *Ehrlichia canis* infection in dogs. *J. Vet. Intern. Med.* 21 (6), 1237–1242.
- Eddlestone, S.M., Gaunt, S.D., Neer, T.M., Boudreault, C.M., Gill, A., Haschke, E., Corstvet, R.E., 2007b. PCR detection of *Anaplasma platys* in blood and tissue of dogs during acute phase of experimental infection. *Exp. Parasitol.* 115 (2), 205–210.
- El-Ashker, M., Hotzel, H., Gwida, M., El-Beskawy, M., Silaghi, C., Tomaso, H., 2015. Molecular biological identification of Babesia, Theileria, and Anaplasma species in cattle in Egypt using PCR assays, gene sequence analysis and a novel DNA microarray. *Vet. Parasitol.* 207 (3–4), 329–334.
- Eremeeva, M., Yu, X., Raoult, D., 1994. Differentiation among spotted fever group rickettsia species by analysis of restriction fragment length polymorphism of PCR-amplified DNA. *J. Clin. Microbiol.* 32, 803–810.
- Eremeeva, M.E., Dasch, G.A., Silverman, D.J., 2003. Evaluation of a PCR assay for quantitation of *Rickettsia rickettsii* and closely related spotted fever group rickettsiae. *J. Clin. Microbiol.* 41 (12), 5466–5472.
- Faburay, B., Geysen, D., Munstermann, S., Bell-Sakyi, L., Jongejan, F., 2007. Longitudinal monitoring of *Ehrlichia ruminantium* infection in Gambian lambs and kids by pCS20 PCR and MAP1-B ELISA. *BMC Infect. Dis.* 7, 85.
- Penollar, F., Raoult, D., 2004. Molecular genetic methods for the diagnosis of fastidious microorganisms. *APMIS* 112 (11–12), 785–807.
- Fournier, P.E., Raoult, D., 2004. Suicide PCR on skin biopsy specimens for diagnosis of rickettsioses. *J. Clin. Microbiol.* 42 (8), 3428–3434.

- Fournier, P.E., Roux, V., Raoult, D., 1998. Phylogenetic analysis of spotted fever group rickettsiae by study of the outer surface protein rOmpA. *Int. J. Syst. Bacteriol.* 48 (Pt 3), 839–849.
- Fournier, P.E., Dumler, J.S., Greub, G., Zhang, J., Wu, Y., Raoult, D., 2003. Gene sequence-based criteria for identification of new rickettsia isolates and description of *Rickettsia heilongjiangensis* sp. nov. *J. Clin. Microbiol.* 41 (12), 5456–5465.
- Friedman, C.S., Andree, K.B., Beauchamp, K.A., Moore, J.D., Robbins, T.T., Shields, J.D., Hedrick, R.P., 2000. 'Candidatus xenohaliotis californiensis', a newly described pathogen of abalone, *Haliotis* spp., along the West Coast of North America. *Int. J. Syst. Evol. Microbiol.* 50 (Pt 2), 847–855.
- Fuxelius, H.H., Darby, A., Min, C.K., Cho, N.H., Andersson, S.G., 2007. The genomic and metabolic diversity of Rickettsia. *Res. Microbiol.* 158 (10), 745–753.
- García-García, J.C., Portillo, A., Núñez, M.J., Santibáñez, S., Castro, B., Oteo, J.A., 2010. A patient from Argentina infected with *Rickettsia massiliae*. *Am. J. Trop. Med. Hyg.* 82 (4), 691–692.
- Gillespie, J.J., Beier, M.S., Rahman, M.S., Ammerman, N.C., Shallom, J.M., Purkayastha, A., Sobral, B.S., Azad, A.F., 2007. Plasmids and rickettsial evolution: insight from *Rickettsia felis*. *PLoS One* 2 (3), e266 (7).
- Gillespie, J.J., Williams, K., Shukla, M., Snyder, E.E., Nordberg, E.K., Ceraul, S.M., Dharmanolla, C., Rainey, D., Soneja, J., Shallom, J.M., Vishnubhat, N.D., Wattam, R., Purkayastha, A., Czar, M., Crasta, O., Subtil, J.C., Azad, A.F., Sobral, B.S., 2008. Rickettsia phylogenomics: unwinding the intricacies of obligate intracellular life. *PLoS One* 3 (4), e2018.
- Goldman, E.E., Breitschwerdt, E.B., Grindem, C.B., Hegarty, B.C., Walls, J.J., Dumler, J.S., 1998. Granulocytic ehrlichiosis in dogs from North Carolina and Virginia. *J. Vet. Intern. Med.* 12, 61–70.
- Gross, L., 1996. How Charles Nicolle of the Pasteur institute discovered that epidemic typhus is transmitted by lice: reminiscences from my years at the Pasteur Institute in Paris. *Proc. Natl. Acad. Sci. U. S. A.* 93 (20), 10539–10540.
- Guillemi, E.C., Ruybal, P., Lia, V., Gonzalez, S., Lew, S., Zimmer, P., Lopez Arias, L., Rodriguez, J.L., Rodriguez, S.Y., Frutos, R., Wilkowsky, S.E., Farber, M.D., 2015. Development of a multilocus sequence typing scheme for the study of *Anaplasma marginale* population structure over space and time. *Infect. Genet. Evol.* 30C, 186–194.
- Henningsson, A.J., Hvidsten, D., Kristiansen, B.E., Matussek, A., Stuen, S., Jenkins, A., 2015. Detection of *Anaplasma phagocytophilum* in *Ixodes ricinus* ticks from Norway using a realtime PCR assay targeting the *Anaplasma* citrate synthase gene gltA. *BMC. Microbiol.* 15, 153.
- Horta, M.C., Labruna, M.B., Pinter, A., Linardi, P.M., Schumaker, T.T., 2007. Rickettsia infection in five areas of the state of São Paulo, Brazil. *Mem. Inst. Oswaldo Cruz* 102, 793–801.
- Huhn, C., Winter, C., Wolfsperger, T., Wüppenhorst, N., Strašek Smrdel, K., Skuballa, J., Pfäffle, M., Petney, T., Silaghi, C., Dyachenko, V., Pantchev, N., Straubinger, R.K., Schaarschmidt-Kiener, D., Ganter, M., Aardema, M.L., von Loewenich, F.D., 2014. Analysis of the population structure of *Anaplasma phagocytophilum* using multilocus sequence typing. *PLoS One* 9 (4), e93725.
- Humair, P.F., Douet, V., Cadena, F.M., Schouls, L.M., Van De Pol, I., Gern, L., 2007. Molecular identification of bloodmeal source in *Ixodes ricinus* ticks using 12S rDNA as a genetic marker. *J. Med. Entomol.* 44, 869–880.
- Inokuma, H., Raoult, D., Brouqui, P., 2000. Detection of *Ehrlichia platys* DNA in brown dog ticks (*Rhipicephalus sanguineus*) in Okinawa island, Japan. *J. Clin. Microbiol.* 38 (11), 4219–4221.
- Inokuma, H., Ohno, K., Onishi, T., Raoult, D., Brouqui, P., 2001a. Detection of ehrlichial infection by PCR in dogs from Yamaguchi and Okinawa prefectures, Japan. *J. Vet. Med. Sci.* 63 (7), 815–817.
- Inokuma, H., Brouqui, P., Drancourt, M., Raoult, D., 2001b. Citrate synthase gene sequence: a new tool for phylogenetic analysis and identification of *Ehrlichia*. *J. Clin. Microbiol.* 39 (9), 3031–3039.
- Inokuma, H., Fujii, K., Okuda, M., Onishi, T., Beaufils, J.P., Raoult, D., Brouqui, P., 2002. Determination of the nucleotide sequences of heat shock operon groESL and the citrate synthase gene (gltA) of *Anaplasma* (*Ehrlichia*) *Platys* for phylogenetic and diagnostic studies. *Clin. Diagn. Lab. Immunol.* 9 (5), 1132–1136.
- Jado, I., Escudero, R., Gil, H., Jiménez-Alonso, M.I., Sousa, R., García-Pérez, A.L., Rodríguez-Vargas, M., Lobo, B., Anda, P., 2006. Molecular method for identification of *Rickettsia* species in clinical and environmental samples. *J. Clin. Microbiol.* 44 (12), 4572–4576.
- Joblet, C., Roux, V., Drancourt, M., Gouvernet, J., Raoult, D., 1995. Identification of *Bartonella* (*Rochalimae*) species among fastidious gramnegative bacteria on the basis of the partial sequence of the citrate-synthase gene. *J. Clin. Microbiol.* 33, 1879–1883.
- Jolley, K.A., Kalmusova, J., Feil, E.J., Gupta, S., Musilek, M., Kriz, P., Maiden, M.C., 2000. Carried meningococci in the Czech Republic: a diverse recombining population. *J. Clin. Microbiol.* 38 (12), 4492–4498.
- Kang, Y.J., Diao, X.N., Zhao, G.Y., Chen, M.H., Xiong, Y., Shi, M., Fu, W.M., Guo, Y.J., Pan, B., Chen, X.P., Holmes, E.C., Gillespie, J.J., Dumler, S.J., Zhang, Y.Z., 2014. Extensive diversity of Rickettsiales bacteria in two species of ticks from China and the evolution of the Rickettsiales. *BMC Evol. Biol.* 14, 167.
- Kent, R.J., 2009. Molecular methods for arthropod bloodmeal identification and applications to ecological and vector-borne disease studies. *Mol. Ecol. Resour.* 9 (1), 4–18.
- Kidd, L., Maggi, R., Diniz, P.P., Hegarty, B., Tucker, M., Breitschwerdt, E., 2008. Evaluation of conventional and real-time PCR assays for detection and differentiation of spotted fever group rickettsia in dog blood. *Vet. Microbiol.* 129 (3–4), 294–303.
- Kirstein, F., Gray, J.S., 1996. A molecular marker for the identification of the zoonotic reservoirs of *Lyme borreliosis* by analysis of the blood meal in its European vector *Ixodes ricinus*. *Appl. Environ. Microbiol.* 62 (11), 4060–4065.
- Kong, F., Gilbert, G.L., 2006. Multiplex PCR-based reverse line blot hybridization assay (mPCR/RLB) a practical epidemiological and diagnostic tool. *Nat. Protoc.* 1 (6), 2668–2680.
- Labruna, M.B., 2009. Ecology of rickettsia in South America. *Ann. N. Y. Acad. Sci.* 1166, 156–166.
- Labruna, M.B., Whitworth, T., Horta, M.C., Bouyer, D.H., McBride, J.W., Pinter, A., Popov, V., Gennari, S.M., Walker, D.H., 2004. Rickettsia species infecting *Amblyomma cooperi* ticks from an area in the state of São Paulo, Brazil, where Brazilian spotted fever is endemic. *J. Clin. Microbiol.* 42 (1), 90–98.
- Lefterova, M.I., Suarez, C.J., Banaei, N., Pinsky, B.A., 2015. Next-generation sequencing for infectious disease diagnosis and management: a report of the association for molecular pathology. *J. Mol. Diagn.* (pii: S1525-1578(15)00172-5).
- Leitner, M., Yitzhaki, S., Rzotkiewicz, S., Keysary, A., 2002. Polymerase chain reaction-based diagnosis of Mediterranean spotted fever in serum and tissue samples. *Am. J. Trop. Med. Hyg.* 67 (2), 166–169.
- Li, J.S., Chu, F., Reilly, A., Winslow, G.M., 2002. Antibodies highly effective in SCID mice during infection by the intracellular bacterium *Ehrlichia chaffeensis* are of picomolar affinity and exhibit preferential epitope and isotype utilization. *J. Immunol.* 169, 1419–1425.
- Loftis, A.D., Massung, R.F., Levin, M.L., 2003. Quantitative real-time PCR assay for detection of *Ehrlichia chaffeensis*. *J. Clin. Microbiol.* 41 (8), 3870–3872.
- Loftis, A.D., Reeves, W.K., Spurlock, J.P., Mahan, S.M., Troughton, D.R., Dasch, G.A., Levin, M.L., 2006. Infection of a goat with a tick-transmitted *Ehrlichia* from Georgia, U.S.A., that is closely related to *Ehrlichia ruminantium*. *J. Vector Ecol.* 31 (2), 213–223.
- LoGiudice, K., Ostfeld, R.S., Schmidt, K.A., Keesing, F., 2003. The ecology of infectious disease: effects of host diversity and community composition on Lyme disease risk. *Proc. Natl. Acad. Sci. U. S. A.* 100 (2), 567–571.
- Lomanno, E., Bertiola, L., Dupasquier, C., Gern, L., 2012. Infections and coinfections of questing *Ixodes ricinus* ticks by emerging zoonotic pathogens in western Switzerland. *Appl. Environ. Microbiol.* 78 (13), 4606–4612.
- Louw, M., Allsopp, M.T., Meyer, E.C., 2005. *Ehrlichia ruminantium*, an emerging human pathogen—a further report. *S. Afr. Med. J.* 95 (12), 948 (950).
- Martínez, M.C., Gutiérrez, C.N., Monger, F., Ruiz, J., Watts, A., Mijares, V.M., Rojas, M.G., Triana-Alonso, F.J., 2008. *Ehrlichia chaffeensis* in child, Venezuela. *Emerg. Infect. Dis.* 14 (3), 519–520.
- Mayer, L.W., Reeves, M.W., Al-Hamdan, N., Sacchi, C.T., Taha, M.K., Ajello, G.W., Schminke, S.E., Noble, C.A., Tondella, M.L., Whitney, A.M., Al-Mazrou, Y., Al-Jefri, M., Mishkhis, A., Sabban, S., Caugant, D.A., Lingappa, J., Rosenstein, N.E., Popovic, T., 2002. Outbreak of W135 meningococcal disease in 2000: not emergence of a new W135 strain but clonal expansion within the electrophoretic type-37 complex. *J. Infect. Dis.* 185 (11), 1596–1605.
- Miranda, J., Contreras, V., Negrete, Y., Labruna, M.B., Máttar, S., 2011. Surveillance of *Rickettsia* sp. infection in capybaras (*Hydrochoerus hydrochaeris*) a potential model of epidemiological alert in endemic areas. *Biomedica* 31 (2), 216–221.
- Miranda, J., Portillo, A., Oteo, J.A., Mattar, S., 2012. *Rickettsia* sp. strain colombianensi (Rickettsiales: Rickettsiaceae): a new proposed *Rickettsia* detected in *Amblyomma dissimile* (Acar: Ixodidae) from iguanas and free-living larvae ticks from vegetation. *J. Med. Entomol.* 49 (4), 960–965.
- Munderloh, U.G., Madigan, J.E., Dumler, J.S., Goodman, J.L., Hayes, S.F., Barlough, J.E., Nelson, C.M., Kurtti, T.J., 1996. Isolation of the equine granulocytic ehrlichiosis agent, *Ehrlichia equi*, in tick cell culture. *J. Clin. Microbiol.* 34 (3), 664–670.
- Murphy, G.L., Ewing, S.A., Whitworth, L.C., Fox, J.C., Kocan, A.A., 1998. A molecular and serologic survey of *Ehrlichia canis*, *E. chaffeensis*, and *E. ewingii* in dogs and ticks from Oklahoma. *Vet. Parasitol.* 79 (4), 325–339.
- Nefedova, V.V., Korenberg, E.I., Kovalevski, I.V., Gorelova, N.B., Vorob'eva, N.N., 2008. Microorganisms of the order Rickettsiales in taiga tick (*Ixodes persulcatus* Sch.) from the Pre-Ural region. *Vestn. Ross. Akad. Med. Nauk* 7, 47–50.
- Nicholson, W.L., Allen, K.E., McQuiston, J.H., Breitschwerdt, E.B., Little, S.E., 2010. The increasing recognition of rickettsial pathogens in dogs and people. *Trends Parasitol.* 26 (4), 205–212.
- Niebylski, M.L., Schrumpf, M.E., Burgdorfer, W., Fischer, E.R., Gage, K.L., Schwan, T.G., 1997. *Rickettsia peacockii* sp. nov., a new species infecting wood ticks, *Dermacentor andersoni*, in western Montana. *Int. J. Syst. Bacteriol.* 47, 446–452.
- Ohashi, N., Rikihisa, Y., Unver, A., 2001. Analysis of transcriptionally active gene clusters of major outer membrane protein multigene family in *Ehrlichia canis* and *E. chaffeensis*. *Infect. Immun.* 69 (4), 2083–2091.
- OIE Terrestrial Manual, 2015. Chapter 2.4.1 Bovine anaplasmosis, Version adopted by the World Assembly of Delegates of the OIE in May 2015.
- Onder, Ö., Shao, W., Kemps, B.D., Lam, H., Brisson, D., 2013. Identifying sources of tick blood meals using unidentified tandem mass spectral libraries. *Nat. Commun.* 4, 1746.
- O'Nion, V.L., Montilla, H.J., Quroollo, B.A., Maggi, R.G., Hegarty, B.C., Tornquist, S.J., Breitschwerdt, E.B., 2015. Potentially novel *Ehrlichia* species in horses, Nicaragua. *Emerg. Infect. Dis.* 21 (2), 335–338.
- Ortranto, D., Dantas-Torres, F., Breitschwerdt, E.B., 2009. Managing canine vector-borne diseases of zoonotic concern: part one. *Trends Parasitol.* 25 (4), 157–163.
- Ortranto, D., Testini, G., Dantas-Torres, F., Latrofa, M.S., Diniz, P.P., de Capriatis, D., Lia, R.P., Mencke, N., Stanneck, D., Capelli, G., Breitschwerdt, E.B., 2010. Diagnosis of canine vector-borne diseases in young dogs: a longitudinal study. *J. Clin. Microbiol.* 48 (9), 3316–3324.
- Paddock, C.D., Childs, J.E., 2003. *Ehrlichia chaffeensis*: a prototypical emerging pathogen. *Clin. Microbiol. Rev.* 16, 37–64.
- Paris, D.H., Blacksell, S.D., Stenos, J., Graves, S.R., Unsworth, N.B., Phetsouvanh, R., Newton, P.N., Day, N.P., 2008. Real-time multiplex PCR assay for detection and differentiation of rickettsiae and orientiae. *Trans. R. Soc. Trop. Med. Hyg.* 102 (2), 186–193.
- Parola, P., Roux, V., Camicas, J.L., Baradji, I., Brouqui, P., Raoult, D., 2000. Detection of ehrlichiae in African ticks by polymerase chain reaction. *Trans. R. Soc. Trop. Med. Hyg.* 94 (6), 707–708.

- Parola, P., Davoust, B., Raoult, D., 2005. Tick- and flea-borne rickettsial emerging zoonoses. *Vet. Res.* 36, 469–492.
- Perez, M., Boodor, M., Zhang, C., Xiong, Q., Rikihisa, Y., 2006. Human infection with *Ehrlichia canis* accompanied by clinical signs in Venezuela. *Ann. N. Y. Acad. Sci.* 1078, 110–117.
- Peter, T.F., Deem, S.L., Barbet, A.F., Norval, R.A.I., Simbi, B.H., Kelly, P.J., Mahan, S.M., 1995. Development and evaluation of PCR assay for detection of low levels of *Cowdria ruminantium* infection in Amblyomma ticks not detected by DNA probe. *J. Clin. Microbiol.* 33, 166–172.
- Philip, R.N., Casper, E.A., Burgdorfer, W., Gerloff, R.K., Hughes, L.E., Bell, E.J., 1978. Serologic typing of rickettsiae of the spotted fever group by micro-immuno-fluorescence. *J. Immunol.* 121 (5), 1961–1968.
- Pichon, B., Egan, D., Rogers, M., Gray, J., 2003. Detection and identification of pathogens and host DNA in unfed host-seeking *Ixodes ricinus* L. (Acar: Ixodidae). *J. Med. Entomol.* 40 (5), 723–731.
- Pichon, B., Rogers, M., Egan, D., Gray, J., 2005. Blood-meal analysis for the identification of reservoir hosts of tick-borne pathogens in Ireland. *Vector Borne Zoonotic Dis.* 5 (2), 172–180.
- Pierce, K.A., Paddock, C.D., Sumner, J.W., Nicholson, W.L., 2009. Pathogen prevalence and blood meal identification in Amblyomma ticks as a means of reservoir host determination for ehrlichial pathogens. *Clin. Microbiol. Infect.* 15 (Suppl. 2), 37–38.
- Pritt, B.S., Sloan, L.M., Johnson, D.K., Munderloh, U.G., Paskewitz, S.M., McElroy, K.M., McFadden, J.D., Binnicker, M.J., Neitzel, D.F., Liu, G., Nicholson, W.L., Nelson, C.M., Franson, J.J., Martin, S.A., Cunningham, S.A., Steward, C.R., Bogumill, K., Bjorgaard, M.E., Davis, J.P., McQuiston, J.H., Warshauer, D.M., Wilhelm, M.P., Patel, R., Trivedi, V.A., Eremeeva, M.E., 2011. Emergence of a new pathogenic *Ehrlichia* species, Wisconsin and Minnesota, 2009. *N. Engl. J. Med.* 365 (5), 422–429.
- Quroollo, B.A., Riggins, D., Comyn, A., Zewde, M.T., Breitschwerdt, E.B., 2014. Development and validation of a sensitive and specific sodB-based quantitative PCR assay for molecular detection of *Ehrlichia* species. *J. Clin. Microbiol.* 52 (11), 4030–4032.
- Randolph, S.E., Gern, L., Nuttall, P.A., 1996. Co-feeding ticks: epidemiological significance for tick-borne pathogen transmission. *Parasitol. Today* 12 (12), 472–479.
- Raoult, D., Fournier, P.E., Fenollar, F., Jensenius, M., Prioe, T., de Pina, J.J., Caruso, G., Jones, N., Laferl, H., Rosenblatt, J.E., Marrie, T.J., 2001. *Rickettsia africae*, a tick-borne pathogen in travelers to sub-Saharan Africa. *N. Engl. J. Med.* 344 (20), 1504–1510.
- Raoult, D., Lakos, A., Fenollar, F., Beytout, J., Brouqui, P., Fournier, P.E., 2002. Spotless rickettsiosis caused by *Rickettsia slovaca* and associated with Dermacentor ticks. *Clin. Infect. Dis.* 34 (10), 1331–1336.
- Rasmussen, H.B., 2012. Restriction Fragment Length Polymorphism Analysis of PCR-amplified fragments (PCR-RFLP) and gel electrophoresis – valuable tool for genotyping and genetic fingerprinting. *Gel Electrophoresis - Principles and Basics*, pp. 315–334.
- Reeves, W.K., Loftis, A.D., Nicholson, W.L., Czarkowski, A.G., 2008. The first report of human illness associated with the Panola Mountain *Ehrlichia* species: a case report. *J. Med. Case Rep.* 2, 139.
- Regnery, R.L., Spruill, C.L., Plikaytis, B.D., 1991. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J. Bacteriol.* 173, 1576–1589.
- Rikihisa, Y., 2010. *Anaplasma phagocytophilum* and *Ehrlichia chaffeensis*: subversive manipulators of host cells. *Nat. Rev. Microbiol.* 8, 328–339.
- Rikihisa, Y., 2011. Mechanisms of obligatory intracellular infection with *Anaplasma phagocytophilum*. *Clin. Microbiol. Rev.* 24, 469–489.
- Roland, J.M., Stuhl, L., Maurin, M., Raoult, D., 2002. Evaluation of antibiotic susceptibilities of three rickettsial species including *Rickettsia felis* by a quantitative PCR DNA assay. *Antimicrob. Agents Chemother.* 46 (9), 2747–2751.
- Romer, Y., Nava, S., Govedic, F., Cicuttin, G., Denison, A.M., Singleton, J., Kelly, A.J., Kato, C.Y., Paddock, C.D., 2014. *Rickettsia parkeri* rickettsiosis in different ecological regions of Argentina and its association with *Amblyomma tigrinum* as a potential vector. *Am. J. Trop. Med. Hyg.* 91 (6), 1156–1160.
- Roux, V., Raoult, D., 2000. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer-membrane protein rOmpB (ompB). *Int. J. Syst. Evol. Microbiol.* 50 (Pt 4), 1449–1455.
- Roux, V., Fournier, P.E., Raoult, D., 1996. Identification of the spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein rOmpA. *J. Clin. Microbiol.* 34, 2058–2065.
- Roux, V., Rydkina, E., Eremeeva, M., Raoult, D., 1997. Citrate synthase gene comparison, a new tool for phylogenetic analysis, and its application for the rickettsiae. *Int. J. Syst. Bacteriol.* 47 (2), 252–261.
- Schouls, L.M., Van De Pol, I., Rijpkema, S.G., Schot, C.S., 1999. Detection and identification of *Ehrlichia*, *Borrelia burgdorferi* sensu lato, and *Bartonella* species in Dutch *Ixodes ricinus* ticks. *J. Clin. Microbiol.* 37 (7), 2215–2222.
- Scorpio, D.G., Akkoyunlu, M., Fikrig, E., Dumler, J.S., 2004. CXCR2 blockade influences *Anaplasma phagocytophilum* propagation but not histopathology in the mouse model of human granulocytic anaplasmosis. *Clin. Diagn. Lab. Immunol.* 11 (5), 963–968.
- Sirigireddy, K.R., Ganta, R.R., 2005. Multiplex detection of *Ehrlichia* and *Anaplasma* species pathogens in peripheral blood by real-time reverse transcriptase-polymerase chain reaction. *J. Mol. Diagn.* 7 (2), 308–316.
- Socolovschi, C., Kernif, T., Raoult, D., Parola, P., 2012. *Borrelia*, *Rickettsia*, and *Ehrlichia* species in bat ticks, France, 2010. *Emerg. Infect. Dis.* 18 (12), 1966–1975.
- Sparagano, O.A., Allsopp, M.T., Mank, R.A., Rijpkema, S.G., Figueroa, J.V., Jongejan, F., 1999. Molecular detection of pathogen DNA in ticks (Acar: Ixodidae): a review. *Exp. Appl. Acarol.* 23 (12), 929–960.
- Stenos, J., Graves, S.R., Unsworth, N.B., 2005. A highly sensitive and specific real-time PCR assay for the detection of spotted fever and typhus group Rickettsiae. *Am. J. Trop. Med. Hyg.* 73 (6), 1083–1085.
- Stich, R.W., Rikihisa, Y., Ewing, S.A., Needham, G.R., Grover, D.L., Jittapalapong, S., 2002. Detection of *Ehrlichia canis* in canine carrier blood and in individual experimentally infected ticks with a p30-based PCR assay. *J. Clin. Microbiol.* 40 (2), 540–546.
- Stoffel, R.T., McClure, J.C., Butcher, M.M., Johnson, G.C., Roland, W., Cheng, C., Sirigireddy, K.R., Ganta, R., Boughan, K., Ewing, S.A., Stich, R.W., 2014. Experimental infection of *Rhipicephalus sanguineus* with *Ehrlichia chaffeensis*. *Vet. Microbiol.* 172 (1–2), 334–338.
- Sumner, J.W., Childs, J.E., Paddock, C.D., 1999. Molecular cloning and characterization of the *Ehrlichia chaffeensis* variable-length PCR target: an antigen-expressing gene that exhibits interstrain variation. *J. Clin. Microbiol.* 37, 1447–1453.
- Tautz, D., Arctander, P., Minelli, A., Thomas, R.H., Vogler, A.P., 2003. A plea for DNA taxonomy. *Trends Ecol. Evol.* 18, 70–74.
- Teshale, S., Geysen, D., Ameni, G., Asfaw, Y., Berkvens, D., 2015. Improved molecular detection of *Ehrlichia* and *Anaplasma* species applied to *Amblyomma* ticks collected from cattle and sheep in Ethiopia. *Ticks Tick Borne Dis.* 6 (1), 1–7.
- Tomassone, L., Nuñez, P., Gürler, R.E., Ceballos, L.A., Orozco, M.M., Kitron, U.D., Farber, M., 2008. Molecular detection of *Ehrlichia chaffeensis* in *Amblyomma parvum* ticks, Argentina. *Emerg. Infect. Dis.* 14 (12), 1953–1955.
- Torioni de Echaide, S., Knowles, D.P., McGuire, T.C., Palmer, G.H., Suarez, C.E., McElwain, T.F., 1998. Detection of cattle naturally infected with *Anaplasma marginale* in a region of endemicity by nested PCR and a competitive enzyme-linked immunosorbent assay using recombinant major surface protein 5. *J. Clin. Microbiol.* 36 (3), 777–782.
- Tzianabos, T., Anderson, B.E., McDade, J.E., 1989. Detection of *Rickettsia rickettsii* DNA in clinical specimens by using polymerase chain reaction technology. *J. Clin. Microbiol.* 27 (12), 2866–2868.
- Vaumourin, E., Gasqui, P., Buffet, J.P., Chapuis, J.L., Pisani, B., Ferquel, E., Vayssié-Taussat, M., Vourc'h, G., 2013. A probabilistic model in cross-sectional studies for identifying interactions between two persistent vector-borne pathogens in reservoir populations. *PLoS One* 8 (6), e66167.
- Wagner, E.R., Bremer, W.G., Rikihisa, Y., Ewing, S.A., Needham, G.R., Unver, A., Wang, X., Stich, R.W., 2004. Development of a p28-based PCR assay for *Ehrlichia chaffeensis*. *Mol. Cell. Probes* 18 (2), 111–116.
- Wen, B., Rikihisa, Y., Mott, J.M., Greene, R., Kim, H.Y., Zhi, N., Couto, G.C., Unver, A., Bartsch, R., 1997. Comparison of nested PCR with immunofluorescent-antibody assay for detection of *Ehrlichia canis* infection in dogs treated with doxycycline. *J. Clin. Microbiol.* 35 (7), 1852–1855.
- Wen, B., Jian, R., Zhang, Y., Chen, R., 2002. Simultaneous detection of *Anaplasma marginale* and a new *Ehrlichia* species closely related to *Ehrlichia chaffeensis* by sequence analyses of 16S ribosomal DNA in *Boophilus microplus* ticks from Tibet. *J. Clin. Microbiol.* 40 (9), 3286–3290.
- Wodecka, B., Rymaszewska, A., Skotarczak, B., 2014. Host and pathogen DNA identification in blood meals of nymphal *Ixodes ricinus* ticks from forest parks and rural forests of Poland. *Exp. Appl. Acarol.* 62 (4), 543–555.
- Yu, X.J., Crocqet-Valdes, P., Culman, L.C., Walker, D.H., 1996. The recombinant 120-kilodalton protein of *Ehrlichia chaffeensis*, a potential diagnostic tool. *J. Clin. Microbiol.* 34, 2853–2855.