

Seed Science Research

<http://journals.cambridge.org/SSR>

Additional services for **Seed Science Research**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Structural aspects of dormancy in quinoa (*Chenopodium quinoa*): importance and possible action mechanisms of the seed coat

Diana Ceccato, Daniel Bertero, Diego Batlla and Beatriz Galati

Seed Science Research / Volume 25 / Issue 03 / September 2015, pp 267 - 275
DOI: 10.1017/S096025851500015X, Published online: 06 May 2015

Link to this article: http://journals.cambridge.org/abstract_S096025851500015X

How to cite this article:

Diana Ceccato, Daniel Bertero, Diego Batlla and Beatriz Galati (2015). Structural aspects of dormancy in quinoa (*Chenopodium quinoa*): importance and possible action mechanisms of the seed coat. Seed Science Research, 25, pp 267-275 doi:10.1017/S096025851500015X

Request Permissions : [Click here](#)

Structural aspects of dormancy in quinoa (*Chenopodium quinoa*): importance and possible action mechanisms of the seed coat

Diana Ceccato^{1*}, Daniel Bertero^{2,3}, Diego Batlla^{2,4} and Beatriz Galati⁵

¹Banco Base de Germoplasma, Instituto de Recursos Biológicos, CIRN, CNIA-INTA, B1686EYR Hurlingham, Buenos Aires, Argentina; ²Instituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a la Agricultura (IFEVA), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) - Universidad de Buenos Aires (UBA), C1417DSE Buenos Aires, Argentina; ³Cátedra de Producción Vegetal, ⁴Cátedra de Cerealicultura and ⁵Cátedra de Botánica General, Facultad de Agronomía, Universidad de Buenos Aires, C1417DSE Buenos Aires, Argentina

(Received 25 September 2014; accepted after revision 5 March 2015; first published online 6 May 2015)

Abstract

Two possible sources of resistance to pre-harvest sprouting were evaluated in quinoa. They showed dormancy at harvest and significant variations in dormancy level in response to environmental conditions experienced during seed development. The aims of this work were to evaluate the importance of seed coats in the regulation of dormancy in this species, to investigate possible mechanisms of action and to assess association of seed coat properties with changes in dormancy level caused by the environment. Accessions Chadmo and 2-Want were grown under field conditions on different sowing dates during 2 years. Seed coats were manipulated and seed germination was evaluated at different temperatures. Seed coat perforation before incubation led to faster dormancy loss in both accessions. This effect decreased with delayed sowing date, and seeds expressed a level of dormancy not imposed by coats. This suggests the presence of embryo dormancy in the genus *Chenopodium*. Seeds of the accession 2-Want had a significantly thinner seed coat at later sowing dates, associated with a decreasing coat-imposed dormancy, but this pattern was not detected in Chadmo. The seed coat acts as a barrier to the release of endogenous abscisic acid (ABA) in quinoa, suggested by the increase in germination and a higher amount of ABA leached from perforated seeds. ABA is able to leach from seeds with an intact seed coat, suggesting that differences in seed coat thickness may allow the leakage of different amounts of ABA. This mechanism may contribute to the observed differences in dormancy level, either between sowing dates or between accessions.

Keywords: *Chenopodium quinoa*, coat-imposed dormancy, embryo-imposed dormancy, pre-harvest sprouting, seed coat, sowing date

Introduction

The cultivation of the Andean seed crop quinoa (*Chenopodium quinoa*) out of its region of origin increases the risk of pre-harvest sprouting. This risk is low in its traditional area of cultivation, where it matures after the end of the rainy season (Geerts *et al.*, 2006), but coincidence with rains around the maturity date is known to affect seed quality (Jacobsen and Bach, 1998). The occurrence of dormancy after physiological maturity was confirmed for two *C. quinoa* accessions, as part of a research project aimed at the identification of resistance sources to pre-harvest sprouting. It was found that environmental conditions experienced by quinoa seeds during their development regulate the level of dormancy present at harvest and their release rate afterwards: high temperatures and long photoperiods produced more dormant seeds than low temperatures and short photoperiods (Ceccato *et al.*, 2011).

Seed dormancy is usually classified depending on the tissues that determine it – as an embryo- or coat-imposed dormancy – and that imposed by the seed coats is perhaps the most commonly studied (Finch-Savage and Leubner-Metzger, 2006; Nonogaki, 2006; Hilhorst, 2007; Linkies and Leubner-Metzger, 2012). Coats impose dormancy through several mechanisms. They can: (1) interfere with water absorption or gas exchange with the environment; (2) impose a mechanical restraint on radicle emergence; (3) prevent the release of inhibitors from the embryo; (4) release inhibitors to the embryo; and/or (5) filter light (Debeaujon *et al.*, 2007).

*Correspondence
Email: ceccato.diana@inta.gob.ar

Environmental effects on seed dormancy level can be exerted through changes in seed coat properties. In some cases, these effects are associated with changes in structural or chemical characteristics, such as seed coat thickness or their polyphenolic content. These kinds of association have been reported in some members of the genus *Chenopodium*. In *C. polyspermum* and *C. album*, seed coat thickness and germination are affected by the photoperiod experienced during development (Jacques, 1968; Karssen, 1970; Pourrat and Jacques, 1975). In particular for *C. album*, darker seed coats (showing higher dormancy) are correlated with the occurrence of longer days (Karssen, 1970). In *C. bonus-henricus*, Dorne (1981) found a negative association between altitude of origin and germination percentage. The latter decreased as the altitude at which the mother plant was grown increased. This decrease in germination was related to an increase of seed coat thickness and polyphenol content. The effect of altitude was associated with average temperature 30 d before harvest and this temperature was positively correlated with germination. Dorne suggested that radiation, which increases in association with altitude, may cause a higher content of oxidizable polyphenols, which determines a lower amount of oxygen reaching the embryo.

The possibility of identifying morphological and physiological processes controlling dormancy in quinoa, and how these traits are affected by the environment, is of paramount importance for the selection of sprouting-resistant genotypes in this species. Based on these considerations, the aims of this work were to establish the importance of seed coats in the regulation of dormancy in quinoa, to reveal possible mechanisms of action and to assess associations with changes in dormancy level caused by the environment.

Materials and methods

Seed production and germination

The seed coat in *C. quinoa* is composed of two cell layers, the endotegmen (the inner layer) and the exotesta (the external one) (Prego *et al.*, 1998; Sukhorukov and Zhang, 2013). Surrounding the seed coat is the pericarp, usually as a discontinuous layer at maturity. Experimental details are as described by Ceccato *et al.* (2011). Briefly, two quinoa accessions, Chadmo and 2-Want, both presenting dormancy at harvest, were cultivated in the field (34°60'S, 58°65'W, Hurlingham, Buenos Aires, Argentina) on three sowing dates during the first experimental year (2 November, 27 December and 13 March) and on two sowing dates during the second experimental year (7 November and 8 February) during 2005–2006 and 2006–2007. Both accessions were obtained from the Germplasm Bank at the US Department of Agriculture,

National Plant Germplasm System at Beltsville, Maryland, where they are identified as PI 614850 and AMES 13737, respectively. Environmental data were obtained from a conventional meteorological station located about 300 m from the experimental site [Instituto de Clima y Agua, Instituto Nacional de Tecnología Agropecuaria (INTA)]. Flowering date was determined as the date of first anthesis in at least 50% of the plants. Seed dry weight and moisture content were determined during seed filling by weighing 50 seeds per replicate before and after drying at 105°C for 24 h in an oven (Ministério da Agricultura e Reforma Agrária, 1992). Physiological maturity (PM) was considered to be complete when dry weight levelled off. Harvest time (H) was determined based on a 20% seed moisture content, which is common practice in the traditional production areas. Four panicles of each accession were sampled starting 15–20 d after flowering (DAF) and at intervals of 5 d until crop harvest. At each harvest time, 35 seeds were placed on filter paper (Schleicher & Schuell 0859, Dassel, Germany) and moistened with 4 ml of distilled water in plastic boxes of 85 mm diameter, with four replicates. Then they were incubated under fluorescent light, with a photoperiod of 8/16 h (light/dark), at 5, 10 or 25°C in an incubator with a temperature variation of $\pm 1^\circ\text{C}$. Seed germination was recorded daily for 15 d, and visible radicle protrusion was used as germination criterion. Seeds used for germination tests were extracted from the middle third of each panicle, and each replicate represents a single plant.

Bois *et al.* (2006) indicated that maximum germination in quinoa occurs between 18 and 23°C. Accessions studied in this work reached 100% of germination in just 2 d of incubation at 25°C when dormancy was not present (data not shown), and therefore 25°C was considered an optimal temperature for germination in subsequent experiments. The following treatments were applied to assess the coat's role in seed germination behaviour: *control* – whole seeds with pericarp; *perforated* – seeds punctured through the pericarp + seed coat in the perisperm area (to avoid embryo damage). This treatment was chosen in replacement of embryo isolation for operational reasons and based on previous observations by Jacques (1968) that a perforation treatment led to the complete release of dormancy in *C. polyspermum*. Ungerminated seeds were tested by the cutting test to rule out dead seeds, and the remaining were considered dormant. Furthermore, random samples of the same seed lot, taken months later, had 100% germination (Ceccato *et al.*, 2011).

Seed coat thickness measurement

Seed coat thickness was measured in seeds of both accessions that were sampled at harvest for each

sowing date in both experimental years. For this purpose seeds were hydrated in warm water with a few drops of detergent, pre-fixed in 2.5% glutaraldehyde in phosphate buffer (pH 7.2) for 2 h and post-fixed in 1.5% osmium tetroxide in the same buffer at 2°C for 3 h, then dehydrated in acetone at increasing concentrations and embedded in Spurr resin. Afterwards, seeds were cut longitudinally with respect to the embryo, with a Reichert–Jung ultramicrotome (Reichert Inc., New York, USA), and thin sections (1 μm thickness) were stained with cresyl violet and mounted in synthetic resin (O'Brien and McCully, 1981). Seed coat and pericarp thickness were measured in a section lateral to the radicle, where endosperm that sheaths the radicle is absent (Prego *et al.*, 1998), under a Wild M20 microscope (Wild Heerbrugg AG, Gais, Switzerland) with a graduated ocular lens. Photomicrographs were taken in a Motic DMWB1-223ASC microscope (Motic Instruments Co., Hong Kong, China) with incorporated digital camera. Ten replicate measurements were performed for each sowing date.

Measurement of abscisic acid leaching and abscisic acid content of seeds

Differences in the amount of abscisic acid (ABA) leaching into the incubation medium between perforated and control seeds (intact seeds) of both accessions were tested for seeds from the first sowing date of the second experimental year. Four replicates of 25 seeds were incubated at 25°C for 8 h in 20-mm plastic boxes containing 300 μl of distilled water. After incubation, the remaining distilled water contained in the boxes of each treatment was stored in a freezer at -20°C until ABA determination. ABA content in dry seeds from the same harvest was measured. For this purpose, whole seeds in four replicates of 25 were ground and water was added [24 ml (mg dry weight) $^{-1}$] for extraction. The amount of ABA was determined by radioimmunoassay, as described by Steinbach *et al.* (1995), using the monoclonal antibody AFR MAC 252 (Quarrie *et al.*, 1988) and tritiated ABA (Amersham Biosciences, Little Chalfont, Bucks, UK) (Rodríguez *et al.*, 2009; Mendiondo *et al.*, 2010; Di Mauro *et al.*, 2012). Each sample was assessed twice. The results presented are the mean value of four (for Chadmo) or three (for 2-Want) biological replicates \pm standard error (SE). Values are expressed either as picograms of ABA per milligram of seed dry weight [pg ABA (mg dw) $^{-1}$] or, as picograms of ABA per microlitre of incubation medium (pg ABA μl^{-1}).

Data analysis

The data were analysed using the InfoStat software (InfoStat Group, University of Córdoba, Argentina). T_{50} was calculated by fitting time-course curves to observed

germination data (germination percentage vs. time in days) using a Gómpertz function (Notivol *et al.*, 2007). Curves were fitted with GraphPad Prism 4 for Windows (GraphPad Software, San Diego California USA, www.graphpad.com) and the fitting value (r^2) was in every case greater than 0.95. Differences in coat thickness between accessions for each sowing date were evaluated by *t*-test for independent samples. The correlation between coat thickness and environmental conditions was analysed using the complete data set, the mean \pm 1 SE is presented for clarity. Differences in amounts of ABA were evaluated by unpaired *t*-test with Welch's correction (which does not assume equal variances) using GraphPad Prism version 5.00 for Windows (GraphPad Software, www.graphpad.com).

Results

Role of the seed coat in dormancy and sowing date effects

The effect of seed coat perforation on seed germination depended on sowing date and incubation temperature. Coat perforation produced a significant increase in the rate of dormancy loss during development for seeds in both accessions belonging to the first sowing date and incubated at 25°C (Fig. 1A, B). This difference was particularly contrasting in Chadmo, in which perforated seeds showed 84% germination at harvest while the control seeds showed no germination (Fig. 1B). For seeds belonging to the late sowing date (March), the effect of seed perforation was not significant in both accessions when seeds were incubated at 25°C. The seeds still expressed a level of dormancy that was not overcome by coat perforation (Fig. 1C, D).

At lower incubation temperatures (10 and 5°C), at the first sowing date, the effect of perforation was still significant in both accessions (Fig. 2A, B). In seeds belonging to the late sowing date a significant level of coat-imposed dormancy was still expressed in 2-Want at 10 and 5°C, overcome by perforation (Fig. 2C). Chadmo seeds expressed strongly reduced germination under low incubation temperatures, which was not overcome by coat perforation (Fig. 2D). The fact that dry afterripened seeds of both accessions achieve maximum germination at 10 and 5°C (Ceccato *et al.*, 2011), indicates that the observed reduction in germination was due to dormancy expression, and not a consequence of non-optimal temperatures for germination.

Seed coat thickness, sowing date and seed dormancy level

In accession 2-Want, seed coat thickness was reduced in late sowings as compared to the earlier ones, in both




Figure 1. Changes in the maximum germination after incubation at 25°C during seed development and maturation in (▲) control (intact seeds with pericarp) and (●) perforated seeds (seeds punctured through the pericarp + seed coat), in accession 2-Want (A and C) and Chadmo (B and D), from the first (A and B) and third sowing dates (C and D) in the first experimental year. In each panel, the left arrow indicates physiological maturity and the right arrow the harvest date. Symbols represent the mean ± 1SE.

experimental years ($P \leq 0.05$, Table 1, Fig. 3); however, no differences were detected for Chadmo. Additionally, independent samples *t*-test showed that the seed coat was significantly thicker in Chadmo as compared to 2-Want for each sowing date ($P \leq 0.01$, Table 1). No consistent variation in pericarp thickness was detected between sowing dates or between accessions (data not shown).

In the first experimental year, seed filling occurred during the months of January and February, March and April, and June for the first, second and third sowing date, respectively. In the second year seed filling occurred between January and February, and April and May for the first and second sowing date, respectively. Differences in seed coat thickness were associated with different environmental conditions (temperature and photoperiod) experienced by the seeds during their development for each sowing date (Fig. 4). The association of seed coat thickness with T_{50} as a dormancy parameter was also significant, for accession 2-Want ($P = 0.0009$; $r^2 = 0.21$; data not shown). T_{50} is the time in days from anthesis for germination at 10°C of harvested seeds to reach 50% (see Materials and methods).

A measurement of whole-seed polyphenol content, based on the assumption that higher polyphenol could lead to higher dormancy and explain variation

between sowing dates, showed an opposite trend to that of seed dormancy (data not shown). So, this hypothesis was provisionally discarded, although more specific and localized measurements are needed.

Interference of seed coat properties with leaching of abscisic acid

In accession Chadmo, ABA content was 2.5 times higher in the incubation medium of perforated seeds compared to that of intact seeds (Fig. 5A). For 2-Want, no significant differences were detected (Fig. 5B). Detectable ABA levels in the incubation medium were observed even in control seeds. The amount of ABA leaching from intact (control) seeds was higher in 2-Want than in Chadmo (Fig. 5A, B), and this was consistent with differences in ABA content of dry seeds between accessions (Fig. 5C).

Discussion

In agreement with previous results in other *Chenopodium* species (Jacques, 1968), the results of perforation treatment indicate that a significant part of dormancy is explained by the action of the seed coat




Figure 2. Changes in the maximum germination after incubation at 10 (closed symbols) and 5°C (open symbols) during seed development and maturation in (▲) control (intact seeds with pericarp), and (●) perforated seeds (seeds punctured through the pericarp + seed coat), in accession 2-Want (A and C) and Chadmo (B and D), from the first (A and B) and third sowing dates (C and D) in the first experimental year. In each panel, the left arrow indicates physiological maturity and the right arrow the harvest date. Symbols represent the mean \pm 1SE.

and pericarp tissues, since a discontinuity in these layers resulted in the acquisition of a high germination capability (Fig. 1A, B). On the other hand, since the pericarp is partially separated from the seeds at harvest time, it can be speculated that coat-imposed dormancy at harvest and afterwards is mainly imposed by the seed coat.

The lack of response to perforation in seeds from the late sowing (March) incubated at 25°C may be explained by an increase in embryo dormancy under this maturation condition (Fig. 1C, D). Generally, high temperatures during seed development on the mother plant decrease dormancy (Fenner, 1991;

Benech-Arnold, 2004; Gualano and Benech-Arnold, 2009). Results obtained in this work suggest that embryo dormancy in quinoa follows this general pattern, therefore in seeds exposed to higher temperatures during development (early sowing date) embryo dormancy was maintained at relatively low levels, while seeds exposed to lower temperatures during development (late sowing date) showed a higher level of embryo dormancy. However, considering dormancy imposed by the seed coats, our results suggest that it responds to environmental conditions in the opposite way: higher in early sowing and lower in late sowings. The reduced effect of perforation

Table 1. Seed coat thickness (μm) of 2-Want and Chadmo seeds, from different sowing dates.

Campaign	Sowing date	2-Want	Chadmo
First year	First; November	14.43 \pm 1.26 ^{a b}	20.00 \pm 1.44 ^a
	Second; December	10.19 \pm 0.86 ^{bc b}	17.31 \pm 0.91 ^a
	Third; March	7.30 \pm 0.56 ^{c b}	17.50 \pm 0.93 ^a
Second year	First; November	12.50 \pm 0.59 ^{ab b}	17.69 \pm 0.75 ^a
	Second; February	8.46 \pm 0.59 ^{c b}	16.92 \pm 1.06 ^a

The data are expressed as the mean \pm standard error; different letters indicate significant differences between sowing dates for each accession (normal letters, $P \leq 0.05$) and between accessions for each sowing date (bold letters, $P \leq 0.01$).




Figure 3. Cross-section of fruit and seed of accession 2-Want (A) and detail of the tissues that composed the seed coat (B). The rectangle indicates the area where measurements of seed coat thickness were performed. emb, embryo; p, perisperm; en, endosperm; per, pericarp; sc, seed coat; tes, testa; teg, tegmen. The scale bars indicate (A) 250 μm and (B) 50 μm .

treatments in late sowing dates is explained by the combination of low coat dormancy and high embryo dormancy (compare Fig. 1A with C for 2-Want, and Fig. 1B with D for Chadmo). This response is similar to that of sunflower, where high temperatures during seed development increase coat (pericarp and seed membranes) imposed dormancy and decrease embryo dormancy (Bodrone, 2014).

Under low incubation temperatures (5 and 10°C) the expression of dormancy is stronger in quinoa seeds (Ceccato *et al.*, 2011). That could explain the fact that coat-imposed dormancy (and perforation effects) was expressed at 5 and 10°C in 2-Want seeds from the late sowing date (even though embryo dormancy was higher than in early sowing dates). Chadmo seeds showed a very low response to perforation, this

might be a consequence of a deeper embryo dormancy expressed under low incubation temperatures in late sowing dates that limits the expression of coat-imposed dormancy in Chadmo seeds. Embryo dormancy was suggested (but not proved) before for *C. album*, as a possible explanation for the lack of response to seed scarification (Williams, 1963). This paper presents firm arguments in support of the existence of embryo dormancy in the genus *Chenopodium*.

The seed coat thickness of 2-Want was reduced due to late sowings in both years, in accordance with the decrease in dormancy level (Table 1). Similar results were reported previously in the genus *Chenopodium* for *C. polyspermum*, *C. album* and *C. bonus-henricus* (Jacques, 1968; Karssen, 1970; Pourrat and Jacques, 1975; Dorne, 1981). In those three species the increased coat thickness




Figure 4. Correlation between the seed coat thickness (μm) at harvest and the mean temperature (A) or photoperiod (B) during seed development for each sowing date. $P < 0.001$. Accession 2-Want. Data are the mean \pm 1SE.




Figure 5. ABA measured in the incubation medium of intact seeds (control) and seeds with perforated coats (perforated), in accessions Chadmo (A) and 2-Want (B); and ABA content in dry seeds at harvest (C). The data are expressed as picograms of ABA per microlitre of incubation medium ($\text{pg ABA } \mu\text{l}^{-1}$; A and B), or as picograms of ABA per milligram of seed dry weight [$\text{pg ABA (mg dw)}^{-1}$; C]. Different letters indicate significant differences ($P < 0.01$) and the bars indicate the mean \pm 1SE.

has been associated with a higher level of dormancy. However, the fact that Chadmo seeds did not respond in that way to sowing date is a warning against generalizations of these patterns to the whole species.

Nevertheless, seed dormancy was associated with exposure of seeds to different environmental conditions during their development. Temperatures and photoperiods experienced during this period are associated with seed dormancy level in quinoa (Ceccato *et al.*, 2011). Seed coat thickness in 2-Want seeds is also correlated with the mean temperature and photoperiod during their development (Fig. 4). Due to the covariance of these factors, it became impossible under field conditions to know which factor (temperature and/or photoperiod) is responsible for the observed variation in seed dormancy and seed coat thickness, and this matter deserves further analysis.

Like quinoa, *C. polyspermum* and *C. album* seeds formed under a short photoperiod had thinner seed coats and lower dormancy (Jacques, 1968; Karssen, 1970; Pourrat and Jacques, 1975). The relative changes in seed coat thickness, and the germination percentage linked to that, were similar to the ones observed in accession 2-Want, but were obtained by exposing plants to a higher difference between photoperiods (10–16 h). In 2-Want, the difference was of ~ 3.5 h. Any temperature effect during seed development on seed dormancy has been tested extensively in other species (e.g. Fenner, 1991; Fonseca and Sanchez, 2000; Benech-Arnold, 2004). In the case of 2-Want, the difference between average temperatures in the first and last sowing date was 11°C in the first and 7°C in the second year for the effective seed-filling period (between the onset of filling and PM). Thus, possible effects of variation in environmental temperature should not be ruled out.

Archaeological studies in eastern North America related differences in coat thickness to the level of

domestication in *C. berlandieri* ssp. *jonesianum*. Thicker coats (mean $34 \mu\text{m}$) were associated with wild or weedy forms, and thinner coats (mean $15.85 \mu\text{m}$) with cultivated forms (Gremillion, 1993a, b). A similar reduction in seed coat thickness (12.8 to $2.4 \mu\text{m}$ average) was linked to domestication of *C. quinoa*, as observed in Chiripa, Bolivia (Bruno, 2005, 2006). These changes were assumed to be the consequence of selection, with the objective of reducing dormancy. This hypothesis, however, was not assessed with viable seeds. According to results of the present work, an association of seed coat thickness with germination per cent between sowing dates was established in one of the two accessions studied. Moreover, Chadmo seeds consistently exhibited a thicker seed coat (Table 1) and were consistently more dormant than 2-Want seeds (Figs 1 and 2). Besides, seed coats measured in accession 2-Want (Table 1) were two times thicker than those reported by Bruno (2005, 2006) in modern seeds from Chiripa, Bolivia. For BO25, an accession native to central Chile, López Fernández (2008) reported an exotesta thickness nearly twice that of another eight cultivars tested ($\sim 25 \mu\text{m}$ vs. $12\text{--}15 \mu\text{m}$), including Chilean and Bolivian accessions. Together, these observations indicate an important variability in coat thickness between accessions of *C. quinoa*. A comparison of dormancy level between accessions with contrasting coat thickness in a range of environments is needed to provide more information about the association between coat thickness and dormancy level in the species.

Less ABA leached from Chadmo seeds upon imbibition than from 2-Want seeds (compare the two controls in Fig. 5A and B), which suggests that deeper dormancy in Chadmo seeds can be explained partially by less ABA leaching.

More ABA leached from Chadmo seeds upon perforation; therefore, the seed coat plays a stronger role in ABA leaching, and the greater thickness may also

play a role. The role of the seed coat as a barrier for ABA diffusion has been reported for several species (Wang *et al.*, 1995; Bianco *et al.*, 1997; Ren and Kermode, 1999; Feurtado *et al.*, 2008). Interestingly, the dry 2-Want seeds have weaker dormancy, yet a higher ABA content. Apparently, ABA content in dry seeds is not necessarily directly associated with dormancy level. On the other hand, the possibility of differences in the ability of seeds to synthesize ABA *de novo*, or to catabolize ABA during incubation, should not be neglected.

Among other probable action mechanisms of coats in the regulation of dormancy, interference with the absorption of water was considered unlikely, since a very fast imbibition was found when seeds were incubated in water and 100% germination was reached post-maturation (Ceccato *et al.*, 2011). Besides, any mechanical restraint to radicle emergence would be unlikely since a small hole in the middle of the seed (far from the micropyle) was sufficient to remove dormancy, although the possibility of an indirect mechanism cannot be dismissed.

Conclusions

It was suggested that embryo dormancy is present in the genus *Chenopodium*. Coats affect dormancy in quinoa, and the seed coat tissue is largely responsible for this effect. The contribution of embryo dormancy to whole-seed dormancy is expressed under conditions in which coat-imposed dormancy is reduced.

Environmental conditions during seed development affect the level of dormancy and relative contribution of the seed coats and the embryo to seed dormancy level. Those experienced at early sowing dates (higher temperatures and longer photoperiods) lead to an increase in coat and a decrease in embryo dormancy, while those experienced at late sowing dates (lower temperatures and shorter photoperiods) lead to a decrease in coat and an increase in embryo dormancy.

The obstruction of ABA release to the medium by the seed coats appears to have a central role in dormancy maintenance. This was revealed by germination responses to perforation treatments in both accessions, and the concomitant increase in ABA levels in the incubation medium. Consistently, more ABA is able to leak to the medium from seeds with a thin seed coat. The thickness of the seed coat seems to explain variation in seed dormancy between accessions, while its variation among environments was consistent only for one of the two evaluated accessions.

Acknowledgements

We thank Gabriela Zarlavsky for histological sections, Silvina Enciso for ABA determinations, Martín León

for technical assistance, Roberto Benech-Arnold and IFEVA (Institute for Physiological and Ecological Research Applied to Agriculture – UBA – CONICET) for providing us with working facilities and INTA Base Bank of Germplasm for help in experiments.

Financial support

This work was supported by a fellowship from INTA (National Institute of Agricultural Technology), Argentina.

Conflict of interest

None.

References

- Benech-Arnold, R.L. (2004) Inception, maintenance and termination of dormancy in grain crops: physiology, genetics and environmental control. pp. 169–198 in Benech-Arnold, R.L.; Sánchez, R.A. (Eds) *Handbook of seed physiology: Applications to agriculture*. Binghamton, New York, The Haworth Press.
- Bianco, J., Garello, G. and Le Page Degivry, M.T. (1997) *De novo* ABA synthesis and expression of seed dormancy in a gymnosperm: *Pseudotsuga menziesii*. *Plant Growth Regulation* **21**, 115–119.
- Bodrone, M.P. (2014) Cambios en el nivel de dormición de semillas de girasol en función del ambiente térmico explorado durante la etapa de desarrollo-maduración de los frutos y el almacenaje post-cosecha. MSc thesis, Facultad de Agronomía, Universidad de Buenos Aires, Argentina.
- Bois, J.P., Winkel, T., Lhomme, J., Rafailac, J.P. and Rocheteau, A. (2006) Response of some Andean cultivars of quinoa (*Chenopodium quinoa* Willd.) to temperature: effects on germination, phenology, growth and freezing. *European Journal of Agronomy* **25**, 299–308.
- Bruno, M.C. (2005) Domesticado o silvestre? Resultados de la investigación de semillas de *Chenopodium* Chiripa, Bolivia (1500–100 A.C.). *Textos Antropológicos* **15**, 39–50.
- Bruno, M.C. (2006) A morphological approach to documenting the domestication of *Chenopodium* in the Andes. pp. 32–45 in Zeder, M.; Bradley, D.; Emshwiller, E.; Smith, B. (Eds) *Documenting domestication: New genetic and archaeological paradigms*. California, University of California Press.
- Ceccato, D.V., Bertero, H.D. and Batlla, D. (2011) Environmental control of dormancy of quinoa (*Chenopodium quinoa*) seeds. Two potential genetic resources for pre-harvest sprouting tolerance. *Seed Science Research* **21**, 133–141.
- Debeaujon, I., Lepiniec, L., Pourcel, L. and Routaboul, J.M. (2007) Seed coat development and dormancy. pp. 25–43 in Bradford, K.J.; Nonogaki, H. (Eds) *Seed development, dormancy and germination*. Oxford, UK, Blackwell Publishing.
- Di Mauro, M.F., Iglesias, M.J., Arce, D.P., Valle, E.M., Benech-Arnold, R.L., Tsuda, K., Yamazaki, K., Casalongué, C.A. and Godoy, A.V. (2012) MBF1s

- regulate ABA-dependent germination of Arabidopsis seed. *Plant Signaling & Behavior* **7**, 188–192.
- Dorne, A.J.** (1981) Variation in seed germination inhibition of *Chenopodium bonus-henricus* in relation to altitude of plant growth. *Canadian Journal of Botany* **59**, 1893–1901.
- Fenner, M.** (1991) The effects of the parent environment on seed germinability. *Seed Science Research* **1**, 75–84.
- Feurtado, J.A., Ren, C., Ambrose, S.J., Cutler, A.J., Ross, A.R.S., Abrams, S.R. and Kermode, A.R.** (2008) The coat-enhanced dormancy mechanism of western white pine (*Pinus monticola* Dougl. ex D. Don) seeds is mediated by abscisic acid homeostasis and mechanical restraint. *Seed Science and Technology* **36**, 283–300.
- Finch-Savage, W.E. and Leubner-Metzger, G.** (2006) Seed dormancy and the control of germination. *New Phytologist* **171**, 501–523.
- Fonseca, A.E. and Sánchez, R.A.** (2000) Efecto de la temperatura durante el llenado de grano sobre la germinación de semillas de girasol (*Helianthus annuus* L.). pp. 216–217 in *Resúmenes de la XXIII Reunión Argentina de Fisiología Vegetal*, 27–30 November, Córdoba, Argentina.
- Geerts, S., Raes, D., García, M., Del Castillo, C. and Buytaert, W.** (2006) Agro-climatic suitability mapping for crop production in the Bolivian Altiplano: A case study for quinoa. *Agricultural and Forest Meteorology* **139**, 399–412.
- Gremillion, K.J.** (1993a) Crop and weed in prehistoric eastern North America: the *Chenopodium* example. *American Antiquity* **58**, 496–509.
- Gremillion, K.J.** (1993b) The evolution of seed morphology in domesticated *Chenopodium*: an archaeological case study. *Journal of Ethnobiology* **13**, 149–169.
- Gualano, N.A. and Benech-Arnold, R.L.** (2009) Predicting pre-harvest sprouting susceptibility in barley: looking for 'sensitivity windows' to temperature throughout grain filling in various commercial cultivars. *Field Crops Research* **114**, 35–44.
- Hilhorst, H.W.M.** (2007) Definitions and hypotheses of seed dormancy. pp. 50–71 in Bradford, K.J.; Nonogaki, H. (Eds) *Seed development, dormancy and germination*. Oxford, UK, Blackwell Publishing.
- Jacobsen, S.E. and Bach, A.P.** (1998) The influence of temperature on seed germination rate in quinoa (*Chenopodium quinoa* Willd.). *Seed Science and Technology* **26**, 515–523.
- Jacques, R.** (1968) Action de la lumière par l'intermédiaire du phytochrome sur la germination, la croissance et le développement de *Chenopodium polyspermum* L. *Physiologie Végétale* **6**, 137–164.
- Karssen, C.M.** (1970) The light promoted germination of the seeds of *Chenopodium album* L. III. Effect of the photoperiod during growth and development of the plants on the dormancy of the produced seeds. *Acta Botanica Neerlandica* **19**, 81–94.
- Linkies, A. and Leubner-Metzger, G.** (2012) Beyond gibberellins and abscisic acid: how ethylene and jasmonates control seed germination. *Plant Cell Reports* **31**, 253–270.
- López Fernández, M.P.** (2008) Longevidad de las semillas de nueve cultivares de *Chenopodium quinoa* Willd., procedentes de regiones contrastantes: Ecuación de la viabilidad y rol de las cubiertas seminales. Biology degree thesis, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires.
- Mendiondo, G.M., Leymarie, J., Farrant, J.M., Corbineau, F. and Benech-Arnold, R.L.** (2010) Differential expression of abscisic acid metabolism and signalling genes induced by seed-covering structures or hypoxia in barley (*Hordeum vulgare* L.) grains. *Seed Science Research* **20**, 69–77.
- Ministério da Agricultura e Reforma Agrária.** (1992) *Regras para análise de sementes*. Brasília, Secretaria Nacional de Defesa Agropecuária, Departamento Nacional de Defesa Vegetal, Coordenação de Laboratório Vegetal.
- Nonogaki, H.** (2006) Seed germination—the biochemical and molecular mechanism. *Breeding Science* **56**, 93–105.
- Notivol, E., García-Gil, M.R., Alía, R. and Savolainen, O.** (2007) Genetic variation of growth rhythm traits in the limits of a latitudinal cline in Scots pine. *Canadian Journal of Forestry Research* **37**, 540–551.
- O'Brien, T.P. and McCully, M.E.** (1981) *The study of plant structure. Principles and selected methods*. Melbourne, Termarcarphi.
- Pourrat, Y. and Jacques, R.** (1975) The influence of photoperiodic conditions received by the mother plant on morphological and physiological characteristics of *Chenopodium polyspermum* L. seeds. *Plant Science Letters* **4**, 273–279.
- Prego, I., Maldonado, S. and Otegui, M.** (1998) Seed structure and localization of reserves in *Chenopodium quinoa*. *Annals of Botany* **82**, 481–488.
- Quarrie, S.A., Whitford, P.N., Appleford, N.E., Wang, T.L., Cook, S.K., Henson, I.E. and Loveys, B.R.** (1988) A monoclonal antibody to (S)-abscisic acid: its characterisation and use in a radioimmunoassay for measuring abscisic acid in crude extracts of cereal and lupin leaves. *Planta* **173**, 330–339.
- Ren, C. and Kermode, A.R.** (1999) Analyses to determine the role of the megagametophyte and other seed tissues in dormancy maintenance of yellow cedar (*Chamaecyparis nootkatensis*) seeds: morphological, cellular and physiological changes following moist chilling and during germination. *Journal of Experimental Botany* **50**, 1403–1419.
- Rodríguez, M.V., Mendiondo, G.M., Maskin, L., Gudesblat, G.E., Iusem, N.D. and Benech-Arnold, R.L.** (2009) Expression of ABA signaling genes and ABI5 protein levels in imbibed *Sorghum bicolor* caryopses with contrasting dormancy and at different developmental stages. *Annals of Botany* **104**, 975–985.
- Steinbach, H.S., Benech-Arnold, R.L., Kristof, G., Sánchez, R.A. and Marcucci-Poltri, S.** (1995) Physiological basis of pre-harvest sprouting resistance in *Sorghum bicolor* (L.) Moench. ABA levels and sensitivity in developing embryos of sprouting-resistant and -susceptible varieties. *Journal of Experimental Botany* **46**, 701–709.
- Sukhorukov, A. and Zhang, M.** (2013) Fruit and seed anatomy of *Chenopodium* and related genera (Chenopodioideae, Chenopodiaceae/Amaranthaceae): implications for evolution and taxonomy. *PLoS ONE* **8**, e61906.
- Wang, M., Heimovaara-Dijkstra, S. and Van Duijn, B.** (1995) Modulation of germination of embryos isolated from dormant and non dormant barley grains by manipulation of endogenous abscisic acid. *Planta* **195**, 586–592.
- Williams, J.T.** (1963) Biological flora of the British Isles. *Chenopodium album* L. *List of Britain Vascular Plants* **154**, 711–725.