

About the first record of the genus *Chonecolea* (Chonecoleaceae, Marchantiophyta) in Paraguay and its differences to related species

Jorge R. FLORES ^{a,b,c*}, Guillermo M. SUÁREZ ^{a,b,c}
& M. Elena REINER-DREHWALD ^d

^a Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET),
Argentina

^b Fundación Miguel Lillo, Miguel Lillo 205, San Miguel de Tucumán,
Tucumán CP 4000, Argentina

^c Facultad de Ciencias Naturales e Instituto Miguel Lillo, Miguel Lillo 251,
San Miguel de Tucumán, Tucumán CP 4000, Argentina

^d Albrecht-von-Haller-Institute of Plant Sciences, Department of Systematics,
Biodiversity and Evolution of Plants, Untere Karspüle 2,
37073 Göttingen, Germany


Abstract – *Chonecolea doellingeri* (Nees) Grolle, whose South American record is restricted to Argentina and Brazil, is reported for the first time for Paraguay. Additionally, its differences to related species are discussed and SEM microphotographs are provided.

Bryophytes / *Chonecolea doellingeri* / Liverworts / Marchantiophyta / Paraguay


The bryophyte flora of Paraguay is poorly known (Söderstrom et al., 2013, Ellis et al., 2014). During a field trip to Paraguay in 2012, some interesting liverworts were found. Among these, a foliose liverwort genus, *Chonecolea* Grolle was never before recorded for Paraguay. This genus, as originally defined by Grolle (1956), included a single species, *Chonecolea doellingeri* (Nees) Grolle, known from Brazil, Argentina, U.S.A. (Florida) and Australia (e.g., Schuster, 1980; Engel, 1980 [as *Clasmatocolea doellingeri*]; Ramsay et al., 1993; Drehwald, 1995; Drehwald & Reiner-Drehwald, 1996; Brown & Coveny, 1999; Gradstein & Costa, 2003; Yano & Luizi-Ponzo, 2006; Hässel de Menéndez & Rubies, 2009). Subsequently, Grolle and Váña (in Váña, 1980) described *Chonecolea andina* Grolle & Váña, based on collections of E. & P. Hegewald from Peru. A further taxon, *Chonecolea acutiloba* (Schiffn.) R.M.Schust., was registered for Brazil and Argentina (Schuster, 1980; Drehwald, 1995); although this last species was not accepted by Yano & Luizi-Ponzo (2006). Thus, three species are currently distributed in the Neotropics.

* Corresponding author: jrflores@conicet.gov.ar

After an extensive study of the collected material using light and scanning electron microscopy, we were able to identify the sample as *Chonecolea doellingeri*. This species is characterized by its small size (< 1 mm wide), a pale green stem, succubous, bilobed leaves reaching the dorsal midline of the stem, and rounded to acute lobes with entire margins. The leaf cells of *Chonecolea doellingeri* are 16.25-20.0 μm long \times 12.5-15.0 μm wide, thin-walled and with smooth cuticle. The sporophyte is covered by a campanulate perianth with a wide, 3-4 plicate mouth (Figs 1-3). *Chonecolea doellingeri* is often growing in dry or open habitats, on the base of trunks, e. g. of palm trees (Schuster, 1980; Drehwald, 1995; Yano & Luiz-Ponzo, 2006). *Chonecolea* is frequently associated with Lejeuneaceae species (Yano



Figs 1-3. *Chonecolea doellingeri* (Nees) Grolle: **1.** Mature plant aspect. **2.** Perianth and sporophyte with the open capsule. **3.** Leaf cells detail, showing the thin-walled cells. Scale bar: 1 and 2: 250 μm ; 3: 25 μm .



Figs 4-5. *Chonecolea doellingeri* (Nees) Grolle: **4.** Sporophyte, with the open capsule, intermixed with filamentous algae (arrow). **5.** Areolate spores in proximal and distal view.

& Luizi-Ponzo, 2006). In addition, the specimen from Paraguay is intermingled with filamentous algae (Fig. 4).

Yano & Luizi-Ponzo (2006) discussed the distribution of *C. doellingeri* in Brazil, and described its spore morphology (Figs 4-5). From the complete geographical range of *Chonecolea* in Brazil, Mato Grosso do Sul is the nearest location to Cerro Yaguarón (Paraguay). The spore size from Brazilian specimens ranged from 11.5 to 12.6 μm diameter (equatorial view), whereas the material from Paraguay had spores with a diameter of up to 20 μm.

Chonecolea andina can be differentiated from *C. doellingeri* on the base of its narrow-acute leaf sinus, unequal leaf lobes, underleaves present on sterile shoots, and by spore ornamentation (Váña, 1980). In addition, both species have a different ecology. In Paraguay, *Chonecolea doellingeri* was found on trunks in humid, subtropical Chaco forest at 168 m, while *C. andina* grows on soil and rocks in dry, alpine environments between 2690 and 4300 m (Váña, 1980).

Another taxon which resembles *C. doellingeri* is *Cylindrocolea* R.M.Schust. Two species of this genus are listed for Brazil (Gradstein & Pinheiro da Costa, 2003) that grow in environments similar to that of *C. doellingeri* in Paraguay. Although *Cylindrocolea* species are likely confused with *Chonecolea*, several morphological characters separate these two genera. In the first place, *Chonecolea* plants are pale to light green while those of *Cylindrocolea* are brownish to reddish; also, leaf insertion reaches stem midline in *C. doellingeri* but it does not in *Cylindrocolea*. The sporophyte of *C. doellingeri* is covered by a relatively short perianth with truncate mouth which does not fully surpass the capsule level. In *Cylindrocolea*, by contrast, perianths extensively overpass the capsule level and the mouth is often strongly crenate with thick walled cells. Oil bodies are finely granular in *Cylindrocolea* but in *Chonecolea* they are homogeneous. *Cylindrocolea* infrequently produces gemmae and shows also vegetative reproduction by caducous perianths. Further on, *Chonecolea doellingeri* is almost exclusively growing as epiphyte whereas *Cylindrocolea* usually grows on soil and rock.

To date, only a single locality of *Chonecolea* is known from Paraguay. However, additional field work is necessary to investigate if the genus is more widespread in the region.

Specimen examined: PARAGUAY. Dpto Paraguari: Cerro Yaguarón, 25°34'01''S, 57°17'35''W, 168 m, 04-08-2012, G. Suárez, M. Dematteis, E. Mesa Torres & A. Vega 1480 (CTES, LIL).

Acknowledgments. The authors thanks J. Váña for providing helpful comments and suggestions to improve the manuscript. This work was founded by CONICET (PIP 0078) and PICT1838.

REFERENCES

- BROWN E.A. & COVENY R.G., 1999 — *Chonecolea doellingeri* (Nees) Grolle (Chonecoleaceae; Hepaticae) – a new record for Australia. *Haussknechtia Beiheft* 9: 57-60.
- DREHWALD U., 1995 — Epiphytische Pflanzengesellschaften in NO-Argentinien. *Dissertationes botanicae* 250: 1-175, Fotos 1-10.
- DREHWALD U. & REINER-DREHWALD M.E., 1996 — Orthotricho-Frullanietum bonariensis, una nueva asociación epifita de Buenos Aires, Argentina. *Cryptogamie, Bryologie-Lichénologie* 17 (4): 279-286.
- ELLIS L.T., AFONINA O.M., ASTHANA A.K., GUPTA R., SAHU V., NATH V., BATAN N., BEDNAREK-UCHYRA H., BENITEZ A., ERZBERGER P., FEDOSOV V.E., GÓRSKI P., GRADSTEIN S.R., GREMMEN N., HALLINGBÄCK T., HAGSTRÖM M., KÖCKINGER H., LÉBOUVIER M., MEINUNGER L., NÉMETH C., NOBIS M., NOWAK A., ÖZDEMİR T., PANTOVIĆ J., SABOVLJEVIĆ A., SABOVLJEVIĆ M.S., PAWLIKOWSKI P., PLÁSEK V., CÍHAL L., SAWICKI J., SÉRGIO C., MINISTRO P., GARCIA C.A., SMITH V.R., STEFANUT S., STOW S., SUÁREZ G.M., FLORES J.R., THOUVENOT L., VÁÑA J., VAN ROOY J., ZANDER R.H., 2014 — New National and Regional Bryophytes records, 39. *Journal of bryology* 36 (3): 223-244.
- ENGEL J.J., 1980 — A monograph of *Clasmatocolea* (Hepaticae). *Fieldiana botany*, New Series, 3: 1-229.
- GRADSTEIN S.R. & PINHEIRO da COSTA D., 2003 — The Hepaticae and Anthocerotae of Brazil. *Memoirs of the New York botanical garden* 87: 1-318.
- GROLLE R. 1956 — Revision der *Clasmatocolea*-Arten. *Revue bryologique et lichénologique* 25: 288-303.
- HÄSSEL DE MENÉNDEZ G.G. & RUBIES M.F., 2009 — Catalogue of Marchantiophyta and Anthocerotophyta of southern South America. *Nova Hedwigia, Beiheft* 134.
- SCHUSTER R.M., 1980 — *The Hepaticae and Anthocerotae of North America east of the hundredth meridian. vol. 4*. New York, Columbia University Press.
- RAMSAY H.P., COVENY R.G., BROWN E.A. & BROOKS A.K., 1993 — Bryophytes of the Royal Botanic Gardens, Government House and the Domain, Sydney. *Cunninghamia* 3(1): 215-229.
- SÖDERSTRÖM L., HAGBORG A., VON KONRAT M. & SENECA A., 2013 — The Guaraní Land – Checklist of Hornworts (Anthocerotophyta) and Liverworts (Marchantiophyta) of Paraguay. *Polish botanical journal* 58(1): 267-277.
- VÁÑA J., 1980 — Some new South and Central American Hepatics. *The journal of the Hattori botanical laboratory* 48: 225-234.
- YANO O. & LUIZI-PONZO A.P., 2006 — *Chonecolea doellingeri* (Chonecoleaceae, Hepaticae), taxonomia e distribuição geográfica no Brasil. *Acta botanica brasílica* 20(4): 783-788.